

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Clinical Packet  
May 4, 2022**

**Table of Contents**

<b>Helpful Hints/Reference Document</b> .....	2
<b>External Criteria</b>	
Antihypertensive Agents.....	4
Alzheimer’s Agents .....	4
<b>Agenda</b> .....	6
<b>Pharmacotherapy Class Re-Reviews</b>	
Pharmacotherapy Review of Central Alpha-Agonists.....	7
Pharmacotherapy Review of Direct Vasodilators.....	43
Pharmacotherapy Review of Peripheral Adrenergic Inhibitors .....	90
Pharmacotherapy Review of Hypotensive Agents, Miscellaneous .....	91
Pharmacotherapy Review of Alpha-Adrenergic Blocking Agents.....	96
Pharmacotherapy Review of Beta-Adrenergic Blocking Agents .....	165
Pharmacotherapy Review of Dihydropyridines.....	336
Pharmacotherapy Review of Calcium-Channel Blocking Agents, Miscellaneous.....	468
Pharmacotherapy Review of Angiotensin-Converting Enzyme Inhibitors .....	542
Pharmacotherapy Review of Angiotensin II Receptor Antagonists .....	677
Pharmacotherapy Review of Mineralocorticoid (Aldosterone) Receptor Antagonists .....	827
Pharmacotherapy Review of Renin Inhibitors.....	922
Pharmacotherapy Review of Loop Diuretics .....	983
Pharmacotherapy Review of Potassium-Sparing Diuretics .....	1036
Pharmacotherapy Review of Thiazide Diuretics .....	1083
Pharmacotherapy Review of Thiazide-Like Diuretics.....	1158
Pharmacotherapy Review of Vasopressin Antagonists .....	1208
Pharmacotherapy Review of Diuretics, Miscellaneous .....	1234
Pharmacotherapy Review of Alzheimer’s Agents.....	1235

**Pharmacy and Therapeutics (P&T) Committee**  
**Helpful Hints/Reference Document**  
**P&T Charge**

As defined by §22-6-122

The Medicaid Pharmacy and Therapeutics (P&T) Committee shall review and recommend classes of drugs to the Medicaid Commissioner for inclusion in the Medicaid Preferred Drug Plan. Class means a therapeutic group of pharmaceutical agents approved by the FDA as defined by the American Hospital Formulary Service.

The P&T Committee shall develop its preferred drug list recommendations by considering the clinical efficacy, safety and cost effectiveness of a product. Within each covered class, the Committee shall review and recommend drugs to the Medicaid Commissioner for inclusion on a preferred drug list. Medicaid should strive to insure any restriction on pharmaceutical use does not increase overall health care costs to Medicaid.

The recommendations of the P&T Committee regarding any limitations to be imposed on any drug or its use for a specific indication shall be based on sound clinical evidence found in labeling, drug compendia and peer reviewed clinical literature pertaining to use of the drug. Recommendations shall be based upon use in the general population. Medicaid shall make provisions in the prior approval criteria for approval of non-preferred drugs that address needs of sub-populations among Medicaid beneficiaries. The clinical basis for recommendations regarding the PDL shall be made available through a written report that is publicly available. If the recommendation of the P&T Committee is contrary to prevailing clinical evidence found in labeling, drug compendia and/or peer-reviewed literature, such recommendation shall be justified in writing.

**Preferred Drug List/Program Definitions**

**Preferred Drug:** Listed on the Agency's Preferred Drug Lists and will not require a prior authorization (PA).

**Preferred with Clinical Criteria:** Listed on the Agency's Preferred Drug Lists but will require a prior authorization. Clinical criteria must be met in order to be approved.

**Non Preferred Drug:** Covered by the Agency, if it is determined and supported by medical records to be medically necessary, but will require a PA.

**Non Covered Drug:** In accordance with Medicaid Drug Amendments contained in the Omnibus Budget Reconciliation Act of 1990 (OBRA 90 federal legislation), the Agency has the option to not cover (or pay for) some drugs. Alabama Medicaid does not cover/pay for the following:

- Drugs used for anorexia, weight loss or weight gain, with the exception of those specified by the Alabama Medicaid Agency
- Drugs used to promote fertility with the exception of those specified by the Alabama Medicaid Agency
- Drugs used for cosmetic purposes or hair growth
- Over-the-counter/non prescription drugs, with the exception of those specified by the Alabama Medicaid Agency
- Covered outpatient drugs when the manufacturer requires as a condition of sale that associated test and/or monitoring services be purchased exclusively from the manufacturer or designee
  - DESI (Drug Efficacy Study Implementation [less than effective drugs identified by the FDA]) and IRS (Identical, Related and Similar [drugs removed from the market]) drugs which may be restricted in accordance with Section 1927(d) (2) of the Social Security Act
  - Agents when used for the symptomatic relief of cough and colds except for those specified by the Alabama Medicaid Agency
  - Prescription vitamin and mineral products, except prenatal vitamins and fluoride preparations and others as specified by the Alabama Medicaid Agency
  - Agents when used for the treatment of sexual or erectile dysfunction, unless authorized for pulmonary hypertension.

(From Alabama Medicaid Agency Administrative Code, Chapter 16 and Alabama Medicaid Agency Provider Billing Manual, Chapter 27.)

**Prior Authorization (PA):** Process that allows drugs that require approval prior to payment to be reimbursed for an individual patient. Drugs may require PA if they are preferred with clinical criteria, are non-preferred status, or if they required PA prior to the PDL.

Medicaid may require prior authorization for generic drugs only in instances when the cost of the generic product is significantly greater than the net cost of the brand product in the same AHFS therapeutic class or when there is a clinical concern regarding safety, overuse or abuse of the product.

Although a product may require PA, the product is considered a covered product and Medicaid will pay for the product only once the PA has been approved.

**Override:** Process where drugs require approval prior to payment to be reimbursed for an individual patient if the claim falls outside a predetermined limit or criteria. Overrides differ from PA in that drugs or drug classes that require an override will automatically allow payment of the drug unless something on the claim hits a predetermined limit or criteria. The different types of overrides include:

- Accumulation Edit
- Brand Limit Switchover
- Dispense As Written Override
- Early Refill
- Ingredient Duplication
- Maintenance Supply Opt Out
- Maximum Unit/Max Cost Limitations
- Short Acting Opioid Naïve Override
- Therapeutic Duplication

**Electronic PA (EPA):** The EPA system checks patient-specific claims history to determine if pharmacy and medical PA requirements are met at the Point-of-Sale claim submission for a non-preferred drug. If it is determined that all criteria are met and the request is approved, the claim will pay and no manual PA request will be required. Electronic PA results in a reduction in workload for providers because the claim is electronically approved within a matter of seconds with no manual PA required.

#### **Prior Authorization Criteria Definitions**

**Appropriate Diagnosis:** Diagnosis(es) that justifies the need for the drug requested. Diagnosis(es) or ICD-10 code(s) may be used. Use of ICD-10 codes provides specificity and legibility and will usually expedite review.

**Prior Treatment Trials:** Prior authorization requires that two (2) prescribed generic, OTC or brand name drugs have been utilized unsuccessfully relative to efficacy and/or safety within six (6) months prior to requesting the PA. The PA request must indicate that two (2) generic, OTC or other brand drugs have been utilized for a period of at least thirty (30) days each (14 days for Triptans, 3 days for EENT Vasoconstrictor Agents), **unless** there is an adverse/allergic response or contraindication. If the prescribing practitioner feels there is a medical reason for which the patient should not be on a generic, OTC or brand drug or drug trial, medical justification may be submitted in lieu of previous drug therapy. One prior therapy is acceptable in those instances when a class has only one preferred agent, either generic, OTC, or brand.

**Stable Therapy:** Allows for approval of a PA for patients who have been determined to be stable on a medication (same drug, same strength) for a specified timeframe and who continue to require therapy. Medications paid for through insurance, private pay or Medicaid are also counted toward the requirement. Providers will be required to document this information on the PA request form and note the program or method through which the medication was dispensed.

**Medical Justification:** An explanation of the reason the drug is required and any additional information necessary. Medical justification is documentation to support the physician's choice of the requested course of treatment. Documentation from the patient record (history and physical, tests, past or current medication/treatments, patient's response to treatment, etc) illustrates and supports the physician's request for the drug specified. For example, if a recommended therapy trial is contraindicated by the patient's condition or a history of allergy to a first-line drug, and the physician wants to order a non-preferred drug, documentation from the patient record would support that decision. In addition, medical justification may include peer reviewed literature to support the use of a non-preferred medication.

## **External Criteria**

### **Antihypertensive Agents**

#### **Appropriate Diagnosis**

- The patient must have an appropriate diagnosis supported by documentation in the patient record.

#### **Prior Treatment Trials**

- The patient must also have failed 30-day treatment trials with at least two prescribed and preferred antihypertensive agents in this class, either generic, OTC, or brand, within the past 6 months or have a documented allergy or contraindication to all preferred agents in this class.
- To meet these prior usage requirements, drugs within this specific classification must be judged against others in the same class (AHFS specific). For example, to qualify for a non-preferred beta-blocker, the patient must have met prior usage requirements of 30-day treatment trials with two other preferred beta-blockers, either generic, OTC, or brand.
- For fixed-dose combination products containing drugs from 2 or more different subclasses, prior therapies must include at least 2 prescribed and preferred agents from the respective subclasses.
- For BiDil<sup>®</sup>, in lieu of prior usage requirements, approval may be obtained for adjunctive therapy to standard heart failure therapy (including a diuretic, angiotensin converting enzyme inhibitor or angiotensin II receptor antagonist, and beta-blocker) in self-identified black patients.
- For Hemangeol<sup>®</sup>, patients must be five weeks to five months of age when initiating treatment. Hemangeol<sup>®</sup> may only be approved for the treatment of proliferating infantile hemangioma requiring systemic therapy.
- For Jynarque<sup>®</sup>, patients must have documented ALT, AST and bilirubin values before initiating treatment. Jynarque<sup>®</sup> may only be approved to slow kidney function decline in adults at risk of rapidly progressing autosomal dominant polycystic kidney disease.
- For Samsca<sup>®</sup>, patients must have a documented serum sodium <125 mEq/L, or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction. Patients must also have documentation of being initiated on Samsca<sup>®</sup> in an inpatient setting.

#### **Stable Therapy**

- Approval may be given for those who have documented stable therapy on the requested medication for 60 consecutive days or greater.

#### **Medical Justification**

- Medical justification may include peer reviewed literature, medical record documentation, or other information specifically requested.

#### **PA Approval Timeframes**

- Approval may be given for up to 12 months.

#### **Electronic Prior Authorization (EPA)**

- Antihypertensive agents are included in the electronic PA program.

#### **Verbal PA Requests**

- Not Applicable



## **Alzheimer's Agents**

### **Appropriate Diagnosis**

- The patient must have an appropriate diagnosis supported by documentation in the patient record.

### **Prior Treatment Trials**

- The patient must also have failed 30-day treatment trials with at least one other prescribed and preferred Alzheimer's agent in this class, either generic, OTC, or brand, within the past 6 months, or have a documented allergy or contraindication to all preferred agents in this class.

### **Stable Therapy**

- Stable therapy for this class is defined as a 90-day or greater timeframe. Approval may be given for those who have documented stable therapy on the requested medication for 90 consecutive days or greater.

### **Medical Justification**

- Medical justification may include peer-reviewed literature, medical record documentation, or other information specifically requested.

### **PA Approval Timeframes**

- Approval may be given for up to 12 months.

### **Electronic Prior Authorization (EPA)**

- Alzheimer's agents are included in the electronic PA program.

### **Verbal PA Requests**

- PA requests that meet prior usage requirement for approval may be accepted verbally.

# AGENDA

## ALABAMA MEDICAID AGENCY PHARMACY AND THERAPEUTICS (P&T) COMMITTEE

May 4, 2022  
1:00 p.m. – 3:00 p.m.

- 
1. Opening remarks.....Chair
  2. Approval of February 9, 2022 P&T Committee Meeting minutes.....Chair
  3. Pharmacy program update.....Alabama Medicaid
  4. Oral presentations by manufacturers/manufacturers' representatives  
(prior to each respective class review)
  5. Pharmacotherapy class re-reviews.....University of Massachusetts Clinical Pharmacy Services
    - Mineralocorticoid (Aldosterone) Receptor Antagonists – AHFS 243220
    - Central Alpha-Agonists – AHFS 240816
    - Direct Vasodilators – AHFS 240820
    - Peripheral Adrenergic Inhibitors – AHFS 240832
    - Hypotensive Agents, Miscellaneous – AHFS 240892
    - Alpha-Adrenergic Blocking Agents – AHFS 242000
    - Beta-Adrenergic Blocking Agents – AHFS 242400
    - Dihydropyridines – AHFS 242808
    - Calcium-Channel Blocking Agents, Miscellaneous – AHFS 242892
    - Angiotensin-Converting Enzyme Inhibitors – AHFS 243204
    - Angiotensin II Receptor Antagonists – AHFS 243208
    - Renin Inhibitors – AHFS 243240
    - Loop Diuretics – AHFS 402808
    - Potassium-Sparing Diuretics – AHFS 402816
    - Thiazide Diuretics – AHFS 402820
    - Thiazide-Like Diuretics – AHFS 402824
    - Vasopressin Antagonists – AHFS 402828
    - Diuretics, Miscellaneous – AHFS 402892
    - Alzheimer's Agents
      - *Parasympathomimetic (Cholinergic) Agents – AHFS Class 120400 (current brands to be included: Aricept®, Exelon®, and Razadyne® ER only)*
      - *Central Nervous System Agents, Miscellaneous – AHFS Class 289200 (current brands to be included: Aduhelm®, Namenda®, Namenda® XR, and Namzaric® only)*
  6. New business
  7. Results of voting announced.....Chair
  8. Next meeting dates
    - August 10, 2022
    - November 9, 2022
    - February 8, 2023
    - May 3, 2023
  9. Adjourn

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Central Alpha-Agonists  
AHFS Class 240816  
May 4, 2022**

**I. Overview**

Drugs to treat hypertension are among the most frequently prescribed pharmacologic agents. The incidence of hypertension increases with age, and the proper selection of an antihypertensive agent is an important issue.<sup>1,2</sup> While a multitude of neurohormonal, renal, and vascular mechanisms have been proposed as contributors to hypertension, no specific cause can be assigned in most cases.<sup>3,4</sup> Antihypertensive agents are separated into broad classes depending on which aspect of blood pressure regulation they affect: sodium and water balance, the sympathetic nervous system, resistance from vascular smooth muscle, or the renin-angiotensin-aldosterone system (RAAS).<sup>5</sup> Most patients will require therapy with more than one agent to achieve adequate blood pressure control. When monotherapy fails to achieve the blood pressure goal, then a second agent from a different class should be added to the treatment regimen.<sup>1</sup>

The central alpha-agonists are approved for the treatment of hypertension. They lower blood pressure primarily through stimulation of  $\alpha_2$ -adrenergic receptors in the central nervous system (CNS). This action inhibits sympathetic vasomotor centers, causing decreased sympathetic outflow from the CNS and an associated increase in vagal tone. Sympathetic activity is reduced while parasympathetic activity is increased. This leads to a reduction in total peripheral resistance, systolic and diastolic blood pressure, baroreceptor reflexes, heart rate, and cardiac output.<sup>3,6-10</sup> Plasma renin activity is also affected by the central  $\alpha$ -agonists, but the relationship between this and their hypotensive effects has not been fully elucidated. Chronic central  $\alpha$ -agonist use is associated with sodium and fluid retention, which may require concomitant diuretic therapy.<sup>3</sup> Methyldopa is available in combination with a thiazide diuretic. Thiazide diuretics inhibit the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride.<sup>6-10</sup>

The central  $\alpha$ -agonists that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the products are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Central Alpha-Agonists Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Clonidine	tablet, transdermal patch	Catapres-TTS®*	clonidine, Catapres-TTS®*
Guanfacine	tablet	N/A	guanfacine
Methyldopa	tablet	N/A	methyldopa
Methyldopate	injection^	N/A	methyldopate

\*Generic is available in at least one dosage form or strength.

^Product is primarily administered in an institution.

PDL=Preferred Drug List

N/A=Not available

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the central  $\alpha$ -agonists are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Central Alpha-Agonists**

Clinical Guideline	Recommendations
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)<sup>1</sup></b></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic blood pressure <math>&lt; 140</math> mm Hg.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 140</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)<sup>11</sup></b></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g</li> </ul>

Clinical Guideline	Recommendations
	<p>alcohol/standard drink). Avoid binge drinking.</p> <ul style="list-style-type: none"> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week. Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR &lt;45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> &gt;4.5</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>mmol/L.</p> <ul style="list-style-type: none"> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indication for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>12</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq 100</math> mmHg or average systolic blood pressure (SBP) measurements of <math>\geq 160</math> mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq 90</math> mmHg or for average SBP readings <math>\geq 140</math> mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should</li> </ul>

Clinical Guideline	Recommendations
	<p>be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</p> <ul style="list-style-type: none"> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt; 40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal</li> </ul>



Clinical Guideline	Recommendations
	<p>function loss, and acute pulmonary edema.</p> <ul style="list-style-type: none"> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index</li> </ul>

Clinical Guideline	Recommendations
	<p>and maternal diabetes.</p> <ul style="list-style-type: none"> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>13</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not</li> </ul>

Clinical Guideline	Recommendations
	<p>tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</p> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV</li> </ul>

Clinical Guideline	Recommendations
	<p>infusion is recommended.</p> <ul style="list-style-type: none"> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive</li> </ul>

Clinical Guideline	Recommendations
	<p>treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</p> <ul style="list-style-type: none"> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>14</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant</li> </ul>

Clinical Guideline	Recommendations
	<p>hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</p> <ul style="list-style-type: none"> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>15</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>16</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors</li> </ul>

Clinical Guideline	Recommendations
	<p>(RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</p> <ul style="list-style-type: none"> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><b>BP management in kidney transplant recipients</b></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>• The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><b>BP management in children with CKD</b></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>17</sup></p>	<p><b>Hypertension/blood pressure control</b></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq 140/90</math>) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq 180/110</math> and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events..</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math></li> </ul>

Clinical Guideline	Recommendations
	<p>mg/g creatinine and/or estimated glomerular filtration rate &lt;60 mL/min/1.73 m<sup>2</sup>.</p> <ul style="list-style-type: none"> <li>Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (&lt;30 mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate &lt;30 mL/min/1.73 m<sup>2</sup>.</li> <li>Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American College of Cardiology/American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>18</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) ≥130 mmHg or an average diastolic blood pressure (DBP) of ≥80 mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of ≥10% and an average SBP of ≥130 mmHg or an average DBP of ≥80 mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of ≥140 mmHg or a DBP of ≥90 mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of ≥10%, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>Adults with SIHD and hypertension (BP ≥130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago</li> </ul>

Clinical Guideline	Recommendations
	<p>and have angina.</p> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of</li> </ul>

Clinical Guideline	Recommendations
	<p>a thiazide diuretic plus ACE inhibitor, is useful.</p> <ul style="list-style-type: none"> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal <math>&lt;130/80</math> mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal <math>&lt;130</math> mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP <math>&lt;140</math> mmHg and a DBP <math>&lt;90</math> mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal <math>&lt;130/80</math> mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target <math>&lt;130/80</math> mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal <math>&lt;130</math> mmHg is</li> </ul>

Clinical Guideline	Recommendations
	<p>recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</p> <ul style="list-style-type: none"> <li>For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to <math>&lt; 140</math> mmHg during the first hour and to <math>&lt; 120</math> mmHg in aortic dissection.</li> <li>For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>In patients with planned elective major surgery and SBP <math>\geq 180</math> mmHg or DBP <math>\geq 110</math> mmHg, deferring surgery may be considered.</li> <li>For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the central  $\alpha$ -agonists are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Central Alpha-Agonists<sup>6-10</sup>**

Indication	Single Entity Agents		
	Clonidine	Guanfacine	Methyldopa
Treatment of hypertension	✓ *	✓ *	✓

\*Alone or in combination with other antihypertensive agents.

†This fixed combination drug is not indicated for the initial therapy of hypertension.

#### IV. Pharmacokinetics

The pharmacokinetic parameters of the central  $\alpha$ -agonists are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Central Alpha-Agonists<sup>7</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Clonidine	Oral: 75 to 100 TD: 60	20 to 40	Liver	Renal (40 to 60) Feces (22)	Oral: 22 TD: 12 to 13
Guanfacine	80	70	Liver	Renal (50)	17
Methyldopa	25 to 50	Negligible (% not reported)	Liver	Renal (70) Feces (30 to 50)	1.7

TD=transdermal

#### V. Drug Interactions

Significant drug interactions with the central  $\alpha$ -agonists are listed in Table 5.

**Table 5. Significant Drug Interactions with the Central Alpha-Agonists<sup>7</sup>**

Generic Name(s)	Interaction	Mechanism
Central $\alpha$ -agonists (clonidine)	Beta-adrenergic blockers	The severity of rebound hypertension associated with abrupt withdrawal of clonidine may be greater in patients taking $\beta$ -adrenergic blockers. This combination has also been reported to cause paradoxical hypertension. The mechanism of this interaction is unknown.
Central $\alpha$ -agonists (clonidine)	Tricyclic antidepressants	The antihypertensive effectiveness of clonidine may be decreased. Tricyclic antidepressants may also worsen the rebound reactions, such as hypertension and tachycardia, from abrupt clonidine withdrawal. The mechanism of this interaction is unknown.
Central $\alpha$ -agonists (clonidine)	Diltiazem, verapamil	Sinus bradycardia, AV block and severe hypotension may occur with coadministration of clonidine and diltiazem/verapamil. The mechanism of this interaction is unknown.
Central $\alpha$ -agonists (guanfacine)	Conivaptan	Concurrent use of conivaptan and guanfacine may result in increased guanfacine exposure due to CYP3A4 inhibition.
Central $\alpha$ -agonists (methyldopa)	Sympathomimetics	The coadministration of methyldopa and sympathomimetics may result in an increased pressor response, possibly resulting in hypertension.
Central $\alpha$ -agonists (methyldopa)	Entacapone	Concurrent use of entacapone and methyldopa may result in an increased risk of tachycardia, hypertension, and arrhythmias.
Central $\alpha$ -agonists (methyldopa)	Monoamine oxidase inhibitors	Metabolites of methyldopa stimulate release of endogenous catecholamines that are usually metabolized by MAOIs, thereby leading to excessive sympathetic stimulation.

## VI. Adverse Drug Events

The most common adverse drug events reported with the central  $\alpha$ -agonists are listed in Table 6. Abrupt discontinuation may cause nervousness, palpitations, headache, perspiration, nausea, and agitation. In some cases, sudden discontinuation may cause potentially dangerous rebound hypertension.<sup>6,7</sup>

**Table 6. Adverse Drug Events (%) Reported with the Central Alpha-Agonists<sup>6-10</sup>**

Adverse Events	Single Entity Agents			
	Clonidine Oral	Clonidine Transdermal	Guanfacine	Methyldopa
<b>Cardiovascular</b>				
Angina	-	-	-	✓
Arrhythmia	✓	-	-	-
Atrioventricular block	✓	-	-	-
Bradycardia	✓	-	≤3	✓
Carotid sinus sensitivity	-	-	-	✓
Chest pain	<1	-	≤3	-
Congestive heart failure	✓	-	-	✓
Edema	-	-	-	✓
Electrocardiogram abnormalities	✓	-	-	-
Hypotension	-	-	-	✓
Myocarditis	-	-	-	✓
Orthostatic hypotension	3	-	-	✓
Palpitations	✓	-	≤3	-
Pericarditis	-	-	-	✓
Peripheral edema	-	-	-	>10
Reynaud's phenomenon	✓	-	-	-
Syncope	✓	-	-	<1
Tachycardia	✓	-	-	-
<b>Central Nervous System</b>				
Agitation	✓	-	-	-
Amnesia	-	-	≤3	-
Anxiety	✓	-	-	1 to 10
Bell's palsy	-	-	-	✓
Confusion	-	-	≤3	-
Delirium	✓	-	-	-
Decreased mental acuity	-	-	-	✓
Delusional perception	✓	-	-	✓
Depression	✓	-	≤3	1 to 10
Dizziness	16	2	12 to 15	✓

Adverse Events	Single Entity Agents			
	Clonidine Oral	Clonidine Transdermal	Guanfacine	Methyldopa
Drowsiness	33	12	-	1 to 10
Fatigue	4	6	2 to 10	-
Hallucinations	<1	-	-	-
Headache	1	5	3 to 13	1 to 10
Insomnia	5	2	≤3	-
Involuntary movements	-	-	-	✓
Lightheadedness	-	-	-	✓
Lethargy	-	3	-	-
Nervousness	3	1	-	-
Nightmares	✓	-	-	✓
Paresthesia	✓	-	-	✓
Parkinsonism	-	-	-	✓
Restlessness	✓	-	-	-
Sedation	10	3	-	✓
Sleep disturbances	✓	-	-	-
Somnolence	-	-	5 to 39	-
Weakness	10	-	2 to 7	✓
<b>Dermatological</b>				
Allergic contact sensitization	-	5	-	-
Alopecia	✓	-	-	-
Angioedema	✓	-	-	-
Blanching	-	1	-	-
Burning	-	3	-	-
Contact dermatitis	-	19	-	-
Dermatitis	-	-	≤3	-
Edema	3	3	-	-
Erythema	-	15 to 50	-	-
Excoriation	-	3	-	-
Exfoliative dermatitis	-	-	-	✓
Hives	✓	-	-	-
Hyperpigmentation	-	5	-	-
Lupus-like syndrome	-	-	-	✓
Morbilliform or macro papular eruptions	-	1	-	-
Pruritus	7	15 to 50	≤3	-
Purpura	-	-	≤3	-
Rash	✓	-	-	✓



Adverse Events	Single Entity Agents			
	Clonidine Oral	Clonidine Transdermal	Guanfacine	Methyldopa
Sweating	-	-	≤3	<1
Throbbing	-	3	-	-
Toxic epidermal necrolysis	-	-	-	✓
Urticaria	✓	<1	-	-
Vasculitis	-	-	-	✓
Vesiculation	-	7	-	-
<b>Endocrine and Metabolic</b>				
Breast enlargement	-	-	-	✓
Erectile dysfunction	✓	-	-	-
Gynecomastia	✓	-	-	✓
Hyperprolactinemia	-	-	-	✓
Impotence	3	2	3 to 7	✓
Lactation	-	-	-	✓
Pancreatitis	-	-	-	✓
Sexual dysfunction	3	2	≤3	✓
Sodium retention	-	-	-	<1
Weight gain	✓	-	-	-
<b>Gastrointestinal</b>				
Abdominal Pain	✓	-	≤3	-
Anorexia	1	-	-	-
Colitis	-	-	-	✓
Constipation	10	1	2 to 15	✓
Diarrhea	-	-	≤3	✓
Distention	-	-	-	✓
Dry mouth	40	25	10 to 54	1 to 10
Dry throat	-	2	-	-
Dyspepsia	-	-	≤3	-
Dysphagia	-	-	≤3	-
Flatus	-	-	-	✓
Nausea	5	1	≤3	✓
Pseudo-obstruction	✓	-	-	-
Parotitis	✓	-	-	-
Salivary gland pain	✓	-	-	-
Sialadenitis	-	-	-	✓
Sore tongue	-	-	-	✓
Taste alteration	-	1	≤3	-

Adverse Events	Single Entity Agents			
	Clonidine Oral	Clonidine Transdermal	Guanfacine	Methyldopa
Vomiting	5	-	-	✓
Weight gain	1	-	-	✓
<b>Genitourinary</b>				
Micturition difficulties	✓	-	-	-
Nocturia	✓	-	-	-
Testicular disorder	-	-	∩	-
Urinary incontinence	-	-	∩	<1
Urinary retention	1	-	-	-
<b>Hematologic</b>				
Bone marrow depression	-	-	-	✓
Eosinophilia	-	-	-	✓
Granulocytopenia	-	-	-	✓
Hemolytic anemia	-	-	-	✓
Leukopenia	-	-	-	✓
Positive antinuclear antibody test	-	-	-	✓
Positive Rheumatoid factor test	-	-	-	✓
Positive Coombs test	✓	-	-	✓
Thrombocytopenia	✓	-	-	✓
<b>Hepatic</b>				
Cholestasis	-	-	-	<1
Cirrhosis	-	-	-	<1
Hepatitis	✓	-	-	✓
Jaundice	-	-	-	✓
<b>Laboratory Test Abnormalities</b>				
Blood urea nitrogen increased	-	-	-	✓
Creatinine phosphokinase increased	✓	-	-	-
Hyperglycemia	✓	-	-	-
Liver function test abnormalities	✓	-	-	✓
<b>Musculoskeletal</b>				
Arthralgia	-	-	-	✓
Hypokinesia	-	-	∩	-
Leg cramps	✓	-	∩	-
Myalgia	✓	-	-	✓
<b>Respiratory</b>				
Dyspnea	-	-	∩	<1
Rhinitis	-	-	∩	-

Adverse Events	Single Entity Agents			
	Clonidine Oral	Clonidine Transdermal	Guanfacine	Methyldopa
<b>Other</b>				
Blurred vision	✓	-	-	-
Dry eyes	✓			
Conjunctivitis	-	-	∞	-
Drug fever	-	-	-	1 to 10
Fever	✓	-	-	-
Iritis	-	-	∞	-
Malaise	1	-	∞	-
Nightmares	<1	-	-	-
Paresis	-	-	∞	-
Paresthesia	-	-	∞	-
Tinnitus	-	-	∞	-
Vision disturbance	-	-	∞	-
Withdrawal syndrome	✓	-	-	-

- ✓ Percent not specified
- Event not reported

## VII. Dosing and Administration

The usual dosing regimens for the central  $\alpha$ -agonists are listed in Table 7.

**Table 7. Usual Dosing Regimens for the Central Alpha-Agonists<sup>6-10</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Clonidine	<p><u>Hypertension:</u> Tablet: initial, 0.1 mg twice daily; maintenance, 0.1 to 0.6 mg/day in two divided doses; maximum, 2.4 mg/day</p> <p>Transdermal: initial, 0.1 mg patch once weekly; maintenance, 0.1 to 0.3 mg patch once weekly; maximum, two of the 0.3 mg patches once weekly</p>	Safety and effectiveness in pediatric patients have not been established in adequate and well-controlled trials.	<p>Tablet: 0.1 mg 0.2 mg 0.3 mg</p> <p>Transdermal patch: 0.1 mg/24 hours 0.2 mg/24 hours 0.3 mg/24 hours</p>
Guanfacine	<p><u>Hypertension:</u> Tablet: initial, 1 mg once daily at bedtime; maintenance, 1 to 2 mg once daily; maximum, 3 mg once daily</p>	Safety and efficacy in children under 12 have not been established.	Tablet: 1 mg 2 mg
Methyldopa	<p><u>Hypertension:</u> Tablet: initial, 250 mg 2 to 3 times daily; maintenance, 500 to 2,000 mg daily in two divided doses; maximum dose, 3 g daily</p>	<p>There are no well-controlled clinical trials in pediatric patients. Information on dosing in pediatric patients is supported by evidence from published literature regarding the treatment of hypertension in pediatric patients.</p> <p><u>Hypertension:</u> Tablet: initial, 10 mg/kg/day in 2 to 4 divided doses; maintenance, titrate up or down until adequate response achieved; maximum, 65 mg/kg/day or 3 g daily, whichever is less</p>	Tablet: 250 mg 500 mg

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the central  $\alpha$ -agonists are summarized in Table 8.

**Table 8. Comparative Clinical Trials with the Central Alpha-Agonists**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Lilja M et al. <sup>19</sup> (1991)  Clonidine 0.1 mg tablets BID  vs  clonidine 0.2 mg transdermal patch QD	DB, DD, PC, RCT, XO  Patients with mild to moderate HTN	N=16  12 weeks	Primary: Change from baseline in supine SBP, standing SBP and heart rate  Secondary: Difference in primary endpoints between oral and transdermal clonidine	Primary: Clonidine transdermal patch reduced both supine SBP and DBP by 13/7 mm Hg (P<0.01 and P<0.01) and heart rate by 9 bpm (P<0.01). Oral clonidine reduced only supine SPB by 11 mm Hg (P<0.01).  In a standing position, clonidine transdermal patch reduced SBP and DBP by 14/9 mm Hg (P<0.01 and P<0.01) and heart rate by 9 bpm (P<0.01). Oral clonidine reduced only standing heart rate by 8 bpm (P<0.05).  Secondary: There were no differences reported in primary endpoints between clonidine transdermal patch and oral clonidine (P value not reported).
Houston et al. <sup>20</sup> (1993)  Clonidine transdermal 0.1 to 0.3 mg QD plus nifedipine 60 mg QD (single entity products)  vs  nifedipine 60 mg QD	OL, PC, PRO  Male and nonpregnant female patients between 18 and 75 years of age with mild to moderate HTN and inadequate response to nifedipine	N=42  8 weeks	Primary: Change in seated DBP to less than 90 mmHg at 8 weeks  Secondary: Not reported	Primary: Patients on combination therapy experienced a reduction of 16/14 mmHg in the mean seated blood pressure vs placebo (P<0.01) with mean seated blood pressure of 127/87 mmHg.  A reduction of 5/10 mmHg in the mean seated blood pressure was seen with combination therapy vs nifedipine monotherapy (P<0.01).  A reduction of 18/12 mmHg in the mean standing blood pressure was seen with combination therapy vs placebo (P<0.01).  A reduction of 9/9 mmHg in the mean standing blood pressure was seen with combination therapy vs nifedipine monotherapy (P<0.01).  Secondary: Not reported
Krieger et al. <sup>21</sup> (2018) ReHOT	OL, RCT  Patients with resistant	N=162  12 weeks	Primary: BP control (determined by office BP<140/90)	Primary: Compared with the spironolactone group, the clonidine group presented similar rates of achieving the primary end point (20.5 vs 20.8%, respectively; RR, 1.01; 95% CI, 0.55 to 1.88; P=1.00).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Clonidine 0.1 mg BID (could be titrated to 0.2 or 0.3 mg BID)</p> <p>vs</p> <p>spironolactone 12.5 mg QD (could be titrated to 25 or 50 mg/ day)</p>	<p>hypertension (no office and ambulatory BP monitoring control, despite treatment with 3 drugs, including a diuretic, for 12 weeks)</p>		<p>and ambulatory 24-hour mean BP &lt;130/80)</p> <p>Secondary: BP control by each evaluation method, absolute BP reduction</p>	<p>Secondary: Secondary end point analysis showed similar office BP (33.3 vs 29.3%) and ambulatory BP monitoring (44 vs 46.2%) control for spironolactone and clonidine, respectively. However, spironolactone promoted greater decrease in 24-hour systolic and diastolic BP and diastolic daytime ambulatory BP than clonidine.</p>
<p>Boyles et al.<sup>22</sup> (1984)</p> <p>Methyldopa 250 to 800 mg/day and HCTZ 25 to 100 mg/day</p> <p>vs</p> <p>HCTZ 25 to 100 mg/day</p>	<p>OL, RCT</p> <p>Patients ≥59 years with isolated systolic HTN</p>	<p>N=21</p> <p>18 weeks</p>	<p>Primary: Changes in blood pressure from baseline</p> <p>Secondary: Not reported</p>	<p>Primary: At two weeks standing blood pressure fell from a mean of 166/90 mmHg at baseline to 164/88 mmHg with HCTZ monotherapy.</p> <p>At four weeks standing blood pressure fell from a mean of 164/88 mmHg at the end of the two week HCTZ monotherapy period to 145/81 mmHg at two weeks with combination therapy.</p> <p>At 18 weeks standing blood pressure fell from a mean of 166/90 mmHg at baseline to 132/80 mmHg with combination therapy.</p> <p>Secondary: Not reported</p>
<p>Channick et al.<sup>23</sup> (1981)</p> <p>Methyldopa 250 mg/day and HCTZ 15 mg/day</p> <p>vs</p> <p>chlorthalidone 50 mg/day and reserpine 0.25 mg/day</p>	<p>OL, RCT</p> <p>Patients with HTN</p>	<p>N=56</p> <p>12 weeks</p>	<p>Primary: Efficacy of blood pressure lowering to goal DBP ≤90 mmHg</p> <p>Secondary: Adverse effects</p>	<p>Primary: Goal DBP of ≤90 mmHg was reached in 91% of the chlorthalidone and reserpine group vs 55% in the methyldopa and HCTZ group (P&lt;0.001).</p> <p>Secondary: The incidence of adverse effects was 31% with chlorthalidone and reserpine vs 64% with methyldopa and HCTZ (P&lt;0.02).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Finnerty et al.<sup>24</sup> (1979)</p> <p>Methyldopa 500 mg to 2,000 mg QD</p> <p>vs</p> <p>reserpine 0.125 mg to 0.25 mg QD</p> <p>vs</p> <p>propranolol 80 mg to 320 mg QD</p> <p>All patients received hydro-flumethiazide* 50 or 100 mg QD.</p>	<p>RCT, SB</p> <p>Patients with HTN unresponsive to hydroflumethiazide monotherapy</p>	<p>N=59</p> <p>9 weeks</p>	<p>Primary: Percentage of patients achieving a DBP &lt;90 mm Hg</p> <p>Secondary: Not reported</p>	<p>Primary: At trial endpoint, 20 patients (100%) receiving reserpine, 13 of the 19 patients (68.4%) receiving methyldopa and 16 of the 20 patients (80%) receiving propranolol achieved a DBP &lt;90 mm Hg (mean reductions and P values not reported).</p> <p>Secondary: Not reported</p>
<p>Fernandez et al.<sup>25</sup> (1980)</p> <p>Methyldopa 750 mg/day</p> <p>vs</p> <p>chlorothiazide 450 mg/day</p> <p>vs</p> <p>methyldopa and chlorothiazide 250-150 mg/day* (fixed-dose combination)</p>	<p>DB, PC, RCT, XO</p> <p>Patients with uncomplicated HTN</p>	<p>N=44</p> <p>4 weeks</p>	<p>Primary: Blood pressure lowering efficacy</p> <p>Secondary: Adverse effects</p>	<p>Primary: No significant differences in supine blood pressure for any treatment compared to placebo was observed (P value not reported). However, upright SBP, DBP and mean blood pressure were significantly lower with methyldopa and methyldopa and chlorothiazide compared to placebo (P&lt;0.05 for all).</p> <p>Secondary: Adverse effects were reported as infrequent (P value not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
product)  vs  placebo				
Materson et al. <sup>26</sup> (1990)  Hydralazine 25, 50 or 100 mg BID  vs  methyldopa 250, 500 or 1,000 mg BID  vs  metoprolol 50, 100 or 200 mg BID  vs  reserpine 0.05, 0.10 or 0.25 mg QD  All patients received HCTZ 25 to 100 mg QD.	DB, MC, RCT  Men ≥60 years with HTN not currently receiving antihypertensive therapy and DBP 90 to 114 mm Hg and SBP <240 mm Hg or a DBP <100 mm Hg and a SBP <240 mm Hg if currently taking antihypertensive therapy and the blood pressure criteria was met after ≥2 weeks without medication	N=690  12 months	Primary: The average reduction in SBP and DBP, the number of patients achieving the goal blood pressure, the average change in heart rate  Secondary: The rates of drug intolerances, adverse effects	Primary: Across all four treatments, there was an additional average reduction in BP of 13.1/10.6 mm Hg. The average reduction in SBP from baseline to endpoint for hydralazine, methyldopa, metoprolol and reserpine were -11.5±10.1 (P<0.001), -15.0±13.7 (P<0.001), -13.0±15.4 (P<0.001) and -12.7±11.5 (P<0.001), respectively. There was no significant difference in SBP reductions among the different treatments (P=0.43). The average reduction in DBP from baseline to endpoint for hydralazine, methyldopa, metoprolol and reserpine were -11.3±5.9 (P<0.001), -10.6±6.3 (P<0.001), -10.6±6.7 (P<0.001) and -9.8±6.3 (P<0.001), respectively. There was no significant difference in DBP reductions among the different treatments (P=0.59).  The average change in heart rate from baseline to endpoint for hydralazine, methyldopa, metoprolol and reserpine were 1.4±10.5 (P value not significant), -1.6±9.3 (P value not significant), 15.9±11.9 (P<0.05) and -7.9±10.7 (P<0.05), respectively. There was a significant difference in change in heart rate among the different treatments (P<0.001).  The percentage of patients achieving the goal blood pressure at endpoint with hydralazine, methyldopa, metoprolol and reserpine were 85.3, 81.7, 76.9 and 72.3%, respectively (P=0.28).  Secondary: Drug intolerance, defined as adverse effects prompting dose reduction or discontinuation, was present in 23.3% of patients not achieving goal blood pressure compared to 2.8% of those who did (P<0.001). This was significant with hydralazine, methyldopa and metoprolol, but not with reserpine.  There were 27 (10%) treatment discontinuations due to adverse effects (hydralazine [n=3], methyldopa [n=8], metoprolol [n=9] and reserpine



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>[n=7]). There were two treatment discontinuations with methyldopa and one with reserpine due to depression.</p> <p>The overall incidence of volunteered moderate or severe adverse effects, not prompting treatment discontinuation, was significantly greater (P&lt;0.01) with methyldopa (31%) and hydralazine (25%) compared to reserpine (15%) or metoprolol (9%).</p>
<p>McAreevey et al.<sup>27</sup> (1984)</p> <p>Hydralazine 12.5 mg QD up to 100 mg BID</p> <p>vs</p> <p>labetalol 200 mg QD up to 1,600 mg BID</p> <p>vs</p> <p>methyldopa 125 mg QD up to 1,000 mg BID</p> <p>vs</p> <p>prazosin 0.5 mg QD up to 10 mg BID</p> <p>vs</p> <p>placebo</p> <p>Minoxidil as add on therapy was given to</p>	<p>DB, PG, RCT</p> <p>Patients with inadequately controlled HTN while receiving atenolol 100 mg/day and bendrofluazide* 5 mg/day</p>	<p>N=238</p> <p>6 months</p>	<p>Primary: Comparative safety and efficacy, target blood pressure &lt;140/95 mm Hg</p> <p>Secondary: Not reported</p>	<p>Primary: Target blood pressure was reached in 25% of patients receiving hydralazine, 23% of patients receiving minoxidil, 19% of patients receiving prazosin, 17% of patients receiving methyldopa and zero percent of patients receiving placebo (P values not reported).</p> <p>Labetalol had the highest withdrawal rate compared to the other treatments with 78% (P&lt;0.05). Minoxidil had the second highest withdrawal rate with 57% (P&lt;0.05), due to fluid retention. There were no significant differences in withdrawal rates among the other treatments.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>men only.</p> <p>Doses were titrated upward at 2-week intervals until target BP or maximum dose was reached.</p>				

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, QD=once daily

Study design abbreviation: DB=double-blind, DD=double-dummy, MC=multicenter, OL=open-label, PC=placebo-controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, SB=single blind, XO=cross over

Miscellaneous abbreviations: DBP=diastolic blood pressure, HCTZ=hydrochlorothiazide, HTN=hypertension, SBP=systolic blood pressure

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of ‘\$’ signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Central Alpha-Agonists**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Clonidine	tablet, transdermal patch	Catapres-TTS®*	\$\$\$\$\$	\$ to \$\$
Guanfacine	tablet	N/A	N/A	\$
Methyldopa	tablet	N/A	N/A	\$

\*Generic is available in at least one dosage form or strength.  
HCTZ=hydrochlorothiazide, N/A=not available

**X. Conclusions**

The central alpha-agonists are approved for the treatment of hypertension, and all of the agents are available in a generic formulation.<sup>6-10</sup> There are several national and international organizations that have published guidelines on the treatment of hypertension. Most of the guidelines do not address the use of the central  $\alpha$ -agonists.<sup>1,11-18</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension. According to the National Heart, Lung, and Blood Institute’s Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium

channel blockers).<sup>1</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>1,11-18</sup> Methyldopa is safe and effective to use during pregnancy.<sup>12-13</sup>

There are limited head-to-head studies with the central  $\alpha$ -agonists. Clinical trials have compared combination therapy to monotherapy. In these studies, the more aggressive treatment regimen lowered systolic and diastolic blood pressure to a greater extent than the less-intensive treatment regimen.<sup>20,22</sup> There does not appear to be any difference in efficacy with the oral or transdermal formulations of clonidine.<sup>19</sup> According to treatment guidelines, most patients will need more than one antihypertensive agent to achieve blood pressure goals.<sup>1,11-18</sup> Certain guidelines note that that fixed combination antihypertensive medications can favor compliance and simplify medication regimens.<sup>12</sup> However, there are no prospective, randomized-controlled trials that have demonstrated better clinical outcomes with any central  $\alpha$ -agonist fixed-dose combination product compared to the coadministration of the individual components as separate formulations.

The most common adverse events reported with the central  $\alpha$ -agonists include dizziness, drowsiness, dry mouth, and somnolence. Abrupt discontinuation may cause nervousness, palpitations, headache, perspiration, nausea, and agitation. In some cases, sudden discontinuation may cause potentially dangerous rebound hypertension.<sup>6-10</sup>

There is insufficient evidence to support that one brand central alpha-agonist is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand central alpha-agonists within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand central alpha-agonist is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
2. Kaplan NM. Hypertension in the population at large. In: Kaplan NM, Victor RG, Flynn JT. *Kaplan's clinical hypertension*. 11<sup>th</sup> ed. Philadelphia (PA): Lippincott, Williams, and Wilkins; 2015.
3. Saseen JJ, MacLaughlin EJ. Chapter 13: Hypertension. In: DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, editors. *Pharmacotherapy: a pathophysiologic approach*. 10th edition. New York (NY): McGraw-Hill; 2017. <http://accesspharmacy.mhmedical.com/book.aspx?bookid=1861>.
4. Victor RG. Systemic hypertension: mechanisms and diagnosis. In: Mann DL, Zipes DP, Libby P, Bonow RO, eds. *Braunwald's heart disease: a textbook of cardiovascular medicine*. 10th ed. Philadelphia (PA): Saunders Elsevier, 2015.
5. Benowitz NL. Antihypertensive agents. In: Katzung BG, editor. *Basic and clinical pharmacology*. 14th ed. New York (NY): McGraw-Hill; 2018.
6. *Lexicomp Online Database [database on the Internet]*. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
7. Micromedex<sup>®</sup> Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
8. Catapres<sup>®</sup> [package insert]. Ridgefield (CT): Boehringer Ingelheim Pharmaceuticals; 2016 Aug.
9. Catapres TTS<sup>®</sup> [package insert]. Ridgefield (CT): Boehringer Ingelheim Pharmaceuticals; 2016 Aug.
10. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
11. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
12. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
13. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
14. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
15. American Diabetes Association. Cardiovascular disease and risk management & Microvascular complications and foot care. Sec. 10-11. In *Standards of Medical Care in Diabetes-2019*. *Diabetes Care* 2019; 42(Suppl. 1): S103–S138.
16. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
17. American Diabetes Association Professional Practice Committee. *Standards of Medical Care in Diabetes—2022*. *Diabetes Care* 2022;45(Suppl. 1).
18. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
19. Lilja M, Juustila H, Sarna S, Jounela AJ. Transdermal and oral clonidine. *Ann Med*. 1991;23(3):265-9.
20. Houston MC, Hays L. Transdermal clonidine as an adjunct to nifedipine-GITS therapy in patients with mild-to-moderate hypertension. *Am Heart J*. 1993;126:918-23.
21. Krieger EM, Drager LF, Giorgi DMA, Pereira AC, Barreto-Filho JAS, Nogueira AR, et al. Spironolactone Versus Clonidine as a Fourth-Drug Therapy for Resistant Hypertension: The ReHOT Randomized Study (Resistant Hypertension Optimal Treatment). *Hypertension*. 2018 Apr;71(4):681-690.
22. Boyles PW. Hydrochlorothiazide plus methyldopa in the treatment of isolated systolic hypertension in the elderly. *Clin Ther*. 1984;6:793-9.

23. Channick BJ, Kessler WB, Marks AD et al. A comparison of chlorthalidone –reserpine and hydrochlorothiazide –methyldopa as step 2 therapy for hypertension. *Clin Ther.* 1981; 4: 175-83.
24. Finnerty FA Jr, Gyftopoulos A, Berry C, McKenney A. Step 2 regimens in hypertension: an assessment. *JAMA.* 1979 Feb 9;241(6):579-81.
25. Fernandez PG, Zachariah PK, Bryant DG et al. Antihypertensive efficacy of alpha-methyldopa, chlorothiazide and Supres (alpha-methyldopa-chlorothiazide). *Can Med Assoc J.* 1980; 123:284-7.
26. Materson BJ, Cushman WC, Goldstein G, et al. Treatment of hypertension in the elderly: I. Blood pressure and clinical changes: results of a Department of Veterans Affairs cooperative study. *Hypertension.* 1990 Apr;15(4):348-60.
27. McAreavey D, Ramsey LE, Latham L, McLaren AD, Lorimer AR, Reid JL, et al. Third drug trial: comparative study of antihypertensive agents added to treatment when blood pressure remains uncontrolled by a beta blocker plus thiazide diuretic. *Br Med J (Clin Res Ed).* 1984 Jan 14;288(6411):106-11.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Direct Vasodilators  
AHFS Class 240820  
May 4, 2022**

**I. Overview**

The direct vasodilators are approved for the treatment of heart failure and hypertension.<sup>1-4</sup> Hydralazine and minoxidil interfere with calcium movement within the vascular smooth muscle, which is responsible for initiating and maintaining the contractile state. They exert a peripheral vasodilating effect through a direct relaxation of vascular smooth muscle. This leads to decreased arterial blood pressure, decreased peripheral vascular resistance, as well as an increase in heart rate, stroke volume, and cardiac output. Hydralazine is available as a single entity product, as well as in combination with isosorbide dinitrate. Isosorbide dinitrate enters vascular smooth muscle and is converted to nitric oxide, which results in dilatation of peripheral arteries and veins. Dilatation of the veins promotes peripheral pooling of blood and decreases venous return to the heart. Dilatation of the arteries reduces systemic vascular resistance, systolic arterial pressure, and mean arterial pressure.<sup>1,2</sup> The exact mechanism of action of the fixed-dose combination product containing isosorbide dinitrate and hydralazine in the treatment of heart failure has not been established.<sup>3</sup>

The direct vasodilators that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. Hydralazine and minoxidil are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Direct Vasodilators Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Hydralazine	injection, tablet	N/A	hydralazine
Minoxidil	tablet	N/A	minoxidil
Nitroprusside	injection <sup>^</sup>	Nitropress <sup>®*</sup> , Nipride <sup>®</sup>	none
<b>Combination Products</b>			
Isosorbide dinitrate and hydralazine	tablet	BiDil <sup>®</sup>	none

\*Generic is available in at least one dosage form or strength.

<sup>^</sup>Product is primarily administered in an institution.

PDL=Preferred Drug List

N/A=Not available

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the direct vasodilators are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Direct Vasodilators**

Clinical Guideline	Recommendations
Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b> <sup>5</sup>	<ul style="list-style-type: none"> <li>Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic</li> </ul>

Clinical Guideline	Recommendations
	<p>blood pressure &lt;140 mm Hg.</p> <ul style="list-style-type: none"> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>6</sup></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><u>Pharmacological Treatment</u></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment               <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment               <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy               <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indication for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>7</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk</li> </ul>

Clinical Guideline	Recommendations
	<p>factors.</p> <ul style="list-style-type: none"> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in</li> </ul>

Clinical Guideline	Recommendations
	<p>combination therapy.</p> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt;40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (<math>&lt;40\%</math>) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium <math>&lt;5.2</math> mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg</li> </ul>

Clinical Guideline	Recommendations
	<p>and DBP of &lt;80 mmHg.</p> <ul style="list-style-type: none"> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90th percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and</li> </ul>

Clinical Guideline	Recommendations
	<p>retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>8</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug</li> </ul>

Clinical Guideline	Recommendations
	<p>treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</p> <ul style="list-style-type: none"> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of</li> </ul>

Clinical Guideline	Recommendations
	<p>nitroprusside should be considered in case of emergency (pre-eclampsia).</p> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection</li> </ul>



Clinical Guideline	Recommendations
	<p>fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</p> <ul style="list-style-type: none"> <li>• In patients with HF<sub>r</sub>EF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HF<sub>p</sub>EF, BP treatment threshold and target values should be the same as for HF<sub>r</sub>EF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>9</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>

Clinical Guideline	Recommendations
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>10</sup></p>	<ul style="list-style-type: none"> <li>To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group: <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>11</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>12</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq 140/90</math>) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq 180/110</math> and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery</li> </ul>

Clinical Guideline	Recommendations
	<p>disease.</p> <ul style="list-style-type: none"> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events..</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for</li> </ul>

Clinical Guideline	Recommendations
	<p>the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (&lt;30 mg/g creatinine), and normal estimated glomerular filtration rate.</p> <ul style="list-style-type: none"> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate &lt;30 mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American College of Cardiology/American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults</b> (2017)<sup>13</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>• Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) ≥130 mmHg or an average diastolic blood pressure (DBP) of ≥80 mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of ≥10% and an average SBP of ≥130 mmHg or an average ≥80 mmHg.</li> <li>• Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of ≥140 mmHg or a DBP of ≥90 mmHg.</li> <li>• Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>• For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of ≥10%, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>• Adults with SIHD and hypertension (BP ≥130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension</li> </ul>

Clinical Guideline	Recommendations
	<p>should be &lt;130 mmHg.</p> <ul style="list-style-type: none"> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce</li> </ul>

Clinical Guideline	Recommendations
	<p>the risk of recurrent stroke and other vascular event.</p> <ul style="list-style-type: none"> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to <math>&lt;140</math> mmHg during the first hour and to <math>&lt;120</math> mmHg in aortic dissection.</li> <li>For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>In patients with planned elective major surgery and SBP <math>\geq 180</math> mmHg or DBP <math>\geq 110</math> mmHg, deferring surgery may be considered.</li> <li>For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)<sup>14</sup></p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>In patients with LVEF <math>\leq 40\%</math>, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three β-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is &gt;30 mL/min/1.73 m<sup>2</sup> and serum potassium is &lt;5.0 mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at &lt;5.5 mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: A)</li> </ul>

Clinical Guideline	Recommendations
	<p>B)</p> <ul style="list-style-type: none"> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><b>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</b></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed</li> </ul>

Clinical Guideline	Recommendations
	<p>medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</p> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>American College of Cardiology Foundation/American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes (2014)</b><sup>15</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation &lt;90%, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications <ul style="list-style-type: none"> <li>○ Nitrates <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACE if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>○ Beta-adrenergic blockers <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval <math>&gt;0.24</math> second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</p> <ul style="list-style-type: none"> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> <ul style="list-style-type: none"> <li>○ Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval <math>&gt;0.24</math> seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</li> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> </li> <li>○ Other anti-ischemic interventions <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> </li> <li>● Inhibitors of renin-angiotensin-aldosterone system <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF <math>&lt;0.40</math> and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF <math>&lt;0.40</math> who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine <math>&gt;2.5</math> mg/dL in men or <math>&gt;2.0</math> mg/dL in women) or hyperkalemia (<math>K &gt;5.0</math> mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF <math>&lt;0.40</math>, diabetes mellitus, or heart failure.</li> </ul> </li> <li>● Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include: <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatid or tirofiban.</li> <li>▪ Fibrinolytic therapy in patients with definite NSTEMI-ACS</li> </ul> </li> </ul> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>● Antiplatelet agents <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatid, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> <li>● Anticoagulant therapy <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>● Timing of CABG in relation to use of antiplatelet agents</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>● Medications at discharge <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</li> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post-NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> </li> <li>● Late hospital and post-hospital oral antiplatelet therapy <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation (2020)</b><sup>16</sup></p>	<p><u>Pharmacological treatment of ischemia</u></p> <ul style="list-style-type: none"> <li>• Sublingual or intravenous nitrates and early initiation of <math>\beta</math>-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>• Continuation of chronic <math>\beta</math>-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>• Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>• In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> <li>• A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds. <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors</li> </ul>



Clinical Guideline	Recommendations
	<p>during PCI.</p> <ul style="list-style-type: none"> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul> <p><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li>• If at low bleeding risk (HAS-BLED <math>\leq 2</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED <math>\geq 3</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED <math>\geq 3</math> and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>17</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a</li> </ul>

Clinical Guideline	Recommendations
	<p>dilated LV in NYHA Class III–IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</p> <ul style="list-style-type: none"> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered</li> </ul>

Clinical Guideline	Recommendations
	<p>in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</p> <ul style="list-style-type: none"> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g., with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the direct vasodilators are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Direct Vasodilators<sup>1-4</sup>**

Indication	Single Entity Agents		Combination Products
	Hydralazine	Minoxidil	Isosorbide Dinitrate and Hydralazine
<b>Heart Failure</b>			
Treatment of heart failure as an adjunct to standard therapy in self-identified black patients to improve survival, to prolong time to hospitalization for heart failure, and to improve patient-reported functional status			✓
<b>Hypertension</b>			
Treatment of essential hypertension	✓ *		
Treatment of hypertension		✓ †	
Treatment of severe essential hypertension	✓ ‡		

\*Tablet: Alone or as an adjunct.

†Because of the potential for serious adverse effects, minoxidil tablet is only indicated for the treatment of hypertension that is symptomatic or associated with target organ damage and is not manageable with maximum therapeutic doses of a diuretic plus two other antihypertensive drugs. At the present time use in milder degrees of hypertension is not recommended because the benefit-risk relationship has not been defined.

‡Injection: When the drug cannot be given orally or when there is an urgent need to lower blood pressure.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the direct vasodilators are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Direct Vasodilators<sup>2</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Hydralazine	38 to 50	88 to 90	Liver, significant (%)	Renal (3 to 14) Feces (3 to 12)	3 to 5

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
			not reported)		
Minoxidil	90 to 100	Insignificant (% not reported)	Liver (90)	Renal (90) Feces (3)	4.2
<b>Combination Products</b>					
Isosorbide dinitrate and hydralazine	H: 10 to 26 I: ~25	H: 88 to 90 I: 28	H: Liver, significant (% not reported) I: Liver, significant (% not reported)	H: Not reported I: Not reported	H: 3 to 5 I: 2

H=hydralazine, I=isosorbide dinitrate

## V. Drug Interactions

Significant drug interactions with the direct vasodilators are listed in Table 5.

**Table 5. Significant Drug Interactions with the Direct Vasodilators<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
Nitrates and nitrites	Phosphodiesterase type 5 inhibitors	Sildenafil may potentiate the hypotensive effects of nitrates. The use of these agents in combination is contraindicated.
Nitrates and nitrites	Avanafil	Concurrent use of avanafil and isosorbide dinitrate may result in potentiation of hypotensive effects.
Nitrates and nitrites	Riociguat	Concurrent use of riociguat and nitrates may result in increased risk of hypotension.

## VI. Adverse Drug Events

The most common adverse drug events reported with the direct vasodilators are listed in Table 6. The boxed warning for minoxidil is listed in Table 7.

**Table 6. Adverse Drug Events (%) Reported with the Direct Vasodilators<sup>1-4</sup>**

Adverse Events	Single Entity Agents		Combination Products
	Hydralazine	Minoxidil	Isosorbide Dinitrate and Hydralazine
<b>Cardiovascular System</b>			
Angina pectoris	✓	✓	16
Cardiovascular collapse	-	-	✓
Crescendo angina	-	-	✓
Electrocardiogram changes	-	60	-
Flushing	✓	-	✓
Heart failure	-	✓	-
Hypotension	-	-	8
Orthostatic hypotension	✓	-	✓
Pallor	-	-	✓
Palpitations	✓	-	4
Paradoxical pressor response	✓	-	✓
Peripheral edema	✓	7	✓
Pericardial effusion with tamponade	-	3	-
Pericarditis	-	✓	-
Postural hypotension	-	✓	✓
Rebound hypertension	-	-	✓
Shock	-	-	✓

Adverse Events	Single Entity Agents		Combination Products
	Hydralazine	Minoxidil	Isosorbide Dinitrate and Hydralazine
Syncope	-	-	✓
Tachycardia	✓	✓	2
Vascular collapse	✓	-	✓
Ventricular tachycardia	-	-	4
<b>Central Nervous System</b>			
Anxiety	✓	-	✓
Asthenia	✓	-	✓
Chills	✓	-	✓
Depression	✓	-	✓
Disorientation	✓	-	✓
Dizziness	✓	-	32
Fever	✓	-	✓
Headache	✓	-	50
Lightheadedness	-	-	✓
Psychotic reaction	✓	-	✓
Restlessness	-	-	✓
<b>Dermatological</b>			
Alopecia	-	-	1
Hypertrichosis	-	80	-
Pruritus	✓	-	✓
Rash	✓	✓	✓
Stevens-Johnson syndrome	-	✓	-
Urticaria	✓	-	✓
<b>Endocrine and Metabolic</b>			
Breast tenderness	-	✓	-
Fluid and electrolyte imbalance	-	✓	-
Hyperglycemia	-	-	4
Hyperlipidemia	-	-	3
<b>Gastrointestinal</b>			
Anorexia	✓	-	✓
Bowel incontinence	-	-	✓
Constipation	✓	-	✓
Diarrhea	✓	-	✓
Nausea	✓	✓	10
Paralytic ileus	✓	-	✓
Vomiting	✓	✓	4
Weight gain	-	✓	-
Xerostomia	-	-	✓
<b>Genitourinary</b>			
Blood urea nitrogen increased	-	✓	-
Dysuria	✓	-	✓
Impotence	✓	-	✓
Serum creatine increased	-	✓	-
Urinary incontinence	-	-	✓
<b>Hematological</b>			
Agranulocytosis	✓	-	✓
Eosinophilia	✓	-	✓
Erythrocyte count reduced	✓	✓	✓
Hematocrit decreased	-	✓	-
Hemoglobin decreased	✓	✓	✓
Hemolytic anemia	✓	-	✓
Leukopenia	✓	✓	✓

Adverse Events	Single Entity Agents		Combination Products
	Hydralazine	Minoxidil	Isosorbide Dinitrate and Hydralazine
Methemoglobinemia	-	-	✓
Thrombocytopenia	✓	✓	✓
<b>Hepatic</b>			
Alkaline phosphatase increased	-	✓	-
Cholecystitis	-	-	1
<b>Musculoskeletal</b>			
Arthralgia	-	-	1
Muscle cramps	✓	-	✓
Myalgia	-	-	1
Paresthesia	-	-	4
Peripheral neuritis	✓	-	✓
Rheumatoid arthritis	✓	-	✓
Tendon disorder	-	-	1
Tremor	✓	-	✓
Weakness	✓	-	14
<b>Ocular</b>			
Blurred vision	-	-	✓
Conjunctivitis	✓	-	✓
Lacrimation	✓	-	✓
<b>Respiratory</b>			
Bronchitis	-	-	8
Dyspnea	✓	-	✓
Nasal congestion	✓	-	✓
Pulmonary edema	-	✓	-
Rhinitis	-	-	4
Sinusitis	-	-	4
<b>Other</b>			
Allergic reactions	-	-	1
Angioedema	-	-	1
Diaphoresis	✓	-	1
Drug-induced lupus-like syndrome	✓	-	✓

✓ Percent not specified  
- Event not reported

**Table 7. Boxed Warning for Minoxidil<sup>1</sup>**

<b>WARNING</b>
<p>Minoxidil may produce serious adverse effects. It can cause pericardial effusion, occasionally progressing to tamponade, and it can exacerbate angina pectoris. Reserve for hypertensive patients who do not respond adequately to maximum therapeutic doses of a diuretic and two other antihypertensive agents.</p> <p>In experimental animals, minoxidil caused several kinds of myocardial lesions and other adverse cardiac effects.</p> <p>Administer under close supervision, usually concomitantly with a <math>\beta</math>-adrenergic blocking agent, to prevent tachycardia and increased myocardial workload. Usually, it must be given with a diuretic, frequently one acting in the ascending limb of the loop of Henle to prevent serious fluid accumulation. When first administering minoxidil, hospitalize and monitor patients with malignant hypertension and those already receiving guanethidine to avoid too rapid or large orthostatic decreases in blood pressure.</p>

## VII. Dosing and Administration

The usual dosing regimens for the direct vasodilators are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Direct Vasodilators<sup>1-4</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Hydralazine	<u>Essential hypertension:</u> Injection, tablet: initial, 10 mg four times daily for the first two to four days, followed by 25 mg four times daily for the balance of the first week, then for the second and subsequent weeks, increase dosage to 50 mg four times daily; maintenance, adjust dosage to the lowest effective levels	<u>Essential hypertension:</u> Safety and effectiveness in pediatric patients have not been established in controlled clinical trials, although there is experience with the use of hydralazine in pediatric patients.  Injection, tablet: initial, 0.75 mg/kg/day administered in four divided doses; maintenance, dosage may be increased gradually over the next three to four weeks; maximum, 7.5 mg/kg or 200 mg/day	Injection: 20 mg/mL  Tablet: 10 mg 25 mg 50 mg 100 mg
Minoxidil	<u>Hypertension:</u> Tablet: initial, 5 mg/day and increase gradually every three days; maintenance, 10 to 40 mg daily in single or divided doses; maximum, 100 mg/day	<u>Hypertension:</u> Tablet: initial, 0.2 mg/kg/day; maintenance, 0.25 to 1 mg/kg/day; maximum, 50 mg/day	Tablet: 2.5 mg 10 mg
<b>Combination Products</b>			
Isosorbide dinitrate and hydralazine	<u>Heart failure:</u> Tablet: initial, 20-37.5 mg three times daily; maximum, 40-75 mg (two tablets) three times daily	The safety and effectiveness have not been established in children.	Tablet: 20-37.5 mg



## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the direct vasodilators are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Direct Vasodilators**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Heart Failure</b>				
Unverferth et al. <sup>18</sup> (1983)  Hydralazine 225 mg/day  vs  ISDN 160 mg/day  vs  hydralazine and ISDN (individual agents)  vs  placebo	DB, PC, RCT  Patients with idiopathic dilated cardiomyopathy were evaluated to determine the hemodynamic and morphologic effects of vasodilator therapy	N=49  3 months	Primary: Echocardiographic percent change of left ventricular diameter, the systolic time intervals ratio of PEP/LVET, the pulmonary capillary wedge pressure, mean pulmonary artery pressure, pulmonary vascular resistance, cardiac index, and SVR  Secondary: Not reported	Primary: For the percent change in left ventricular diameter and PEP/LVET, a significant improvement with hydralazine and combination therapy (P<0.05) was seen compared to ISDN alone or placebo.  Significant decrease with ISDN and combination therapy vs placebo or hydralazine alone (P<0.05) was seen for pulmonary capillary wedge pressure, mean pulmonary artery pressure, and the pulmonary vascular resistance.  Hydralazine resulted in a decrease in SVR and increase in cardiac index from 2.5±0.4 to 3.1±0.4 L/min/m <sup>2</sup> vs placebo or ISDN alone (P<0.05).  Combination therapy resulted in a decrease in SVR and cardiac index increased from 2.3±0.4 to 3.1±0.4 L/min/m <sup>2</sup> (P<0.01).  There was no improvement in SVR or cardiac index with ISDN alone or with placebo.  Myocardial cell diameter decreased from 25.4±3.1 microns at baseline to 23.1±3.8 microns with hydralazine (P<0.05). Combination therapy decreased its cell diameter from 23.9±3.7 to 22.2±2.2 microns (P<0.05).  There was no change in the myocardial cell diameter seen in patients treated with ISDN alone or with placebo.  Secondary: Not reported
Taylor <sup>19</sup> (2005) A-HeFT	DB, MC, PC, RCT  African American	N=1,050  6 to 18 months	Primary: Composite score (all-cause	Primary: Mortality in the fixed-dose ISDN and hydralazine group was 6.2% compared to 10.2% in the placebo group (P=0.02).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
ISDN and hydralazine 60-112.5 mg/day in 3 divided doses, titrated up to ISDN and hydralazine 120-225 mg/day in 3 divided doses (fixed-dose combination product)  vs  placebo	patients with moderate-to-severe symptomatic heart failure, classified NYHA class III to IV heart failure with dilated ventricles and low ejection fractions		mortality, first hospitalization for heart failure, and quality of life at 6 months as measured by the Minnesota Living with Heart Failure questionnaire)  Secondary: Not reported	Survival was increased by 43% in the active treatment arm (HR, 0.57; P=0.02).  The composite score and all individual components of the composite score were significantly and positively impacted by treatment with ISDN and hydralazine (primary composite score P=0.01, death from any cause P=0.02, first hospitalization for heart failure P=0.001, change in quality of life score at 6 months P=0.02).  The study was prematurely terminated in as a result of the significantly improved survival in the ISDN and hydralazine group.  Secondary: Not reported
Taylor et al. <sup>20</sup> (2004) A-HeFT  ISDN and hydralazine 20-37.5 mg TID, increased to ISDN and hydralazine 40-75 mg TID (fixed-dose combination product)  vs  placebo	DB, MC, PC, RCT  Patients ≥18 years of age, self-identified as of African descent, with NYHA class III or IV heart failure on standard therapy for ≥3 months and evidence of left ventricular dysfunction within the prior 6 months	N=1,050  18 months (mean duration of follow-up was 10 months)	Primary: A composite score made up of weighted values for death from any cause, a first hospitalization for heart failure, and quality of life changes  Secondary: Individual components of the primary composite score	Primary: From a range of possible scores of -6 to 2 for the composite endpoint, patients in the active treatment group achieved a significantly better score of -0.1±1.9 compared -0.5±2.0 in the placebo group (P=0.01).  Secondary: There was a significantly higher mortality rate in the placebo group compared to the ISDN and hydralazine group (6.2 vs 10.2%; P=0.02). Survival was increased by 43% in the active treatment group (HR, 0.57; P=0.02). This led to the early termination of the trial.  Compared to the placebo group, the rate of first hospitalization for heart failure was significantly reduced in the ISDN and hydralazine group (16.4 vs 24.4%; P=0.001).  There was a significant improvement in quality of life scores found with the ISDN and hydralazine group when compared to the placebo group (-5.6±20.6 vs -2.7±21.2; P=0.02).
Taylor et al. <sup>21</sup> (2007) A-HeFT	Post-hoc analysis of A-HeFT	N=1,050  Mean duration	Primary: Cause specific mortality, event	Primary: Cardiovascular deaths were significantly reduced in the treatment group compared to the placebo group (5.0 vs 8.5%; P=0.027). Pump failure

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ISDN and hydralazine 20-37.5 mg TID, increased to ISDN and hydralazine 40-75 mg TID (fixed-dose combination product)</p> <p>vs</p> <p>placebo</p>	<p>Patients <math>\geq</math>18 years of age, self-identified as of African descent, with NYHA class III or IV heart failure on standard therapy for <math>\geq</math>3 months and evidence of left ventricular dysfunction within the prior 6 months</p>	<p>of follow-up was 18 months</p>	<p>free survival (time to either death or first hospitalization and time to first hospitalization for heart failure)</p> <p>Secondary: Subgroup analysis</p>	<p>death was also significantly reduced (75%) compared to the placebo group (0.8 vs 3.0%; <math>P=0.012</math>). There were no significant differences between the groups for other causes of death.</p> <p>In the treatment group event-free survival (death or first hospitalization for heart failure) was significantly improved compared to the placebo group (HR, 0.63; 95% CI, 0.49 to 0.81; <math>P&lt;0.001</math>).</p> <p>The time to first hospitalization for heart failure was also significantly reduced (HR, 0.61; 95% CI, 0.46 to 0.80; <math>P&lt;0.001</math>).</p> <p>Secondary: A consistent beneficial effect was seen in the treatment sub groups (age, sex, baseline BP, history of chronic renal insufficiency, presence of diabetes, cause of heart failure, and baseline medication use) on primary composite score and event-free survival.</p>
<p>Anand et al.<sup>22</sup> (2014) A-HeFT</p> <p>ISDN and hydralazine 20-37.5 mg TID, increased to ISDN and hydralazine 40-75 mg TID (fixed-dose combination product)</p> <p>vs</p> <p>placebo</p>	<p>Post-hoc analysis of A-HeFT</p> <p>Patients <math>\geq</math>18 years of age, self-identified as of African descent, with NYHA class III or IV heart failure on standard therapy for <math>\geq</math>3 months and evidence of left ventricular dysfunction within the prior 6 months</p>	<p>N=1,050</p> <p>Mean duration of follow-up was 18 months</p>	<p>Primary: Mortality, all hospitalizations including recurrences (first hospitalizations only, all hospitalizations including recurrences, 30-day all-cause readmission rates)</p> <p>Secondary: Not reported</p>	<p>Primary: During a median follow-up of 450 days, 86 (8.2%) patients died. The cumulative mortality was significantly lower (HR, 0.57; 95% CI, 0.37 to 0.89; <math>P=0.013</math>) in the treatment group vs the placebo group.</p> <p>When deaths were analyzed as a competing risk for first hospitalizations, the effect (HR) of treatment was 0.88 (95% CI, 0.72 to 1.06; <math>P=0.18</math>) on hospitalization for any cause and 0.61 (95% CI, 0.47 to 0.80; <math>P&lt;0.001</math>) on heart failure hospitalizations.</p> <p>The use of fixed-dose combination product was associated with a significant 25% reduction in all hospitalizations for any cause (HR, 0.75; 95% CI, 0.63 to 0.91; <math>P=0.003</math>) and a 34% reduction in all heart failure hospitalizations (HR, 0.66; 95% CI, 0.52 to 0.83; <math>P&lt;0.001</math>).</p> <p>Of the subjects who had at least one admission for heart failure and were discharged alive, 29 of 123 (23.6%) in the placebo group and 12 of 81 (14.8%) in the combination product group were readmitted for any cause <math>&lt;</math>30 days of being discharged from their first hospitalization for heart failure. This reduction in the 30-day all-cause readmissions by the combination product was not statistically significant.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Yancy et al. <sup>23</sup> (2007) A-HeFT  ISDN and hydralazine 20-37.5 mg TID, increased to ISDN and hydralazine 40-75 mg TID (fixed-dose combination product)  vs  placebo	ES, OL  Patients previously enrolled in A-HeFT with NYHA class I to IV heart failure symptoms while receiving background therapy and satisfying the A-HeFT inclusion criteria	N=158  12 months or until ISDN and hydralazine approved by the FDA	Primary: Compliance with study drug, safety, tolerability  Secondary: Change in NYHA association class, death, hospitalization for heart failure	Primary: Compliance in the treatment group averaged 87±25%, with no significant difference when compared to the placebo group.  There were no significant differences in adverse events between the groups.  Secondary: No significant difference was seen in hospitalizations from heart failure according to randomization.  The greatest improvement in heart failure symptoms occurred in NYHA class III (at baseline) compared to other classes (P<0.001).  Overall most patients were unchanged with 24% showing improved NYHA class and 9% showing a worsening.
Cohn et al. <sup>24</sup> (1986) V-HeFT I  Hydralazine 300 mg/day plus ISDN 160 mg/day (individual agents, concurrent therapy)  vs  prazosin 20 mg/day  vs  placebo	AC, DB, PC, RCT  Men with impaired cardiac function and reduced exercise tolerance on digoxin and a diuretic	N=642  3 years	Primary: Mortality  Secondary: Effect on left ventricular function	Primary: There was a 34% risk reduction in mortality by two years in the hydralazine plus ISDN group compared to placebo (P<0.028).  Cumulative mortality rates of 25.6 and 36.2% were observed in the hydralazine plus ISDN group at 2 and 3 years respectively, compared to 34.3 and 46.9% in the placebo group. The results found in the prazosin group were similar to placebo.  Secondary: A significant increase in the left ventricular ejection fraction was reported at eight weeks and one year in the hydralazine plus ISDN treatment group, but not in either the prazosin or placebo groups.
Cohn et al. <sup>25</sup>	AC, DB, MC, RCT	N=804	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1991) V-HeFT II</p> <p>Hydralazine 300 mg/day plus ISDN 160 mg/day</p> <p>vs</p> <p>enalapril 20 mg/day</p>	<p>Men between the ages of 18 and 75 years with chronic heart failure receiving digoxin and diuretic therapy</p>	<p>2 years</p>	<p>Mortality</p> <p>Secondary: Peak oxygen consumption during exercise, LVEF</p>	<p>Mortality after two years was significantly lower in the group treated with enalapril (18%) than hydralazine plus isosorbide dinitrate (25%; P=0.016), and overall mortality tended to be lower (P=0.08).</p> <p>The lower mortality in the enalapril arm was attributable to a reduction in the incidence of sudden death, and this beneficial effect was more prominent in patients with less severe symptoms (NYHA class I or II).</p> <p>Secondary: Peak oxygen consumption during exercise was increased only by hydralazine plus isosorbide dinitrate (P&lt;0.05).</p> <p>While LVEF increased with both regimens during the two years after randomization, LVEF increased more (P&lt;0.05) during the first 13 weeks in the hydralazine plus isosorbide dinitrate group.</p>
<p>Mullens et al.<sup>26</sup> (2009)</p> <p>Isosorbide dinitrate and hydralazine (I/H) added to an ACE inhibitor or angiotensin receptor blockers</p> <p>vs</p> <p>ACE inhibitor or angiotensin receptor blockers</p> <p>Titration of oral drugs was aimed to wean off parental therapy and based on maintaining a target mean</p>	<p>PRO</p> <p>Patients ≥18 years of age with advanced decompensate heart failure with a cardiac index &lt;2.2 L/min/m<sup>2</sup> who were admitted to the hospital for intensive medical therapy</p>	<p>N=239</p> <p>Mean 26.3 months</p>	<p>Primary: All-cause mortality, cardiac transplantation, and first readmission for heart failure after index hospitalization discharge</p> <p>Secondary: Not reported</p>	<p>Primary: Patients receiving I/H had lower all-cause mortality (34 vs 41%; OR, 0.65; 95% CI, 0.43 to 0.99, P=0.04) and lower all-cause mortality/heart failure rehospitalization (70% vs 85%; OR, 0.72; 95% CI, 0.54 to 0.97; P=0.03) compared to the control group. There was no difference in overall cardiac transplantation or heart failure rehospitalization rates among the treatment groups.</p> <p>The improved outcomes in the I/H group was independent of race; however, there was a trend toward improved outcomes in African-Americans (all-cause mortality for whites in the I/H group, OR 0.66; 95% CI, 0.4 to 0.98; P=0.05; all-cause mortality for African-Americans in the I/H group, OR 0.44; 95% CI, 0.23 to 0.85; P=0.01).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
arterial pressure of 65 to 70 mm Hg and/or systolic blood pressure >85 mm Hg				
<b>Hypertension</b>				
Johnson et al. <sup>27</sup> (1983)  Minoxidil 5 to 40 mg/day as add-on therapy  vs  hydralazine 25 to 200 mg/day as add-on therapy	DB, RCT  Patients with normal renal function receiving HCTZ or propranolol (doses unknown) with DBP >95 mmHg	N=36  28 weeks	Primary: Percentage of patients with DBP <90 mmHg at weeks 4 and 28  Secondary: Not reported	Primary: There were greater response rates (DBP <90 mmHg) with minoxidil (69%) vs hydralazine (35%) at week four.  At week 28, there were greater response rates (DBP <90 mmHg) with minoxidil (55%) vs hydralazine (40%).  Secondary: Not reported
Bevan et al. <sup>28</sup> (1993)  Captopril (unknown dose)  vs  hydralazine (unknown dose)  vs  nifedipine (unknown dose)  vs  placebo	DB, PC, RCT  Patients with inadequately controlled HTN, despite treatment with atenolol 100 mg/day and bendrofluazide* 5 mg/day	N=160  12 weeks	Primary: Comparative antihypertensive, biochemical, adverse effects  Secondary: Not reported	Primary: Mean supine blood pressure changes: captopril 13.4/10.3 mmHg, hydralazine 15.0/10.0 mmHg, and nifedipine 16.8/8.1 mmHg (differences not significant).  Erect blood pressure changes were similar; target blood pressure (<140/95 mmHg) was achieved in 33% with captopril, 29% with hydralazine, 17% with nifedipine, and 10% with placebo.  Compared to other agents, captopril increased serum potassium (value not reported; P=0.01).  Mean changes in serum cholesterol: captopril -0.2 mmol/L, hydralazine -0.8 mmol/L, nifedipine -0.2 mmol/L, and placebo 0.2 mmol/L (P<0.001).  Side effects did not differ significantly between the groups. Withdrawal rates: captopril 15%, hydralazine 24%, nifedipine 22%, and placebo 3% (P=0.04).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Julien et al.<sup>29</sup> (1990)</p> <p>Captopril 150 to 300 mg/day</p> <p>vs</p> <p>minoxidil 7.5 to 30 mg/day</p>	<p>DB, PG, RCT</p> <p>Male patients with left ventricular hypertrophy and essential HTN with DBP &gt;95 mmHg who were taking metoprolol 200 mg/day and furosemide 80 mg/day</p>	<p>N=34</p> <p>6 months</p>	<p>Primary: Blood pressure changes and left ventricular hypertrophy changes as seen on electrocardiogram</p> <p>Secondary: Not reported</p>	<p>Primary: Blood pressure decreased significantly in both groups; captopril (163/102 to 135/89 mmHg) and minoxidil (160/99 to 137/87 mmHg; P&lt;0.001).</p> <p>Electrocardiogram criteria for left ventricular hypertrophy improved with captopril only with a decrease in intraventricular septum, posterior wall, and left ventricular mass (17.4 to 15.9 mm; P&lt;0.05, 14.5 to 13.4 mm; P&lt;0.05 and 236 to 198 g/m<sup>2</sup>; P&lt;0.001, respectively). No changes on electrocardiogram criteria with minoxidil.</p> <p>Secondary: Not reported</p>
<p>McAreevey et al.<sup>30</sup> (1984)</p> <p>Hydralazine 12.5 mg QD up to 100 mg BID</p> <p>vs</p> <p>labetalol 200 mg QD up to 1,600 mg BID</p> <p>vs</p> <p>methyldopa 125 mg QD up to 1,000 mg BID</p> <p>vs</p> <p>prazosin 0.5 mg QD up to 10 mg BID</p> <p>vs</p>	<p>DB, PG, RCT</p> <p>Patients with inadequately controlled HTN while receiving atenolol 100 mg/day and bendrofluazide* 5 mg/day</p>	<p>N=238</p> <p>6 months</p>	<p>Primary: Comparative safety and efficacy, target blood pressure &lt;140/95 mm Hg</p> <p>Secondary: Not reported</p>	<p>Primary: Target blood pressure was reached in 25% of patients receiving hydralazine, 23% of patients receiving minoxidil, 19% of patients receiving prazosin, 17% of patients receiving methyldopa and zero percent of patients receiving placebo (P values not reported).</p> <p>Labetalol had the highest withdrawal rate compared to the other treatments with 78% (P&lt;0.05). Minoxidil had the second highest withdrawal rate with 57% (P&lt;0.05), due to fluid retention. There were no significant differences in withdrawal rates among the other treatments.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>placebo</p> <p>Minoxidil as add-on therapy was given to men only.</p>				
<p>Materson et al.<sup>31</sup> (1990)</p> <p>Hydralazine 25, 50 or 100 mg BID</p> <p>vs</p> <p>metoprolol 50, 100 or 200 mg BID</p> <p>vs</p> <p>methyldopa 250, 500 or 1,000 mg BID</p> <p>vs</p> <p>reserpine 0.05, 0.10 or 0.25 mg QD</p> <p>All patients received HCTZ 25 to 100 mg QD.</p>	<p>DB, MC, RCT</p> <p>Men <math>\geq 60</math> years with HTN not currently receiving antihypertensive therapy with a DBP 90 to 114 mm Hg and a SBP <math>&lt; 240</math> mm Hg; or a DBP <math>&lt; 100</math> mm Hg and a SBP <math>&lt; 240</math> mm Hg if currently taking antihypertensive therapy and the blood pressure criteria was met after <math>\geq 2</math> weeks without medication</p>	<p>N=690</p> <p>12 months</p>	<p>Primary: Average reduction in SBP, DBP, the number of patients achieving the goal blood pressure and the average change in heart rate</p> <p>Secondary: Rates of drug intolerances and incidence of adverse effects</p>	<p>Primary: A total of 269 patients were uncontrolled with HCTZ therapy alone and were randomized to receive hydralazine (n=68), methyldopa (n=71), metoprolol (n=65), or reserpine (n=65).</p> <p>A total of 213 of the 269 patients achieved goal blood pressure with the addition of one of four therapies was added to HCTZ and entered the 6 month maintenance phase; 186 patients completed the maintenance phase.</p> <p>Across all four add-on therapies, there was an additional average reduction in blood pressure of 13.1/10.6 mm Hg. The average reduction in SBP (mm Hg)<math>\pm</math>SD from baseline to endpoint for hydralazine, methyldopa, metoprolol, and reserpine add-on therapies was: <math>-11.5\pm 10.1</math> (P<math>&lt;0.001</math>), <math>-15.0\pm 13.7</math> (P<math>&lt;0.001</math>), <math>-13.0\pm 15.4</math> (P<math>&lt;0.001</math>), and <math>-12.7\pm 11.5</math> (P<math>&lt;0.001</math>), respectively. There was no statistically significant difference in SBP reductions among the different groups (P=0.43).</p> <p>The average reduction in DBP (mm Hg)<math>\pm</math>SD from baseline to endpoint for hydralazine, methyldopa, metoprolol, and reserpine add-on therapies was: <math>-11.3\pm 5.9</math> (P<math>&lt;0.001</math>), <math>-10.6\pm 6.3</math> (P<math>&lt;0.001</math>), <math>-10.6\pm 6.7</math> (P<math>&lt;0.001</math>), and <math>-9.8\pm 6.3</math> (P<math>&lt;0.001</math>), respectively. There was no statistically significant difference in DBP reductions among the different groups (P=0.59).</p> <p>The average change in heart rate (beats per minute) <math>\pm</math>SD from baseline to endpoint for hydralazine, methyldopa, metoprolol, and reserpine add-on therapies was: <math>1.4\pm 10.5</math> (P value not significant), <math>-1.6\pm 9.3</math> (P value not significant), <math>15.9\pm 11.9</math> (P<math>&lt;0.05</math>), and <math>-7.9\pm 10.7</math> (P<math>&lt;0.05</math>), respectively. There was a statistically significant difference in change in heart rate among the different groups (P<math>&lt;0.001</math>).</p> <p>The percentage of patients achieving the goal blood pressure at endpoint in</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>the hydralazine, methyldopa, metoprolol, and reserpine groups was: 85.3, 81.7, 76.9, and 72.3%, respectively (P=0.28).</p> <p>Secondary: Drug intolerance, defined as adverse effects prompting dose reduction or discontinuation, was present in 23.3% of those not achieving goal blood pressure compared to 2.8% of those achieving the goal blood pressure (P&lt;0.001). This was statistically significant in the hydralazine, methyldopa, and metoprolol groups, but not the reserpine group.</p> <p>There were 27 (10%) study terminations due to adverse drug events: hydralazine (n=3), methyldopa (n=8), metoprolol (n=9), and reserpine (n=7). There were 2 study terminations in the methyldopa-treated group and 1 in the reserpine group due to depression.</p> <p>The overall incidence of volunteered moderate or severe adverse effects, not prompting study termination was significantly greater (P&lt;0.01) with methyldopa (31%) and hydralazine (25%) compared to reserpine (15%) or metoprolol (9%).</p>

\*Synonym for bendroflumethiazide.

Drug regimen abbreviations: BID=twice daily, QD=once daily, TID=three times daily

Study abbreviations: AC=active controlled, DB=double blind, ES=extended study, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial

Miscellaneous abbreviations: ACE=angiotensin converting enzyme, CI=confidence interval, DBP=diastolic blood pressure, FDA=Food and Drug Administration, HCTZ=hydrochlorothiazide, HR=hazard ratio, HTN=hypertension, ISDN=isosorbide dinitrate, LVEF=left ventricular ejection fraction, LVET=left ventricular ejection time, NYHA=New York Heart Association, PEP=pre-ejection period, SBP=systolic blood pressure, SD=standard deviation, SVR=systemic vascular resistance

### Additional Evidence

#### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

## IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Direct Vasodilators**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Hydralazine	injection, tablet	N/A	N/A	\$
Minoxidil	tablet	N/A	N/A	\$
<b>Combination Products</b>				
Isosorbide dinitrate and hydralazine	tablet	BiDil®	\$\$\$\$\$	N/A

\*Generic is available in at least one dosage form or strength.

N/A=not available

## X. Conclusions

Hydralazine and minoxidil are approved for the treatment of hypertension, and both agents are available in a generic formulation.<sup>1,2,4</sup> There are several national and international organizations that have published guidelines on the treatment of hypertension. Most of the guidelines do not provide recommendations on the use of the oral direct vasodilators.<sup>5-11</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension. According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs, β-

blockers, calcium channel blockers).<sup>5</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>5-11</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>5-11</sup> Clinical trials have demonstrated that hydralazine and minoxidil are effective for the treatment of hypertension when added to existing therapy in patients whose blood pressure is inadequately controlled. There are limited head-to-head trials comparing the direct vasodilators.<sup>27-31</sup> These agents are associated with several potentially severe adverse effects, which limits their use in the treatment of hypertension.<sup>1,2</sup>

Hydralazine and isosorbide dinitrate (administered as single entity products) have been used off-label to treat heart failure for many years. The combination of these agents has been shown to reduce mortality compared to placebo in patients receiving standard therapy with digoxin and diuretics.<sup>24</sup> However, when hydralazine and isosorbide dinitrate were directly compared to an ACE inhibitor, mortality was significantly lower in the ACE inhibitor group.<sup>25</sup> Treatment guidelines for the management of heart failure currently recommend the use of hydralazine and an oral nitrate in patients who do not tolerate an ACE inhibitor or ARB.<sup>13-17</sup> The fixed-dose combination of isosorbide dinitrate and hydralazine is FDA-approved for the treatment of heart failure as an adjunct to standard therapy in self-identified black patients.<sup>3</sup> In the A-HeFT trial, the use of this combination product improved mortality, prolonged time to hospitalization for heart failure, and improved functional status compared to placebo. The patients in this trial were also receiving standard heart failure therapy prior to enrollment (ACE inhibitors, angiotensin II receptor antagonists,  $\beta$ -blockers, diuretics, digoxin, spironolactone).<sup>3,19-22</sup> The Heart Failure Society of America and the American College of Cardiology Foundation/American Heart Association recommend the use of the fixed-dose combination of isosorbide dinitrate and hydralazine in African American patients with NYHA functional class III or IV heart failure who are on a standard regimen including an ACE inhibitor (or ARB) and a  $\beta$ -blocker.<sup>14</sup> Both hydralazine and isosorbide dinitrate are available generically; however, generic hydralazine is not available in a strength equivalent to the fixed-dose combination product.<sup>1-4</sup>

Therefore, all brand direct vasodilators within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use. The fixed-dose combination of isosorbide dinitrate and hydralazine (BiDil<sup>®</sup>) should be available through the medical justification portion of the prior authorization process as an adjunct to standard heart failure therapy in self-identified black patients.

## **XI. Recommendations**

No brand direct vasodilator is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. *Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.*
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. BiDil® [package insert]. Atlanta (GA): Arbor Pharmaceuticals, LLC; 2019 Nov.
4. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
5. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
6. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
7. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
8. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
9. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
10. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
11. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
12. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
13. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
14. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
15. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
16. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>.
17. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.

18. Unverferth DV, Mehegan JP, Magorien RD, Unverferth BJ, Leier CV. Regression of myocardial cellular hypertrophy with vasodilator therapy in chronic congestive heart failure associated with idiopathic dilated cardiomyopathy. *Am J Cardiol.* 1983 May 1;51(8):1392-8.
19. Taylor AL. The African American Heart Failure Trial: a clinical trial update. *Am J Cardiol.* 2005 Oct 10;96(7B):44-8.
20. Taylor AL, Ziesche S, Yancy C, Carson P, D'Agostino R Jr, Ferdinand K, et al. African-American Heart Failure Trial Investigators. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *N Engl J Med.* 2004 Nov 11;351(20):2049-57.
21. Taylor AL, Ziesche S, Yancy CW, Carson P, Ferdinand K, Taylor M, et al; African-American Heart Failure Trial Investigators. Early and sustained benefit on event-free survival and heart failure hospitalization from fixed-dose combination of isosorbide dinitrate/hydralazine: consistency across subgroups in the African-American Heart Failure Trial. *Circulation.* 2007 Apr 3;115(13):1747-53.
22. Anand IS, Win S, Rector TS, Cohn JN, and Taylor AL. Effect of fixed-dose combination of isosorbide dinitrate and hydralazine on all hospitalizations and on 30-day readmission rates in patients with heart failure: results from the African-American Heart Failure Trial. *Circ Heart Fail.* 2014 Sep;7(5):759-765.
23. Yancy CW, Ghali JK, Braman VM, Sabolinski ML, Worcel M, Archambault WT, et al. Evidence for the continued safety and tolerability of fixed-dose isosorbide dinitrate/hydralazine in patients with chronic heart failure (the extension to African-American Heart Failure Trial). *Am J Cardiol.* 2007 Aug 15;100(4):684-9.
24. Cohn JN, Archibald DG, Ziesche S, et al. Effect of vasodilator therapy on mortality in chronic congestive heart failure: Results of a Veterans Administration Cooperative Study (VHEFT-I). *N Engl J Med.* 1986; 314:1547-52.
25. Cohn JN, Ziesche S, Johnson G, Cobb F, Francis G, Tristani F, et al. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med.* 1991 Aug 1;325(5):303-10.
26. Mullens W, Abrahams Z, Francis GS, et al. Usefulness of Isosorbide Dinitrate and Hydralazine as add-on therapy in patients discharged for advanced decompensated heart failure. *Am J Cardiol* 2009;103:1113-9.
27. Johnson BF, Black HR, Becker R, Weiner B, Angeletti F. A Comparison of minoxidil and hydralazine in non-azotemic hypertensives. *J Hypertens.* 1983 Jun;1(1):103-7.
28. Bevan EG, Pringle SD, Waller PC, Herrick AL, Findlay JG, Murray GD, et al. Comparison of captopril, hydralazine, and nifedipine as third drug in hypertensive patients. *J Hum Hypertens.* 1993 Feb;7(1):83-8.
29. Julien J, Dufloux MA, Prasquier R, Chatellier G, Menard D, Plouin PF, et al. Effects of captopril and minoxidil on left ventricular hypertrophy in resistant hypertensive patients: A 6 month double-blind comparison. *J Am Coll Cardiol.* 1990 Jul;16(1):137-42.
30. McAreavey D, Ramsey LE, Latham L, McLaren AD, Lorimer AR, Reid JL, et al. Third drug trial: comparative study of antihypertensive agents added to treatment when blood pressure remains uncontrolled by a beta blocker plus thiazide diuretic. *Br Med J (Clin Res Ed).* 1984 Jan 14;288(6411):106-11.
31. Materson BJ, Cushman WC, Goldstein G, et al. Treatment of hypertension in the elderly: I. Blood pressure and clinical changes: results of a Department of Veterans Affairs cooperative study. *Hypertension.* 1990 Apr;15(4):348-60.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Peripheral Adrenergic Inhibitors  
AHFS Class 240832  
May 4, 2022**

**I. Overview**

In 2016, reserpine was discontinued. Currently, there are no drugs classified by AHFS as peripheral adrenergic inhibitors.

**II. Conclusions**

There are no drugs available in the peripheral adrenergic inhibitor class (AHFS Class 240832).

**III. Recommendations**

No brand peripheral adrenergic inhibitor is recommended for preferred status. Alabama Medicaid should continue to include AHFS Class 240832 in the PDL screening process. If new outpatient peripheral adrenergic inhibitors are added, it is recommended that this class be re-reviewed at that time.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Hypotensive Agents, Miscellaneous  
AHFS Class 240892  
May 4, 2022**

**I. Overview**

Mecamylamine was one of the first oral antihypertensive agents, introduced in the mid-1950s under the trade name Inversine<sup>®</sup>. It was withdrawn from the market in 2009 due to increased competition of antihypertensive drugs and decreasing use of the agent. In March 2013, mecamylamine was issued FDA approval and re-entered the market under the name of Vecamyl<sup>®</sup>.<sup>1,2</sup> Mecamylamine, a ganglionic blocker and secondary amine, inhibits acetylcholine at the autonomic ganglia. This causes blood vessel dilation and an increase in peripheral blood flow resulting in a decrease in blood pressure. Additionally, it blocks central nicotinic cholinergic receptors. Mecamylamine use has diminished due to its ganglionic side effects at antihypertensive doses.<sup>1,3,4</sup>

The miscellaneous hypotensive agents that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. This class was last reviewed in February 2020.

**Table 1. Miscellaneous Hypotensive Agents Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Mecamylamine	tablet	Vecamyl <sup>®</sup>	none

PDL=Preferred Drug List

**II. Evidence-Based Medicine and Current Treatment Guidelines**

The miscellaneous hypotensive agents are not included in the treatment guidelines and there are no specific recommendations for this drug.

**III. Indications**

Food and Drug Administration (FDA)-approved indications for the miscellaneous hypotensive agents are listed in Table 2. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 2. FDA-Approved Indications for the Miscellaneous Hypotensive Agents<sup>3</sup>**

Indication	Mecamylamine
Management of moderately severe to severe essential hypertension and in uncomplicated cases of malignant hypertension.	✓

**IV. Pharmacokinetics**

The pharmacokinetic parameters for miscellaneous hypotensive agents are summarized in Table 3.

**Table 3. Pharmacokinetic Parameters of the Miscellaneous Hypotensive Agents<sup>4</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Elimination (%)	Half-life (hours)	Active Metabolites
Mecamylamine	Not reported*	Not reported	Renal (100)	24	Not reported

\*It is noted that mecamylamine is readily absorbed from the gastrointestinal tract

## V. Drug Interactions

There are no reported drug interactions of major or moderate significance with the miscellaneous hypotensive agent, mecamylamine, although the package insert states that patients receiving antibiotics and sulfonamides generally should not be treated with ganglion blockers.<sup>3,5</sup> Additionally, the action of mecamylamine may be amplified by anesthesia, other antihypertensive agents, and alcohol. The mechanism of these interactions and specific drug agents are not specified.<sup>3</sup>

## VI. Adverse Drug Events

The most common adverse reactions reported with the miscellaneous hypotensive agents are noted in Table 4.

**Table 4. Adverse Events (%) Reported with the Miscellaneous Hypotensive Agents<sup>3</sup>**

Adverse Events	Mecamylamine
<b>Cardiovascular</b>	
Orthostatic dizziness	✓
Postural hypotension	✓
Syncope	✓
<b>Central Nervous System</b>	
Choreiform movements	✓
Convulsions	✓
Mental aberrations	✓
Paresthesias	✓
Tremor	✓
<b>Gastrointestinal</b>	
Anorexia	✓
Constipation	✓
Dry mouth	✓
Glossitis	✓
Ileus	✓
Nausea	✓
Vomiting	✓
<b>Respiratory</b>	
Fibrosis	✓
Interstitial pulmonary edema	✓
<b>Urogenital</b>	
Decreased libido	✓
Impotence	✓
Urinary retention	✓
<b>Other</b>	
Blurred vision	✓
Dilated pupils	✓
Fatigue	✓
Sedation	✓
Weakness	✓

## VII. Dosing and Administration

The usual dosing regimens for the miscellaneous hypotensive agents are summarized in Table 5.

**Table 5. Usual Dosing for the Miscellaneous Hypotensive Agents<sup>3</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
-----------------	------------------	----------------------	--------------



Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Mecamylamine	<p><u>Management of moderately severe to severe essential hypertension and in uncomplicated cases of malignant hypertension:</u>                      Tablet: initial, 2.5 mg twice daily (titrate in increments of 2.5 mg with at least two day intervals); average, 25 mg daily, in three divided doses (some patients may respond to as little as 2.5 mg daily; however, some patients may require two to four doses or greater in severe cases when consistent control is difficult to achieve); partial tolerance may develop requiring daily dosage increases</p>	Safety and effectiveness have not been established.	Tablet: 2.5 mg

### VIII. Effectiveness

A thorough literature search from 1966 to the present failed to retrieve any clinical studies evaluating the safety and effectiveness of mecamylamine for the treatment of hypertension. The initial clinical trials were conducted in the 1950s. These trials established the drug’s efficacy and side effect profile in patients with severe hypertension.<sup>1</sup>

#### Additional Evidence

##### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

##### Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

##### Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

### IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of ‘\$’ signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Miscellaneous Hypotensive Agents**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Mecamylamine	tablet	Vecamyl <sup>®</sup>	\$\$\$\$\$	N/A

\*Generic is available in at least one dosage form or strength.  
 N/A=not available

**X. Conclusions**

Although the clinical literature reports that mecamylamine is effective for the management of moderate-to-severe hypertension, its clinical utility is minimal due to its adverse events profile and the availability of newer and more effective agents. Current hypertension treatment guidelines do not mention mecamylamine as a first-line or alternative agent for the treatment of hypertension. Therefore, all brand miscellaneous hypotensive agents within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

**XI. Recommendations**

No brand miscellaneous hypotensive agent is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine cost effective products and possibly designate one or more preferred brands.

## **XII. References**

1. Young JM, Shytle RD, Sanberg PR, George TP. Mecamylamine: new therapeutic uses and toxicity/risk profile. *Clin Ther.* 2001 Apr;23(4):532-65.
2. Manchester Announces FDA Approval of Vecamyl™ [press release on the Internet]. Fort Collins (CO): Manchester Pharmaceuticals; 2013 May 1 [cited 2015 Mar 25]. Available from: <http://www.evaluategroup.com/Universal/View.aspx?type=Story&id=487170&sectionID=&isEPVantage=no>
3. Vecamyl® [Package Insert]. New York (NY): Vyera Pharmaceuticals, LLC; 2018 Jul.
4. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2021 [cited 2021 Jan]. Available from: <http://www.thomsonhc.com/>.
5. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Alpha-Adrenergic Blocking Agents  
AHFS Class 242000  
May 4, 2022**

**I. Overview**

The alpha-adrenergic blocking agents are approved for the treatment of benign prostatic hyperplasia (BPH) and hypertension.<sup>1-6</sup> However, the use of these agents for the treatment of hypertension is somewhat limited due to adverse events. They can cause postural hypotension, reducing the standing systolic blood pressure by more than 10 mm Hg. Syncope with sudden loss of consciousness can also occur, especially with the first few doses, rapid dose increases, or the addition of another antihypertensive agent to the treatment regimen. Unlike diuretics and  $\beta$ -adrenergic blocking agents,  $\alpha$ -adrenergic blocking agents do not adversely affect lipids. They have been shown to reduce total cholesterol by 3 to 5% and triglycerides by 3 to 4%, as well as increase high-density lipoprotein cholesterol.<sup>7</sup> The  $\alpha$ -adrenergic blocking agents are more commonly used to relieve symptoms of BPH, which is characterized by an enlargement of the prostate gland. BPH is associated with lower urinary tract symptoms, such as frequent daytime urination, nocturia, a sensation of incomplete bladder emptying, and a hesitant, weak, or intermittent urinary stream.<sup>8,9</sup>

The  $\alpha$ -adrenergic blocking agents competitively inhibit postsynaptic  $\alpha_1$ -adrenergic receptors, which are classified into three subtypes:  $\alpha_{1A}$ ,  $\alpha_{1B}$ , and  $\alpha_{1D}$ .<sup>10-13</sup> These receptors are located in the smooth muscle cell membrane of the peripheral blood vessels, as well as in various nonvascular smooth muscle and non-muscular tissues.<sup>11-15</sup> The  $\alpha$ -adrenergic blocking agents lower blood pressure by acting peripherally to dilate the blood vessels. They also cause rapid relaxation of smooth muscle in the bladder neck, prostate capsule, and prostatic urethra.<sup>14,16</sup>

The  $\alpha$ -adrenergic blocking agents that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the products are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Alpha-Adrenergic Blocking Agents Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Doxazosin	extended-release tablet, tablet	Cardura <sup>®*</sup> , Cardura XL <sup>®</sup>	doxazosin
Prazosin	capsule	Minipress <sup>®*</sup>	prazosin
Terazosin	capsule	N/A	terazosin

\*Generic is available in at least one dosage form or strength.

PDL=Preferred Drug List

N/A=Not available

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the  $\alpha$ -adrenergic blocking agents are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Alpha-Adrenergic Blocking Agents**

Clinical Guideline	Recommendations
Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b> <sup>17</sup>	<ul style="list-style-type: none"> <li>Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to</li> </ul>

Clinical Guideline	Recommendations
	<p>lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic blood pressure <math>&lt; 140</math> mm Hg.</p> <ul style="list-style-type: none"> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 140</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>18</sup></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio <math>&lt; 0.5</math> is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week. Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and</li> </ul>

Clinical Guideline	Recommendations
	<p>mindfulness or meditation introduced into the daily routine.</p> <ul style="list-style-type: none"> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><u>Pharmacological Treatment</u></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment               <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment               <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy               <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p><b>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children</b></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence</li> </ul>

Clinical Guideline	Recommendations
(2020) <sup>19</sup>	<p>of macrovascular target organ damage or other independent cardiovascular risk factors.</p> <ul style="list-style-type: none"> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both</li> </ul>

Clinical Guideline	Recommendations
	<p>agents may be used in patients with certain comorbid conditions or in combination therapy.</p> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt;40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (<math>&lt;40\%</math>) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium <math>&lt;5.2</math> mmol/L, an eGFR</li> </ul>



Clinical Guideline	Recommendations
	<p>≤30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</p>
	<p><u>Treatment of hypertension in association with stroke</u></p>
	<ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul>
	<p><u>Treatment of hypertension in association with LVH</u></p>
	<ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul>
	<p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p>
	<ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul>
	<p><u>Treatment of hypertension in association with renovascular disease</u></p>
	<ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul>
	<p><u>Treatment of hypertension in association with diabetes mellitus</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption,</li> </ul>

Clinical Guideline	Recommendations
	<p>prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>20</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential</li> </ul>

Clinical Guideline	Recommendations
	<p>antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</p> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HF<sub>r</sub>EF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HF<sub>p</sub>EF, BP treatment threshold and target values should be the same as for HF<sub>r</sub>EF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>21</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if</li> </ul>

Clinical Guideline	Recommendations
	<p>there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</p> <ul style="list-style-type: none"> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension</li> </ul>

Clinical Guideline	Recommendations
	taking the optimal tolerated doses of four drugs, seek specialist advice.
<p>American College of Cardiology/American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>22</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>• Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>• Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>• Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>• For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>• Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be <math>&lt; 130</math> mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP <math>&lt; 130/80</math> mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal</li> </ul>

Clinical Guideline	Recommendations
	<p>&lt;130/80 mmHg may be reasonable.</p> <ul style="list-style-type: none"> <li>For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> <li>For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>In patients with planned elective major surgery and SBP <math>\geq</math>180 mmHg or DBP <math>\geq</math>110 mmHg, deferring surgery may be considered.</li> <li>For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>23</sup></p>	<ul style="list-style-type: none"> <li>To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>

Clinical Guideline	Recommendations
<p>Kidney Disease Improving Clinical Outcomes Group: <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>24</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>25</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (≥140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure ≥180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>All hypertensive patients with diabetes should monitor their blood pressure at</li> </ul>

Clinical Guideline	Recommendations
	<p>home.</p> <ul style="list-style-type: none"> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of <math>&lt;130/80</math> mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;15\%</math>), treat to blood pressure target of <math>&lt;140/90</math>.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;120/80</math>, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic</li> </ul>

Clinical Guideline	Recommendations
	<p>kidney disease (CKD).</p> <ul style="list-style-type: none"> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium–glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium–glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin–to–creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin–to–creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin–to–creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Urological Association: <b>Management of Benign Prostatic Hyperplasia/ Lower Urinary Tract Symptoms (2021)</b><sup>8</sup></p>	<p><u>Medical therapy</u></p> <ul style="list-style-type: none"> <li>• Offer one of the following <math>\alpha</math>-blockers as a treatment option for patients with bothersome, moderate to severe lower urinary tract symptoms/benign prostatic hyperplasia (LUTS/BPH): alfuzosin, doxazosin, silodosin, tamsulosin, or terazosin.</li> <li>• When prescribing an <math>\alpha</math>-blocker for the treatment of LUTS/BPH, the choice of <math>\alpha</math>-blocker should be based on patient age and comorbidities, and different adverse event profiles (e.g., ejaculatory dysfunction, changes in blood pressure).</li> <li>• When initiating <math>\alpha</math>-blocker therapy, patients with planned cataract surgery should be informed of the associated risks and be advised to discuss these risks with their ophthalmologists.</li> <li>• For the purpose of symptom improvement, 5<math>\alpha</math>-reductase inhibitor (5-ARI) monotherapy should be used as a treatment option in patients with LUTS/BPH with prostatic enlargement as judged by a prostate volume of <math>&gt; 30</math>cc on imaging, a prostate specific antigen (PSA) <math>&gt; 1.5</math>ng/dL, or palpable prostate enlargement on digital rectal exam (DRE).</li> <li>• 5-ARIs alone or in combination with <math>\alpha</math>-blockers are recommended as a</li> </ul>

Clinical Guideline	Recommendations
	<p>treatment option to prevent progression of LUTS/BPH and/or reduce the risks of urinary retention and need for future prostate-related surgery.</p> <ul style="list-style-type: none"> <li>• Before starting a 5-ARI, clinicians should inform patients of the risks of sexual side effects, certain uncommon physical side effects, and the low risk of prostate cancer.</li> <li>• Clinicians may consider 5-ARIs as a treatment option to reduce intraoperative bleeding and peri- or postoperative need for blood transfusion after transurethral resection of the prostate (TURP) or other surgical intervention for BPH.</li> <li>• For patients with LUTS/BPH irrespective of comorbid erectile dysfunction (ED), 5mg daily tadalafil should be discussed as a treatment option.</li> <li>• 5-ARI in combination with an <math>\alpha</math>-blocker should be offered as a treatment option only to patients with LUTS associated with demonstrable prostatic enlargement as judged by a prostate volume of &gt; 30cc on imaging, a PSA &gt;1.5ng/dL, or palpable prostate enlargement on DRE.</li> <li>• Anticholinergic agents, alone or in combination with an <math>\alpha</math>-blocker, may be offered as a treatment option to patients with moderate to severe predominant storage LUTS.</li> <li>• <math>\beta</math>-3-agonists in combination with an <math>\alpha</math>-blocker may be offered as a treatment option to patients with moderate to severe predominate storage LUTS.</li> <li>• Clinicians should not offer the combination of low-dose daily 5mg tadalafil with <math>\alpha</math>-blockers for the treatment of LUTS/BPH as it offers no advantages in symptom improvement over either agent alone.</li> <li>• Physicians should prescribe an oral <math>\alpha</math>-blocker prior to a voiding trial to treat patients with acute urinary retention (AUR) related to BPH.</li> <li>• Patients newly treated for AUR with <math>\alpha</math>-blockers should complete at least three days of medical therapy prior to attempting trial without a catheter (TWOC). Clinicians should inform patients who pass a successful TWOC for AUR from BPH that they remain at increased risk for recurrent urinary retention.</li> </ul>
<p>European Association of Urology: <b>Non-neurogenic Male LUTS (2021)</b><sup>9</sup></p>	<p><u>Pharmacological treatment</u></p> <ul style="list-style-type: none"> <li>• Offer <math>\alpha</math>1-blockers to men with moderate-to-severe lower urinary tract symptoms (LUTS).</li> <li>• <math>\alpha</math>1-blockers are effective in reducing urinary symptoms (IPSS) and increasing the peak urinary flow rate (Qmax) compared with placebo.</li> <li>• Alfuzosin, terazosin and doxazosin showed a statistically significant increased risk of developing vascular-related events compared with placebo.</li> <li>• Alfuzosin, doxazosin, tamsulosin or terazosin exposure has been associated with an increased risk of intra-operative floppy iris syndrome (IFIS).</li> <li>• Ejaculatory dysfunction is significantly more common with <math>\alpha</math>1-blockers than with placebo, particularly with more selective <math>\alpha</math>1-blockers such as tamsulosin and silodosin.</li> <li>• Use 5<math>\alpha</math>-reductase inhibitors in men who have moderate-to-severe LUTS and an increased risk of disease progression (e.g., prostate volume &gt; 40 mL).</li> <li>• Counsel patients about the slow onset of action of 5<math>\alpha</math>-reductase inhibitors.</li> <li>• Use muscarinic receptor antagonists in men with moderate-to-severe LUTS who mainly have bladder storage symptoms.</li> <li>• Do not use antimuscarinic overactive bladder medications in men with a post-void residual volume &gt; 150 mL.</li> <li>• Use <math>\beta</math>-3 agonists in men with moderate-to-severe LUTS who mainly have bladder storage symptoms.</li> <li>• Use phosphodiesterase type 5 inhibitors in men with moderate-to-severe LUTS with or without erectile dysfunction.</li> <li>• Offer combination treatment with an <math>\alpha</math>1-blocker and a 5<math>\alpha</math>-reductase inhibitor to men with moderate-to-severe LUTS and an increased risk of disease progression (e.g., prostate volume &gt; 40 mL).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>Use combination treatment of a <math>\alpha 1</math>-blocker with a muscarinic receptor antagonist in patients with moderate-to-severe LUTS if relief of storage symptoms has been insufficient with monotherapy with either drug. Do not prescribe combination treatment in men with a post-void residual volume &gt; 150 mL.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the  $\alpha$ -adrenergic blocking agents are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Alpha-Adrenergic Blocking Agents<sup>3-6</sup>**

Indication	Doxazosin	Prazosin	Terazosin
<b>Benign Prostatic Hyperplasia</b>			
Treatment of signs and symptoms of benign prostatic hypertension	✓		
Treatment of symptomatic benign prostatic hyperplasia			✓
<b>Hypertension</b>			
Treatment of hypertension	✓ * (immediate-release)	✓ *	✓ *

\*Alone or in combination with other antihypertensive agents.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the  $\alpha$ -adrenergic blocking agents are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Alpha-Adrenergic Blocking Agents<sup>2</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
Doxazosin	IR: 65 ER: 54 to 59	98	Liver, extensive (% not reported)	Renal (9) Feces (63)	IR: 22 ER: 15 to 19
Prazosin	56 to 63	92 to 97	Liver, extensive (% not reported)	Renal (<1) Feces, extensive (% not reported)	2 to 3
Terazosin	90	90 to 94	Liver, extensive (% not reported)	Renal (40) Feces (55 to 60)	9 to 12

ER=extended-release, IR=immediate-release

### V. Drug Interactions

Major drug interactions with the  $\alpha$ -adrenergic blocking agents are listed in Table 5.

**Table 5. Major Drug Interactions with the Alpha-Adrenergic Blocking Agents<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
Alpha-adrenergic blocking agents (doxazosin, prazosin, terazosin)	Asenapine	Concurrent use of asenapine and alpha-adrenergic antagonists may result in additive hypotensive effect.



Generic Name(s)	Interaction	Mechanism
Alpha-adrenergic blocking agents (doxazosin, prazosin, terazosin)	Phosphodiesterase type 5 Inhibitors	Hypotension may occur when alpha-blockers and phosphodiesterase type 5 inhibitors are co-administered. Alpha-blockers and phosphodiesterase type 5 inhibitors may exert additive pharmacologic activity.
Alpha-adrenergic blocking agents (doxazosin)	Boceprevir	Concurrent use of boceprevir and doxazosin may result in increased doxazosin exposure.

## VI. Adverse Drug Events

The most common adverse drug events reported with the  $\alpha$ -adrenergic blocking agents are listed in Table 6. These agents can cause marked hypotension and syncope with sudden loss of consciousness with the first few doses. This “first-dose” effect can be minimized by administration of the first dose at bedtime. Hypotension and syncope can also occur with dose increases, addition of other antihypertensives, and therapy interruptions. The elderly are more at risk for this adverse reaction.

**Table 6. Adverse Drug Events (%) Reported with the Alpha-Adrenergic Blocking Agents<sup>1-6</sup>**

Adverse Events	Doxazosin	Prazosin	Terazosin
<b>Cardiovascular</b>			
Angina	<1	✓	-
Arrhythmia	1	-	<1
Atrial fibrillation	-	-	<1
Bradycardia	<1	✓	-
Chest pain	1 to 2	-	<1
Edema	3 to 4	1 to 4	-
Flushing	1	✓	-
Hypotension	1 to 2	✓	-
Myocardial infarction	<1	-	-
Orthostatic hypotension	<2	1 to 4	1 to 4
Palpitations	1 to 2	5	≤4
Peripheral edema	-	-	1 to 6
Peripheral ischemia	<1	-	-
Syncope	1 to 2	1 to 4	≤1
Tachycardia	<1	<1	≤2
Vasodilation	-	-	<1
<b>Central Nervous System</b>			
Abnormal thinking	<1	-	-
Agitation	<1	-	-
Amnesia	<1	-	-
Anxiety	1	-	<1
Ataxia	1	-	-
Cerebrovascular accident	<1	-	-
Confusion	<1	-	-
Decreased energy	-	7	-
Depersonalization	<1	-	-
Depression	1	1 to 4	-
Dizziness	5 to 19	10	9 to 19
Drowsiness	-	8	-
Emotional lability	<1	-	-
Fatigue	8 to 12	-	-
Fever	<1	-	<1
Hallucinations	-	<1	-
Headache	5 to 14	8	1 to 16

Adverse Events	Doxazosin	Prazosin	Terazosin
Hypertonia	1	-	-
Insomnia	1	✓	<1
Kinetic disorders	1	-	-
Migraine	<1	-	-
Nervousness	2	1 to 4	-
Paranoia	<1	-	-
Paresis	<1	-	-
Paresthesia	≤1	<1	≤3
Somnolence	1 to 5	-	4 to 5
Stroke	<1	-	-
Vertigo	2 to 4	1 to 4	1
<b>Dermatological</b>			
Alopecia	-	<1	-
Lichen planus	-	<1	-
Pallor	<1	-	-
Rash	1	1 to 4	<1
Pruritus	1	<1	<1
Urticaria	<1	<1	-
<b>Endocrine and Metabolic</b>			
Breast pain	<1	-	-
Gout	<1	-	<1
Gynecomastia	<1	✓	-
Pancreatitis	-	<1	-
<b>Gastrointestinal</b>			
Abdominal pain	2	<1	<1
Anorexia	<1	-	-
Appetite decreased	<1	-	-
Cholestasis	<1	-	-
Constipation	1	1 to 4	<1
Diarrhea	2	1 to 4	<1
Dyspepsia	1 to 2	-	<1
Fecal incontinence	<1	-	-
Flatulence	1	-	<1
Gastroenteritis	<1	-	-
Nausea	1 to 3	5	2 to 4
Vomiting	<1	1 to 4	<1
Xerostomia	2	1 to 4	<1
<b>Genitourinary</b>			
Hematuria	<1	-	-
Impotence	1	<1	≤2
Libido decreased	-	-	<1
Micturition abnormality	<1	-	-
Nocturia	<1	-	-
Polyuria	2	-	<1
Priapism	<1	<1	<1
Renal calculus	<1	-	-
Sexual dysfunction	2	-	-
Urinary frequency	-	1 to 4	-
Urinary incontinence	1	<1	<1
Urinary tract infection	1	-	<1
<b>Hematologic</b>			
Leukopenia	<1	-	-
Neutropenia	<1	-	-
Purpura	<1	-	-

Adverse Events	Doxazosin	Prazosin	Terazosin
Thrombocytopenia	<1	-	<1
<b>Hepatic</b>			
Jaundice	<1	-	-
Liver function tests increased	<1	<1	-
<b>Laboratory Test Abnormalities</b>			
Hypokalemia	<1	-	-
<b>Musculoskeletal</b>			
Arthralgia	1	✓	<1
Arthritis	1	-	<1
Back pain	2 to 3	-	≤2
Extremity pain	-	-	<1
Joint disorder	-	-	<1
Muscle cramps	1	-	-
Muscle weakness	1	-	7 to 11
Myalgia	1	-	<1
Neck pain	-	-	<1
Pain	2	✓	-
Shoulder pain	-	-	<1
Weakness	<1	7	-
<b>Respiratory</b>			
Bronchitis	-	-	<1
Bronchospasm	<1	-	-
Cough	-	-	<1
Dyspnea	1 to 3	1 to 4	2 to 3
Epistaxis	1	1 to 4	<1
Hepatitis	<1	-	-
Nasal congestion	-	1 to 4	2 to 6
Pharyngitis	-	-	<1
Respiratory disorder	1	-	-
Respiratory tract infection	5	-	-
Rhinitis	3	-	<1
Sinusitis	-	-	≤3
<b>Special Senses</b>			
Abnormal vision	1 to 2	-	<1
Blurred vision	-	1 to 4	≤2
Cataracts	-	<1	-
Conjunctivitis	1	-	<1
Hypoesthesia	<1	-	-
Intraoperative floppy iris syndrome	<1	<1	<1
Pigmentary mottling and serous retinopathy	-	<1	-
Sclera reddened	-	1 to 4	-
Tinnitus	1	<1	<1
Parosmia	<1	-	-
<b>Other</b>			
Allergic reaction	<1	✓	<1
Anaphylaxis	-	-	<1
Diaphoresis	1	✓	<1
Facial edema	1	-	<1
Infection	<1	-	-
Influenza-like symptoms	1	-	≤2
Lymphadenopathy	<1	-	-
Rigors	<1	-	-
Vasculitis	-	✓	-

✓ Percent not specified

- Event not reported

## VII. Dosing and Administration

The usual dosing regimens for the  $\alpha$ -adrenergic blocking agents are listed in Table 7. Treatment should be initiated at bedtime and at the lowest dose to minimize the likelihood of the “first-dose” effect. Dosages should be titrated up slowly to achieve the desired response. If therapy is interrupted for more than a few days, the initial dosing regimen and titration schedule should be reinstated. Other antihypertensive agents should be added cautiously to reduce the risk of developing significant hypotension.<sup>1-6</sup>

**Table 7. Usual Dosing Regimens for the Alpha-Adrenergic Blocking Agents<sup>1-6</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Doxazosin	<p><u>Benign prostatic hyperplasia:</u> Extended-release: initial, 4 mg once daily; maintenance, 4 to 8 mg daily; maximum, 8 mg/day</p> <p>Tablet: initial, 1 mg once daily; maintenance, 1 to 8 mg once daily; maximum:, 8 mg/day</p> <p><u>Hypertension:</u> Tablet: initial, 1 mg once daily; maintenance, 1 to 16 mg once daily; maximum, 16 mg/day</p>	Safety and efficacy in children have not been established.	<p>Extended-release tablet: 4 mg 8 mg</p> <p>Tablet: 1 mg 2 mg 4 mg 8 mg</p>
Prazosin	<p><u>Hypertension:</u> Capsule: initial, 1 mg two to three times a day; maintenance, 6 to 15 mg/day in divided doses; maximum, 40 mg/day</p>	Safety and efficacy in children have not been established.	<p>Capsule: 1 mg 2 mg 5 mg</p>
Terazosin	<p><u>Benign prostatic hyperplasia:</u> Capsule: initial, 1 mg at bedtime; maintenance, 1 to 10 mg/day; maximum, 20 mg/day</p> <p><u>Hypertension:</u> Capsule: initial, 1 mg at bedtime; maintenance, 1 to 20 mg once daily; maximum, 20 mg/day</p>	Safety and efficacy in children have not been established.	<p>Capsule: 1 mg 2 mg 5 mg 10 mg</p>

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the  $\alpha$ -adrenergic blocking agents are summarized in Table 8.

**Table 8. Comparative Clinical Trials with the Alpha-Adrenergic Blocking Agents**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Benign Prostatic Hyperplasia</b>				
<p>Lee et al.<sup>26</sup> (2011)</p> <p><math>\alpha</math>-adrenergic blocking agent</p> <p>vs</p> <p>no <math>\alpha</math>-adrenergic blocking agent</p> <p>All patients were receiving finasteride.</p> <p>Patients were divided into 2 groups based on treatment pattern (<math>\alpha</math>-adrenergic blocking agent monotherapy vs <math>\alpha</math>-adrenergic blocking agent combined with finasteride) and further divided into 4 subgroups based on severity of storage symptoms (IPSS</p>	<p>MC, RETRO</p> <p>Patients <math>\geq 50</math> years of age with LUTS consistent with moderate to severe BPH</p>	<p>N=1315</p> <p>4 years</p>	<p>Primary: Prostate volume, PSA, IPSS, <math>Q_{max}</math></p> <p>Secondary: Not reported</p>	<p>Primary: All groups showed significant improvements in IPSS total scores, IPSS voiding subscores and QOL at one year (P values not reported). Total IPSS from baseline to year four decreased by -11.5 in group IV compared to -0.18 in group I (P&lt;0.001), -6.1 in group II (P=0.97) and -2.6 in group III (P=0.031). However, IPSS storage subscores only improved in patients with high (<math>\geq 6</math>) storage subscores at baseline (P value not reported). After one year, prostate volume and PSA were reduced by 21.3 and 47.0%, respectively, in the combination groups compared to an increase of 9 and 18%, respectively, in the monotherapy groups (P&lt;0.001 for both).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>storage domain score <math>\geq 6</math> vs <math>&lt; 6</math>).</p> <p>Group I was classified as monotherapy and storage scores <math>&lt; 6</math>, group II as monotherapy and storage scores <math>\geq 6</math>, group III as combination therapy and storage scores <math>&lt; 6</math> and group IV as combination therapy and storage scores <math>\geq 6</math>.</p>				
<p>Demir et al.<sup>27</sup> (2009)</p> <p>Doxazosin 4 mg QD</p> <p>Patients were grouped into 2 groups according to self-reported erectile status: patients who reported the presence of erectile dysfunction (group I) and patients who reported the absence of erectile dysfunction (group</p>	<p>RETRO</p> <p>Males &gt;40 years of age who had been in a steady sexual relationship for the past 6 months and were admitted to urology clinics with complaints of BPH</p>	<p>N=64</p> <p>6 weeks</p>	<p>Primary: Not reported</p> <p>Secondary: Not reported</p>	<p>Primary Endpoints: Not reported</p> <p>Secondary Endpoints: Not reported</p> <p>Mean reductions in total IPSS and quality of life compared to baseline were <math>-7.7 \pm 6.1</math> and <math>1.5 \pm 1.5</math> (<math>P=0.006</math> and <math>P=0.024</math>, respectively). Treatment with doxazosin also resulted in significant improvements in <math>Q_{max}</math> over baseline (<math>3.2 \pm 4.6</math> mL/s; <math>P=0.002</math>). Both groups exhibited significant improvements in IPSS and quality of life scores over baseline (<math>P&lt;0.001</math> for both). Improvements in LUTS appeared to be numerically greater in group II; however, quality of life was the only parameter for which a significant improvement was seen compared to group I (<math>-1.0 \pm 1.8</math> vs <math>-1.9 \pm 1.1</math>, for groups I and II respectively; <math>P=0.018</math>).</p> <p>Mean International Index of Erectile Function erectile function domain scores increased in group I and slightly decreased in group II when compared to baseline. Mean changed of other International Index of</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
II)				<p>Erectile Function domains were not significant in either group. When stratified according to erectile dysfunction severity, the mean changes in International Index of Erectile Function erectile function domain scores over baseline were: <math>4.3 \pm 6.0</math>, <math>0.3 \pm 5.3</math>, <math>-1.2 \pm 1.6</math> in those participants with severe, moderate, and mild erectile dysfunction, respectively.</p> <p>No serious adverse events were observed during the treatment course in either group.</p>
<p>Sun et al.<sup>28</sup> (abstract) (2010)</p> <p>Doxazosin SR 4 mg QD</p> <p>At week 4, subjects who achieved an increase in <math>Q_{max} \geq 3</math> mL/s and a <math>\geq 30\%</math> reduction in the total IPSS continued on doxazosin SR 4 mg for the remaining 4 weeks; all other subjects were titrated up to 8 mg QD.</p>	<p>OL, PM</p> <p>Taiwanese males with BPH</p>	<p>N=80</p> <p>8 weeks</p>	<p>Primary: Change from baseline <math>Q_{max}</math> and IPSS</p> <p>Secondary: Safety</p>	<p>Primary Endpoints: Baseline <math>Q_{max}</math> and IPSS were <math>10.7 \pm 3.4</math> mL/s and <math>20.6 \pm 5.4</math>, respectively. At week eight, a significant increase from baseline in <math>Q_{max}</math> of <math>3.3 \pm 4.6</math> mL/s (95% CI, 2.2 to 4.4, <math>P &lt; 0.001</math>) and a significant decrease in total IPSS of <math>-8.9 \pm 7.0</math> (95% CI, -10.5 to -7.3; <math>P &lt; 0.001</math>) was observed.</p> <p>Secondary Endpoints: The most common treatment-related adverse event was dizziness.</p>
<p>Kirby et al.<sup>29</sup> (2001)</p> <p>Doxazosin vs</p>	<p>2 DB, MC, PG, RCT</p> <p>Men 50 to 80 years of age with BPH</p>	<p>N=1,475</p> <p>17 weeks</p>	<p>Primary: Change from baseline in IPSS and <math>Q_{max}</math></p> <p>Secondary: Sexual function,</p>	<p>Primary: A 45% significant decrease from baseline in IPSS was attained with both formulations of doxazosin compared to a 34% decrease with placebo after 13 weeks (<math>P &lt; 0.001</math> vs placebo). Doxazosin SR was as effective as doxazosin in improving IPSS with a least squares mean difference of 0.07 (SEM, 0.28; 95% CI, -0.47 to 0.61; <math>P = 0.799</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
doxazosin SR  vs  placebo  Comparison with placebo was evaluated in 1 of the 2 trials.			tolerability	Effect on Q <sub>max</sub> was also comparable between the two doxazosin formulations; a least square mean difference of 0.19 (SEM, 0.23; 95% CI, -0.27 to 0.64; P=0.426) was reported. The improvements were significantly greater compared to placebo (P<0.001 for both).  Secondary: Only the non-PC trial evaluated sexual function. Both formulations of doxazosin demonstrated modest but significant improvements in sexual function from baseline as measured by the IIEF (P≤0.001 for doxazosin SR and P<0.05 for doxazosin).  Forty one percent of patients receiving doxazosin SR, 54% of patients receiving doxazosin and 39% of patients receiving placebo experienced adverse events (P<0.001 for differences among treatments). Headache, dizziness, respiratory tract infections and asthenia were the most frequently reported side effects of active treatment.
Keten et al. <sup>30</sup> (2015)  Doxazosin XL 4 mg (group 1)  vs  doxazosin XL 8 mg (group 2)  Patients with an inadequate response to 4 mg treatment were switched to 8 mg after one month (group 1b)	PRO  Patients aged >45 years, with a total PSA <4 ng/mL, IPSS of >7, and Q <sub>max</sub> ≤15 mL/s	N=162  4 months	Primary: IPSS, Q <sub>max</sub> , quality of life (QoL) score  Secondary: Not reported	Primary: From the time of presentation to the first month follow-up, the IPSS and QoL values had decreased more in group 2 compared with group 1 (P <sub>IPSS</sub> =0.028, P <sub>Q<sub>max</sub></sub> =0.206, P <sub>QoL</sub> =0.038, and P <sub>PVR</sub> =0.070).  The comparison of the patients in Group 1b who used 4 and 8 mg doxazosin XL for one month showed that during the use of 4 mg doxazosin XL, the change in the IPSS was 1.3 ± 1.3 units, and during the use of 8 mg doxazosin XL, the change was 3.6 ± 2.5 units (P<0.001). The change in the Q <sub>max</sub> values for 4 and 8 mg doxazosin XL was found to be 1.6 ± 1.8 and 3.2 ± 2.7 mL/s, respectively (P=0.019). For the 4 mg doxazosin XL, the QoL values increased by 0.4 ± 0.6 units, and during the use of 8 mg doxazosin XL, the QoL values decreased by 1.8 ± 0.7 units (P<0.001).  No difference was found concerning the adverse reactions between the groups (P>0.05).  Secondary: Not reported
Samli et al. <sup>31</sup>	RCT, XO	N=50	Primary:	Primary:



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(2004) Doxazosin 8 mg QD  vs terazosin 10 mg QD	Men with LUTS associated with BPH	3 months	Change from baseline in IPSS and Qmax  Secondary: Not reported	The proportion of patients who showed improvement in both IPSS and Qmax were 44 and 40% of patients receiving doxazosin and Terazosin, respectively. After three months, both treatments resulted in a significantly increased Qmax (P<0.001) and a significantly decreased IPSS (P<0.01).  The number of patients who did not show improvement and had to switch to the other treatment was 19. Of these patients, two showed improvement in both IPSS and Qmax, two showed improvement in IPSS only and 15 did not show any improvement.  Secondary: Not reported
Kaplan et al. <sup>32</sup> (1997) Doxazosin 4 to 8 mg QD  vs terazosin 5 to 10 mg QD	OL, PRO  Men >80 years of age with BPH	N=36  6 months	Primary: Change from baseline in Qmax and AUA SS  Secondary: Not reported	Primary: There was significant improvement in Qmax (P<0.008) and AUA SS (P<0.01) with both treatments.  There were small, nonsignificant decreases in blood pressure with both treatments.  Secondary: Not reported
Kaplan et al. <sup>33</sup> (1995) Doxazosin 4 mg QD in the morning  vs doxazosin 4 mg QD in the evening  vs terazosin 5 mg QD in the morning	RCT  Men without HTN and symptomatic prostatism	N=43  4 to 17 months	Primary: Changes from baseline in Boyarsky symptom score, Qmax and blood pressure; adverse events  Secondary: Not reported	Primary: There were significant improvements in Boyarsky symptom scores and Qmax with all four treatments (P<0.05), with no significant differences between the treatments (P values not reported).  Adverse events were significantly decreased with evening doses (P<0.05).  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs terazosin 5 mg QD in the evening				
Bozlu et al. <sup>34</sup> (2004)  Doxazosin 4 mg QD  vs  terazosin 5 mg QD  vs  alfuzosin 2.5 mg TID  vs  tamsulosin 0.4 mg QD	RETRO  Patients with LUTS suggestive of BPH with and without diabetes	N=281  6 months	Primary: Change from baseline in IPSS, bother score, Qmax and PVR  Secondary: Not reported	Primary: Doxazosin, terazosin and alfuzosin significantly improved IPSS, bother scores, Qmax and PVR compared to baseline (P<0.001). IPSS and bother scores were significantly improved in diabetic patients compared to nondiabetic patients (P<0.01).  There was no significant differences among the treatments in the improvement rates of any of the parameters (P>0.05).  Secondary: Not reported
Xue et al. <sup>35</sup> (2007)  Doxazosin SR 4 mg QD  vs  tamsulosin 0.2 mg QD	RCT  Chinese men with confirmed BPH	N=117  8 weeks	Primary: Efficacy, safety  Secondary: Not reported	Primary: Both treatments significantly improved the IPSS (total, irritative subscore, and obstructive subscore; P=0.001 for all) and Qmax (P=0.001). Other differences between groups were not statistically significant.  Secondary: Not reported
Rahardjo et al. <sup>36</sup> (2006)	MC, OL, RCT  Patients with LUTS	N=101  6 weeks	Primary: Changes from baseline in IPSS,	Primary: The total IPSS decreased significantly with both tamsulosin and doxazosin compared to baseline (P<0.001), with tamsulosin being associated with a

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Doxazosin 2 mg QD  vs  tamsulosin 0.2 mg QD	due to BPH		Qmax, average urinary flow rate and residual urine; safety  Secondary: Not reported	significant decrease compared to doxazosin (P=0.036).  Qmax, average urinary flow rate and residual urine significantly improved with tamsulosin only (P<0.001, P<0.001, and P<0.05, respectively).  There were no significant differences in SBP or DBP with tamsulosin; however, doxazosin resulted in significant differences in SBP (P<0.01) but not in DBP (P value not reported) at the end of the study.  Tamsulosin was well tolerated; only three patients (six percent) receiving tamsulosin reported an adverse event (dizziness), while 11 patients (22%) with doxazosin reported an adverse event (dizziness), one of whom withdrew from the trial.  Secondary: Not reported
Pompeo et al. <sup>37</sup> (2006)  Doxazosin SR 4 mg QID  vs  tamsulosin 0.4 mg QID	DB, DD, RCT  Brazilian patients with BPH	N=165  12 week	Primary: Absolute and percentage change from baseline in symptoms measured by IPSS  Secondary: Quality of life question from the IPSS and questions six and seven of the SFAQ	Primary: Doxazosin and tamsulosin improved IPSS with no significant differences between the two after 12 weeks. During weeks four to eight, tamsulosin demonstrated a slower improvement (P<0.001) in IPSS compared to doxazosin.  Secondary: The proportion of satisfied patients did not change over the course of the trial with doxazosin, while it did change significantly between weeks four and eight with tamsulosin (P=0.006); suggesting that a change for the better was observed earlier with doxazosin. After 12 weeks, the proportion of patients with little or no difficulty at ejaculation (question six of SFAQ) was significantly higher with doxazosin (P=0.019). Both treatments were well tolerated.
Cao et al. <sup>38</sup> (2016)  Doxazosin SR 4 mg QD  vs	OL, PRO, RCT  Men 50 to 80 years of age with newly diagnosed symptoms of BPH	N=192  12 weeks	Primary: IPSS  Secondary: Quality of life and Qmax	Primary: After six weeks of treatment, LUTS were improved in both groups, as seen by the change in IPSS from baseline. There was no significant difference between groups in terms of total IPSS (P=0.86). At 12 weeks, greater improvement of total IPSS was observed in the doxazosin group than in the tamsulosin group (P<0.001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
tamsulosin 0.2 mg QD  Both groups received tolterodine ER 4 mg QD				Secondary: After six weeks treatment there was no significant difference between the groups in Qmax (P=0.19). However, quality of life was significantly improved in the doxazosin group compared to the tamsulosin group (P=0.01). At 12 weeks Qmax showed greater improvement in the doxazosin group (14.1±1.6 mL/s) than in the tamsulosin group (13.5±2.1 mL/s; P=0.03). In addition, quality of life (2.5±0.67 vs 3.1±0.7; P<0.001) of the doxazosin group was more improved (lower score represents better quality of life).
Johnson et al. <sup>39</sup> (2007)  Doxazosin 2, 4, or 8 mg QD  vs  finasteride 5 mg QD  vs  doxazosin 2, 4, or 8 mg QD plus finasteride 5 mg QD  vs  placebo	PC, RCT  Men with LUTS suggestive of BPH	N=3,047  4 years	Primary: Efficacy (mean reduction in self-reported nightly nocturia at 1 and 4 years)  Secondary: Not reported	Primary: The number of men reporting one or more episodes of nocturia who finished ≥12 months of the trial came to a total of 2,583. Mean nocturia was similar with all treatments at baseline. Mean nocturia was reduced after one year by 0.35, 0.40, 0.54 and 0.58 with placebo, finasteride, doxazosin and combination therapy, respectively. Reductions with doxazosin and combination therapy were significantly greater compared to placebo (P<0.05).  After four years, nocturia was also significantly reduced in patients receiving doxazosin and combination therapy (P<0.05 vs placebo). In men >70 years of age (n=495) all treatments significantly reduced nocturia after one year (Finasteride, 0.29; Doxazosin, 0.46 and combination therapy, 0.42) compared to placebo (0.11; P<0.05 for all).  Secondary: Not reported
Crawford et al. <sup>40</sup> (2006)  Doxazosin 4 to 8 mg QD  vs	PC, RCT  Men with LUTS suggestive of BPH	N=737  4 years	Primary: Time to overall clinical progression of BPH (either a confirmed ≥4 point increase in AUA	Primary: The rate of overall clinical progression of BPH events with placebo was 4.5 per 100 person-years, for a cumulative incidence (among men who had at least four years of follow up data) of 17%.  The risk of BPH progression was significantly greater with placebo with a baseline TPV ≥31 mL compared to a baseline TPV <31 mL (P<0.0001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>finasteride 5 mg QD</p> <p>vs</p> <p>doxazosin 4 to 8 mg QD plus finasteride 5 mg QD</p> <p>vs</p> <p>placebo</p>			<p>SS, acute urinary retention, incontinence, renal insufficiency or recurrent urinary tract infection)</p> <p>Secondary: Not reported</p>	<p>The risk of BPH progression was significantly greater with placebo with a baseline PSA <math>\geq 1.6</math> ng/dL compared to a baseline PSA <math>&lt; 1.6</math> ng/dL (P=0.0009).</p> <p>The risk of BPH progression was significantly greater with placebo with a baseline Qmax <math>&lt; 10.6</math> mL/second compared to a baseline Qmax <math>\geq 10.6</math> mL/second (P=0.011)</p> <p>The risk of BPH progression was significantly greater with placebo with a baseline PVR <math>\geq 39</math> mL compared to a baseline PVR <math>&lt; 39</math> mL (P=0.0008).</p> <p>The risk of BPH progression was significantly greater with placebo with baseline age <math>\geq 62</math> years compared to those aged <math>&lt; 62</math> years (P=0.0002).</p> <p>Secondary: Not reported</p>
<p>Kaplan et al.<sup>41</sup> (2006)</p> <p>Doxazosin 4 to 8 mg QD</p> <p>vs</p> <p>finasteride 5 mg QD</p> <p>vs</p> <p>doxazosin 4 to 8 mg QD plus finasteride 5 mg QD</p> <p>vs</p>	<p>PC, RCT</p> <p>Men with LUTS suggestive of BPH</p>	<p>N=3,047</p> <p>4 years</p>	<p>Primary: Overall clinical progression of BPH (either a confirmed <math>\geq 4</math> point increase in AUA SS, acute urinary retention, incontinence, renal insufficiency or recurrent urinary tract infection)</p> <p>Secondary: Need for invasive therapy for BPH, change from baseline in AUA SS and Qmax</p>	<p>Primary: In patients with a small prostate (baseline TPV <math>&gt; 25</math> mL) combination therapy was no better than doxazosin for decreasing the risk of clinical progression of BPH and need for invasive therapy as well as improving AUA SS and Qmax. However, in patients with a moderate sized (25 to <math>&gt; 40</math> mL) or enlarged (<math>\geq 40</math> mL) gland, combination therapy led to a clinical benefit in these outcomes that was “superior” to that of doxazosin or finasteride (P<math>&lt; 0.05</math>).</p> <p>Secondary: In men with baseline TPV <math>&lt; 25</math> mL, there was no difference in the risk of invasive therapy for combination therapy relative to doxazosin or finasteride. However, in the baseline TPV subgroups of 25 to <math>&lt; 40</math> and <math>\geq 40</math> mL there was a significant and marked percent risk decrease in invasive therapy, of around 60 to 80%, for combination therapy compared to doxazosin (P<math>&lt; 0.05</math>).</p> <p>In men with baseline TPV <math>&lt; 25</math> mL, the improvement after four years in AUA SS for combination therapy relative to doxazosin was not different, whereas the improvement for combination therapy compared to finasteride</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo				was significantly different in favor of combination therapy ( $P<0.05$ ).  In the baseline TPV subgroups of 25 to $<40$ and $\geq 40$ mL, the improvement in AUA SS with combination therapy was significantly better than that for doxazosin and finasteride ( $P<0.05$ ).
Kaplan et al. <sup>42</sup> (2008)  Doxazosin 4 to 8 mg/day  vs  finasteride 5 mg/day  vs  doxazosin 4 to 8 mg/day and finasteride 5 mg/day  vs  placebo	DB, PC, RCT  Men $\geq 50$ years of age with an AUA SS of 8 to 30 and a Qmax of 4 to 15 ml/second with a voided volume of $\geq 125$ mL	N=3,047  Mean 4.5 years	Primary: TPV  Secondary: Not reported	Primary: Long-term treatment with finasteride alone or in combination with doxazosin led to a consistent reduction in TPV of approximately 25% compared to placebo in men with a relatively small prostate (baseline TPV less than 25 mL and 25 to 30 mL) as well as those with a moderate size (greater than 30 to less than 40 mL) or enlarged prostate (40 mL or greater).  Secondary: Not reported
Kirby et al. <sup>43</sup> (2003) PREDICT  Doxazosin 1 to 8 mg QD  vs  finasteride 5 mg QD	DB, MC, PC, PRO, RCT  Men 50 to 80 years of age with BPH and an enlarged prostate	N=1,095  52 weeks	Primary: Change from baseline in Qmax and IPSS  Secondary: Tolerability	Primary: Doxazosin ( $3.6\pm 0.3$ mL/second) and combination therapy ( $3.8\pm 0.3$ mL/second) were associated with a significantly greater improvement in Qmax after one year compared to finasteride ( $1.8\pm 0.3$ mL/second; $P\leq 0.0001$ ) or placebo ( $1.4\pm 0.3$ mL/second; $P\leq 0.0001$ ). There were no differences between doxazosin and combination therapy or finasteride and placebo (P values not reported).  Similar results were found with total IPSS. Again, doxazosin ( $3.6\pm 0.3$ mL/second) and combination therapy ( $3.8\pm 0.3$ mL/second) caused a significantly greater improvement in score over finasteride alone ( $1.8\pm 0.3$ mL/second; $P<0.01$ ) or placebo ( $1.4\pm 0.3$ mL/second; $P\leq 0.0001$ ). There

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  doxazosin 1 to 8 mg QD plus finasteride 5 mg QD  vs  placebo				were no differences between doxazosin and combination therapy or finasteride and placebo (P values not reported).  Secondary: Doxazosin use increased the risk of asthenia, dizziness and hypotension, while impotence was reported most frequently with combination therapy.
Fwu et al. <sup>44</sup> (2013) MTOPS  Doxazosin 4 or 8 mg QD  vs  finasteride 5 mg QD  vs  doxazosin 4 or 8 mg QD plus finasteride 5 mg QD  vs  placebo	DB, MC, RCT  Patients ≥50 years of age with an AUA-SS of 8 to 30, a Qmax of 4 to 15 mL/second, and a minimum voided volume of 125 mL	N=2,872  4 years	Primary: Change in quality of life (QoL) using Benign Prostatic Hyperplasia Impact Index (BII), IPSS-QoL, and annually by the Outcomes Study Short-Form 36 (MOS-SF-36)  Secondary: Not reported	Primary: Changes in the MOS-SF-36 scores from baseline to year four of follow-up by treatment group demonstrated a statistically significant reduction (worsening) for the subscales of physical functioning, role limitations due to physical problems, bodily pain, general health perception, and vitality in all treatment groups, except bodily pain in men assigned to finasteride. No significant change was observed in mental health in any treatment group. The subscale with the greatest reduction was role limitations due to physical problems. The decrease was greatest in the placebo group (−8.83, 95% CI, −12.09 to −5.58) and least in finasteride (−6.97, −10.19 to −3.74).  The differences in changes for MOS-SF-36 subscales and summary scores between drug groups and placebo group were not statistically significant. Similarly, neither significant differences nor important effect sizes of the subscales and summary scores were observed when drug groups were compared with each other at year four.  Compared with men assigned to placebo, men assigned to doxazosin and combination experienced a statistically significant improvement in the BII at year four. Men assigned to each of the drug groups also experienced a significant improvement in the IPSS-QoL compared with those assigned to placebo.  Secondary: Not reported
Djavan et al. <sup>45</sup> (1999)	MA	N=6,333 (PC trials)	Primary: Changes from	Primary: There was no difference in efficacy among the four treatments. Alfuzosin

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Doxazosin vs terazosin vs alfuzosin vs tamsulosin</p>	<p>Men with LUTS suggestive of benign prostatic obstruction</p>	<p>N=507 (comparative trials)</p>	<p>baseline in total symptom score and maximum urinary flow rate, tolerability</p> <p>Secondary: Not reported</p>	<p>(IR 2.5 mg TID), alfuzosin (SR 5 mg BID), terazosin (5 to 10 mg/day), doxazosin (4 to 8 mg/day) and tamsulosin (0.4 mg/day) all produced comparable improvements in LUTS and Qmax (P values not reported). The total symptom score improved by 30 to 40% and the Qmax by 16 to 25%.</p> <p>Alfuzosin and tamsulosin were better tolerated than terazosin and doxazosin. Alfuzosin and tamsulosin had similar withdrawal rates as placebo. With terazosin and doxazosin, an additional 4 to 10% of patients withdrew from due to intolerability (P value not reported).</p> <p>Tamsulosin had less effect on blood pressure than alfuzosin (P value not reported). Tamsulosin also caused less symptomatic orthostatic hypotension than terazosin (P value not reported).</p> <p>Secondary: Not reported</p>
<p>Nickel et al.<sup>46</sup> (2008)</p> <p>Doxazosin 4 to 8 mg/day vs terazosin 1 to 10 mg/day vs alfuzosin 10 mg/day vs tamsulosin 0.4 mg/day</p>	<p>MA</p> <p>Men with BPH</p>	<p>26 trials</p> <p>4 weeks to 4.5 years</p>	<p>Primary: Vascular-related adverse events with <math>\alpha</math>1-adrenergic blockers including dizziness, hypotension, or syncope</p> <p>Secondary: Efficacy based on change from baseline of Qmax and change from baseline of AUA SI or IPSS</p>	<p>Primary: Treatment with <math>\alpha</math>1-adrenergic blockers was associated with a significant increase in the development of a vascular-related adverse event compared to placebo (OR, 2.54; 95% CI, 2.00 to 3.23; P&lt;0.0001).</p> <p>There was a higher risk of developing the primary composite end-point compared to placebo for alfuzosin (P=0.005), terazosin (P&lt;0.0001), doxazosin (P&lt;0.0001), and doxazosin SR (P&lt;0.0001).</p> <p>Secondary: Alpha-1-adrenergic blockers improved Q<sub>max</sub> by 1.32 mL/min compared to placebo (95% CI, 1.07 to 1.57; P&lt;0.0001).</p> <p>The WMD in AUA SI/IPSS for all <math>\alpha</math>1-adrenergic blockers was -1.92 points compared to placebo (95% CI, -2.71 to -1.14); P&lt;0.0001).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo				
MacDonald et al. <sup>47</sup> (2005)  Doxazosin, tamsulosin, or finasteride  vs  alfuzosin  vs  alfuzosin plus finasteride or placebo	SR (11 trials)  Men with symptomatic BPH	N=3,901  4 to 26 weeks	Primary: Change from baseline in IPSS  Secondary: Change from baseline in Qmax and urinary symptom scores, adverse effects, incidence of treatment discontinuation	Primary: In the two trials comparing alfuzosin to other $\alpha$ -adrenergic blocking agents, doxazosin demonstrated the greatest improvement in IPSS (WMD, 1.70; 95% CI, 0.76 to 1.64; P=0.05). One trial evaluated alfuzosin vs finasteride or alfuzosin plus finasteride. Alfuzosin, both alone or in combination, significantly improved LUTS compared to finasteride. When compared to placebo, alfuzosin demonstrated a greater improvement in the IPSS with a WMD of -1.8 points (95% CI, -2.49 to -1.11).  Secondary: No difference was found among $\alpha$ -adrenergic blocking agents in Qmax, while alfuzosin and tamsulosin (0.4 mg) demonstrated similar improvements in Boyarsky symptom scores.  Alfuzosin, finasteride and combination treatment all had similar changes in Qmax; however, a subgroup analysis showed greater improvement in patients with obstruction with alfuzosin and combination therapy over finasteride.  Qmax was 2.6 mL/second (10 to 54%) with alfuzosin vs 1.1 mL/second with placebo (2 to 29%). Alfuzosin demonstrated benefit over placebo in the mean urinary symptom score with a WMD of -0.90 point (95% CI, -0.94 to -0.87).  The incidences of adverse events as well as withdrawal rates were comparable among $\alpha$ -adrenergic blocking agents. Vasodilatory effects were similar with alfuzosin, finasteride and combination therapy, whereas impotence occurred significantly more often with finasteride and in combination therapy. Discontinuation of treatment was higher with alfuzosin than finasteride and lower with alfuzosin compared to combination therapy. Dizziness was the most frequently reported side effect with alfuzosin compared to placebo. Postural hypotension, syncope and somnolence were reported in less than two percent of patients treated

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				with alfuzosin, but more often than with placebo. Withdrawal rates were similar between treatments.
<p>Tsujii et al.<sup>48</sup> (2000)</p> <p>Prazosin 0.5 to 1 mg BID</p> <p>vs</p> <p>terazosin 0.5 to 1 mg BID</p> <p>vs</p> <p>tamsulosin 0.1 to 0.2 QD</p>	<p>RCT, XO</p> <p>Patients with symptomatic BPH</p>	<p>N=121</p> <p>4 weeks</p>	<p>Primary: Changes from baseline in symptom score, Qmax, average urinary flow rate, PVR and blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: Terazosin was associated with a significant improvement in four out of nine symptoms compared to tamsulosin (P&lt;0.05).</p> <p>There were significant increases in Qmax with prazosin, and in average urinary flow rate with tamsulosin (P&lt;0.05 for both).</p> <p>There were no significant changes in PVR with any of the treatments.</p> <p>Significant reductions in blood pressure were observed in the hypertensive patients with prazosin, terazosin and tamsulosin (P&lt;0.05 for all). In the normotensive patients, no significant changes in blood pressure were observed with any of the treatments.</p> <p>Secondary: Not reported</p>
<p>Tsai et al.<sup>49</sup> (2007)</p> <p>Group A: Terazosin (generic) 1 to 4 mg QD for 6 weeks (Period 1), followed by terazosin (brand Hytrin®*) 1 to 4 mg QD for 6 weeks (Period 2)</p> <p>vs</p> <p>Group B: Terazosin (brand Hytrin®*) 1 to 4</p>	<p>OL, RCT</p> <p>Adult men in Taiwan newly diagnosed with symptomatic BPH who had not previously received treatment for BPH</p>	<p>N=53</p> <p>13 weeks</p>	<p>Primary: Change from baseline in IPSS, tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: After two and six weeks, no significant between-product differences were found in mean (SD) decreases from baseline in IPSS total score (generic, 2.46 [0.84] and 2.46 [1.00], respectively; branded, 1.56 [0.60] and 2.87 [0.71]) (P=0.29). After six weeks, the between-product difference in mean (SD) increase from baseline in Qmax was nonsignificant (generic, 2.36 [0.90] mL/s; branded, 2.03 [0.62] mL/s) (P=0.72).</p> <p>A total of 86 treatment emergent adverse events were reported (45 with the generic drug; 41 with the branded drug), all of which were considered by the investigator as non-serious except for one case of acute epididymitis, which occurred with the generic drug. The most common adverse events reported with the generic and branded formulations were dizziness (7/48 [14.6%] and 10/50 [20.0%], respectively) and peripheral edema (1/48 [2.1%] and 3/50 [6.0%]). No significant differences in the prevalence of adverse events were found between the two treatments.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD for 6 weeks (Period 1), followed by terazosin (generic) 1 to 4 mg QD for six weeks (Period 2)</p>				
<p>Yang et al.<sup>50</sup> (2007)</p> <p>Terazosin 2 mg QD for 1 week</p> <p>Those patients with continued LUTS after the initial treatment were allocated randomly to: terazosin 2 mg QD for 6 weeks</p> <p>vs</p> <p>terazosin 2 mg QD plus tolterodine 2 mg BID for 6 weeks</p>	<p>RCT</p> <p>Patients diagnosed with LUTS due to BPH</p>	<p>N=69</p> <p>7 weeks</p>	<p>Primary: Change from baseline in IPSS</p> <p>Secondary: Not reported</p>	<p>Primary: IPSS was significantly improved with both treatments after the initial first week, and the reduction of IPSS with combination therapy was significantly greater compared to terazosin (P&lt;0.01). A decrease in urgency, frequency and nocturia were the main contributory factors causing the reduction of IPSS with combination therapy. Differences in Qmax and residual urine from baseline were noted with both treatments, but there was no difference between the treatments (P values not reported).</p> <p>The incidence of adverse effects with combination therapy was higher compared to terazosin. The most commonly reported adverse effects were mouth dryness, which is associated with anticholinergic drugs such as tolterodine.</p> <p>Secondary: Not reported</p>
<p>Dong et al.<sup>51</sup> (2009)</p> <p>Terazosin</p> <p>vs</p> <p>tamsulosin</p>	<p>MA (12 trials)</p> <p>Patients with BPH</p>	<p>N=2,816</p> <p>4 weeks</p>	<p>Primary: IPSS, quality of life, Qmax, Qave, residual volume, prostate volume, adverse effects</p> <p>Secondary:</p>	<p>Primary: After four weeks of treatment, tamsulosin demonstrated a significant improvement in IPSS compared to terazosin (WMD, -1.24; 95% CI, -1.98 to -0.51; P=0.0009).</p> <p>There was no significant difference in quality of life between the treatment groups (WMD, -0.04; 95% CI, -0.16 to 0.24), Qmax (WMD, -0.38; 95% CI, -1.18 to 0.41), Qave (WMD, -0.39; 95% CI, -0.84 to 0.06), residual volume (WMD, -4.32; 95% CI, -10.96 to 2.33), or prostate volume</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Not reported	(WMD, -0.28; 95% CI, -3.37 to 2.81).  Fewer patients experienced dizziness (RR, 0.38; 95% CI, 0.30 to 0.48), severe hypotension (RR, 0.16; 95% CI, 0.04 to 0.68), and dry mouth (RR, 0.14; 95% CI, 0.03 to 0.77) with tamsulosin compared to patients receiving terazosin.  Secondary: Not reported
Lepor et al. <sup>52</sup> (1996)  Terazosin 1 to 10 mg QD  vs  finasteride 5 mg QD  vs  finasteride 5 mg QD plus terazosin 1 to 10 mg QD  vs  placebo	DB, MC, RCT  Men 45 to 80 years of age with symptomatic BPH	N=1,229  1 year	Primary: Change from baseline in AUA SS and Qmax  Secondary: Not reported	Primary: A significantly greater reduction in symptom scores were observed with terazosin and combination therapy compared to finasteride and placebo (6.1, 6.2, 3.2 and 2.6 points respectively; P<0.001 for terazosin vs finasteride, combination therapy vs finasteride, terazosin vs placebo and combination therapy vs placebo). There was no difference in scores noted between terazosin and combination therapy (P=1.00) or finasteride and placebo (P=0.63).  Terazosin and combination therapy was also associated with a greater increase in Qmax compared to finasteride or placebo (2.7, 3.2, 1.6 and 1.4 mL/second). Differences between finasteride and terazosin, finasteride and combination therapy, combination therapy and placebo and terazosin and placebo all reached statistical significance (P<0.001 for all comparisons), whereas the difference between terazosin and combination therapy (P=0.15) and finasteride and placebo (P=0.07) did not.  Secondary: Not reported
Liu et al. <sup>53</sup> (2009)  Terazosin 2 mg/day  vs	DB, PG, RCT  Men ≥50 years of age with Stage 1 or 2 essential HTN (SBP 140 to 180 mm Hg and/or DBP 90 to 110 mm Hg)	N=360  28 days	Primary: Reduction in the total and sub-scores of the IPSS and blood pressure  Secondary: Not reported	Primary: Treatment with terazosin and amlodipine monotherapy led to a similar reduction in the total IPSS (6.7 vs 6.9). There were no significant difference in the reduction in the bladder outlet obstruction sub-score (4.0 vs 4.1), OAB sub-score (2.9 vs 2.6), or quality of life score (1.1 vs 1.2) with amlodipine compared to terazosin.  Treatment with terazosin and amlodipine led to a greater reduction in the

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amlodipine 5 mg/day</p> <p>vs</p> <p>terazosin 2 mg/day and amlodipine 5 mg/day</p>	<p>and with LUTS (IPSS <math>\geq 10</math>)</p>			<p>QOL score (1.4 vs 1.1, <math>P &lt; 0.05</math>) compared to amlodipine monotherapy. There was no significant difference in the reduction in the total IPSS (7.8), bladder outlet obstruction sub-score (4.8), or OAB sub-score (3.2) with terazosin and amlodipine compared to amlodipine alone or terazosin alone.</p> <p>The rate of the responders (defined as patients with a reduction of 40% or more in the total IPSS, bladder outlet obstruction sub-score, OAB sub-score, or quality of life score or total IPSS of <math>&lt; 8</math>) were similar between the amlodipine group (36.1, 41.2, 46.2, and 33.6%, respectively) and terazosin group (39.3, 46.2, 39.3, and 41.0%, respectively). The rate of responders in the OAB sub-score was significantly greater in the terazosin and amlodipine group than in the terazosin group (53.8 vs 39.3%, <math>P &lt; 0.05</math>). The rate of responders in the quality of life score was significantly greater in the terazosin + amlodipine group than in the amlodipine group (47.1 vs 33.6%, <math>P &lt; 0.05</math>).</p> <p>The mean reduction in SBP and DBP was greater with amlodipine than terazosin (21.8/10.0 vs 11.9/6.5 mm Hg, <math>P &lt; 0.01</math>). The greatest reduction in SBP and DBP (25.2/12.6 mm Hg) occurred in the terazosin and amlodipine group (<math>P &lt; 0.01</math> vs terazosin and <math>P &lt; 0.05</math> vs amlodipine).</p> <p>The rates of blood pressure control were greater in the amlodipine group (63.9%) and the terazosin and amlodipine group (73.1%) than in the terazosin group (36.8%, both <math>P &lt; 0.001</math>).</p> <p>Secondary: Not reported</p>
<p>Wilt et al.<sup>54</sup> (2000)</p> <p>Terazosin</p> <p>vs</p> <p>other <math>\alpha</math>-adrenergic blocking agents,</p>	<p>SR (17 trials)</p> <p>Men with symptomatic benign prostatic obstruction</p>	<p>N=5,151</p> <p>4 to 52 weeks</p>	<p>Primary: Change from baseline in urological symptom scale scores</p> <p>Secondary: Urodynamic</p>	<p>Primary: Boyersky symptom score improved by 37% with terazosin and by 15% with placebo. AUA SS scores improved by 38% with terazosin compared to 20% with finasteride and 17% with placebo. Terazosin was comparable to tamsulosin (40 and 43%, respectively) in improving IPSS (P values not reported).</p> <p>Secondary: The improvement in Qmax reported with terazosin (22%) was similar to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
finasteride, finasteride plus terazosin or placebo			measures, adverse effects	other $\alpha$ -adrenergic blocking agents, but higher compared to finasteride (15%) and placebo (11%). Side effects, including dizziness, asthenia, headache and postural hypotension, occurred more often with terazosin compared to placebo. Rates of discontinuation with terazosin were higher than other $\alpha$ -adrenergic blocking agents, but similar to finasteride and placebo.
Wilt et al. <sup>55</sup> (2002)  Other $\alpha$ -adrenergic blocking agents, Permixon <sup>®*</sup> or placebo  vs  tamsulosin 0.2 to 0.8 mg QD	SR (14 trials)  Men with BPH and LUTS	N=4,122  4 to 26 weeks	Primary: Change from baseline in urological symptom scale scores  Secondary: Changes from baseline in Qmax, adverse effects	Primary: The WMD in the Boyarsky symptom score for tamsulosin compared to placebo was -1.1 points (95% CI, -1.49 to -0.72) or a 12% improvement with 0.4 mg and -1.6 points (95% CI, -2.3 to -1.0) or a 16% improvement with 0.8 mg.  Secondary: The WMD in Qmax was 1.1 mL/second with both tamsulosin 0.4 and 0.8 mg (95% CI, 0.59 to 1.51 with 0.4 mg; 95% CI, 0.65 to 1.48 with 0.8 mg).  Tamsulosin was reported to be as effective as other $\alpha$ -adrenergic blocking agents or Permixon <sup>®</sup> in the improvement of LUTS and Qmax.  Dizziness, rhinitis and abnormal ejaculation occurred significantly more often with tamsulosin than placebo. The rates of adverse events and withdrawal increased with higher doses of tamsulosin. Terazosin was associated with a higher rate of discontinuation than low dose tamsulosin.
<b>Hypertension</b>				
Hayduk et al. <sup>56</sup> (1987)  Doxazosin 1 to 16 mg QD  vs  terazosin 1 to 20 mg QD	DB, MC, RCT  Patients with high blood pressure	N=55  14 weeks	Primary: Proportion of patients achieving blood pressure success and normalization (blood pressure $\leq$ 90 mm Hg), safety  Secondary: Not reported	Primary: Blood pressure success was higher with doxazosin compared to terazosin (73 vs 64%; P value not reported).  Blood pressure normalization was higher with doxazosin compared to terazosin (65 vs 57%; P value not reported).  The incidence of treatment-related side effects was higher with terazosin compared to doxazosin (39 vs 30%; P value not reported).  Secondary: Not reported
Torvik et al. <sup>57</sup>	DB	N=172	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(1986) Doxazosin 1 to 16 mg QD vs prazosin 0.5 to 10 mg BID vs placebo	Patients with essential HTN	12 weeks	Changes from baseline in blood pressure, heart rate, and plasma lipid profiles  Secondary: Not reported	Doxazosin and prazosin both produced significant reductions in blood pressure compared to placebo (P<0.05 to P<0.005).  There was no significant difference between the three treatments in changes in plasma lipid profiles or heart rate (P values not reported). There was a significant baseline reduction in TG only with doxazosin (P<0.05).  Secondary: Not reported
Fukiyama et al. <sup>58</sup> (1991) Doxazosin vs prazosin	DB, MC, RCT  Patients with essential HTN	N=126  12 weeks	Primary: Changes from baseline in blood pressure and heart rate  Secondary: Not reported	Primary: There was no significant difference between the two treatments in reductions in blood pressure (P=0.7826); however, both treatments produced significant reductions from baseline (P<0.001).  No significant changes in heart rate were observed with either treatment (P value not reported).  Secondary: Not reported
DePlanque et al. <sup>59</sup> (1991) Doxazosin QD vs prazosin BID	DB, DD, PG  Patients with mild or moderate essential HTN not adequately controlled by diuretics and $\beta$ -blockers	N=43  14 weeks	Primary: Changes from baseline in blood pressure, heart rate and serum lipid levels; calculated CHD risk using the Framingham equation  Secondary: Not reported	Primary: There was no difference between the two treatments in changes in SBP (P value not significant), heart rate (P value not significant) or serum lipid levels (P value not reported). Doxazosin was associated with a significantly greater reduction in standing (P=0.01) and supine (P=0.04) DBP compared to prazosin.  At the end of the trial, 84.2 and 56.5% of patients receiving doxazosin and prazosin achieved therapeutic success (P value not reported).  Doxazosin (P=0.02) was associated with a greater reduction from baseline in the calculated risk of CHD compared to prazosin (P value not significant).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Deger et al. <sup>60</sup> (1986)  Prazosin BID  vs  terazosin QD  vs  placebo	DB, MC, PC  Patients with mild to moderate HTN	N=174  14 weeks	Primary: Changes from baseline in blood pressure and heart rate  Secondary: Not reported	Primary: Terazosin was associated with a significant reduction in supine and standing DBP compared to placebo (P≤0.05). Prazosin was not associated with a significant reduction in supine DBP, but was associated with a significant reduction in mean standing DBP compared to placebo (P values not reported).  There was no difference in the changes in heart rate between the two treatments (P value not reported).  Secondary: Not reported
Ruoff et al. <sup>61</sup> (1986)  <u>Study 1:</u> Prazosin  vs  terazosin  vs  placebo  <u>Study 2:</u> Terazosin  vs  HCTZ  <u>Study 3:</u> Terazosin and	DB, PG, RCT  Patients with mild to moderate HTN	Study 1 N=54  Study 2 N=37  Study 3 N=28	Primary: Blood pressure, pulse rate, body weight, laboratory tests, physical examinations, ECG  Secondary: Not reported	Primary: Study 1- There was no significant difference in blood pressure changes between the terazosin and prazosin treatment groups.  Study 2- HCTZ produced a significantly greater reduction in supine DBP compared to terazosin. There were no significant differences in standing blood pressure between the HCTZ and terazosin treatment groups.  Study 3- There were no significant differences in blood pressure between the treatment groups.  The drug treatments did not produce significant changes in pulse rates, body weights, laboratory test results, physical examinations, or electrocardiograms.  Secondary: Not reported



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
HCTZ  vs  prazosin and HCTZ				
Neaton et al. <sup>62</sup> (1993) TOMHS  Doxazosin 2 to 4 mg QD  vs  chlorthalidone 15 to 30 mg QD  vs  acebutolol 400 mg QD  vs  amlodipine 5 mg QD  vs  enalapril 5 to 10 mg QD  vs  placebo	DB, MC, PC, RCT  Patients with mild HTN (DBP <100 mm Hg)	N=902  4.4 years	Primary: Blood pressure, quality of life, side effects, blood lipid levels and analysis of other serum components, echocardiographic changes, and incidence of cardiovascular events  Secondary: Not reported	Primary: There was a significant reduction in blood pressure in all the active treatment groups compared to placebo (-15.9 vs -9.1 mm Hg for SBP and -12.3 vs -8.6 mm Hg for DBP; P<0.0001).  There were no major differences in blood pressure lowering between the 5 active treatment groups (P=0.10).  TC was significantly reduced more in the doxazosin group than in the amlodipine, chlorthalidone, and placebo groups (P<0.01). The reduction in LDL-C was significantly more in doxazosin group than in the amlodipine, chlorthalidone, and placebo groups. Reduction in TG was significantly larger with the doxazosin, enalapril, and amlodipine groups than acebutolol group (P<0.01).  The lowest level of fasting insulin was observed with doxazosin; fasting insulin was lower than placebo in all drug groups.  Secondary: Not reported
Liebson et al. <sup>63</sup>	DB, PC, RCT	N=844	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1995) TOMHS</p> <p>Doxazosin</p> <p>vs</p> <p>chlorthalidone</p> <p>vs</p> <p>acebutolol</p> <p>vs</p> <p>amlodipine</p> <p>vs</p> <p>enalapril</p> <p>vs</p> <p>placebo</p>	<p>Patients with mild HTN</p>	<p>4 years</p>	<p>Changes in blood pressure and pulse, changes in left ventricular mass from baseline to end of study period as assessed by ECG</p> <p>Secondary: Not reported</p>	<p>All drug treatment groups showed significantly greater reduction of blood pressure compared to placebo (mean decrease of 16/12 vs 9/9 mm Hg; P&lt;0.001).</p> <p>Pulse rate decreased by 10 bpm for the acebutolol group compared to 1 to 3 bpm for the other treatment groups.</p> <p>All drug treatment groups and the placebo group showed significant decreases (10 to 15%) in left ventricular mass. The chlorthalidone group showed the largest decrease in left ventricular mass at 34 g compared to 24 to 27 g for the other treatment groups.</p> <p>Secondary: Not reported</p>
<p>Brown et al.<sup>64</sup> (1995)</p> <p><u>Study A:</u> Doxazosin, followed by amlodipine, followed by doxazosin and amlodipine</p> <p>vs</p>	<p>DB, RCT, XO</p> <p>Patients with moderate or severe HTN</p>	<p>N=24</p> <p>18 weeks</p>	<p>Primary: Blood pressure and heart rate, foot volume as measure of edema, plasma noradrenaline concentration</p> <p>Secondary: Not reported</p>	<p>Primary: <u>Study A:</u> The decrease in blood pressure was significantly greater than the sum of the blood pressure falls at the end of the single drug treatment periods. The reduction in blood pressure was greater with amlodipine than doxazosin (P&lt;0.01). The reduction in blood pressure was greater with combination than amlodipine (P&lt;0.001).</p> <p>No significant changes in heart rate were observed. One subject developed ankle edema. The plasma noradrenaline concentration did not change significantly during the single drug treatment periods, but doubled at the end of the combination treatment period (P&lt;0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p><u>Study B:</u> Enalapril, followed by amlodipine, followed by enalapril and amlodipine</p>				<p><u>Study B:</u> The reduction in blood pressure was significantly greater with amlodipine than enalapril (P&lt;0.05). The reduction in blood pressure was significantly greater with combination than amlodipine (P&lt;0.05) with the exception of erect blood pressure.</p> <p>No significant changes in heart rate were noted. No significant difference in foot volume was observed between treatments. The plasma noradrenaline was significantly higher than at baseline (P&lt;0.01).</p> <p>Secondary: Not reported</p>
<p>Deary et al.<sup>65</sup> (2002)</p> <p>Doxazosin 1 to 4 mg/day</p> <p>vs</p> <p>amlodipine 5 mg/day</p> <p>vs</p> <p>lisinopril 2.5 to 10 mg/day</p> <p>vs</p> <p>bisoprolol 5 mg/day</p> <p>vs</p> <p>bendroflumethiazide* 2.5</p>	<p>DB, XO</p> <p>Hypertensive patients, aged 18 to 55 years old</p>	<p>N=34</p> <p>42 weeks (6 week treatment of each drug or placebo, then the 7<sup>th</sup> week was a repeat of each patient's most effective, tolerated drug)</p>	<p>Primary: Blood pressure, heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: All drug treatments caused significant decreases in blood pressure.</p> <p>Bendroflumethiazide performed significantly worse (P=0.0016) and bisoprolol performed significantly better (P=0.004) than amlodipine.</p> <p>When the most effective drugs for each patient were tabulated, all drugs included in the study except for bendroflumethiazide, were represented.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day vs placebo				
Hayduk et al. <sup>66</sup> (1987)  <u>Study 1:</u> Doxazosin 1 to 16 mg QD  vs  prazosin 1 to 20 mg BID  vs  HCTZ 25 to 100 mg QD  vs  nadolol 40 to 160 mg QD  vs  atenolol 50 to 100 mg QD  vs  metoprolol 100 to 200 mg BID	DB, MC  Patients with HTN	<u>Study 1:</u> N=903  10 to 24 week trial; therapy continued for up to 62 weeks   <u>Study 2:</u> N=52  12 weeks	Primary: Blood pressure, heart rate  Secondary: Not reported	Primary: Blood pressure lowering effect of doxazosin was similar to that of the other antihypertensive drugs.  There was no significant difference in the heart rate with the doxazosin treated group. The $\beta$ -blockers demonstrated clinically significant bradycardia.  Both doxazosin and terazosin were equally efficacious, but doxazosin was effective at significantly lower doses.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  placebo  <u>Study 2:</u> Doxazosin 16 mg QD  vs  terazosin 20 mg QD				
Trost et al. <sup>67</sup> (1987)  Doxazosin 1 to 16 mg QD  vs  HCTZ 25 to 100 mg QD	DB, MC, PG  Patients with HTN	N=104  6 months	Primary: Blood pressure, serum lipid changes  Secondary: Not reported	Primary: There was no significant difference in the supine and standing blood pressures between the two treatment groups.  There was significantly greater reduction in total TG (P=0.002) and TC concentration (P=0.006) and significantly greater increase in HDL-C:TC (P=0.001) in the doxazosin arm compared to the HCTZ arm.  Secondary: Not reported
Grimm et al. <sup>68</sup> (1996)  Doxazosin 2 to 16 mg  vs  HCTZ 25 to 50 mg	DB, PG, RCT  Patients with HTN	N=107  1 year	Primary: Blood pressure, heart rate, biochemistries, lipids/lipoproteins, quality of life, ECGs, adverse effects  Secondary: Not reported	Primary: There were no significant differences in blood pressure lowering, heart rate, quality of life measures, or serious adverse effects between the two treatment groups.  The doxazosin treated group experienced a more favorable high density lipoprotein /total cholesterol ratio (P≤0.01) compared to the hydrochlorothiazide group.  Both drug treatments showed significant reduction in left ventricular mass (P<0.001) and wall thickness (P<0.05). The left ventricular systolic and diastolic internal dimensions were significantly less in the HCTZ group compared to the doxazosin group.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
Ferrara et al. <sup>69</sup> (1993)  Doxazosin 1 to 16 mg QD  vs  captopril 25 to 150 mg QD	MC, OL, PG  Patients with hypercholesterolemia and HTN	N=224  14 weeks	Primary: Blood pressure (normalized blood pressure defined as standing diastolic pressure $\leq$ 90 mm Hg), serum lipid levels, quality of life  Secondary: Not reported	Primary: Blood pressure was significantly reduced with both drugs (P<0.001).  A total of 73% of the doxazosin group and 67% of the captopril group achieved normalized blood pressure.  Serum TC level was significantly improved with both drugs (P<0.001). The HDL-C concentration was only significantly increased in the doxazosin group (P<0.001).  The calculated 10-year risk for the development of CHD was significantly reduced with both drug treatments (P<0.001).  Secondary: Not reported
Derosa et al. <sup>70</sup> (2005)  Doxazosin 4 mg QD  vs  irbesartan 300 mg QD	DB, PG, RCT  Patients with type 2 diabetes and mild HTN	N=96  1 year	Primary: Blood pressure, glucose metabolism, lipid parameters  Secondary: Not reported	Primary: Blood pressure was significantly reduced in both treatment groups compared to baseline (P<0.01).  Irbesartan was significantly better in lowering blood pressure compared to doxazosin (P<0.05).  Doxazosin significantly reduced glycosylated hemoglobin, fasting plasma glucose, fasting plasma insulin, TC, LDL-C, HDL-C, and TG (P $\leq$ 0.05 for all parameters).  As monotherapy, neither of the drugs achieved adequate blood pressure control.  Secondary: Not reported
Taylor et al. <sup>71</sup> (1988)  Doxazosin 1 to 16 mg QD	DB, PG  Patients with mild or moderate essential HTN	N=67  18 weeks	Primary: Blood pressure (therapeutic success defined as standing DBP $\leq$ 90	Primary: A total of 74% of the doxazosin group achieved therapeutic success compared to 81% of the enalapril group.  Blood pressures were significantly reduced in both groups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs enalapril 10 to 40 mg QD			mm Hg), lipid parameters  Secondary: Not reported	There were no significant changes in the lipid profile observed for either drug.  Secondary: Not reported
Wessels et al. <sup>72</sup> (1991)  Doxazosin QD  vs  enalapril QD	DB, DD, PC, RCT  Patients with mild or moderate essential HTN	N=54  12 weeks	Primary: Blood pressure, heart rate, serum lipid profile, calculated CHD risk  Secondary: Not reported	Primary: Both drugs produced significant reductions in blood pressure (P<0.05).  There was no significant change in heart rate with both drugs.  Doxazosin showed a significant reduction in the total serum cholesterol concentration (P<0.05). Doxazosin also showed a decrease in triglyceride level (P value not significant) and an increase in HDL-C/total cholesterol ratio (P value not significant).  Coronary heart disease risk reduction was significant and greater in the doxazosin group compared to the enalapril group (-27.58 vs -18.49%, P<0.02).  Secondary: Not reported
Hjordahl et al. <sup>73</sup> (1987)  Doxazosin QD  vs  HCTZ QD	DB, RCT  Patients with mild to moderate essential HTN	N=115  24 weeks	Primary: Blood pressure, heart rate, lipid profile, side effects  Secondary: Not reported	Primary: There was no significant difference between treatment groups for blood pressure and heart rate except HCTZ produced significantly greater supine SBP than doxazosin (P=0.04).  There were significant reductions in TC (P=0.006) and total TG (P=0.018) for the doxazosin group.  Eleven patients of the HCTZ group had an abnormally low potassium level and seven of the HCTZ treated group had abnormally high uric acid concentrations.  Secondary: Not reported
Ott et al. <sup>74</sup>	DB, MC, RCT	N=126	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(1987)  Doxazosin 1 to 16 mg QD  vs  atenolol 50 to 100 mg QD	Patients with mild to moderate HTN	20 weeks	Primary: Blood pressure, heart rate  Secondary: Not reported	Primary: There was no significant difference between treatment groups in blood pressure.  Secondary: Both drugs reduced heart rate, but atenolol produced a significantly greater decrease in heart rate than doxazosin (P<0.001).  Secondary: Not reported
Frick et al. <sup>75</sup> (1986)  Doxazosin 1 to 16 mg QD  vs  atenolol 50 to 100 mg QD	DB, DD, MC, RCT  Patients with mild to moderate essential HTN	N=152  1 year	Primary: Blood pressure, heart rate, lipid profile  Secondary: Not reported	Primary: At endpoint, there was greater blood pressure reduction with atenolol than doxazosin. This was statistically significant only in the supine position (P<0.05).  Secondary: Doxazosin reduced the heart rate slightly, while atenolol produced a marked bradycardia (P<0.0001).  HDL-C:TC was raised in the doxazosin group and lowered in the atenolol group (P=0.001). TG levels decreased in the doxazosin group and increased in the atenolol group (-5.0 vs 42.7%; P<0.001).  Secondary: Not reported
Daae et al. <sup>76</sup> (1998)  Doxazosin QD  vs  atenolol QD	DB, MC, PG  Patients with mild to moderate HTN	N=228  1 year followed by a 4-year OL, ES	Primary: Blood pressure, heart rate, lipid profile, calculated risk of developing CHD in 10 years using the Framingham equation  Secondary: Not reported	Primary: Both groups showed similar decreases in blood pressure.  Secondary: The doxazosin-treated group had a significantly greater reduction from baseline in CHD risk than the atenolol-treated group (P<0.05).  TC significantly decreased from baseline in both treatment groups (P≤0.05), with no statistically significant difference between the groups.  HDL-C (P<0.01), the HDL-C:TC (P<0.01), and TG levels (P<0.01) significantly improved in the doxazosin group compared to the atenolol group.  Secondary:



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
Talseth et al. <sup>77</sup> (1991)  Doxazosin (mean dose used: 5.2 mg QD)  vs  atenolol (mean dose used: 66.4 mg QD)	PG, RCT  Patients with mild and moderate HTN	N=164  3 years	Primary: Blood pressure, heart rate, lipids profile, calculated CHD risk using the Framingham equation  Secondary: Not reported	Primary: Both drugs produced similar reductions in blood pressure.  Atenolol produced a significant decrease in heart rate (P<0.05), while doxazosin did not change the heart rate significantly.  Doxazosin significantly reduced TG levels (P<0.001), increased HDL-C levels (P<0.001), and increased the HDL-C:TC (P<0.001) compared to atenolol.  The calculated CHD risk was significantly increased with atenolol (P<0.05) and significantly decreased with doxazosin (P<0.05) from baseline.  Secondary: Not reported
Carruthers et al. <sup>78</sup> (1993)  Doxazosin QD  vs  atenolol QD	RCT  Patients with mild to moderate systemic HTN and normal serum lipid	N=191  24 weeks	Primary: Calculated CAD risk using the Framingham formula  Secondary: Not reported	Primary: Doxazosin treatment produced a significantly greater reduction in CHD risk compared to atenolol (P=0.0074).  The relative risk of CHD was reduced to 0.92 in the atenolol group (P=0.144) and 0.74 in the doxazosin group (P=0.0001) from baseline.  Secondary: Not reported
Searle et al. <sup>79</sup> (1990)  Doxazosin 11 mg (mean dose)  vs  placebo  All patients	DB, MC, RCT  Patients with mild to moderate essential HTN	N=87  12 weeks	Primary: Changes from baseline in blood pressure, heart rate and serum lipids  Secondary: Not reported	Primary: Doxazosin was associated with significant reductions in blood pressure compared to placebo (17.0/12.3 vs 6.2/6.7 mm Hg; P<0.05). The supine blood pressure was decreased by 13.2/9.8 mm Hg with doxazosin compared to 9.2/6.0 mm Hg with placebo (P value not reported).  Only minor, nonsignificant changes in serum lipids and heart rate were observed between the two treatments (P value not reported).  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
received atenolol 100 mg/day.				
Ohta et al. <sup>80</sup> (2007)  Doxazosin 1 to 2 mg QD to BID  Treatment was added to calcium channel blockers, ARBs and ACE inhibitors.	RETRO  Patients with HTN that had poorly controlled blood pressure	N=41  3 months (mean follow up 170 days)	Primary: Changes from baseline in blood pressure and blood chemistry  Secondary: Not reported	Primary: Blood pressure decreased from 152±14/81±12 to 135±14/70±11 mm Hg after the addition of doxazosin at a mean dose of 1.5 mg/day (P<0.001).  When good SBP control was defined as <140 mm Hg, the prevalence of patients with good SBP control increased from 24 to 61% with the addition of doxazosin (P<0.01). Similarly, the prevalence of patients with good DBP control (<90 mm Hg) increased from 78 to 98% (P<0.01).  Patients whose SBP decreased >10 mm Hg (n=25) showed significantly higher baseline SBP, TC and LDL-C compared to those who showed less SBP reduction (<10 mm Hg; P<0.01).  Comparable reductions in blood pressure were obtained between obese patients (BMI ≥25, change in blood pressure at three months: -15±15/-12±9 mm Hg, n=18) and non-obese patients (-14±19/-7±8 mm Hg, n=23).  Secondary: Not reported
de Alvaro et al. <sup>81</sup> (2006) ASOCIA  Doxazosin SR 4 mg QD, increased to 8 mg/day at week 4 if required  Added to entry medication.	MC, PRO  Patients with HTN (>140/>90 mm Hg) on previous antihypertensive medication who were uncontrolled	N=3,631  16 weeks	Primary: Proportion of patients achieving goal blood pressure (<140/<90 mm Hg), adverse events  Secondary: Not reported	Primary: The proportion of patients achieving goal blood pressure after four weeks of add on therapy with doxazosin was 39% and increased to 61% after 16 weeks. SBP and DBP (mean±SEM) decreased, respectively, from 161.6±0.2/95.1±0.1 mm Hg at baseline to 142.2±0.2/84.1±0.1 mm Hg after four weeks (P<0.0001) and to 136.8±0.2/80.6±0.2 mm Hg after 16 weeks (P<0.0001).  Adverse events occurred in 108 patients (3.0%), with 57 (1.6%) related to the study treatment. In 17 patients (0.5%), serious adverse events were described, but only one was related to the study drug.  Secondary: Not reported
Os et al. <sup>82</sup> (2006)	DB, PG, RCT	N=310	Primary: Efficacy, safety	Primary: All groups had a significant decrease in blood pressure at all study visits

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Doxazosin 4 mg QD</p> <p>vs</p> <p>doxazosin 2 mg QD</p> <p>vs</p> <p>doxazosin SR 4 mg QD</p>	<p>Patients 18 to 80 years of age with mild to moderate essential HTN (sitting DBP 95 to 110 mm Hg and SBP &lt;180 mm Hg)</p>	<p>9 weeks</p>	<p>Secondary: Not reported</p>	<p>compared to baseline. The proportion of patients who reached goal sitting DBP (&lt;90 mm Hg) was similar among the three treatment groups, except at week one, when more patients in the doxazosin SR group had obtained the goal compared to those in the doxazosin 2 mg group (40.6 vs 22.3%; P=0.005). The proportion of patients who reached sitting SBP (&lt;140 mm Hg) goal was similar among groups.</p> <p>Adverse event profiles among the groups were similar.</p> <p>Secondary: Not reported</p>
<p>Williams et al.<sup>83</sup> (2015) PATHWAY-2</p> <p>Twelve weeks of once daily treatment with each of spironolactone (25 to 50 mg), bisoprolol (5 to 10 mg), doxazosin modified release (4 to 8 mg), and placebo, in addition to their baseline blood pressure drugs</p>	<p>DB, PC, XO</p> <p>Patients 18 to 79 years of age with seated clinic SBP <math>\geq</math> 140 mmHg (or <math>\geq</math>135 mmHg for patients with diabetes) and home SBP (18 readings over four days) <math>\geq</math>130 mmHg, despite treatment for at least three months with maximally tolerated doses of three drugs (an ACE or ARB, a CCB, and a diuretic)</p>	<p>N=335</p> <p>12 months</p>	<p>Primary: Average home SBP, recorded in the morning and the evening in triplicate, on four consecutive days before study visits</p> <p>Secondary: Clinic SBP, BP control rates, adverse events</p>	<p>Primary: The average reduction in home SBP by spironolactone was significantly greater compared to placebo (-8.70 mmHg; 95% CI, -9.72 to -7.69; P&lt;0.0001), compared to the mean of the other two active treatments (doxazosin and bisoprolol; -4.26; 95% CI, -5.13 to -3.38; P&lt;0.0001), and compared to the individual treatments; versus doxazosin (-4.03; 95% CI, -5.04 to -3.02; P&lt;0.0001) and versus bisoprolol (-4.48; 95% CI, -5.50 to -3.46; P&lt;0.0001).</p> <p>Secondary: The results for seated clinic SBP largely mirror those seen with home SBP except that there was a large placebo effect on clinic BP that was not seen with home BP measurement.</p> <p>Overall 219 (68.9%; 95% CI, 63.6 to 73.8) of 314 patients achieved target home SBP of &lt;135 mmHg. 58% of patients had their BP controlled with spironolactone, which was significantly greater than rates for other treatments (P&lt;0.001 when compared to doxazosin, bisoprolol, and placebo). Most patients who were controlled by doxazosin or bisoprolol had a still greater fall in blood pressure on spironolactone, which was consequently the most effective treatment in almost 60% of patients. This was at least three times the proportion in whom doxazosin or bisoprolol were the most effective.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Materson et al.<sup>84</sup> (1994)</p> <p>Prazosin 4 to 20 mg QD</p> <p>vs</p> <p>HCTZ 12.5 to 50 mg QD</p> <p>vs</p> <p>atenolol 25 to 100 mg QD</p> <p>vs</p> <p>captopril 25 to 100 mg QD</p> <p>vs</p> <p>clonidine 0.2 to 0.6 mg QD</p> <p>vs</p> <p>diltiazem SR 120 to 360 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, RCT</p> <p>Men with DBP of 95 to 109 mm Hg</p>	<p>N=1,292</p> <p>1 year</p>	<p>Primary: Success as defined by DBP ≤95 mm Hg at 1 year</p> <p>Secondary: Not reported</p>	<p>All active treatments were well tolerated with similar low rates of adverse events and withdrawals due to adverse events.</p> <p>Primary: Success rates were 59% for diltiazem, 51% for atenolol, 50% for clonidine, 46% for HCTZ, 42% for captopril, 42% for prazosin, and 25% for placebo (P&lt;0.001 between diltiazem and HCTZ, atenolol and prazosin).</p> <p>The rates of adverse effects leading to termination of treatment were highest with prazosin at 13.8% and clonidine at 10.1%, which was significantly different from captopril at 4.8%, atenolol at 2.2%, HCTZ at 1.1%, diltiazem at 5.5%, and placebo at 6.4%.</p> <p>Successful blood pressure control was highest with diltiazem at 64% in African Americans, highest with captopril at 55% in younger whites, and highest with atenolol at 68% in older whites.</p> <p>Secondary: Not reported</p>
<p>McAreavey et al.<sup>85</sup></p>	<p>DB, PG, RCT</p>	<p>N=238</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1984)</p> <p>Prazosin 0.5 mg QD up to 10 mg BID</p> <p>vs</p> <p>hydralazine 12.5 mg QD up to 100 mg BID</p> <p>vs</p> <p>labetalol 200 mg QD up to 1,600 mg BID</p> <p>vs</p> <p>methyldopa 125 mg QD up to 1,000 mg BID</p> <p>vs</p> <p>placebo</p> <p>Minoxidil as add on therapy was given to men only.</p> <p>Doses were titrated upward at 2 week intervals until target BP or maximum dose</p>	<p>Patients with inadequately controlled HTN while receiving atenolol 100 mg/day and bendrofluazide* 5 mg/day</p>	<p>6 months</p>	<p>Comparative safety and efficacy, target blood pressure &lt;140/95 mm Hg</p> <p>Secondary: Not reported</p>	<p>Target blood pressure was reached in 25% of patients receiving hydralazine, 23% of patients receiving minoxidil, 19% of patients receiving prazosin, 17% of patients receiving methyldopa and zero percent of patients receiving placebo (P values not reported).</p> <p>Labetalol had the highest withdrawal rate compared to the other treatments with 78% (P&lt;0.05). Minoxidil had the second highest withdrawal rate with 57% (P&lt;0.05), due to fluid retention. There were no significant differences in withdrawal rates among the other treatments.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
was reached.				
Chrysant et al. <sup>86</sup> (1986)  Terazosin  vs  placebo	DB, MC, PC, RCT  Patients with inadequate control of essential HTN	N=138  Duration not specified	Primary: Changes from baseline in blood pressure, physical examination and ECG  Secondary: Not reported	Primary: There was a significant mean reduction in supine DBP with the terazosin compared to placebo (7.3 vs 0.6 mm Hg; P<0.05).  There were no significant changes between treatments in physical examinations or ECGs.  Secondary: Not reported
Holtzman et al. <sup>87</sup> (1988)  Terazosin  vs  placebo in combination with atenolol	DB, MC, PC  Patients with HTN	N=92  10 weeks	Primary: Changes from baseline in blood pressure and lipid profiles  Secondary: Not reported	Primary: There was a significant reduction in supine and standing blood pressure (P<0.05), TC (P<0.05) and LDL-C plus VLDL-C (P<0.05) with terazosin.  Secondary: Not reported
Casas et al. <sup>88</sup> (2005)  ACE inhibitor or ARBs compared to other antihypertensive drugs (β-adrenergic blocking agents, α-adrenergic blocking agents, calcium-channel blocking agents, or combinations)  vs	MA (127 trials)  Studies in adults that examined the effect of any drug treatment with a blood pressure lowering action on progression of renal disease	N=not reported  4.2 years (mean)	Primary: Doubling of serum creatinine, and ESRD  Secondary: Serum creatinine, urine albumin excretion and GFR	Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.  A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.  Secondary: Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).  Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ACE inhibitor or ARBs compared to placebo</p> <p>Specific agents and doses were not specified.</p>				<p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>
<b>Outcomes Trials</b>				
<p>ALLHAT<sup>89-91</sup> (2000, 2003, 2004)</p> <p>Doxazosin 2 to 8 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg QD</p>	<p>AC, DB, RCT</p> <p>Patients <math>\geq 55</math> years of age with HTN and <math>\geq 1</math> CHD risk factor</p>	<p>N=24,335</p> <p>3.3 years</p>	<p>Primary: Combined occurrence of CHD death or nonfatal MI</p> <p>Secondary: All-cause mortality, stroke, combined CHD (CHD death, nonfatal MI, revascularization procedures and hospitalized angina), stroke, combined cardiovascular disease (CHD death, nonfatal MI, stroke, revascularization procedures, angina, CHF and PAD)</p>	<p>Primary: There was no difference in risk of the combined primary endpoint between the two treatments (P=0.71).</p> <p>Secondary: There was no difference in risk of all-cause mortality between the two treatments (P=0.71).</p> <p>Compared to chlorthalidone, doxazosin was associated with a significantly higher risk of stroke (RR, 1.19; 95% CI, 1.01 to 1.40; P=0.04) and combined cardiovascular disease (RR, 1.25; 95% CI, 1.17 to 1.33; P&lt;0.001).</p> <p>The risk of CHF doubled with doxazosin compared to chlorthalidone (P&lt;0.001).</p> <p>Doxazosin was associated with a significantly higher risk of angina (RR, 1.16; P&lt;0.001) and coronary revascularization (RR, 1.15; P=0.05).</p> <p>No difference between the two treatments were observed for risk of PAD (RR, 1.07; P=0.50)</p>
<p>Wright et al.<sup>92</sup> (2008) ALLHAT</p>	<p>DB, RCT</p> <p>Hypertensive individuals with and</p>	<p>N=42,418</p> <p>3.2 years (median)</p>	<p>Primary: Fatal CHD or nonfatal MI</p>	<p>Primary: No differences were noted among the four treatment groups, regardless of race or metabolic/cardiomatabolic syndrome status for the primary end point (fatal CHD or nonfatal MI).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Doxazosin vs chlorthalidone vs amlodipine vs lisinopril</p>	<p>without metabolic/ cardiometabolic syndrome</p>	<p>follow-up)</p>	<p>Secondary: Heart failure, combined cardiovascular disease, stroke, ESRD</p>	<p>Secondary: Significantly higher rates of heart failure were consistent across all treatment comparisons in those with metabolic/cardiometabolic syndrome. RRs were 1.50 (95% CI, 1.18 to 1.90), 1.49 (1.17 to 1.90), and 1.88 (1.42 to 2.47) in African American participants and 1.25 (1.06 to 1.47), 1.20 (1.01 to 1.41), and 0.82 (1.51 to 2.19) in non-African American participants for amlodipine, lisinopril, and doxazosin comparisons with chlorthalidone, respectively.</p> <p>Higher rates for combined cardiovascular disease were observed with lisinopril and chlorthalidone (RR, 1.24; 95% CI, 1.09 to 1.40; RR, 1.10; 95% CI, 1.02 to 1.19, respectively) and doxazosin and chlorthalidone comparisons (RR, 1.37; 95% CI, 1.19 to 1.58; RR, 1.18; 95% CI, 1.08 to 1.30, respectively) in African American and non-African American participants with metabolic/cardiometabolic syndrome.</p> <p>Higher rates of stroke were seen in African American participants only (RR, 1.37; 95% CI, 1.07 to 1.76 for the lisinopril and chlorthalidone comparison, and RR, 1.49; 95% CI, 1.09 to 2.03 for the doxazosin and chlorthalidone comparison). African American patients with metabolic/cardiometabolic syndrome also had higher rates of end-stage renal disease (RR, 1.70; 95% CI, 1.13 to 2.55) with lisinopril compared to chlorthalidone.</p>
<p>Dahlöf et al.<sup>93</sup> (2005) ASCOT-BPLA  Amlodipine 5 to 10 mg and if needed perindopril 4 to 8 mg  or  atenolol 50 to 100</p>	<p>MC, RCT  Patients with HTN</p>	<p>N=19,257  5.5 years</p>	<p>Primary: Nonfatal MI and fatal CHD Secondary: Nonfatal MI, and fatal CHD, total coronary endpoint, total cardiovascular events and procedures, all- cause mortality, cardiovascular</p>	<p>Primary: The trial was halted early due to findings that patients on the amlodipine and perindopril regimen had fewer of the primary endpoints (P=0.1052) and lower rates of fatal and nonfatal stroke (P=0.0003), total cardiovascular events and procedures (P&lt;0.0001), all-cause mortality (P=0.025), and incidence of developing diabetes (P&lt;0.0001).</p> <p>There was a greater reduction in blood pressure by an average of 2.7/1.9 mm Hg in the amlodipine-based regimen compared to the atenolol-based regimen.</p> <p>There was no significant difference in the percent of patients (25%) that stopped therapy because of an adverse event between the two treatment</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg and if needed bendroflumethiazide* 1.25 to 2.5 mg</p> <p>If goal blood pressure was still not achieved, doxazosin 4 to 8 mg was added to the regimen.</p>			<p>mortality, fatal and nonfatal stroke, fatal and nonfatal heart failure, silent MI, unstable angina, chronic stable angina, PAD, life-threatening arrhythmias, development of diabetes mellitus, development of renal impairment</p>	<p>groups. However, a significantly greater proportion of patients in the amlodipine-based regimen stopped the trial therapy early because of serious adverse events compared to the atenolol-based regimen (P&lt;0.0001).</p> <p>Secondary: Patients on the amlodipine-perindopril regimen had fewer fatal and nonfatal strokes (P=0.0003), total cardiovascular events and procedures (P&lt;0.0001), and all-cause mortality (P=0.025).</p> <p>Patients on the amlodipine and perindopril regimen had less chance of developing diabetes (P&lt;0.0001).</p>
<p>Chapman et al.<sup>94</sup> (2007) ASCOT-BPLA</p> <p>Atenolol 50 to 100 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); bendroflumethiazide* plus potassium 1.25 to 2.5 mg plus doxazosin were added for additional blood pressure control; if blood pressure remained elevated on the 3 above drugs,</p>	<p>Subanalysis of ASCOT-BPLA evaluating effects of spironolactone on treatment-resistant HTN</p> <p>Patients 40 to 79 years of age with HTN and ≥3 cardiovascular risk factors, with SBP ≥160 mm Hg and/or DBP ≥100 mm Hg (not on antihypertensive therapy) or SBP ≥140 mm Hg and/or DBP ≥90 mm Hg (on antihypertensive therapy)</p>	<p>N=1,411</p> <p>1.3 years</p>	<p>Primary: Change in DBP and SBP, adverse effects</p> <p>Secondary: Not reported</p>	<p>Primary: Spironolactone-treated patients lead to a significant 21.9 mm Hg reduction in SBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 20.8 to 23.0 mm Hg; P&lt;0.001).</p> <p>Spironolactone-treated patients lead to a significant 9.5 mm Hg reduction in DBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 9.0 to 10.1; P&lt;0.001).</p> <p>Spironolactone-treated patients exhibited small but significant decreases in sodium, LDL-C and TC as well as increases in potassium, glucose, creatinine and HDL-C (P&lt;0.05).</p> <p>The most common adverse effect reported in the trial was gynecomastia in men (P value not reported).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
spironolactone 25 mg was added to the regimen  vs  amlodipine 5 to 10 mg titrated to target blood pressure <140/90 mm Hg (or <130/90 mm Hg in diabetic patients); perindopril 4 to 8 mg and doxazosin were added for additional control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen				

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, IR=immediate-release, QD=once daily, QID=four times daily, SR=sustained-release, TID=three times daily

Study design abbreviations: AC=active comparator, DB=double blind, DD=double dummy, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group,

PRO=prospective, RCT=randomized controlled trial, RETRO=retrospective, SR=systematic review, XO=cross over

Miscellaneous abbreviations: AUA-SS=American Urology Association Symptom Score, BPH=benign prostatic hyperplasia, CAD=coronary artery disease, CHD=coronary artery disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, ECG=electrocardiogram, ESRD=end stage renal disease, GFR=glomerular filtration rate, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HTN=hypertension, IIEF=International Index of Erectile Function, IPSS=International Prostate Symptom Score, LDL-C=low-density lipoprotein cholesterol, LUTS=lower urinary tract symptoms, MI=myocardial infarction, OAB=overactive bladder, OR=odds ratio, PAD=peripheral artery disease, PSA=prostate-specific antigen, PVR=post-void residual urine volume, Qave=average urinary flow rate, Qmax=maximum urinary flow rate, RR=relative risk, SBP=systolic blood pressure, SD=standard deviation, SEM=standard error of mean, SFAQ= Sexual Function Abbreviated Questionnaire, TC=total cholesterol, TG=triglycerides, TPV=total prostate volume, VLDL-C=very low-density lipoprotein cholesterol, WMD=weighted mean difference

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 9. Relative Cost of the Alpha-Adrenergic Blocking Agents**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Doxazosin	extended-release tablet, tablet	Cardura <sup>®*</sup> , Cardura XL <sup>®</sup>	\$\$\$\$\$	\$
Prazosin	capsule	Minipress <sup>®*</sup>	\$\$\$\$\$	\$
Terazosin	capsule	N/A	N/A	\$

\*Generic is available in at least one dosage form or strength.

N/A=not available

**X. Conclusions**

The alpha-adrenergic blocking agents are approved for the treatment of benign prostatic hyperplasia (BPH) and hypertension.<sup>1-6</sup> All of the agents are available in a generic formulation. Treatment guidelines on the management of BPH recommend the use of an  $\alpha$ -adrenergic blocking agent or a 5 $\alpha$ -reductase inhibitor in patients with moderate-to-severe symptoms. Alpha-blockers can quickly improve symptoms and flow rate, while the 5 $\alpha$ -reductase inhibitors have the potential for long-term reduction in prostate volume.<sup>8,9</sup> Available data suggests that the combination is also effective. Clinical trials have demonstrated similar efficacy among the various  $\alpha$ -adrenergic blocking agents for the treatment of BPH.<sup>26-55</sup>

There are several national and international organizations that have published guidelines on the treatment of hypertension. Most of the guidelines do not address the use of the  $\alpha$ -adrenergic blocking agents.<sup>17-25</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension. According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>17</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>17-22</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>17-23</sup>

Several clinical trials have demonstrated that the  $\alpha$ -adrenergic blocking agents effectively lower blood pressure when administered as monotherapy or in combination with other antihypertensive agents. Comparative studies have demonstrated similar efficacy when the  $\alpha$ -blockers were directly compared to each other, as well as when they were compared to ACE inhibitors,  $\beta$ -blockers, calcium-channel blocking agents and thiazide-type diuretics.<sup>56-88</sup> The ALLHAT trial evaluated the effects of doxazosin on cardiovascular morbidity and mortality. Treatment with doxazosin increased the risk of stroke and cardiovascular events; however, it provided other benefits including improvements in insulin resistance and lipid parameters.<sup>89-92</sup>

There is insufficient evidence to support that one brand alpha-adrenergic blocking agent is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand alpha-adrenergic blocking agents within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand alpha-adrenergic blocking agent is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. Cardura® [package insert]. New York (NY): Pfizer Inc; 2021 Jun.
4. Cardura XL® [package insert]. New York, NY: Pfizer Inc; 2020 Dec.
5. Minipress® [package insert]. New York, NY: Pfizer Inc; 2016 Jul.
6. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
7. Kasiske BL, Ma JZ, Kalil RS, et al. Effects of antihypertensive therapy on serum lipids. *Ann Intern Med*. 1995;122:133-41.
8. Lerner LB, McVary KT, Barry MJ, Kumar Das A, Gandhi MC, Kaplan SA, et al. Management of lower urinary tract symptoms attributed to benign prostatic hyperplasia: AUA Guideline. *J Urol* 2021; 206: 818.
9. Gravas S, Cornu JN, Gacci M, Gratzke C, Herrmann TRW, Mamoulakis C, et al, Guidelines Associates. Management of Non-neurogenic Male LUTS. EAU Guideline. ISBN 978-94-92671-13-4. EAU Guidelines Office, Arnhem, The Netherlands. 2021.
10. Frishman WH. Alpha and beta-adrenergic blocking drugs. In: Cardiovascular pharmacotherapeutics. 3rd ed. Frishman WH, Sica DA, editors. Minneapolis (MN): Cardiotext Publishing; 2011; 57-86.
11. Schwinn DA, Roehrborn CG. A<sub>1</sub> adrenoceptor subtypes and lower urinary tract symptoms. *Int J Urol*. 2008 Mar;15(3):193-9.
12. Cheng TO. Should patients with benign prostatic hypertrophy stop taking doxazosin in the light of the ALLHAT study? *Int J Cardiol*. 2006;107:275-6.
13. Berecek KH, Carey RM. Adrenergic and dopaminergic receptors and actions. In: (Izzo JL Jr [ed.]). Hypertension primer: the essentials of high blood pressure. Dallas (TX): American Heart Association;2008: 39-43.
14. Fulton B, Wagstaff AJ, Sorkin EM. Doxazosin: an update of its clinical pharmacology and therapeutic applications in hypertension and benign prostatic hyperplasia. *Drugs*. 1995;49(2):295-320.
15. Maclaren R, Mueller SW, Dasta JF. Use of vasopressors and inotropes in the pharmacotherapy of shock. In: DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, editors. Pharmacotherapy: a pathophysiologic approach. 10th edition. New York (NY): McGraw-Hill; 2017.
16. Cunningham GR, Kadmon D. Medical treatment of benign prostatic hyperplasia. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA, 2019.
17. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
18. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
19. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086
20. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
21. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
22. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
23. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010;56:780-800.

24. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int.* 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003
25. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
26. Lee JY, Lee SH, Kim SJ, Kim CS, Lee HM, Kim CI, et al. Change in International Prostate Symptom storage subscore after long-term medical therapy in BPH patients: finasteride and alpha-blocker combination therapy in men with moderate-to-severe LUTS/BPH in Korea. *Urology.* 2011 Jan;77(1):171-6.
27. Demir O, Ozdemir I, Bozkurt O, Aslan G, Esen AA. The effect of  $\alpha$ -blocker therapy on erectile functions in patients with lower urinary tract symptoms due to benign prostate hyperplasia. *Asian Journal of Andrology.* 2009;11:716-22.
28. Sun GH, Tsui KH, Wu TT, Chang CH, Cheng CL, Schou M. Efficacy and safety of the doxazosin gastrointestinal therapeutic system for the treatment of benign prostate hyperplasia. *Kaohsiung J Med Sci.* 2010;26(10):532-9.
29. Kirby RS, Andersen M, Gratzke P, Dahlstrand C, Hoe K. A combined analysis of double-blind trials of the efficacy and tolerability of doxazosin-gastrointestinal therapeutic system, doxazosin standard and placebo in patients with benign prostatic hyperplasia. *BJU Int.* 2001;87:192-200.
30. Keten T, Aslan Y, Balci M, et al. Determination of the efficiency of 8 mg doxazosin XL treatment in patients with an inadequate response to 4 mg doxazosin XL treatment for benign prostatic hyperplasia. *Urology.* 2015 Jan;85(1):189-194.
31. Samli MM, Dincel C. Terazosin and doxazosin in the treatment of BPH: results of a randomized study with crossover in non-responders. *Urol Int.* 2004;73:125-9.
32. Kaplan SA, Te AE, Ikeguchi E, et al. The treatment of benign prostatic hyperplasia with alpha blockers in men over the age of 80 years. *Br J Urol.* 1997;80:875-9.
33. Kaplan SA, Soldo KA, Olsson CA. Terazosin and doxazosin in normotensive men with symptomatic prostatism: a pilot study to determine the effect of dosing regimen on efficacy and safety. *Eur Urol.* 28(3):223-8, 1995.
34. Bozlu M, Ulusoy E, Cayan S, et al. A comparison of four different  $\alpha 1$ -blockers in benign prostatic hyperplasia patients with and without diabetes. *Scand J Urol Nephrol.* 2004;38:391-5.
35. Xue Z, Zhang Y, Ding Q, et al. Doxazosin gastrointestinal therapeutic system versus tamsulosin for the treatment of benign prostatic hyperplasia: a study in Chinese patients. *Int J Urol.* 2007 Feb;14(2):118-22.
36. Rahardjo D, Soebadi D, Sugandi S, et al. Efficacy and safety of tamsulosin hydrochloride compared to doxazosin in the treatment of Indonesian patients with lower urinary tract symptoms due to benign prostatic hyperplasia. *Int J Urol.* 2006 Nov;13(11):1405-9.
37. Pompeo AC, Rosenblatt C, Bertero E, et al. A randomised, double-blind study comparing the efficacy and tolerability of controlled-release doxazosin and tamsulosin in the treatment of benign prostatic hyperplasia in Brazil. *Int J Clin Pract.* 2006 Oct;60(10):1172-7.
38. Cao Y, Wang Y, Guo L, Yang X, Chen T, Niu H. A Randomized, Open-Label, Comparative Study of Efficacy and Safety of Tolterodine Combined with Tamsulosin or Doxazosin in Patients with Benign Prostatic Hyperplasia. *Med Sci Monit.* 2016 Jun 4;22:1895-902.
39. Johnson TM 2nd, Burrows PK, Kusek JW, et al. The effect of doxazosin, finasteride and combination therapy on nocturia in men with benign prostatic hyperplasia. *J Urol.* 2007 Nov;178(5):2045-50.
40. Crawford ED, Wilson SS, McConnell JD, et al. Baseline factors as predictors of clinical progression of benign prostatic hyperplasia in men treated with placebo. *J Urol.* 2006 Apr;175(4):1422-6.
41. Kaplan SA, McConnell JD, Roehrborn CG, et al. Combination therapy with doxazosin and finasteride for benign prostatic hyperplasia in patients with lower urinary tract symptoms and a baseline total prostate volume of 25 ml or greater. *J Urol.* 2006 Jan;175(1):217-20.
42. Kaplan SA, Roehrborn CG, McConnell JD, et al. Long-term treatment with finasteride results in a clinically significant reduction in total prostate volume compared to placebo over the full range of baseline prostate sizes in men enrolled in the MTOPS trial. *J Urol* 2008;180:1030-2.
43. Kirby RS, Roehrborn C, Boyle P, Bartsch G, Jardin A, Cary MM, et al. Efficacy and tolerability of doxazosin and finasteride, alone or in combination, in treatment of symptomatic benign prostatic hyperplasia: the prospective European doxazosin and combination therapy (PREDICT) trial. *Urology.* 2003;61(1):119-26.
44. Fwu CW, Eggers PW, Kaplan SA, Kirkali Z, Lee JY, Kusek JW. Long-term effects of doxazosin, finasteride and combination therapy on quality of life in men with benign prostatic hyperplasia. *J Urol.* 2013 Jul;190(1):187-193.

45. Djavan B, Marberger M. A meta-analysis on the efficacy and tolerability of  $\alpha_1$ -adrenoceptor antagonists in patients with lower urinary tract symptoms suggestive of benign prostatic obstruction. *Eur Urol*. 1999;36:1-13.
46. Nickel JC, Sander S, Moon TD, et al. A meta-analysis of the vascular-related safety profile and efficacy of alpha-adrenergic blockers for symptoms related to benign prostatic hyperplasia. *Int J Clin Pract* 2008;62:1547-59.
47. MacDonald R, Wilt TJ. Alfuzosin for treatment of lower urinary tract symptoms compatible with benign prostatic hyperplasia: a systematic review of efficacy and adverse effects. *Urology*. 2005;66:780-8.
48. Tsujii T. Comparison of prazosin, terazosin and tamsulosin in the treatment of symptomatic benign prostatic hyperplasia: a short-term open, randomized multicenter study. *Int J Urol*. 2000 Jun;7(6):199-205.
49. Tsai YS, Lan SK, Ou JH, et al. Effects of branded versus generic terazosin hydrochloride in adults with benign prostatic hyperplasia: a randomized, open-label, crossover study in Taiwan. *Clin Ther*. 2007 Apr;29(4):670-82.
50. Yang Y, Zhao XF, Li HZ, et al. Efficacy and safety of combined therapy with terazosin and tolterodine for patients with lower urinary tract symptoms associated with benign prostatic hyperplasia: a prospective study. *Chin Med J (Engl)*. 2007 Mar 5;120(5):370-4.
51. Dong Z, Wang Z, Yang K, et al. Tamsulosin versus terazosin for benign prostatic hyperplasia: a systematic review. *Syst Biol Reprod Med* 2009;55:129-36.
52. Lepor H, Williford WO, Barry MJ, Brawer MK, Dixon CM, Gormley G, et al. The efficacy of terazosin, finasteride, or both in benign prostatic hyperplasia. *N Eng J Med*. 1996;355:533-9.
53. Liu H, Liu P, Mao G, et al. Efficacy of combined amlodipine/terazosin therapy in male hypertensive patients with lower urinary tract symptoms: a randomized, double-blind clinical trial. *Urology* 2009;74:130-6.
54. Wilt T, Howe RW, Rutks I, MacDonald R. Terazosin for benign prostatic hyperplasia. *Cochrane Database of Systemic Reviews* 2000, Issue 1. Art. No.: CD003851. DOI: 10.1002/14651858. CD003851.
55. Wilt T, MacDonald R, Rutks I. Tamsulosin for benign prostatic hyperplasia. *Cochrane Database of Systemic Reviews* 2002, Issue 4. Art No.: CD002081. DOI: 10.1002/14651858. CD002081.
56. Hayduk K and Schneider HT. Antihypertensive effects of doxazosin in systemic hypertension and comparison with terazosin. *Am J Cardiol*. 1987;59:95G-8G.
57. Torvik D and Madsbu HP. Multicentre 12-week double-blind comparison of doxazosin, prazosin and placebo in patients with mild to moderate essential hypertension. *Br J Clin Pharmacol*. 1986;21 Suppl 1:69S-75S.
58. Fukiyama K, Omae T, Limura O, et al. A double-blind comparative study of doxazosin and prazosin in the treatment of essential hypertension. *Am Heart J*. 1991;121:317-22.
59. DePlanque BA. A double-blind comparative study of doxazosin and prazosin when administered with beta-blockers or diuretics. *Am Heart J*. 1991;121:304-11.
60. Deger G. Comparison of the safety and efficacy of once-daily terazosin versus twice-daily prazosin for the treatment of mild to moderate hypertension. *Am J Med*. 1986 May 23;80(5B):62-7.
61. Ruoff G. Comparative trials of terazosin with other antihypertensive agents. *Am J Med*. 1986;80(5B):42-8.
62. Neaton JD, Grimm RH, Prineas RJ, et al. Treatment of mild hypertension study. Final results. *JAMA*. 1993;270(6):713-24.
63. Liebson PR, Grandits GA, Dianzumba S, et al. Comparison of five antihypertensive monotherapies and placebo for change in left ventricular mass in patients receiving nutritional-hygienic therapy in the treatment of mild hypertension study (TOMHS). *Circulation*. 1995;91:698-706.
64. Brown MJ and Dickerson JEC. Alpha-blockade and calcium antagonism: an effective and well-tolerated combination for the treatment of resistant hypertension. *J Hypertens*. 1995;13:701-7.
65. Deary AJ, Schumann AL, Murfet H, et al. Double-blind, placebo-controlled crossover comparison of five classes of antihypertensive drugs. *J Hypertens*. 2002 Apr;20(4):771-7.
66. Hayduk K. Efficacy and safety of doxazosin in hypertension therapy. *Am J Cardiol*. 1987;59:35G-39G.
67. Trost BN, Weidmann P, Riesen W, et al. Comparative effects of doxazosin and hydrochlorothiazide on serum lipids and blood pressure in essential hypertension. *Am J Cardiol*. 1987;59:99G-104G.
68. Grimm RH, Flack JM, Schoenberger JA, et al. Alpha-blockade and thiazide treatment of hypertension: a double-blind randomized trial comparing doxazosin and hydrochlorothiazide. *Am J Hypertens*. 1996;9:445-54.
69. Ferrara LA, Di Marino L, Russo O, et al. Doxazosin and captopril in mildly hypercholesterolemic hypertensive patients: the doxazosin-captopril in hypercholesterolemic hypertensives study. *Hypertension*. 1993;21:97-104.
70. Derosa G, Cicero AFG, Gaddi A, et al. Effects of doxazosin and irbesartan on blood pressure and metabolic control in patients with type 2 diabetes and hypertension. *J Cardiovasc Pharmacol*. 2005;45(6):599-604.
71. Taylor SH, Lee PS, Sharma SK. A comparison of doxazosin and enalapril in the treatment of mild and moderate essential hypertension. *Am Heart J*. 1988;116:1820-5.

72. Wessels F. Double-blind comparison of doxazosin and enalapril in patients with mild or moderate essential hypertension. *Am Heart J.* 1991;121:299-303.
73. Hjortdahl P, von Krogh H, Daae L, et al. A 24-week multicenter double-blind study of doxazosin and hydrochlorothiazide in patients with mild to moderate essential hypertension. *Acta Med Scand.* 1987;221(5):427-34.
74. Ott P, Storm TL, Krusell LR, et al. Multicenter, double-blind comparison of doxazosin and atenolol in patients with mild to moderate hypertension. *Am J Cardiol.* 1987;59(14):73G-77G.
75. Frick MH, Halttunen P, Himanen P, et al. A long-term double-blind comparison of doxazosin and atenolol in patients with mild to moderate essential hypertension. *Br J Clin Pharmacol.* 1986;21(Supp 1):55S-62S.
76. Daae LN, Westlie L. A 5-year comparison of doxazosin and atenolol in patients with mild-to-moderate hypertension: effects on blood pressure, serum lipids, and coronary heart disease risk. *Blood Press.* 1998;7:39-45.
77. Talseth T, Westlie L, Daae L. Doxazosin and atenolol as monotherapy in mild and moderate hypertension: a randomized, parallel study with a three-year follow-up. *Am Heart J.* 1991;121:280-5.
78. Carruthers G, Dessain P, Fodor G, et al. Comparative trial of doxazosin and atenolol on cardiovascular risk reduction in systemic hypertension. *Am J Cardiol.* 1993;71(7):575-81.
79. Searle M, Dathan R, Dean S, et al. Doxazosin in combination with atenolol in essential hypertension: a double-blind placebo-controlled multicentre trial. *Eur J Clin Pharmacol.* 1990;39:299-300.
80. Ohta Y, Tsuchihashi T, Onaka U, et al. Usefulness of the alpha1-blocker doxazosin as a third-line antihypertensive drug. *Hypertens Res.* 2007 Apr;30(4):301-6.
81. de Alvaro F, Hernández-Presa MA; ASOCIA Study. Effect of doxazosin gastrointestinal therapeutic system on patients with uncontrolled hypertension: the ASOCIA Study. *J Cardiovasc Pharmacol.* 2006 Feb;47(2):271-6.
82. Os I. Comparison of doxazosin GITS and standard doxazosin in the treatment of high blood pressure. *Int J Clin Pharmacol Ther.* 2006 Mar;44(3):99-106.
83. Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G, et al. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial.
84. Materson BJ, Reda DJ, Cushman WC, et al. Single-drug therapy for hypertension in men – A comparison of six antihypertensive agents with placebo. *N Engl J Med.* 1994;330(23):914-21.
85. McAreavey D, Ramsey LE, Latham L, McLaren AD, Lorimer AR, Reid JL, et al. Third drug trial: comparative study of antihypertensive agents added to treatment when blood pressure remains uncontrolled by a beta blocker plus thiazide diuretic. *Br Med J (Clin Res Ed).* 1984 Jan 14;288(6411):106-11.
86. Chrysant SG. Experience with terazosin administered in combination with other antihypertensive agents. *Am J Med.* 1986;80:5(B):55-61.
87. Holtzman JL, Kaihlanen PM, Rider JA, et al. Concomitant administration of terazosin and atenolol for the treatment of essential hypertension. *Arch Intern Med.* 1988;148(3):539-43.
88. Casas JP, Chua W, Loukogeorgakis S, Vallance P, Smeeth L, Hingorani AD, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet.* 2005 Dec 10;366:2026-33.
89. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major cardiovascular events in hypertensive patients randomized to doxazosin vs chlorthalidone. The antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *JAMA.* 2000;283:1967-75.
90. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Diuretic versus [alpha]-blocker as first-step antihypertensive therapy: final results from the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *JAMA.* 2003;42(3):239-46.
91. Barzilay JI, Davis BR, Bettencourt J, et al. Cardiovascular outcomes using doxazosin vs chlorthalidone for the treatment of hypertension in older adults with and without glucose disorders: a report from the ALLHAT study. *J Clin Hypertens.* 2004;6(3):116-25.
92. Wright JT, Harris-Haywood S, Pressel S, et al. Clinical outcomes by race in hypertensive patients with and without the metabolic syndrome: Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Arch Intern Med.* 2008 Jan 28;168(2):207-17.
93. Dahlöf B, Sever PS, Poulter NR, et al. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomized controlled trial. *Lancet.* 2005;366(9489):895-906.
94. Chapman N, Dobson J, Wilson S, Dahlöf B, Sever PS, Wedel H, Poulter NR; Anglo-Scandinavian Cardiac Outcomes Trial Investigators. Effect of spironolactone on blood pressure in subjects with resistant hypertension. *Hypertension.* 2007 Apr;49(4):839-45.



**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Beta-Adrenergic Blocking Agents  
AHFS Class 242400  
May 4, 2022**

**I. Overview**

The beta-adrenergic blocking agents ( $\beta$ -blockers) are approved for the treatment of angina, arrhythmias, essential tremor, heart failure, hypertension, hypertrophic aortic stenosis, migraine prophylaxis, myocardial infarction, and pheochromocytoma.<sup>1-22</sup> Additionally, propranolol is the first agent Food and Drug Administration-approved for the treatment of proliferating infantile hemangioma requiring systemic therapy.<sup>12,24</sup> The  $\beta$ -blockers differ with regards to their adrenergic-receptor blocking, membrane stabilizing and intrinsic sympathomimetic activities, as well as lipophilicity.<sup>1-23</sup> There are at least three distinct types of  $\beta$  receptors distributed throughout the body ( $\beta_1$ ,  $\beta_2$ , and  $\beta_3$ ).  $\beta_1$  receptors are located predominantly in the heart and kidneys.  $\beta_2$  receptors are located in the lungs, gastrointestinal tract, liver, uterus, vascular smooth muscle, and skeletal muscle.  $\beta_3$ -receptors are located in fat cells.  $\beta$ -blockers primarily exert their effects through a blockade of  $\beta_1$  and  $\beta_2$  receptor subtypes. Agents that have a greater affinity for  $\beta_1$  receptors are considered to be cardioselective. These agents may be safer in patients with asthma, chronic obstructive pulmonary disease, and peripheral vascular disease because they produce less inhibition of  $\beta_2$  receptors, which mediate vasoconstriction and bronchospasm. Cardioselectivity is dose dependent; therefore,  $\beta_2$  blockade can occur at higher doses with these agents.<sup>23,25</sup> Carvedilol and labetalol also block  $\alpha$ -adrenergic receptors, which would be expected to reduce peripheral vascular resistance to a greater extent than other  $\beta$ -blockers.<sup>23,25</sup>

The  $\beta$ -blockers are available as single entity agents, as well as fixed-dose combination products. Each of the combination products contains a thiazide-type diuretic. The thiazide-type diuretics inhibit the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride.<sup>1,2,17-22</sup>

The  $\beta$ -adrenergic blocking agents that are included in this review are listed in Table 1 and comparative information on cardioselectivity is highlighted in Table 2. This review encompasses all dosage forms and strengths. All of the agents are available in a generic formulation, with the exception of penbutolol. This class was last reviewed in February 2020.

**Table 1. Beta-Adrenergic Blocking Agents Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Acebutolol	capsule	N/A	acebutolol
Atenolol	tablet	Tenormin <sup>®*</sup>	atenolol
Betaxolol	tablet	N/A	betaxolol
Bisoprolol	tablet	N/A	bisoprolol
Carvedilol	extended-release capsule, tablet	Coreg <sup>®*</sup> , Coreg CR <sup>®*</sup>	carvedilol
Esmolol	injection <sup>^</sup>	Brevibloc <sup>®*</sup>	none
Labetalol	injection <sup>^</sup> , tablet	N/A	labetalol
Metoprolol	extended-release capsule, extended-release tablet, injection, tablet	Kaspargo Sprinkle <sup>®</sup> , Lopressor <sup>®*</sup> , Toprol-XL <sup>®*</sup>	metoprolol
Nadolol	tablet	Corgard <sup>®*</sup>	nadolol
Nebivolol	tablet	Bystolic <sup>®*</sup>	Bystolic <sup>®</sup> , nebivolol
Penbutolol	tablet	Levatol <sup>®†</sup>	none
Pindolol	tablet	N/A	pindolol
Propranolol	extended-release capsule, injection, solution, tablet	Hemangeol <sup>®</sup> , Inderal LA <sup>®*</sup> , Inderal XL <sup>®</sup> , InnoPran XL <sup>®</sup>	propranolol
Sotalol	tablet, solution	Betapace <sup>®*</sup> , Betapace	sotalol

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
		AF <sup>®*</sup> , Sotylize <sup>®</sup>	
Timolol	tablet	N/A	timolol
<b>Combination Products</b>			
Atenolol and chlorthalidone	tablet	Tenoretic <sup>®*</sup>	atenolol and chlorthalidone
Bisoprolol and hydrochlorothiazide	tablet	Ziac <sup>®*</sup>	bisoprolol and hydrochlorothiazide
Metoprolol and hydrochlorothiazide	tablet	N/A	metoprolol and hydrochlorothiazide
Nadolol and bendroflumethiazide	tablet	N/A	nadolol and bendroflumethiazide

\*Generic is available in at least one dosage form or strength.

^Product is primarily administered in an institution.

†Product was discontinued by manufacturer, but remains active in pharmacy system.

PDL=Preferred Drug List

N/A=Not available

**Table 2. Selected Pharmacologic Properties of the Beta-Adrenergic Blocking Agents<sup>1-22</sup>**

Generic Name(s)	Adrenergic-Receptor Blocking Activity	Membrane Stabilizing Activity	Intrinsic Sympathomimetic Activity
Acebutolol	$\beta_1^*$	+†	+
Atenolol	$\beta_1^*$	0	0
Betaxolol	$\beta_1^*$	+	0
Bisoprolol	$\beta_1^*$	0	0
Carvedilol	$\alpha_1 - \beta_1 - \beta_2$	++	0
Labetalol	$\alpha_1 - \beta_1 - \beta_2$	0	+
Metoprolol	$\beta_1^*$	0†	0
Nadolol	$\beta_1 - \beta_2$	0	0
Nebivolol	$\beta_1^*$	0	0
Penbutolol	$\beta_1 - \beta_2$	0	+
Pindolol	$\beta_1 - \beta_2$	+	++
Propranolol	$\beta_1 - \beta_2$	++	0
Sotalol	$\beta_1 - \beta_2$	0	0
Timolol	$\beta_1 - \beta_2$	0	0

0=none; +=low; ++=moderate; +++ =high

\*Inhibits  $\beta_2$  receptors (bronchial and vascular) at higher doses.

†Detectable only at doses much greater than required for  $\beta$  blockade.

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the  $\beta$ -adrenergic blocking agents are summarized in Table 3.

**Table 3. Treatment Guidelines Using the Beta-Adrenergic Blocking Agents**

Clinical Guideline	Recommendations
European Society of Cardiology: <b>Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)<sup>26</sup></b>	<p><b>Pharmacological management of stable coronary artery disease (CAD) patients</b></p> <ul style="list-style-type: none"> <li>The two aims of the pharmacological management of stable CAD patients are to obtain relief of symptoms and to prevent CV events.</li> <li>Optimal medical treatment indicates at least one drug for angina/ischaemia relief plus drugs for event prevention.</li> <li>It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>It is indicated to review the patient's response soon after starting therapy.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin, consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</li> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief: <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a beta-blocker and/or a non-DHP-calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</li> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</li> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischaemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and beta-blockers avoided.</li> </ul> </li> <li>• Event prevention: <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><u>Treatment in patients with microvascular angina</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>• <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>• Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>• ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>• Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul> <p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>• Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy (DAPT).</li> <li>• Aspirin is recommended for elective stenting.</li> <li>• Clopidogrel is recommended for elective stenting.</li> <li>• Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>• GP IIb/IIIa antagonists should be considered for bailout situation only.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Platelet function testing or genetic testing may be considered in specific or high risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>• Prasugrel or ticagrelor may be considered in specific high-risk situations of elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</li> <li>• Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>• Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>• Prasugrel or ticagrelor is not recommended in low-risk elective stenting.</li> <li>• After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low</li> <li>• Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months</li> </ul> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all revascularized patients receive a secondary prevention and be scheduled for follow-up visit.</li> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after second generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/ American College of Cardiology Foundation/ American Heart Association/ American Association for Thoracic Surgery/ Preventive Cardiovascular Nurses Association/ Society of Thoracic Surgeons:  <b>Management of</b></p>	<p><u>Medical therapy to prevent myocardial infarction (MI) and death in patients with stable ischemic heart disease (IHD)</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</li> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable</li> </ul>

Clinical Guideline	Recommendations
<p><b>Stable Ischemic Heart Disease (2012)</b><sup>27</sup></p>	<p>IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</p> <ul style="list-style-type: none"> <li>• Patients should receive an annual influenza vaccine.</li> </ul> <p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> <li>• Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/ American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes (2014)</b><sup>28</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation &lt;90%, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications <ul style="list-style-type: none"> <li>○ Nitrates <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>○ Beta-adrenergic blockers <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>blockade (e.g., PR interval &gt;0.24 second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</li> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> </li> <li>○ Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval &gt;0.24 seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</p> <ul style="list-style-type: none"> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> <ul style="list-style-type: none"> <li>○ Other anti-ischemic interventions <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> </li> </ul> <ul style="list-style-type: none"> <li>• Inhibitors of renin-angiotensin-aldosterone system <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF &lt;0.40 and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF &lt;0.40 who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine &gt;2.5 mg/dL in men or &gt;2.0 mg/dL in women) or hyperkalemia (K &gt;5.0 mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF &lt;0.40, diabetes mellitus, or heart failure.</li> </ul> </li> <li>• Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include: <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>part of initial antiplatelet therapy. Preferred options are eptifibatide or tirofiban.</p> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet agents <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatide, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> <li>• Anticoagulant therapy <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>• Timing of coronary artery bypass grafting (CABG) in relation to use of antiplatelet agents <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> </li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>• Medications at discharge <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</p> <ul style="list-style-type: none"> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post-NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> <ul style="list-style-type: none"> <li>● Late hospital and post-hospital oral antiplatelet therapy <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation (2020)</b><sup>29</sup></p>	<p><u>Pharmacological treatment of ischemia</u></p> <ul style="list-style-type: none"> <li>● Sublingual or intravenous nitrates and early initiation of <math>\beta</math>-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>● Continuation of chronic <math>\beta</math>-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>● Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>● In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>● Aspirin is recommended for all patients without contraindications at an initial oral</li> </ul>



Clinical Guideline	Recommendations
	<p>loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</p> <ul style="list-style-type: none"> <li>• A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds. <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of</li> </ul>

Clinical Guideline	Recommendations
	<p data-bbox="532 205 821 233">parenteral anticoagulation.</p> <p data-bbox="488 264 1373 323"><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul data-bbox="488 327 1414 1415" style="list-style-type: none"> <li data-bbox="488 327 1390 422">• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li data-bbox="488 426 1414 541">• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li data-bbox="488 546 1382 604">• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li data-bbox="488 609 1414 703">• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li data-bbox="488 707 1328 766">• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li data-bbox="488 770 1382 829">• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li data-bbox="488 833 1295 892">• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li data-bbox="488 896 1382 1012">• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li data-bbox="488 1016 1373 1110">• If at low bleeding risk (HAS-BLED <math>\leq 2</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li data-bbox="488 1115 1409 1230">• If at high bleeding risk (HAS-BLED <math>\geq 3</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li data-bbox="488 1234 1377 1329">• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED <math>\geq 3</math> and low risk of stent thrombosis).</li> <li data-bbox="488 1333 1373 1360">• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li data-bbox="488 1365 1300 1415">• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p data-bbox="488 1451 1227 1478"><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul data-bbox="488 1482 1414 1906" style="list-style-type: none"> <li data-bbox="488 1482 1414 1577">• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li data-bbox="488 1581 1373 1675">• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li data-bbox="488 1680 1349 1738">• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li data-bbox="488 1743 1409 1837">• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li data-bbox="488 1841 1401 1906">• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>

Clinical Guideline	Recommendations
<p>American College of Cardiology/ American Heart Association: <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>30</sup></p>	<p><u>Routine medical therapies: <math>\beta</math>-blockers</u></p> <ul style="list-style-type: none"> <li>• Oral <math>\beta</math>-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral <math>\beta</math>-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li>• <math>\beta</math>-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>• Patients with initial contraindications to the use of <math>\beta</math>-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>• It is reasonable to administer intravenous <math>\beta</math>-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia.</li> </ul> <p><u>Routine medical therapies: Renin-Angiotensin-Aldosterone System Inhibitors</u></p> <ul style="list-style-type: none"> <li>• An angiotensin-converting enzyme (ACE) inhibitor should be administered within the first 24 hours to all patients with STEMI with anterior location, HF, or ejection fraction (EF) <math>\leq</math>40%, unless contraindicated.</li> <li>• An angiotensin receptor blocker (ARB) should be given to patients with STEMI who have indications for but are intolerant of ACE inhibitors.</li> <li>• An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq</math>40% and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>• High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>• It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>31</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of STEMI</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>• Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>• A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>• In patients with an indication for oral anticoagulation, oral anticoagulants are indicated in addition to antiplatelet therapy</li> <li>• In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the estimated risk of recurrent coronary events and bleeding).</li> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age <math>\geq</math>50 years, and at least one of the following</li> </ul>

Clinical Guideline	Recommendations
	<p>risk factors: age <math>\geq 65</math> years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR <math>&lt; 60</math> mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</p> <ul style="list-style-type: none"> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with <math>\beta</math>-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with <math>\beta</math>-blockers is indicated in patients with heart failure or left ventricular dysfunction, LVEF <math>&lt; 40\%</math> unless contraindicated.</li> <li>• Intravenous <math>\beta</math>-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> <li>• Intravenous <math>\beta</math>-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of <math>&lt; 1.8</math> mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C <math>&gt; 1.8</math> mmol/L (<math>&gt; 70</math> mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction <math>\leq 40\%</math> and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
<p>American college of Cardiology/ American Heart Association: <b>Guideline on the Primary Prevention of Cardiovascular disease (2019)</b><sup>32</sup></p>	<p><u>Top 10 messages for the primary prevention of cardiovascular disease</u></p> <ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</li> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened</li> </ul>

Clinical Guideline	Recommendations
	<p>beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</p> <ul style="list-style-type: none"> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</li> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be <math>&lt; 130/80</math> mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul> <p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a decision is made for statin therapy, a moderate-intensity statin should be recommended.</li> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) patients, LDL-C levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</li> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (<math>5\%</math> to <math>&lt; 7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> </li> <li>• In patients at borderline risk (<math>5\%</math> to <math>&lt; 7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul> <p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>• In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include: <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>• In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>• In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>• In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>• In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than 130/80 mm Hg.</li> <li>• In adults with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering medication are recommended.</li> <li>• In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul> <p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>• All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>• To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>• In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>• In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: <b>2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)</b><sup>33</sup></p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><u>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</u></p> <ul style="list-style-type: none"> <li>For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may</li> </ul>



Clinical Guideline	Recommendations
	<p>increase risk of mortality. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may</li> </ul>

Clinical Guideline	Recommendations
	<p>be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</p> <ul style="list-style-type: none"> <li>Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>34</sup></p>	<p><b>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</b></p> <ul style="list-style-type: none"> <li>An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Thromboembolism prophylaxis (e.g., with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>American Heart Association/ American College of Cardiology/ Heart Rhythm Society: <b>2019</b> <b>AHA/ACC/HRS Focused update of the 2014</b> <b>AHA/ACC/ARS Guideline for the Management of Patients with Atrial Fibrillation (2019)</b><sup>35</sup></p>	<p><u>Recommendations for risk-based anticoagulant therapy:</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>• In patients with atrial fibrillation (AF), anticoagulant therapy should be individualized based on shared decision-making after discussion of the absolute and relative risks of stroke, bleeding and the patient’s values and preferences (Level of Evidence: C).</li> <li>• NOACs (dabigatran, rivaroxaban, apixaban, and edoxaban) are recommended over warfarin in NOAC-eligible patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) (Level of Evidence: A).</li> <li>• Selection of anticoagulant therapy should be based on the risk of thromboembolism irrespective of whether the AF patten is paroxysmal, persistent, or permanent (Level of Evidence: B).</li> <li>• In patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve), the CHA<sub>2</sub>DS<sub>2</sub>-VASc score is recommended for assessment of stroke risk (Level of Evidence: B). For patients with AF who have mechanical heart valves, warfarin is recommended. (Level of Evidence: B).</li> <li>• Selection of anticoagulant therapy should be based on the risk of thromboembolism, irrespective of whether the AF pattern is paroxysmal, persistent, or permanent.</li> <li>• For patients with AF and an elevated CHA<sub>2</sub>DS<sub>2</sub>-VASc score <math>\geq 2</math> in men or <math>\geq 3</math> in women, oral anticoagulants are recommended. Options include warfarin (Level of Evidence: A), dabigatran, rivaroxaban, apixaban, or edoxaban. (Level of Evidence: B).</li> <li>• Among patients treated with warfarin, the INR should be determined at least weekly during initiation of anticoagulant therapy and at least monthly when anticoagulation (INR in range) is stable (Level of Evidence: A)</li> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) who are unable to maintain a therapeutic INR level with warfarin, use of a non-vitamin K oral anticoagulant (NOAC) is recommended (Level of Evidence: C-EO).</li> <li>• Re-evaluation of the need for and choice of anticoagulant therapy at periodic intervals is recommended to reassess stroke and bleeding risks (Level of Evidence: C).</li> <li>• Renal function should be evaluated prior to initiation of a NOAC and should be re-evaluated when clinically indicated and at least annually (Level of Evidence: B-NR).</li> <li>• For patients with atrial flutter, anticoagulant therapy is recommended according to the same risk profile used for AF (Level of Evidence: C).</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 0 in men or 1 in women, it is reasonable to omit antithrombotic therapy (Level of Evidence: B).</li> <li>• For patients with AF with a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of <math>\geq 2</math> in men or <math>\geq 3</math> in women and who have end-stage chronic kidney disease (creatinine clearance <math>&lt; 15</math> mL/min) or who are on dialysis, it is reasonable to prescribe warfarin (INR 2.0 to 3.0) or apixaban for oral anticoagulation (Level of Evidence: B).</li> </ul> <p><b>Class IIb</b></p> <ul style="list-style-type: none"> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 1 in men and 2 in women, prescribing an oral anticoagulant to reduce thromboembolic stroke risk may be</li> </ul>

Clinical Guideline	Recommendations
	<p>considered. (Level of Evidence: C-LD).</p> <ul style="list-style-type: none"> <li>For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and moderate-to-severe chronic kidney disease with elevated CHA<sub>2</sub>DS<sub>2</sub>-VASc, treatment with reduced doses of direct thrombin or factor Xa inhibitors may be considered (e.g., dabigatran, rivaroxaban, apixaban, or edoxaban. (Level of Evidence: C).</li> </ul> <p><b>Class III: No Benefit</b></p> <ul style="list-style-type: none"> <li>In patients with AF and end-stage CKD or on dialysis, the direct thrombin inhibitor dabigatran or the factor Xa inhibitors rivaroxaban or edoxaban are not recommended because of the lack of evidence from clinical trials that benefit exceeds risk (Level of Evidence: C-EO)</li> </ul> <p><b>Class III: Harm</b></p> <ul style="list-style-type: none"> <li>The direct thrombin inhibitor, dabigatran, should not be used in patients with AF and a mechanical heart valve (Level of Evidence: B).</li> </ul> <p><u>Recommendations for Interruption and Bridging Anticoagulation</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>Bridging therapy with UFH or LMWH is recommended for patients with AF and a mechanical heart valve undergoing procedures that require interruption of warfarin. Decisions regarding bridging therapy should balance the risks of stroke and bleeding (Level of Evidence: C).</li> <li>For patients with AF without mechanical heart valves who require interruption of warfarin or newer anticoagulants for procedures, decisions about bridging therapy (LMWH or UFH) should balance the risks of stroke and bleeding and the duration of time a patient will not be anticoagulated (Level of Evidence: B-R).</li> <li>Idarucizumab is recommended for the reversal of dabigatran in the event of life-threatening bleeding or an urgent procedure (Level of Evidence: B-NR)</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>Andexanet alfa can be useful for the reversal of rivaroxaban and apixaban in the event of life-threatening or uncontrolled bleeding (Level of Evidence: B-NR).</li> </ul> <p><u>Recommendations for Thromboembolism Prevention:</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>For patients with AF or atrial flutter of 48-hour duration or longer, or when the duration of AF is unknown, anticoagulation with warfarin (INR 2.0 to 3.0), a factor Xa inhibitor, or direct thrombin inhibitor is recommended for at least three weeks prior to and four weeks after cardioversion, regardless of the CHA<sub>2</sub>DS<sub>2</sub>-VASc score and the method used to restore sinus rhythm (Level of Evidence: B).</li> <li>For patients with AF or atrial flutter of more than 48 hours duration that requires immediate cardioversion for hemodynamic instability, anticoagulation should be initiated as soon as possible and continued for at least four weeks after cardioversion unless contraindicated (Level of Evidence: C).</li> <li>After cardioversion for AF of any duration, the decision about long-term anticoagulation therapy should be based on the thromboembolic risk profile and bleeding risk profile (Level of Evidence: C-EO)</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>For patients with AF or atrial flutter of less than 48-hour duration and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of ≥2 in men or ≥3 in women, administration of, heparin, a factor Xa or direct thrombin inhibitor, is recommended as soon as possible before cardioversion, followed by long-term anticoagulation therapy (Level of Evidence: B-NR).</li> <li>After cardioversion for AF of any duration, the decision regarding long-term anticoagulation therapy should be based on the thromboembolic risk profile (Level of Evidence: C-EO).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of 48-hour duration or longer or of unknown duration who have not been anticoagulated for the preceding three weeks, it is reasonable to perform a TEE prior to cardioversion and proceed with cardioversion if no LA thrombus is identified, including in the LAA, provided that anticoagulation is achieved before TEE and maintained after cardioversion for at least four weeks (Level of Evidence: B).</li> <li>• For patients with AF or atrial flutter of 48-hour duration or longer, or when the duration of AF is unknown, anticoagulation with dabigatran, rivaroxaban, or apixaban is reasonable for at least three weeks prior to and four weeks after cardioversion (Level of Evidence: C).</li> </ul> <p>Class IIb</p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of less than 48 hours' duration with a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 0 in men or 1 in women, administration of heparin, a factor Xa inhibitor, or a direct thrombin inhibitor, versus no anticoagulant therapy, may be considered before cardioversion, without the need for post cardioversion oral anticoagulation (Level of Evidence: B-NR).</li> </ul> <p><u>Recommendations for pharmacological cardioversion</u></p> <p>Class I</p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of 48 hours' duration or longer, or when the duration of AF is unknown, anticoagulation with warfarin (INR 2.0 to 3.0), a factor Xa inhibitor, or direct thrombin inhibitor is recommended for at least 3 weeks before and at least 4 weeks after cardioversion, regardless of the CHA<sub>2</sub>DS<sub>2</sub>-VASc score or the method (electrical or pharmacological) used to restore sinus rhythm (Level of Evidence: B-R).</li> <li>• For patients with AF or atrial flutter of more than 48 hours' duration or unknown duration that requires immediate cardioversion for hemodynamic instability, anticoagulation should be initiated as soon as possible and continued for at least 4 weeks after cardioversion unless contraindicated (Level of Evidence: C).</li> <li>• After cardioversion for AF of any duration, the decision about long-term anticoagulation therapy should be based on the thromboembolic risk profile and bleeding risk profile (Level of Evidence: C-EO).</li> </ul> <p><u>Recommendations for AF complicating ACS</u></p> <p>Class I</p> <ul style="list-style-type: none"> <li>• For patients with ACS and AF at increased risk of systemic thromboembolism (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater), anticoagulation is recommended unless the bleeding risk exceeds the expected benefit (Level of Evidence: B-R).</li> <li>• Urgent direct-current cardioversion of new-onset AF in the setting of ACS is recommended for patients with hemodynamic compromise, ongoing ischemia, or inadequate rate control (Level of Evidence: C).</li> <li>• Intravenous beta blockers are recommended to slow a rapid ventricular response to AF in patients with ACS who do not display HF, hemodynamic instability, or bronchospasm (Level of Evidence: C).</li> </ul> <p>Class IIa</p> <ul style="list-style-type: none"> <li>• If triple therapy (oral anticoagulant, aspirin, and P2Y<sub>12</sub> inhibitor) is prescribed for patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone percutaneous coronary intervention (PCI) with stenting for ACS, it is reasonable to choose clopidogrel in preference to prasugrel (Level of Evidence: B-NR).</li> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with a P2Y<sub>12</sub> inhibitor (clopidogrel or ticagrelor) and dose-adjusted vitamin K antagonist is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with P2Y<sub>12</sub> inhibitors (clopidogrel) and low-dose rivaroxaban 15 mg daily is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with a P2Y<sub>12</sub> inhibitor (clopidogrel) and dabigatran 150 mg twice daily is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> </ul> <p>Class IIb</p> <ul style="list-style-type: none"> <li>• If triple therapy (oral anticoagulant, aspirin, and P2Y<sub>12</sub> inhibitor) is prescribed for patients with AF who are at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) and who have undergone PCI with stenting (drug eluting or bare metal) for ACS, a transition to double therapy (oral anticoagulant and P2Y<sub>12</sub> inhibitor) at 4 to 6 weeks may be considered (Level of Evidence: B-R).</li> <li>• Administration of amiodarone or digoxin may be considered to slow a rapid ventricular response in patients with ACS and AF associated with severe LV dysfunction and HF or hemodynamic instability (Level of Evidence: C).</li> <li>• Administration of amiodarone or digoxin may be considered to slow a rapid ventricular response in patients with ACS and AF associated with severe LV dysfunction and HF or hemodynamic instability (Level of Evidence: C).</li> </ul> <p><u>Recommendations for Device Detection of AF and Atrial Flutter</u></p> <p>Class I</p> <ul style="list-style-type: none"> <li>• In patients with cardiac implantable electronic devices (pacemakers or implanted cardioverter-defibrillators), the presence of recorded atrial high-rate episodes (AHREs) should prompt further evaluation to document clinically relevant AF to guide treatment decisions (Level of Evidence: B-NR).</li> </ul> <p>Class IIa</p> <ul style="list-style-type: none"> <li>• In patients with cryptogenic stroke (i.e., stroke of unknown cause) in whom external ambulatory monitoring is inconclusive, implantation of a cardiac monitor (loop recorder) is reasonable to optimize detection of silent AF (Level of Evidence: B-R).</li> </ul> <p><u>Recommendations for Weight loss</u></p> <p>Class I</p> <ul style="list-style-type: none"> <li>• For overweight and obese patients with AF, weight loss, combined with risk factor modification, is recommended (Level of Evidence: B-R).</li> </ul>
<p>National Institute for Health and Care Excellence: <b>Atrial Fibrillation: Diagnosis and Management (2021)</b><sup>36</sup></p>	<p><u>Rate control</u></p> <ul style="list-style-type: none"> <li>• Offer rate control as the first-line treatment strategy for atrial fibrillation except in people:             <ul style="list-style-type: none"> <li>○ whose atrial fibrillation has a reversible cause</li> <li>○ who have heart failure thought to be primarily caused by atrial fibrillation</li> <li>○ with new-onset atrial fibrillation</li> <li>○ with atrial flutter whose condition is considered suitable for an ablation strategy to restore sinus rhythm</li> <li>○ for whom a rhythm-control strategy would be more suitable based on clinical judgement.</li> </ul> </li> <li>• Offer either a standard β-blocker (that is, a β-blocker other than sotalol) or a rate-limiting calcium channel blocker (CCB) (diltiazem or verapamil) as initial monotherapy to people with atrial fibrillation (AF) unless the person has the features described above. Base the choice of drug on the person's symptoms, heart rate, comorbidities and preferences.</li> <li>• Consider digoxin monotherapy for initial rate control for people with</li> </ul>



Clinical Guideline	Recommendations
	<p>non-paroxysmal atrial fibrillation if the person does no or very little physical exercise or if other rate-limiting drug options are ruled out because of comorbidities or the person's preferences.</p> <ul style="list-style-type: none"> <li>• If monotherapy does not control symptoms, and if continuing symptoms are thought to be due to poor ventricular rate control, consider combination therapy with any two of the following: a <math>\beta</math>-blocker, diltiazem, and digoxin.</li> <li>• Do not offer amiodarone for long-term rate control.</li> </ul> <p><u>Rhythm control</u></p> <ul style="list-style-type: none"> <li>• Consider pharmacological and/or electrical rhythm control for people with AF whose symptoms continue after heart rate has been controlled or for whom a rate-control strategy has not been successful.</li> </ul> <p><u>Drug treatment for long-term rhythm control</u></p> <ul style="list-style-type: none"> <li>• Assess the need for drug treatment for long-term rhythm control, taking into account the person's preferences, associated comorbidities, risks of treatment, and likelihood of recurrence of AF.</li> <li>• Do not offer class 1c antiarrhythmic drugs such as flecainide or propafenone to people with known ischaemic or structural heart disease.</li> <li>• If drug treatment for long-term rhythm control is needed, consider a standard <math>\beta</math>-blocker as first-line treatment unless there are contraindications.</li> <li>• If <math>\beta</math>-blockers are contraindicated or unsuccessful, assess the suitability of alternative drugs for rhythm control, taking comorbidities into account.</li> <li>• Dronedarone is recommended as an option for the maintenance of sinus rhythm after successful cardioversion in people with paroxysmal or persistent atrial fibrillation:             <ul style="list-style-type: none"> <li>○ Whose AF is not controlled by first-line therapy (usually including <math>\beta</math>-blockers), that is, as a second-line treatment option and after alternative options have been considered AND</li> <li>○ Who have at least one of the following cardiovascular risk factors:                 <ul style="list-style-type: none"> <li>▪ Hypertension requiring drugs of at least two different classes.</li> <li>▪ Diabetes mellitus.</li> <li>▪ Previous TIA, stroke, or systemic embolism.</li> <li>▪ Left atrial diameter of 50 mm or greater, OR</li> <li>▪ Age <math>\geq 70</math> years, AND</li> </ul> </li> <li>○ Who do not have left ventricular systolic dysfunction, AND</li> <li>○ Who do not have a history of, or current, heart failure.</li> </ul> </li> <li>• People who do not meet the criteria above who are currently receiving dronedarone should have the option to continue treatment until they and their clinicians consider it appropriate to stop.</li> <li>• Consider amiodarone for people with left ventricular impairment or heart failure.</li> <li>• Do not offer class 1c antiarrhythmic drugs such as flecainide or propafenone to people with known ischemic or structural heart disease.</li> <li>• Where people have infrequent paroxysms and few symptoms, or where symptoms are induced by known precipitants (such as alcohol, caffeine), a 'no drug treatment' strategy or a 'pill-in-the-pocket' strategy should be considered and discussed with the person.</li> </ul> <p><u>Preventing postoperative atrial fibrillation</u></p> <ul style="list-style-type: none"> <li>• In people having cardiothoracic surgery:             <ul style="list-style-type: none"> <li>○ reduce the risk of postoperative atrial fibrillation by offering one of the following: amiodarone; a standard beta-blocker (that is, a beta-blocker other than sotalol); a rate-limiting calcium-channel blocker (diltiazem or verapamil).</li> <li>○ do not offer digoxin.</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In people having cardiothoracic surgery who are already on beta-blocker therapy, continue this treatment unless contraindications develop (such as postoperative bradycardia or hypotension).</li> <li>• Do not start statins in people having cardiothoracic surgery solely to prevent postoperative atrial fibrillation.</li> <li>• In people having cardiothoracic surgery who are already on statins, continue this treatment. For further advice on statins for the prevention of cardiovascular disease, see NICE's guideline on cardiovascular disease: risk assessment and reduction.</li> </ul>
<p>American Association for Thoracic Surgery: <b>2014 AATS Guidelines for the Prevention and Management of Peri-Operative Atrial Fibrillation and Flutter (POAF) for Thoracic Surgical Procedures (2014)</b><sup>37</sup></p>	<p><u>Recommended prevention strategies for all postoperative atrial fibrillation (POAF) patients</u></p> <ul style="list-style-type: none"> <li>• Patients taking <math>\beta</math>-blockers prior to thoracic surgery should continue them in the postoperative period to avoid <math>\beta</math>-blockade withdrawal.</li> <li>• Intravenous magnesium supplementation may be considered to prevent postoperative AF when serum magnesium level is low or it is suspected that total body magnesium is depleted.</li> <li>• Digoxin should not be used for prophylaxis against AF.</li> </ul> <p><u>Recommended prevention strategies for intermediate to high-risk POAF patients</u></p> <ul style="list-style-type: none"> <li>• It is reasonable to administer diltiazem to those patients with preserved cardiac function who are not taking <math>\beta</math>-blockers preoperatively in order to prevent POAF.</li> <li>• It is reasonable to consider the postoperative administration of amiodarone to reduce the incidence of POAF for intermediate and high risk patients undergoing pulmonary resection.</li> <li>• Postoperative administration of intravenous amiodarone may be considered to prevent POAF in patients undergoing esophagectomy.</li> <li>• Atorvastatin may be considered to prevent POAF for statin naïve patients scheduled for intermediate and high risk thoracic surgical procedures.</li> </ul> <p><u>Rate control recommendations for patients with new onset POAF</u></p> <ul style="list-style-type: none"> <li>• Intravenous administration of <math>\beta</math>-blockers (e.g., esmolol or metoprolol) or nondihydropyridine calcium channel blockers (diltiazem or verapamil) is recommended to achieve rate control (heart rate <math>\leq 110</math> bpm) for patients who develop POAF with rapid ventricular response.</li> <li>• Caution should be used with patients with hypotension, left ventricular (LV) dysfunction, or heart failure.</li> <li>• Combination use of atrioventricular (AV) nodal blocking agents, such as <math>\beta</math>-blockers (e.g., esmolol or metoprolol), nondihydropyridine calcium channel antagonists (e.g., diltiazem or verapamil), or digoxin, can be useful to control heart rates when a single agent fails to control rates of POAF. The choice should be individualized and doses modified to avoid bradycardia.</li> <li>• For patients with hypotension, heart failure or LV dysfunction, or when other measures are unsuccessful or contraindicated, intravenous amiodarone can be useful for control of heart rate. Amiodarone could result in conversion to sinus rhythm, and if it is initiated after 48 hours of AF, both a transesophageal echocardiography (TEE) when possible, to rule out left atrial/LA appendage (LA/LAA) thrombus, and full anticoagulation should be considered.</li> <li>• For patients with heart failure, LV dysfunction or hypotension, intravenous digoxin may be considered for rate control of POAF.</li> <li>• For patients with ventricular preexcitation (i.e., Wolff-Parkinson-White syndrome) and POAF, use of AV nodal blocking agents, such as <math>\beta</math>-blockers (e.g., esmolol or metoprolol), intravenous amiodarone, nondihydropyridine calcium channel antagonists (e.g., diltiazem or verapamil), or digoxin, should be avoided.</li> </ul> <p><u>Recommendations for the use of antiarrhythmic drugs for pharmacologic cardioversion</u></p>

Clinical Guideline	Recommendations
	<p data-bbox="488 205 589 233"><u>of POAF</u></p> <ul style="list-style-type: none"> <li data-bbox="488 237 1377 327">• Restoration of sinus rhythm with pharmacologic cardioversion is reasonable in patients with symptomatic, hemodynamically stable POAF. Intravenous amiodarone can be useful for pharmacologic cardioversion of POAF.</li> <li data-bbox="488 331 1414 390">• It is reasonable to administer antiarrhythmic medications in an attempt to maintain sinus rhythm for patients with recurrent or refractory POAF.</li> <li data-bbox="488 394 1360 485">• Amiodarone, sotalol, flecainide, propafenone, or dofetilide can be useful to maintain sinus rhythm in patients with POAF, depending on underlying heart disease, renal status and other comorbidities.</li> <li data-bbox="488 489 1406 638">• Flecainide or propafenone may be considered for pharmacologic cardioversion of POAF and maintenance of sinus rhythm if the patient has had no prior history of myocardial infarction, coronary artery disease, impaired LV function, significant LV hypertrophy, or valvular heart disease that is considered moderate or greater. These agents may need to be combined with an AV nodal blocking agent.</li> <li data-bbox="488 642 1403 821">• Intravenous ibutilide or procainamide may be considered for pharmacologic conversion of POAF for patients with structural heart disease and new onset POAF, but no hypotension or manifestations of congestive heart failure. Serum electrolytes and QTc interval must be within a normal range and patients must be closely monitored during and for at least six hours after the infusion if either ibutilide or procainamide.</li> <li data-bbox="488 825 1398 884">• Intravenous ibutilide or procainamide may be considered for patients with POAF and an accessory pathway.</li> <li data-bbox="488 888 1382 1010">• Flecainide and propafenone should not be used to treat POAF in patients with a history of a prior myocardial infarction, coronary artery disease, and/or severe structural heart disease, including severe left ventricular hypertrophy, or significantly reduced left ventricular ejection fraction.</li> <li data-bbox="488 1014 1360 1073">• Dronedarone should not be used for treatment of POAF in patients with heart failure.</li> </ul> <p data-bbox="488 1104 1390 1163"><u>Recommendations for prevention of thromboembolism for patients with stable atrial fibrillation/flutter undergoing direct current cardioversion</u></p> <ul style="list-style-type: none"> <li data-bbox="488 1167 1414 1316">• For stable patients with POAF of 48-hours duration or longer, anticoagulation (with warfarin for INR 2.0 to 3.0, a novel oral anti-coagulant [NOAC] or LMWH) is recommended for at least three weeks prior to and four weeks after cardioversion, regardless of the method (electrical or pharmacological) used to restore sinus rhythm.</li> <li data-bbox="488 1320 1390 1442">• During the first 48 hours after the onset of POAF, the need for anticoagulation before and after direct current (DC) cardioversion may be based on the patient's risk of thromboembolism (CHA<sub>2</sub>DS<sub>2</sub>-VASc score) balanced by the risk of postoperative bleeding.</li> <li data-bbox="488 1446 1390 1591">• For POAF lasting longer than 48 hours, as an alternative to three weeks of therapeutic anticoagulation prior to cardioversion of POAF, it is reasonable to perform TEE in search of thrombus in the LA or LA appendage, preferably with full anticoagulation at the time of TEE in anticipation of DC cardioversion after the TEE.</li> <li data-bbox="488 1596 1414 1774">• For POAF lasting longer than 48 hours in patients who are not candidates for TEE (e.g., post-esophageal surgery), an initial rate control strategy combined with therapeutic anticoagulation using warfarin (aiming for INR 2.0 to 3.0), a direct thrombin inhibitor (e.g. dabigatran), factor Xa inhibitor (e.g. rivaroxaban, apixaban), or LMWH is recommended for at least three weeks prior to and four weeks after cardioversion.</li> <li data-bbox="488 1778 1403 1837">• Anticoagulation recommendations for cardioversion of atrial flutter are similar to those for atrial fibrillation.</li> <li data-bbox="488 1841 1390 1900">• For patients with an identified thrombus, cardioversion should not be performed until a longer period of anticoagulation is achieved (usually at least three weeks)</li> </ul>

Clinical Guideline	Recommendations
	<p>and in accordance with established AF guidelines.</p> <p><u>Management of anticoagulation for new onset POAF</u></p> <ul style="list-style-type: none"> <li>• For the prevention of strokes for patients who develop POAF lasting longer than 48 hours, it is recommended to administer antithrombotic medications similarly to non-surgical patients. Anticoagulation within the first 48-hours of POAF should be considered based on the CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of the patient for stroke weighed against the risk of postoperative bleeding.</li> <li>• New oral anticoagulants (dabigatran, rivaroxaban, apixaban) are reasonable as an alternative to warfarin for patients who do not have a prosthetic heart valve, hemodynamically significant valve disease, and/or severe renal impairment or risk of GI bleeding.</li> <li>• It is reasonable to continue anticoagulation therapy for four weeks after the return of sinus rhythm because of the possibility of slowly resolving impairment of atrial contraction with an associated ongoing risk for thrombus formation and for delayed embolic events.</li> <li>• New oral anticoagulants should be avoided for patients at risk for serious bleeding (including GI bleeding) as they cannot be readily reversed. However, their use may be recommended in situations where achievement of a therapeutic INR with warfarin has proved to be difficult.</li> </ul>
<p>American College of Cardiology/ American Heart Association/ European Society of Cardiology Committee for Practice Guidelines: <b>Guidelines for Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death (2017)</b><sup>38</sup></p>	<p><u>Recommendation for Pharmacological Prevention of Sudden Cardiac Death (SCD)</u> <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with heart failure with reduced ejection fraction (HFrEF, ≤40%), treatment with a β-blocker, a mineralocorticoid receptor antagonist and either an angiotensin-converting enzyme (ACE) inhibitor, an angiotensin-receptor blocker (ARB), or an angiotensin receptor-neprilysin inhibitor (ARNI) is recommended to reduce SCD and all-cause mortality.</li> </ul> <p><u>Recommendations for Autonomic Modulation</u> <u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with symptomatic, non-lifethreatening ventricular arrhythmia (VA), treatment with a β-blocker is reasonable.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with ventricular tachycardia (VT)/ ventricular fibrillation (VF) storm in whom a β-blocker, other antiarrhythmic medications, and catheter ablation are ineffective, not tolerated, or not possible, cardiac sympathetic denervation may be reasonable.</li> </ul> <p><u>Recommendation for Management of Cardiac Arrest</u> <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• Cardiopulmonary resuscitation (CPR) should be performed in patients in cardiac arrest according to published basic and advanced cardiovascular life support algorithms.</li> <li>• In patients with hemodynamically unstable VA that persist or recur after a maximal energy shock, intravenous amiodarone should be administered to attempt to achieve a stable rhythm after further defibrillation.</li> <li>• Patients presenting with VA with hemodynamic instability should undergo direct current cardioversion.</li> <li>• In patients with polymorphic VT or VF with ST-elevation myocardial infarction (MI), angiography with emergency revascularization is recommended.</li> <li>• Patients with a wide-QRS tachycardia should be presumed to have VT if the diagnosis is unclear.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with hemodynamically stable VT, administration of intravenous</li> </ul>

Clinical Guideline	Recommendations
	<p>procainamide can be useful to attempt to terminate VT.</p> <ul style="list-style-type: none"> <li>• In patients with a witnessed cardiac arrest due to VF or polymorphic VT that is unresponsive to CPR, defibrillation, and vasopressor therapy, intravenous lidocaine can be beneficial.</li> <li>• In patients with polymorphic VT due to myocardial ischemia, intravenous <math>\beta</math>-blockers can be useful.</li> <li>• In patients with a recent MI who have VT/VF that repeatedly recurs despite direct current cardioversion and antiarrhythmic medications (VT/VF storm), an intravenous <math>\beta</math>-blocker can be useful.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients in cardiac arrest, administration of epinephrine (1 mg every 3 to 5 minutes) during CPR may be reasonable.</li> <li>• In patients with hemodynamically stable VT, administration of intravenous amiodarone or sotalol may be considered to attempt to terminate VT.</li> </ul> <p><u>Class III: No benefit recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with cardiac arrest, administration of high-dose epinephrine (&gt;1 mg boluses) compared with standard doses is not beneficial.</li> <li>• In patients with refractory VF not related to torsades de pointes, administration of intravenous magnesium is not beneficial.</li> </ul> <p><u>Class III: harm recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with suspected acute myocardial infarction (AMI), prophylactic administration of lidocaine or high-dose amiodarone for the prevention of VT is potentially harmful.</li> <li>• In patients with a wide QRS complex tachycardia of unknown origin, calcium channel blockers (e.g., verapamil and diltiazem) are potentially harmful.</li> </ul> <p><u>Recommendation for Secondary Prevention of SCD in Patients with Ischemic Heart Disease</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with ischemic heart disease, who either survive sudden cardiac arrest (SCA) due to VT/VF or experience hemodynamically unstable VT or stable sustained VT not due to reversible causes, an implantable cardioverter-defibrillator (ICD) is recommended if meaningful survival greater than one year is expected.</li> <li>• Value statement: A transvenous ICD provides intermediate value in the secondary prevention of SCD particularly when the patient's risk of death due to a VA is deemed high and the risk of non-arrhythmic death (either cardiac or noncardiac) is deemed low based on the patient's burden of comorbidities and functional status.</li> <li>• In patients with ischemic heart disease and unexplained syncope who have inducible sustained monomorphic VT on electrophysiological study, an ICD is recommended if meaningful survival of greater than one year is expected.</li> </ul> <p><u>Recommendation for Patients with Coronary Artery Spasm</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with VA due to coronary artery spasm, treatment with maximally tolerated doses of a calcium channel blocker and smoking cessation are indicated to reduce recurrent ischemia and VA.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients resuscitated from SCA due to coronary artery spasm in whom medical therapy is ineffective or not tolerated, an ICD is reasonable if meaningful survival of greater than one year is expected.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients resuscitated from SCA due to coronary artery spasm, an ICD in addition to medical therapy may be reasonable if meaningful survival of greater than one year is expected.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendation for Primary Prevention of SCD in Patients with Ischemic Heart Disease</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with LVEF of 35% or less that is due to ischemic heart disease who are at least 40 days' post-MI and at least 90 days post revascularization, and with NYHA class II or III HF despite guideline-directed management and therapy (GDMT), an ICD is recommended if meaningful survival of greater than one year is expected.</li> <li>• In patients with LVEF of 30% or less that is due to ischemic heart disease who are at least 40 days post-MI and at least 90 days post revascularization, and with NYHA class I HF despite GDMT, an ICD is recommended if meaningful survival of greater than one year is expected.</li> <li>• In patients with nonsustained ventricular tachycardia (NSVT) due to prior MI, LVEF of 40% or less and inducible sustained VT or VF at electrophysiological study, an ICD is recommended if meaningful survival of greater than one year is expected.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In non-hospitalized patients with NYHA class IV symptoms who are candidates for cardiac transplantation or an LVAD, an ICD is reasonable if meaningful survival of greater than one year is expected.</li> </ul> <p><u>Class III: no benefit recommendation</u></p> <ul style="list-style-type: none"> <li>• An ICD is not indicated for NYHA class IV patients with medication-refractory HF who are not also candidates for cardiac transplantation, an LV assist device (LVAD), or a cardiac resynchronization therapy (CRT) defibrillator that incorporates both pacing and defibrillation capabilities.</li> </ul> <p><u>Recommendation for Treatment of Recurrent VA in Patients with Ischemic Heart Disease</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with ischemic heart disease and recurrent VA, with significant symptoms or ICD shocks despite optimal device programming and ongoing treatment with a beta blocker, amiodarone or sotalol is useful to suppress recurrent VA.</li> <li>• In patients with prior MI and recurrent episodes of symptomatic sustained VT, or who present with VT storm and have failed or are intolerant of amiodarone or other antiarrhythmic medications, catheter ablation is recommended.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with ischemic heart disease and ICD shocks for sustained monomorphic VT or symptomatic sustained monomorphic VT that is recurrent, or hemodynamically tolerated, catheter ablation as first-line therapy may be considered to reduce recurrent VA.</li> </ul> <p><u>Class III: Harm recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with prior MI, class IC antiarrhythmic medications (e.g., flecainide and propafenone) should not be used.</li> <li>• In patients with incessant VT or VF, an ICD should not be implanted until sufficient control of the VA is achieved to prevent repeated ICD shocks.</li> </ul> <p><u>Class III: No Benefit recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with ischemic heart disease and sustained monomorphic VT, coronary revascularization alone is an ineffective therapy to prevent recurrent VT.</li> </ul> <p><u>Recommendation for Treatment of Recurrent VA in Patients with Nonischemic Cardiomyopathy (NICM)</u></p> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with NICM and an ICD who experience spontaneous VA or recurrent appropriate shocks despite optimal device programming and treatment with a beta</li> </ul>

Clinical Guideline	Recommendations
	<p>blocker, amiodarone or sotalol can be beneficial.</p> <ul style="list-style-type: none"> <li>• In patients with NICM and recurrent sustained monomorphic VT who fail or are intolerant of antiarrhythmic medications, catheter ablation can be useful for reducing recurrent VT and ICD shocks.</li> </ul> <p><u>Recommendation for Arrhythmogenic Right Ventricular Cardiomyopathy</u> <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In selected first-degree relatives of patients with arrhythmogenic right ventricular cardiomyopathy, clinical screening for the disease is recommended along with genetic counseling and genetic testing, if the proband has a disease causing mutation.</li> <li>• In patients with suspected arrhythmogenic right ventricular cardiomyopathy and VA or electrocardiographic abnormalities, cardiac MRI is useful for establishing a diagnosis and for risk stratification.</li> <li>• In patients with arrhythmogenic right ventricular cardiomyopathy and an additional marker of increased risk of SCD (resuscitated SCA, sustained VT, significant ventricular dysfunction with RVEF or LVEF <math>\leq 35\%</math>), an ICD is recommended if meaningful survival greater than one year is expected.</li> <li>• In patients with arrhythmogenic right ventricular cardiomyopathy and VA, a beta blocker is recommended.</li> <li>• In patients with a clinical diagnosis of arrhythmogenic right ventricular cardiomyopathy, avoiding intensive exercise is recommended.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with clinically diagnosed or suspected arrhythmogenic right ventricular cardiomyopathy, genetic counseling and genetic testing can be useful for diagnosis and for gene-specific targeted family screening.</li> <li>• In patients with arrhythmogenic right ventricular cardiomyopathy and syncope presumed due to VA, an ICD can be useful if meaningful survival greater than one year is expected.</li> <li>• In patients with clinical evidence of arrhythmogenic right ventricular cardiomyopathy but not VA, a beta blocker can be useful.</li> <li>• In patients with arrhythmogenic right ventricular cardiomyopathy and recurrent symptomatic sustained VT in whom a beta blocker is ineffective or not tolerated, catheter ablation with availability of a combined endocardial/epicardial approach can be beneficial.</li> <li>• In patients with suspected arrhythmogenic right ventricular cardiomyopathy, a signal averaged ECG can be useful for diagnosis and risk stratification.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>• In asymptomatic patients with clinical evidence of arrhythmogenic right ventricular cardiomyopathy, an electrophysiological study may be considered for risk stratification.</li> </ul> <p><u>Recommendation for Long QT Syndrome</u> <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with long QT syndrome with a resting QTc greater than 470 ms, a beta blocker is recommended.</li> <li>• In high-risk patients with symptomatic long QT syndrome in whom a beta blocker is ineffective or not tolerated, intensification of therapy with additional medications (guided by consideration of the particular long QT syndrome type), left cardiac sympathetic denervation, and/or an ICD is recommended.</li> <li>• In patients with long QT syndrome and recurrent appropriate ICD shocks despite maximum tolerated doses of a beta blocker, intensification of medical therapy with additional medications (guided by consideration of the particular long QT syndrome type) or left cardiac sympathetic denervation, is recommended.</li> <li>• In patients with clinically diagnosed long QT syndrome, genetic counseling and</li> </ul>

Clinical Guideline	Recommendations
	<p>genetic testing are recommended.</p> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with suspected long QT syndrome, ambulatory electrocardiographic monitoring, recording the ECG lying and immediately on standing, and/or exercise treadmill testing can be useful for establishing a diagnosis and monitoring the response to therapy.</li> <li>In asymptomatic patients with long QT syndrome and a resting QTc less than 470 ms, chronic therapy with a beta blocker is reasonable.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>In asymptomatic patients with long QT syndrome and a resting QTc greater than 500 ms while receiving a beta blocker, intensification of therapy with medications (guided by consideration of the particular long QT syndrome type), left cardiac sympathetic denervation or an ICD may be considered.</li> </ul> <p><u>Class III: harm recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with long QT syndrome, QT prolonging medications are potentially harmful.</li> </ul> <p><u>Recommendation for Catecholaminergic Polymorphic Ventricular Tachycardia</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with catecholaminergic polymorphic ventricular tachycardia, a beta blocker is recommended.</li> <li>In patients with catecholaminergic polymorphic ventricular tachycardia and recurrent sustained VT or syncope, while receiving adequate or maximally tolerated beta blocker, treatment intensification with either combination medication therapy (e.g., beta blocker, flecainide), left cardiac sympathetic denervation, and/or an ICD is recommended.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with catecholaminergic polymorphic ventricular tachycardia and with clinical VT or exertional syncope, genetic counseling and genetic testing are reasonable.</li> </ul> <p><u>Recommendation for short QT Syndrome</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with short QT syndrome who have a cardiac arrest or sustained VA, an ICD is recommended if meaningful survival greater than one year is expected.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with short QT syndrome and recurrent sustained VA, treatment with quinidine can be useful.</li> <li>In patients with short QT syndrome and VT/ VF storm, isoproterenol infusion can be effective.</li> </ul> <p><u>Class IIb recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with short QT syndrome, genetic testing may be considered to facilitate screening of first-degree relatives.</li> </ul> <p><u>Recommendation for VA in the Structurally Normal Heart</u></p> <p><u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with symptomatic premature ventricular complexes (PVCs) in an otherwise normal heart, treatment with a beta blocker or nondihydropyridine calcium channel blocker is useful to reduce recurrent arrhythmias and improve symptoms.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>In patients with symptomatic VA in an otherwise normal heart, treatment with an antiarrhythmic medication is reasonable to reduce recurrent symptomatic arrhythmias and improve symptoms if beta blockers and nondihydropyridine calcium channel blockers are ineffective or not tolerated.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendation for Outflow Tract VA</u>  <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with symptomatic outflow tract VA in an otherwise normal heart for whom antiarrhythmic medications are ineffective, not tolerated, or not the patient's preference, catheter ablation is useful.</li> <li>• In patients with symptomatic outflow tract VT in an otherwise normal heart, a beta blocker or a calcium channel blocker is useful.</li> </ul> <p><u>Recommendation for PVC-Induced Cardiomyopathy</u>  <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• For patients who require arrhythmia suppression for symptoms or declining ventricular function suspected to be due to frequent PVCs (generally &gt;15% of beats and predominately of 1 morphology) and for whom antiarrhythmic medications are ineffective, not tolerated, or not the patient's preference, catheter ablation is useful.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with PVC-induced cardiomyopathy, pharmacological treatment (e.g., beta blocker, amiodarone) is reasonable to reduce recurrent arrhythmias and improve symptoms and LV function.</li> </ul> <p><u>Recommendation for Pregnancy</u>  <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• In mothers with long QT syndrome, a beta blocker should be continued during pregnancy and throughout the postpartum period including in women who are breastfeeding.</li> <li>• In the pregnant patient with sustained VA, electrical cardioversion is safe and effective and should be used with standard electrode configuration.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In pregnant patients needing an ICD or VT ablation, it is reasonable to undergo these procedures during pregnancy, preferably after the first trimester.</li> </ul> <p><u>Recommendation for Medication-Induced Arrhythmias</u>  <u>Class I recommendation</u></p> <ul style="list-style-type: none"> <li>• Administration of digoxin antibodies is recommended for patients who present with sustained VA potentially due to digoxin toxicity.</li> <li>• In patients with recurrent torsades de pointes associated with acquired QT prolongation and bradycardia that cannot be suppressed with intravenous magnesium administration, increasing the heart rate with atrial or ventricular pacing or isoproterenol are recommended to suppress the arrhythmia.</li> <li>• For patients with QT prolongation due to a medication, hypokalemia, hypomagnesemia, or other acquired factor and recurrent torsades de pointes, administration of intravenous magnesium sulfate is recommended to suppress the arrhythmia.</li> <li>• For patients with torsades de pointes associated with acquired QT prolongation, potassium repletion to 4.0 mmol/L or more and magnesium repletion to normal values (e.g., <math>\geq 2.0</math> mmol/L) are beneficial.</li> </ul> <p><u>Class IIa recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients taking sodium channel blockers who present with elevated defibrillation or pacing thresholds, discontinuing the presumed responsible medication or reprogramming the device can be useful to restore effective device therapy.</li> </ul> <p><u>Class III: harm recommendation</u></p> <ul style="list-style-type: none"> <li>• In patients with congenital or acquired long QT syndrome, QT-prolonging medications are potentially harmful.</li> </ul>



Clinical Guideline	Recommendations
<p>European Society of Cardiology: <b>Guidelines on diagnosis and management of hypertrophic cardiomyopathy (2014)</b><sup>39</sup></p>	<ul style="list-style-type: none"> <li>• Patients with symptomatic left ventricular outflow tract obstruction should be treated initially with non-vasodilating <math>\beta</math>-blockers titrated to maximum tolerable dose.</li> <li>• If <math>\beta</math>-blockers alone are ineffective, disopyramide titrated to a maximum tolerated dose (usually 400 to 600 mg/day) may be added.</li> <li>• Verapamil can be used when <math>\beta</math>-blockers are contraindicated or ineffective, but close monitoring is required in patients with severe obstruction (<math>\geq 100</math> mmHg) or elevated pulmonary artery systolic pressures, as it can provoke pulmonary edema.</li> <li>• Nifedipine and other dihydropyridine calcium antagonists are not recommended.</li> <li>• Low-dose loop or thiazide diuretics may be used cautiously to improve dyspnea, but it is important to avoid hypovolemia.</li> <li>• In patients without left ventricular outflow tract obstruction, An ACE inhibitor (or ARB if ACE inhibitor not tolerated) should be considered, in addition to a <math>\beta</math>-blocker, for patients who have an LVEF <math>&lt; 50\%</math>, to reduce the risks of HF hospitalization and premature death.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>40</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic blood pressure <math>&lt; 140</math> mm Hg.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 140</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global</b></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or</li> </ul>

Clinical Guideline	Recommendations
<p><b>Hypertension Practice Guidelines (2020)<sup>41</sup></b></p>	<p>limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</p> <ul style="list-style-type: none"> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is two standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP ≥160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (≥80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq 80</math> years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt; 45</math> mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt; 4.5</math> mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul>
<p>Hypertension Canada: <b>2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)</b><sup>42</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq 100</math> mmHg or average systolic blood pressure (SBP) measurements of <math>\geq 160</math> mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq 90</math> mmHg or for average SBP readings <math>\geq 140</math> mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>● Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>● Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>● If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>● Possible reasons for poor response to therapy should be considered.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt; 40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists</li> </ul>

Clinical Guideline	Recommendations
	<p>(mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</p> <ul style="list-style-type: none"> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\geq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>43</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in</li> </ul>

Clinical Guideline	Recommendations
	<p>proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</p> <ul style="list-style-type: none"> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>44</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium</li> </ul>

Clinical Guideline	Recommendations
	<p>level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</p> <ul style="list-style-type: none"> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>45</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group: <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>46</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a</li> </ul>

Clinical Guideline	Recommendations
	<p>target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</p> <ul style="list-style-type: none"> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><b>BP management in kidney transplant recipients</b></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><b>BP management in children with CKD</b></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>47</sup></p>	<p><b>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</b></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) ≥130 mmHg or an average diastolic blood pressure (DBP) of ≥80 mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of ≥10% and an average SBP of ≥130 mmHg or an average ≥80 mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of ≥140 mmHg or a DBP of ≥90 mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of ≥10%, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><b>Stable Ischemic Heart Disease (SIHD)</b></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Adults with SIHD and hypertension (BP <math>\geq</math>130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be <math>&lt;</math>130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP <math>&lt;</math>130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP <math>&lt;</math>130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal <math>&lt;</math>130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq</math>300 mg/d, or <math>\geq</math>300 mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal <math>&lt;</math>130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP <math>&gt;</math>220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to <math>&lt;</math>140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to <math>&lt;</math>185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be <math>&lt;</math>185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP <math>&gt;</math>140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP <math>&lt; 220/120</math> mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal <math>&lt; 130</math> mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP <math>&lt; 140</math> mmHg and a DBP <math>&lt; 90</math> mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal <math>&lt; 130/80</math> mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target <math>&lt; 130/80</math> mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP ≥180 mmHg or DBP ≥110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>48</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (≥140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure ≥180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at</li> </ul>



Clinical Guideline	Recommendations
	<p>home.</p> <ul style="list-style-type: none"> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of <math>&lt;130/80</math> mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;15\%</math>), treat to blood pressure target of <math>&lt;140/90</math>.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;120/80</math>, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events..</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Academy of Family Physicians: <b>Acute Migraine Headache: Treatment Strategies (2018)</b><sup>49</sup></p>	<p><u>General treatment principles</u></p> <ul style="list-style-type: none"> <li>• Evidence-based guidelines from the United States, Canada, and Europe provide consistent recommendations, including lifestyle modifications, avoidance of triggers, and healthy coping mechanisms.</li> <li>• First-line medication options recommended for acute migraines include nonsteroidal anti-inflammatory drugs (for mild to moderate migraines) or triptans (for moderate to severe migraine).</li> <li>• The choice of triptan should be individualized based on the patient's migraine characteristics and on the route of administration, pharmacokinetics, and cost.</li> <li>• Dopamine antagonist antiemetics are second-line treatments for migraine.</li> <li>• Parenteral dihydroergotamine (DHE 45), magnesium sulfate, valproate (Depacon), and opioids should be reserved for refractory migraine because of adverse effects, weaker evidence of effectiveness, and/or abuse potential.</li> </ul>
<p>American Academy of Family Physicians: <b>Medications for Migraine</b></p>	<ul style="list-style-type: none"> <li>• Preventive therapy should be considered in patients having four or more headaches a month or at least eight headache days a month, significantly debilitating attacks despite appropriate acute management, difficulty tolerating or having a contraindication to acute therapy, medication overuse headache, patient</li> </ul>

Clinical Guideline	Recommendations
<p><b>Prophylaxis (2019)<sup>50</sup></b></p>	<p>preference, or the presence of certain migraine subtypes (i.e., hemiplegic migraine; migraine with brainstem aura; migrainous infarction; or frequent, persistent, or uncomfortable aura symptoms).</p> <ul style="list-style-type: none"> <li>• Divalproex (Depakote), topiramate (Topamax), metoprolol, propranolol, and timolol are effective for migraine prevention and should be offered as first-line treatment.</li> <li>• Petasites (an extract of the butterbur plant) has been established as effective and can be considered for migraine prevention.</li> <li>• Behavioral treatments, such as relaxation training, thermal biofeedback combined with relaxation training, electromyographic biofeedback, and cognitive behavior therapy, are effective options for migraine prevention.</li> <li>• Adding acupuncture to symptomatic treatment decreases the frequency of migraine headaches.</li> <li>• For menstrual migraines, starting preventive therapy before the time of expected migraine onset can help prevent disability and reduce severity.</li> <li>• Frovatriptan and naratriptan are recommended for menstrual-associated migraines.</li> </ul>
<p>American Academy of Neurology/ American Headache Society: <b>Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults (2012)<sup>51</sup></b></p>	<ul style="list-style-type: none"> <li>• The following medications are established as effective and should be offered for migraine prevention: <ul style="list-style-type: none"> <li>○ Antiepileptic drugs (AEDs): divalproex sodium, sodium valproate, topiramate</li> <li>○ <math>\beta</math>-Blockers: metoprolol, propranolol, timolol</li> <li>○ Triptans: frovatriptan for short-term menstrually associated migraine prevention</li> </ul> </li> <li>• The following medications are probably effective and should be considered for migraine prevention: <ul style="list-style-type: none"> <li>○ Antidepressants: amitriptyline, venlafaxine</li> <li>○ <math>\beta</math>-Blockers: atenolol, nadolol</li> <li>○ Triptans: naratriptan, zolmitriptan for short-term menstrually associated migraine prevention</li> </ul> </li> </ul>
<p>European Federation of Neurological Societies: <b>Guideline on the Drug Treatment of Migraine - Revised Report of an European Federation of Neurological Societies Task Force (2009)<sup>52</sup></b></p>	<ul style="list-style-type: none"> <li>• Prophylactic drugs for the treatment of migraine with good efficacy and tolerability and evidence of efficacy are <math>\beta</math>-blockers, calcium-channel blockers, antiepileptic drugs, NSAIDs, antidepressants, and miscellaneous drugs.</li> <li>• The use of all these drugs is based on empirical data rather than on proven pathophysiological concepts.</li> <li>• There is no commonly accepted indication for starting a prophylactic treatment. Prophylactic drug treatment of migraine should be considered and discussed with the patient when 1) the quality of life, business duties, or school attendance are severely impaired; 2) frequency of attacks per month is two or higher; 3) migraine attacks do not respond to acute drug treatment; or 4) frequent, very long, or uncomfortable auras occur.</li> <li>• The recommended drugs of first choice are <math>\beta</math>-blockers (metoprolol or propranolol), calcium-channel blockers (flunarizine), and antiepileptic drugs (valproic acid or topiramate).</li> <li>• Drugs of second choice include amitriptyline, venlafaxine, naproxen, and bisoprolol.</li> <li>• Drugs of third choice include acetylsalicylic acid, gabapentin, magnesium, riboflavin, coenzyme Q10, candesartan, lisinopril, and methylsergide.</li> <li>• <math>\beta</math>-blockers are clearly effective in migraine prophylaxis and very well studied. The best evidence has been obtained for metoprolol and propranolol. Bisoprolol, timolol and atenolol might be effective, but evidence is less convincing compared with propranolol and metoprolol.</li> <li>• The calcium-channel blocker, flunarizine, has been shown to be effective in migraine prophylaxis in several studies.</li> <li>• Valproic acid and topiramate are two antiepileptic drugs with evidence of efficacy in more than one placebo-controlled trial. The efficacy rates are comparable to</li> </ul>

Clinical Guideline	Recommendations
	<p>those of metoprolol, propranolol, and flunarizine. Topiramate is also efficacious in the prophylaxis of chronic migraine and may have some effect in migraine with medication overuse.</p>
<p>National Cancer Institute: <b>Pheochromocytoma and Paraganglioma Treatment (PDQ®) (2021)</b><sup>53</sup></p>	<p><b>Preoperative Medical Preparation</b></p> <ul style="list-style-type: none"> <li>• Surgery is the mainstay of treatment for most patients; however, preoperative medical preparation is critical. <math>\alpha</math>-adrenergic blockade should be initiated at the time of diagnosis and maximized preoperatively to prevent potentially life-threatening cardiovascular complications, which can occur as a result of excess catecholamine secretion during surgery. Complications may include hypertensive crisis, arrhythmia, myocardial infarction, and pulmonary edema.</li> <li>• Phenoxybenzamine (a nonselective <math>\alpha</math>-antagonist) is the usual drug of choice; prazosin, terazosin, and doxazosin (selective <math>\alpha</math>-1-antagonists) are alternative choices. Prazosin, terazosin, and doxazosin are shorter acting than phenoxybenzamine, and therefore, the duration of postoperative hypotension is theoretically less than with phenoxybenzamine; however, there is less overall experience with selective <math>\alpha</math>-1-antagonists than with phenoxybenzamine.</li> <li>• A preoperative treatment period of one to three weeks is usually sufficient.</li> <li>• Resolution of spells and a target low normal blood pressure for age indicate that <math>\alpha</math>-adrenergic blockade is adequate.</li> <li>• During <math>\alpha</math>-adrenergic blockade, liberal salt and fluid intake should be encouraged because volume loading reduces excessive orthostatic hypotension both preoperatively and postoperatively.</li> <li>• If tachycardia develops or if blood pressure control is not optimal with <math>\alpha</math>-adrenergic blockade, a <math>\beta</math>-blocker (e.g., metoprolol or propranolol) can be added, but only after <math>\alpha</math>-blockade.</li> <li>• A <math>\beta</math>-adrenergic blockade must never be initiated before <math>\alpha</math>-blockade; doing so blocks <math>\beta</math>-blocker mediated vasodilation and results in unopposed <math>\alpha</math>-blocker receptor mediated vasoconstriction, which can lead to a life-threatening crisis.</li> </ul> <p><b>Localized Pheochromocytoma Treatment</b></p> <ul style="list-style-type: none"> <li>• The standard treatment option for patients with localized pheochromocytoma is surgery.</li> <li>• Intraoperative hypertension can be controlled with intravenous infusion of phentolamine, sodium nitroprusside, or a short-acting calcium-channel blocker (e.g., nicardipine).</li> <li>• Tumor removal may be followed by a sudden drop in blood pressure that may require rapid volume replacement and intravenous vasoconstrictors (e.g., norepinephrine or phenylephrine).</li> <li>• Postoperatively, patients should remain in a monitored environment for 24 hours.</li> <li>• Postoperative hypotension is managed primarily by volume expansion, and postoperative hypertension usually responds to diuretics.</li> </ul> <p><b>Pheochromocytoma During Pregnancy</b></p> <ul style="list-style-type: none"> <li>• Phenoxybenzamine use is safe in pregnancy, but beta-adrenergic blockers should be initiated only if needed because their use has been associated with intrauterine growth restriction.</li> </ul>
<p>American Academy of Neurology: <b>Practice Parameter: Therapies for Essential Tremor: Report of the Quality Standards Subcommittee of the American Academy</b></p>	<ul style="list-style-type: none"> <li>• Propranolol and primidone are agents that are most commonly used to treat essential tremor (ET).</li> <li>• It is recommended that propranolol, long-acting propranolol, or primidone be offered to patients who want treatment for limb tremor in ET, depending on concurrent medical conditions and potential side effects.</li> <li>• It is recommended that either primidone or propranolol be used as initial therapy to treat limb tremor in ET.</li> <li>• It is recommended that atenolol and sotalol be considered for treatment of limb tremor associated with ET, and propranolol may be considered as a treatment</li> </ul>

Clinical Guideline	Recommendations
<p><b>of Neurology (2005)<sup>54</sup>, Evidence-based guideline update: Treatment of essential tremor (2011 update)<sup>55</sup></b></p> <p><b>Reaffirmed April 2017</b></p>	<p>option for head tremor in patients with ET.</p> <ul style="list-style-type: none"> <li>• Nadolol may be considered a treatment option for limb tremor associated with ET.</li> <li>• Pindolol is not recommended for treatment of limb tremor in ET.</li> <li>• Due to the lack of evidence, a recommendation regarding the use of metoprolol in the treatment of limb tremor in ET cannot be provided.</li> <li>• The combination of primidone and propranolol may be used to treat limb tremor when the use of a single agent does not adequately decrease tremor.</li> <li>• The dosages of propranolol and primidone may need to be increased after 12 months of therapy when treating limb tremor in ET.</li> <li>• Levetiracetam and 3,4-diaminopyridine should not be considered for treatment of limb tremor in ET.</li> <li>• Clinicians may choose not to consider flunarizine for treatment of limb tremor in ET.</li> <li>• The evidence is insufficient to make recommendations regarding the use of pregabalin, zonisamide, or clozapine</li> </ul>
<p>American Academy of Pediatrics: <b>Clinical Practice Guideline for the Management of Infantile Hemangiomas (2019)<sup>56</sup></b></p>	<p><u>Pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Clinicians should use oral propranolol as the first-line agent for Infantile Hemangiomas (IHs) requiring systemic treatment (grade A, strong recommendation)</li> <li>• Clinicians should dose propranolol between 2 and 3 mg/kg per day unless there are comorbidities (e.g., PHACE syndrome) or adverse effects (e.g., sleep disturbance) that necessitate a lower dose (grade A, moderate recommendation).</li> <li>• Clinicians should counsel that propranolol be administered with or after feeding and that doses be held at times of diminished oral intake or vomiting to reduce the risk of hypoglycemia (grade X, strong recommendation)</li> <li>• Clinicians should evaluate patients for and educate caregivers about potential adverse effects of propranolol, including sleep disturbances, bronchial irritation, and clinically symptomatic bradycardia and hypotension (grade X, strong recommendation)</li> <li>• Clinicians may prescribe oral prednisolone or prednisone to treat IHs if there are contraindications or an inadequate response to oral propranolol (grade B, moderate recommendation).</li> <li>• Clinicians may recommend intralesional injection of triamcinolone and/or betamethasone to treat focal, bulky IHs during proliferation or in certain critical anatomic locations (e.g., the lip) (grade B, moderate recommendation)</li> <li>• Clinicians may prescribe topical timolol maleate as a therapy for thin and/or superficial IHs (grade B, moderate recommendation).</li> <li>• Clinicians may recommend surgery and laser therapy as treatment options in managing selected IHs (grade C, moderate recommendation)</li> <li>• Clinicians should educate parents of infants with an IH about the condition, including the expected natural history, and its potential for causing complications or disfigurement (grade X, strong recommendation).Clinicians should educate parents of infants with an IH about the condition, including the expected natural history, and its potential for causing complications or disfigurement (grade X, strong recommendation).</li> </ul>

\*Agent not available in the United States.

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the  $\beta$ -adrenergic blocking agents are noted in Tables 4 and 5. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 4. FDA-Approved Indications for the Beta-Adrenergic Blocking Agents<sup>3-22</sup>**

Indications	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
<b>Angina Pectoris</b>									
Long-term management of angina pectoris		✓ *					✓ †	✓	
<b>Cardiac Arrhythmias</b>									
Management of ventricular premature beats	✓								
<b>Heart Failure</b>									
Mild to severe chronic heart failure of ischemic or cardiomyopathic origin to increase survival and, also, to reduce the risk of hospitalizations					✓				
Stable, symptomatic (NYHA Class II or III) heart failure of ischemic, hypertensive, or cardiomyopathic origin							✓ (succinate)		
<b>Hypertension</b>									
Control of blood pressure in severe hypertension						✓ (injection)			
Hypertension	✓ ‡	✓ ‡	✓ ‡	✓ ‡	✓ ‡	✓ ‡ (tablet)	✓ ‡	✓ ‡	✓ ‡
<b>Myocardial Infarction</b>									
Hemodynamically stable patients with definite or suspected acute myocardial infarction to reduce cardiovascular mortality		✓					✓ (tartrate)		
Reduce cardiovascular mortality in clinically stable patients who have survived the acute phase of a myocardial infarction and have a left ventricular ejection fraction of $\leq 40\%$ (with or without symptomatic heart failure)					✓				

\*Due to coronary atherosclerosis.

†Metoprolol succinate: To reduce angina attacks and to improve exercise tolerance.

‡May be used in combination with other antihypertensive agents.

NYHA=New York Heart Association

**Table 5. FDA-Approved Indications for the Beta-Adrenergic Blocking Agents (continued)<sup>3-22</sup>**

Indications	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendroflu-methiazide	
<b>Angina Pectoris</b>										
Angina pectoris			✓ * (Inderal LA <sup>®</sup> , tablet)							
<b>Cardiac Arrhythmias</b>										
Abolish tachyarrhythmias due to			✓							

Indications	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendroflu-methiazide	
excessive catecholamine action during anesthesia when other measure fail			(injection)							
Control ventricular rate in patients with atrial fibrillation and a rapid ventricular response			✓ (tablet)							
Control ventricular rate in life-threatening digitalis-induced arrhythmias			✓ (injection)							
Documented ventricular arrhythmias, such as sustained ventricular tachycardia, that in the judgement of the physician are life-threatening				✓ † (Betapace®, Sotylize®)						
Maintenance of normal sinus rhythm in patients with symptomatic atrial fibrillation/atrial flutter who are currently in sinus rhythm				✓ † (Betapace AF®, Sotylize®)						
Persistent premature ventricular extrasystoles that impair the well-being of the patient and do not respond to conventional measures			✓ (injection)							
Short-term treatment of supraventricular tachycardia, including Wolff-Parkinson-White syndrome and thyrotoxicosis, to decrease ventricular rate			✓ (injection)							
<b>Hypertension</b>										
Hypertension		✓ ‡	✓ ‡ (oral§)		‡	✓	✓	✓ ¶	✓	
Mild to moderate arterial hypertension	✓ ‡									
<b>Hypertrophic Subaortic Stenosis</b>										
Improves NYHA functional class in symptomatic patients with hypertrophic subaortic stenosis			✓ (Inderal LA®, tablet)							
<b>Myocardial Infarction</b>										
Reduce cardiovascular mortality in patients who have survived the acute phase of myocardial infarction and are clinically stable			✓ (tablet)							
Reduce cardiovascular mortality and reinfarction in patients who have survived the acute phase of myocardial infarction and are clinically stable					✓					
<b>Other</b>										
Adjunct to $\alpha$ -adrenergic blockade to control blood pressure and reduce			✓ (tablet)							

Indications	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendroflu-methiazide	
symptoms of catecholamine-secreting tumors										
Familial or hereditary essential tremor			✓ (tablet)							
Treatment of proliferating infantile hemangioma requiring systemic therapy			✓ (Hemangeol®)							
Prophylaxis of migraine headache			✓ (Inderal LA®, tablet)		✓					

\*Angina pectoris due to coronary atherosclerosis to decrease angina frequency and increase exercise tolerance.

†Intravenous sotalol can substitute for oral sotalol in patients who are unable to take sotalol orally.

‡May be used in combination with other antihypertensive agents.

§Inderal LA® and propranolol tablet are not indicated in the management of hypertensive emergencies.

|| Not indicated for initial treatment of hypertension.

¶Dutoprol® may be used in combination with other antihypertensive agents. Lopressor HCT® is not indicated for initial treatment of hypertension.

HCTZ=hydrochlorothiazide, NYHA=New York Heart Association



#### IV. Pharmacokinetics

The pharmacokinetic parameters of the  $\beta$ -adrenergic blocking agents are listed in Table 6. The lipophilic properties vary among the agents. The higher the lipid solubility, the higher the potential to cross the blood brain barrier and increase the risk of central nervous system adverse events, including dizziness and drowsiness.<sup>32,34</sup>

**Table 6. Pharmacokinetic Parameters of the Beta-Adrenergic Blocking Agents<sup>2</sup>**

Generic Name(s)	Bio-availability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)	Lipid Solubility
<b>Single Entity Agents</b>						
Acebutolol	40	10 to 26	Liver (% not reported)	Renal (30 to 40) Bile (3 to 8) Feces (56)	3 to 4	Low
Atenolol	46 to 60	6 to 16	Not reported	Renal (40 to 50) Feces (50)	6 to 7	Low
Betaxolol	78 to 90	50 to 60	Liver, extensive (% not reported)	Renal (>80)	14 to 22	Low
Bisoprolol	80 to 94	30 to 36	Liver (50)	Renal (50) Feces (<2)	9 to 12	Low
Carvedilol	21 to 35	95 to 98	Liver, extensive (% not reported)	Renal (16) Feces (60)	7 to 10	Moderate
Labetalol	25	50	Liver, extensive (% not reported)	Renal (55 to 60) Feces (50)	5 to 8	Moderate
Metoprolol	50 to 77	12	Liver, extensive (% not reported)	Renal (95)	3 to 7	Moderate
Nadolol	20 to 40	28 to 30	None	Renal (25) Feces (77)	20 to 24	Low
Nebivolol	12 to 96	98	Liver, extensive (% not reported)	Renal (<1) Feces (13 to 44)	12 to 19	High
Penbutolol	100	80 to 98	Liver, extensive (% not reported)	Renal (90)	5	High
Pindolol	95	40 to 60	Liver (60 to 65)	Renal (35 to 40) Feces (6 to 9)	3 to 4	Moderate
Propranolol	30 to 70	93	Liver (50 to 70)	Renal (<1)	3 to 6	High
Sotalol	90 to 100	0	Liver, minor	Renal (66 to 88)	7 to 12	Low
Timolol	61	<10	Liver (80)	Renal (20)	2 to 4	Low-Moderate
<b>Combination Products</b>						
Atenolol and chlorthalidone	50/65	16/75	Not reported/ Liver (% not reported)	Renal (40 to 50) Feces (50)/ Renal (60)	6 to 7/ 40 to 60	Low/not reported
Bisoprolol and HCTZ	80/ 50 to 75	30/ 40 to 68	Liver (50)/ not reported	Renal (50) Feces (<2)/ Renal (>95)	9 to 12/ 6 to 15	Low/not reported
Metoprolol and HCTZ	Not reported	12/68	Liver, extensive (% not reported)/ not reported	Renal (95)/ Renal (72 to 97)	3 to 7/ 10 to 17	Moderate/ not reported
Nadolol and bendro-flumethiazide	30	Not reported	Not Reported	Not reported	20 to 24/ 3	Not reported

HCTZ=hydrochlorothiazide

## V. Drug Interactions

Major drug interactions with the beta-adrenergic blocking agents ( $\beta$ -blockers) are listed in Table 7.

**Table 7. Major Drug Interactions with the Beta-Adrenergic Blocking Agents<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
$\beta$ -blockers (acebutolol, atenolol, betaxolol, bisoprolol, carvedilol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Verapamil	May be synergistic or additive effects. Verapamil may inhibit oxidative metabolism of certain $\beta$ -blockers. Additive QT interval prolongation is possible with sotalol.
$\beta$ -blockers (nadolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Epinephrine	Nonselective $\beta$ blockade allows $\alpha$ -receptor effects of epinephrine to predominate. Increasing vascular resistance leads to a rise in blood pressure and reflex bradycardia.
$\beta$ -blockers (nadolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Sympathomimetics	Nonselective $\beta$ -blockers may block the action of beta-agonists, potentially resulting in severe bronchospasm in asthmatics.
Thiazides (hydrochlorothiazide, chlorthalidone, bendroflumethiazide)	Lithium	Decreased lithium clearance may occur with thiazide use. This may lead to increased serum lithium levels and possibly lithium toxicity. Monitor plasma lithium levels and symptoms of toxicity, and adjust the dose as needed.
Thiazides (hydrochlorothiazide, bendroflumethiazide)	Dofetilide	Thiazide diuretics may induce hypokalemia which may increase the risk of torsades de pointes. The coadministration of dofetilide with a thiazide diuretic is contraindicated.
$\beta$ -blockers (sotalol)	Bepiridil	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.
$\beta$ -blockers (sotalol)	Chloroquine	Prolonged QT interval and cardiac arrhythmias are a potential when sotalol and chloroquine are coadministered.
$\beta$ -blockers (sotalol)	Class IA, IC, III Antiarrhythmic Agents	Class IA, IC, and III antiarrhythmics and sotalol may cause additive pharmacologic and adverse cardiovascular effects when co- administered.
$\beta$ -blockers (sotalol)	Dofetilide	The risk of cardiovascular toxicity, including torsades de pointes, may be increased by co-administration of dofetilide and sotalol. Pharmacologic effects of dofetilide and sotalol on electrical conduction of the heart may be additive.
$\beta$ -blockers (sotalol)	Dronedarone	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.
$\beta$ -blockers (sotalol)	Droperidol	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.
$\beta$ -blockers (sotalol)	Fluconazole	Coadministration of fluconazole and sotalol may increase the risk of potentially fatal cardiac arrhythmias (torsades de pointes), especially in seriously ill patients and/or patients receiving high dose fluconazole.
$\beta$ -blockers (sotalol)	Haloperidol	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.

Generic Name(s)	Interaction	Mechanism
β-blockers (sotalol)	Maprotiline	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.
β-blockers (sotalol)	Methadone	Prolongation of the QT interval with possible development of cardiac arrhythmias, including torsades de pointes, should be considered when sotalol is co-administered with methadone.
β-blockers (sotalol)	Nilotinib	Additive QT prolongation may occur during coadministration of nilotinib and sotalol.
β-blockers (sotalol)	Pentamidine	Prolongation of the QT interval with possible development of cardiac arrhythmias, including torsades de pointes, should be considered when sotalol is co-administered with pentamidine.
β-blockers (sotalol)	Perflutren	Additive QT interval prolongation may occur during coadministration of perflutren and sotalol.
β-blockers (sotalol)	Phenothiazines	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility when sotalol and phenothiazines are co-administered.
β-blockers (sotalol)	Phosphodiesterase type 5 Inhibitors	Phosphodiesterase type 5 inhibitors and sotalol may cause additive adverse effects when co-administered. Prolonged QT interval with the potential for cardiac arrhythmias may occur.
β-blockers (sotalol)	Pimozide	Sotalol and pimozide may cause additive adverse effects when co-administered. Cardiovascular toxicity, including torsades de pointes, may occur due to additive QT-interval prolongation.
β-blockers (sotalol)	Quinolones	The rare occurrence of arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility.
β-blockers (sotalol)	Serotonin Receptor Antagonists Antiemetics	The risk of QT-interval prolongation and cardiac arrhythmias caused by serotonin receptor antagonist antiemetics may be increased by co-administration of sotalol.
β-blockers (sotalol)	Tetrabenazine	Additive QT prolongation may occur during coadministration of tetrabenazine and sotalol.
β-blockers (sotalol)	Tyrosine Kinase Receptor Inhibitor	Additive QT interval prolongation is a possibility when tyrosine kinase receptor inhibitors are coadministered with sotalol.
β-blockers (sotalol)	Ziprasidone	Arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility when sotalol and ziprasidone are co-administered.
β-blockers (acebutolol, atenolol, betaxolol, bisoprolol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Clonidine	B-blocker inhibition of β <sub>2</sub> receptor mediated vasodilation leaves peripheral α <sub>2</sub> -receptor mediated vasoconstriction unopposed to clonidine stimulation.
β-blockers (acebutolol, atenolol, betaxolol, carvedilol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Diltiazem	Additive AV nodal blockade may lead to synergistic bradycardia
β-blockers (acebutolol, atenolol, betaxolol, bisoprolol, carvedilol,	Flecainide	Unknown mechanism. Combination may result in additive bradycardia and cardiac arrest

Generic Name(s)	Interaction	Mechanism
metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, timolol)		
β-blockers (acebutolol, atenolol, betaxolol, carvedilol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Nonsteroidal Anti-inflammatory Drugs	NSAIDs may inhibit renal prostaglandin synthesis, allowing unopposed pressor systems to produce hypertension.
β-blockers (acebutolol, atenolol, betaxolol, carvedilol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Quinazolines	Unknown mechanism. Additive vasodilation may increase risk of hypotension, specifically orthostatic hypotension. Generally occurs with the addition of prazosin to chronic β-blocker therapy, not β-blocker added to chronic prazosin therapy
β-blockers (bisoprolol, carvedilol, nadolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Insulin	β-blockers blunt sympathetic mediated responses to hypoglycemia.
β-blockers (atenolol, carvedilol, metoprolol, nadolol, pindolol, propranolol, sotalol)	Lidocaine	Reduced hepatic lidocaine metabolism and possibly a minor component of diminished hepatic blood flow.
β-blockers (bisoprolol, carvedilol, metoprolol, pindolol, propranolol, timolol)	Cimetidine	Cimetidine may reduce hepatic first-pass extraction, decrease liver blood flow, and inhibit hepatic metabolism of β-blockers.
β-blockers (nadolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Meglitinides	Unknown mechanism. Possible increase in hypoglycemic activity of meglitinides.
β-blockers (nadolol, penbutolol, pindolol, propranolol, sotalol, timolol)	Theophyllines	Pharmacologic antagonism. B-blockers may reduce the n-demethylation of theophylline.
β-blockers (atenolol, carvedilol, metoprolol, propranolol, timolol)	Quinidine	Oxidative metabolism of certain β-blockers may be inhibited by quinidine.
β-blockers (carvedilol, metoprolol, nebivolol, propranolol, timolol)	Terbinafine	Terbinafine inhibits CYP2D6 and may result in increased plasma concentrations of certain β-blockers.
β-blockers (carvedilol, metoprolol, propranolol, timolol)	Diphenhydramine	Inhibition of CYP2D6-mediated β-blocker metabolism may decrease the metabolism of certain β-blockers resulting in excessive cardiovascular effects.
β-blockers (metoprolol, nebivolol, propranolol, timolol)	Serotonin Reuptake Inhibitors	Inhibition of CYP2D6 enzyme may decrease the metabolism of metoprolol resulting in excessive pharmacologic activity.
β-blockers (metoprolol, propranolol, sotalol)	Amiodarone	Additive pharmacologic effects of both drugs may result in severe bradycardia, hypotension, or cardiac arrest. Possible additive QT interval prolongation with sotalol and amiodarone.
β-blockers (pindolol, propranolol, sotalol)	Phenothiazines	Chlorpromazine may inhibit the first-pass hepatic metabolism of propranolol and increase its pharmacologic effects. Certain β-blockers may inhibit the metabolism of phenothiazines increasing the risk for cardiac side effects, including torsades

Generic Name(s)	Interaction	Mechanism
		de pointes.
$\beta$ -blockers (carvedilol, metoprolol, propranolol)	Rifamycins (rifabutin, rifampin, rifapentine)	Possible decrease in oral bioavailability of carvedilol resulting in first-pass metabolism.
$\beta$ -blockers (carvedilol, metoprolol, propranolol)	Thiamines	Hyperthyroidism appears to cause increased clearance of $\beta$ -blockers with a high extraction ration. This may be the result of increased liver blood flow, first-pass metabolism and volume of distribution.
Thiazide diuretics (HCTZ, chlorthalidone, bendroflumethiazide)	Diazoxide	The combination of diazoxide with a thiazide diuretic may lead to hyperglycemia though an unknown mechanism; therefore the combination should be avoided. When used together, blood and urine glucose levels should be frequently monitored, and dosage reductions may be required.
Thiazide diuretics (HCTZ, chlorthalidone, bendroflumethiazide)	Digitalis glycosides	Thiazide diuretics may induce electrolyte disturbances which may predispose patients to digitalis-induced arrhythmias. Measure plasma levels of potassium and magnesium, supplement low levels, and use dietary sodium restriction or potassium-sparing diuretics to prevent further losses.
$\beta$ -blockers (metoprolol, propranolol)	Hydralazine	Hydralazine increases systemic availability of some $\beta$ -blockers, probably by transient increase in splanchnic blood flow and decreasing first-pass hepatic metabolism.
$\beta$ -blockers (metoprolol, propranolol)	Propafenone	Propafenone increases plasma $\beta$ -blocker level by decreasing first-pass metabolism and reducing systemic clearance. Both drugs are oxidized by the hepatic CYP450 system, and propafenone appears to inhibit the metabolism of the $\beta$ -blocker.
$\beta$ -blockers (atenolol)	Ampicillin	The bioavailability of atenolol may be decreased by impaired gastrointestinal absorption induced by ampicillin.
$\beta$ -blockers (carvedilol)	Cyclosporine	Unknown mechanism. Carvedilol may increase plasma concentrations of cyclosporine and dose reduction may be required.
$\beta$ -blockers (carvedilol)	Digoxin	Carvedilol may increase digoxin bioavailability. Possible additive depression of myocardial conduction and decreased renal tubular digoxin secretion.
$\beta$ -blockers (labetalol)	Inhalation anesthetics	Additive myocardial depressant effects possibly resulting in excessive hypotension.
$\beta$ -blockers (propranolol)	Mefloquine	Additive slowing of cardiac conduction possibly resulting in lengthening of the QT interval
$\beta$ -blockers (propranolol)	Triptans	Unknown mechanism. Possible inhibition of triptan metabolism (monoamine oxidase-A) by propranolol resulting in enhanced pharmacologic effects and plasma concentrations.
$\beta$ -blockers (sotalol)	Cisapride	Prolongation of the QT interval with possible development of cardiac arrhythmias, including torsades de pointes, should be considered when cisapride is co-administered with sotalol.
$\beta$ -blockers (sotalol)	H1 Antagonists	The rare occurrence of arrhythmias resulting from the potential for additive QT prolongation should

Generic Name(s)	Interaction	Mechanism
		be considered as a possibility when sotalol and H-1 antagonists are coadministered.
β-blockers (sotalol)	Iloperidone	Prolonged QT interval and cardiac arrhythmias are a potential when sotalol and iloperidone are used concomitantly.
β-blockers (sotalol)	Macrolides	The rare occurrence of arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility when sotalol and macrolides are coadministered.
β-blockers (sotalol)	Mefloquine	Co-administration of mefloquine and sotalol may cause cardiovascular toxicity, including electrocardiographic abnormalities such as QT interval prolongation
β-blockers (sotalol)	Mibefradil	Co-administration of sotalol and mibefradil may cause cardiovascular toxicity.
β-blockers (sotalol)	Paliperidone	Prolongation of the QT interval with possible development of cardiac arrhythmias, including torsades de pointes, should be considered when paliperidone is co-administered with sotalol.
β-blockers (sotalol)	Propafenone	The rare occurrence of arrhythmias resulting from the potential for additive QT prolongation should be considered when sotalol and propafenone are coadministered.
β-blockers (sotalol)	Saquinavir	Coadministration of sotalol with saquinavir/ritonavir may be associated arrhythmias due to potential additive effects on prolongation of the QT interval.
β-blockers (sotalol)	Tricyclic Antidepressants	The rare occurrence of arrhythmias resulting from the potential for additive QT prolongation should be considered as a possibility when tricyclic antidepressants and sotalol are coadministered.
β-blockers (acebutolol)	Ceritinib	Bradycardia causing agents may enhance the bradycardic effect of ceritinib.
β-blockers (acebutolol, atenolol, betaxolol, bisoprolol, carvedilol, labetalol, metoprolol, nadolol, nebivolol, pindolol, propranolol, timolol)	Rivastigmine	Concurrent use may result in additive bradycardic effects.
β-blockers (carvedilol, labetalol, nadolol, pindolol, propranolol, timolol)	Beta2-agonists (non-selective)	Nonselective beta-blockers may diminish the bronchodilatory effect of beta2-agonists
β-blockers (carvedilol)	Topotecan	P-glycoprotein/ABCB1 inhibitors may increase the serum concentration of topotecan
β-blockers (carvedilol)	Vincristine	P-glycoprotein/ABCB1 inhibitors may increase the serum concentration of vincristine
β-blockers (pindolol, propranolol, sotalol)	Thioridazine	Concurrent use may result in increased risk of thioridazine toxicity (e.g., QT prolongation, torsades de pointes, cardiac arrest)
β-blockers (sotalol)	Fingolimod	Concurrent use may result in increased risk of QT interval prolongation, bradycardia, or heart block

CYP=cytochrome P450 isoenzymes, HCTZ=hydrochlorothiazide

## VI. Adverse Drug Events

The most common adverse drug events reported with the  $\beta$ -adrenergic blocking agents are listed in Tables 8 and 9. The boxed warnings for the  $\beta$ -adrenergic blocking agents are listed in Tables 10 through 15.

**Table 8. Adverse Drug Events (%) Reported with the Beta-Adrenergic Blocking Agents<sup>1-22</sup>**

Adverse Events	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
<b>Cardiovascular</b>									
Angina	-	-	<2	-	1 to 6	-	-	-	-
Arrhythmia	-	-	<2	<1	-	-	-	<1	-
Arterial/vascular insufficiency	-	-	-	-	-	-	1	-	<1
Bradycardia	1 to 10	18	6 to 8	<1	2 to 10	<1	2 to 16	1 to 10	≤1
Cardiogenic shock	-	-	-	-	-	-	✓	-	-
Cerebrovascular accident	-	-	-	-	≤4	-	-	-	-
Chest pain	2	1 to 10	2 to 7	1 to 2	-	-	1	<1	≤1
Cold extremities	-	12	2	<1	-	-	1	1 to 10	-
Congestive heart failure	1 to 10	1 to 10	<2	<1	-	<1	1	1 to 10	-
Edema	2	1 to 10	≤2	<1	5 to 6	≤2	-	1 to 10	-
Flushing	-	-	-	<1	-	1	-	-	-
Heart block	✓	1 to 10	<2	-	≤4	<1	5	-	-
Hypertension	-	-	<2	-	≤4	-	-	-	-
Hypotension	2	25	<2	<1	9 to 20	1 to 5	1 to 27	-	-
Myocardial ischemia	-	-	-	-	-	-	-	-	<1
Orthostatic hypotension	-	1 to 10	-	<1	-	-	-	<1	-
Palpitations	✓	-	2	<1	≤4	-	1	1 to 10	-
Peripheral circulation reduced	-	-	-	-	<1	-	-	1 to 10	-
Peripheral edema	-	-	-	-	1 to 7	-	1	-	1
Postural hypotension	-	-	-	-	≤4	-	-	-	-
Rhythm disturbance	-	-	-	<1	-	-	-	-	-
Shortness of breath	-	-	-	-	-	-	✓	-	-
Syncope	-	-	<2	<1	3 to 8	<1	1	-	<1
Ventricular arrhythmias	✓	-	-	-	-	-	-	-	-
<b>Central Nervous System</b>									
Abnormal dreams	2	1 to 10	<1	-	-	-	-	-	-
Anxiety	1 to 10	-	-	<1	-	-	✓	-	-
Concentration decreased	-	-	-	-	<1	-	-	-	-
Confusion	-	1 to 10	-	<1	-	-	✓	<1	-
Depression	2	1 to 10	<1	<1	1 to 10	-	5	1 to 10	-
Diaphoresis	-	-	<2	-	<1	-	-	-	-
Dizziness	6	1 to 10	-	<1	2 to 32	1 to 20	2 to 10	-	2 to 4
Drowsiness	-	1 to 10	-	-	-	-	-	>10	-
Fatigue	11	1 to 10	3 to 10	6 to 8	4 to 24	1 to 11	1 to 10	-	-
Fever	-	-	<2	-	1 to 10	-	-	-	-
Hallucinations	-	<1	<2	<1	-	-	✓	<1	2 to 5
Headache	6	1 to 10	-	<1	5 to 8	2	✓	<1	-

Adverse Events	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
Hyper/hypoesthesia	1 to 10	-	-	1 to 2	1 to 10	-	-	-	-
Insomnia	3	1 to 10	1 to 5	2 to 3	1 to 10	-	✓	>10	6 to 9
Lethargy	-	1 to 10	3	-	-	-	-	-	1
Malaise	-	-	<2	<1	1 to 10	-	-	-	-
Memory loss	-	-	<2	<1	<1	-	✓	-	-
Mental impairment	-	1 to 10	-	-	-	-	-	-	-
Nervousness	-	-	-	<1	<1	-	✓	<1	-
Nightmares/vivid dreams	-	1 to 10	-	-	<1	-	✓	-	-
Paresthesia	-	-	-	<1	-	-	✓	-	-
Psychosis	-	<1	-	-	-	-	-	-	-
Sleep disturbance	-	-	-	<1	-	-	✓	-	-
Somnolence	-	-	-	<1	1 to 10	3	✓	-	-
Vertigo	-	-	-	<1	1 to 10	1 to 2	✓	-	<1
<b>Dermatologic</b>									
Acne	-	-	-	<1	-	-	-	-	-
Alopecia	-	<1	<2	<1	<1	<1	✓	-	-
Dermatitis	-	-	-	<1	-	-	-	-	✓
Eczema	-	-	-	<1	-	-	-	-	-
Erythema multiforme	-	-	-	-	<1	-	-	-	-
Exfoliative dermatitis	-	-	-	-	<1	-	-	-	-
Photosensitivity	-	-	-	-	<1	-	✓	-	-
Pruritus	1 to 10	-	-	<1	<1	1	5	-	<1
Psoriasisiform rash	-	<1	-	<1	-	<1	-	-	-
Psoriasis (exacerbated)	-	-	-	<1	-	-	✓	-	<1
Purpura	-	-	-	<1	-	-	-	-	-
Rash	2	-	1	<1	<1	1	5	-	≤1
Scalp tingling	-	-	-	-	-	≤7	-	-	-
Stevens-Johnson syndrome	-	-	-	-	<1	-	-	-	-
Sweating, excessive	-	-	-	-	-	-	✓	-	-
Systemic lupus erythematosus	✓	-	-	-	-	-	-	-	-
Toxic epidermal necrolysis	-	-	-	-	<1	-	-	-	-
Urticaria	-	-	-	-	-	<1	✓	-	<1
<b>Endocrine and Metabolic</b>									
Diabetes (exacerbated)	-	-	<2	-	1 to 10	-	✓	-	-
Gout	-	-	-	<1	1 to 10	-	-	-	-
Libido decreased	-	-	-	-	-	-	✓	-	-
<b>Gastrointestinal</b>									
Abdominal pain	1 to 10	-	-	<1	1 to 10	-	✓	-	1 to 10
Anorexia	✓	-	<2	-	-	-	-	-	-
Constipation	4	1 to 10	<2	<1	-	-	1	1 to 10	-
Cramping	-	-	-	-	-	-	-	-	-
Diarrhea	4	1 to 10	2	3 to 4	-	-	5	1 to 10	2 to 3
Dyspepsia	4	-	4 to 5	<1	-	≤4	-	-	-
Epigastric distress	-	-	-	-	-	-	-	-	-
Flatulence	3	-	-	-	-	-	1	-	-
Gastritis/gastric irritation	-	-	-	<1	-	-	-	-	-



Adverse Events	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
Gastrointestinal hemorrhage	-	-	-	-	<1	-	-	-	-
Heartburn	-	-	-	-	-	-	1	-	-
Melena	-	-	-	-	1 to 10	-	-	-	-
Nausea	4	1 to 10	2 to 6	2	2 to 9	≤19	1	1 to 10	1 to 3
Pancreatitis	-	-	-	-	<1	-	-	-	-
Peptic ulcer	-	-	-	<1	-	-	-	-	-
Periodontitis	-	-	-	-	1 to 10	-	-	-	-
Retroperitoneal fibrosis	-	-	-	-	-	-	✓	-	-
Stomach discomfort	-	-	-	-	-	-	-	1 to 10	-
Taste disorder	-	-	<2	<1	-	1	✓	-	-
Vomiting	1 to 10	-	<2	1 to 2	1 to 6	≤3	✓	1 to 10	<1
Weight gain	-	-	<2	<1	10 to 12	-	✓	-	-
Xerostomia	✓	-	<2	<1	<1	-	-	-	-
<b>Genitourinary</b>									
Cystitis	-	-	<2	<1	-	-	-	-	-
Diabetes insipidus	-	-	-	-	-	<1	-	-	-
Dysuria	1 to 10	-	<2	-	-	-	-	-	-
Ejaculatory failure	-	-	-	-	-	≤5	-	-	-
Hematuria	-	-	-	-	1 to 10	-	-	-	-
Impotence	1 to 10	1 to 10	-	<1	1 to 10	1 to 4	✓	-	<1
Libido decreased	-	-	<2	<1	<1	-	-	-	-
Micturition (frequency)	3	-	-	-	-	-	-	-	-
Nocturia	1 to 10	-	-	-	-	-	-	-	-
Polyuria	-	-	-	<1	-	-	-	-	-
Sexual ability decreased	-	-	-	-	-	-	-	>10	-
Urinary incontinence	-	-	-	-	<1	-	-	-	-
Urinary retention	✓	-	-	-	-	<1	-	-	-
<b>Hematologic</b>									
Agranulocytosis	-	-	-	-	<1	-	✓	-	-
Anemia (aplastic/hemolytic)	-	-	<2	-	1 to 10	-	-	-	-
Claudication	-	-	-	-	-	-	✓	-	-
Leukopenia	-	-	-	<1	<1	-	-	<1	-
Pancytopenia	-	-	-	-	<1	-	-	-	-
Prothrombin decreased	-	-	-	-	1 to 10	-	-	-	-
Purpura	-	-	<2	-	1 to 10	-	-	-	-
Thrombocytopenia	-	<1	<2	<1	1 to 10	-	✓	<1	1 to 10
<b>Hepatic</b>									
Cholestatic jaundice	-	-	-	-	<1	<1	-	-	-
Hepatic impairment	✓	-	-	-	<1	<1	-	-	-
Hepatitis	-	-	-	-	-	<1	✓	-	-
Increase liver enzymes	-	<1	-	-	-	-	-	-	<1
Transaminases increase	✓	-	<2	<1	1 to 10	4	✓	-	-
<b>Laboratory Test Abnormalities</b>									
Alkaline phosphatase increased	✓	-	-	-	-	-	✓	-	-
Hypercalcemia	-	-	-	-	<1	-	-	-	-
Hypercholesterolemia	-	-	<2	-	1 to 4	-	-	-	1 to 10

Adverse Events	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
Hyperglycemia	-	-	<2	-	-	-	-	-	-
Hyperkalemia	-	-	<2	<1	1 to 10	-	-	-	-
Hypernatremia	-	-	-	-	-	-	-	-	-
Hyperphosphatemia	-	-	-	-	3 to 6	-	-	-	-
Hypertriglyceridemia	-	-	-	<1	1	-	-	-	-
Hyperuricemia	-	-	<2	<1	1 to 10	-	-	-	1 to 10
Hypervolemia	-	-	-	-	≤4	-	-	-	-
Hypoglycemia	-	✓	<2	<1	1 to 10	-	-	-	-
Hyponatremia	-	-	-	-	1 to 10	-	-	-	-
Hypokalemia	-	-	<2	-	1 to 10	-	-	-	-
Lactate dehydrogenase increased	-	-	-	-	-	-	✓	-	-
<b>Musculoskeletal</b>									
Arthralgia	-	-	3 to 5	1 to 10	1 to 6	-	✓	-	-
Arthritis	-	-	-	-	-	-	✓	-	-
Asthenia	-	-	-	≤2	-	-	-	-	-
Back pain	1 to 10	-	-	<1	2 to 7	-	-	-	-
Joint pain	1 to 10	-	-	<1	-	-	-	-	-
Muscle cramps	-	-	<2	<1	1 to 10	-	-	-	-
Muscle pain	-	-	-	<1	-	-	✓	-	-
Muscle spasm	-	-	-	-	-	-	-	-	-
Myalgia	2	-	-	-	-	-	-	-	-
Neuralgia	-	-	<2	-	<1	-	-	-	-
Paresthesia	-	-	-	-	-	≤5	-	-	1 to 10
Peripheral ischemia	✓	-	-	-	-	-	-	-	-
Restlessness	-	-	-	<1	-	-	-	-	-
Tremor	-	-	<2	<1	-	-	-	-	-
Toxic myopathy	-	-	-	-	-	<1	-	-	-
Twitching	-	-	<2	<1	-	-	-	-	-
Weakness	-	-	-	-	7 to 11	1	-	-	1 to 10
<b>Renal</b>									
Blood urea nitrogen increased	-	-	-	<1	≤6	≤8	-	-	1 to 10
Creatinine increase	-	-	-	<1	1 to 10	-	-	-	-
Glycosuria	-	-	-	-	1 to 10	-	-	-	-
Hematuria	-	-	-	1 to 10	-	-	-	-	-
Interstitial nephritis	-	-	-	-	<1	-	-	-	-
Renal colic	-	-	-	<1	-	-	-	-	-
Renal failure/dysfunction	-	-	-	-	1 to 10	-	-	-	<1
<b>Respiratory</b>									
Asthma	-	-	-	<1	<1	-	-	-	-
Bronchitis	-	-	-	<1	-	-	-	-	-
Bronchospasm	-	✓	-	<1	<1	<1	1	1 to 10	<1
Cough	1	-	<2	<1	5 to 8	-	-	-	-
Dyspnea	4	<1	2	1 to 2	>3	2	1 to 3	<1	≤1
Eosinophilic pneumonitis	-	-	-	-	<1	-	-	-	-
Interstitial pneumonitis	-	-	-	-	<1	-	-	-	-
Nasal congestion	-	-	-	-	1	1 to 6	-	-	-

Adverse Events	Single Entity Agents (A-N)								
	Acebutolol	Atenolol	Betaxolol	Bisoprolol	Carvedilol	Labetalol	Metoprolol	Nadolol	Nebivolol
Nasopharyngitis	-	-	-	-	4	-	-	-	-
Pharyngitis	1 to 10	-	2	<1	-	-	-	-	-
Pleurisy	✓	-	-	-	-	-	-	-	-
Pneumonitis	✓	-	-	-	-	-	-	-	-
Pulmonary edema	-	-	-	-	>3	-	-	-	<1
Pulmonary granulomas	✓	-	-	-	-	-	-	-	-
Respiratory failure/distress	-	-	-	-	<1	-	-	-	-
Rhinitis	2	-	-	3 to 4	2	-	✓	-	-
Sinus congestion	-	-	-	-	1	-	-	-	-
Sinusitis	-	-	-	2	-	-	-	-	-
Upper respiratory infection	-	-	-	5	-	-	-	-	-
Wheezing	1 to 10	<1	-	-	-	-	1	-	-
<b>Special Senses</b>									
Abnormal/blurred vision	2	-	-	-	1 to 5	1	✓	-	-
Blepharitis	-	-	<2	-	-	-	-	-	-
Cataract	-	-	<2	-	-	-	-	-	-
Conjunctivitis	1 to 10	-	-	-	-	-	-	-	-
Dry eyes	1 to 10	-	-	-	-	-	✓	-	-
Eye pain	1 to 10	-	-	<1	-	-	-	-	-
Hearing decreased	-	-	<2	<1	<1	-	-	-	-
Lacrimation, abnormal	-	-	-	<1	-	-	-	-	-
Tinnitus	-	-	<2	<1	<1	-	-	-	-
Visual disturbances	-	-	<2	<1	-	-	✓	-	-
<b>Other</b>									
Allergy/allergic reaction	-	-	-	-	1 to 10	-	-	-	-
Anaphylactoid reaction	-	-	-	-	<1	<1	-	-	-
Angioedema	-	-	-	-	-	<1	-	-	<1
Cholecystitis	-	-	-	-	-	-	-	-	-
Cutaneous vasculitis	-	-	-	<1	-	-	-	-	-
Diaphoresis	-	-	-	-	-	≤4	-	-	-
Gangrene	-	-	-	-	-	-	✓	-	-
Hypersensitivity	-	-	-	-	-	<1	-	-	<1
Lupus syndrome	✓	<1	-	-	-	<1	-	-	-
Metabolic acidosis	-	-	<2	-	-	-	-	-	-
Necrotizing angitis	-	-	-	-	-	-	-	-	-
Peyronie's disease	-	<1	<2	<1	-	<1	<1	-	-
Positive antinuclear antibody test	-	<1	5	<1	1 to 10	<1	-	-	-
Tinnitus	-	-	-	-	-	-	✓	-	-

✓ Percent not specified  
-Event not reported

**Table 9. Adverse Drug Events (%) Reported with the Beta-Adrenergic Blocking Agents<sup>1-22</sup>**

Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendo-flumethiazide	
<b>Cardiovascular</b>										
Angina	-	-	✓	-	✓	-	-	-	-	
Arrhythmia	1 to 10	-	-	5	✓	-	<1	-	<1	
Arterial/vascular insufficiency	-	-	✓	-	-	-	-	1	-	
Atrioventricular nodal disturbances	-	-	✓	-	-	-	-	-	-	
Bradycardia	<1	≤2	6	8 to 16	1 to 10	1 to 10	<1	2 to 16	1 to 10	
Cardiac failure/arrest	-	-	-	-	1 to 10	-	-	-	-	
Cardiogenic shock	-	-	✓	-	-	-	-	✓	-	
Chest pain	-	3	2 to 4	3 to 16	-	1 to 10	1 to 2	1	<1	
Cold extremities	<1	≤2	7 to 8	<1	1 to 10	1 to 10	<1	1	1 to 10	
Congestive heart failure	1 to 10	<1	✓	5	-	1 to 10	<1	1	1 to 10	
Edema	<1	10	2	2 to 8	✓	1 to 10	<1	-	1 to 10	
Electrocardiogram abnormal	-	-	-	2 to 7	-	-	-	-	-	
Flushing	-	-	-	-	-	-	<1	-	-	
Heart block	<1	≤2	-	-	✓	1 to 10	-	5	-	
Hypotension	<1	≤2	✓	3 to 6	✓	1 to 10	1 to 10	1 to 27	-	
Myocardial contractility impaired	-	-	✓	-	-	-	<1	<1	-	
Myocardial ischemia	-	-	-	-	-	-	-	-	-	
Orthostatic hypotension	-	-	-	-	-	-	1 to 10	1 to 10	<1	
Palpitations	-	≤1	-	3 to 14	1 to 10	-	<1	1	1 to 10	
Peripheral circulation reduced	-	-	-	3	-	-	-	-	1 to 10	
Peripheral edema	-	-	-	-	-	-	-	1	-	
Rhythm disturbance	-	-	-	-	-	-	<1	-	-	
Shortness of breath	-	-	-	-	-	-	-	✓	-	
Syncope	-	≤2	✓	5	-	-	<1	1	-	
Tachycardia	-	≤2	-	-	-	-	-	-	-	
Torsade de pointes	-	-	-	1 to 4	-	-	-	-	-	
Thrombosis, mesenteric arterial	-	-	✓	-	-	-	-	-	-	
<b>Central Nervous System</b>										
Abnormal dreams	-	-	3	-	-	-	-	-	-	
Amnesia	-	-	✓	-	-	-	-	-	-	
Anxiety	-	-	-	2 to 4	✓	-	<1	✓	-	
Catatonia	-	-	✓	-	-	-	-	-	-	
Cerebral ischemia	-	-	-	-	1 to 10	-	-	-	-	
Cerebral vascular accident	-	-	-	-	1 to 10	-	-	-	-	
Cognitive dysfunction	-	-	✓	-	-	-	-	-	-	
Confusion	<1	-	✓	6	✓	1 to 10	<1	✓	<1	
Depression	1 to 10	-	1 to 3	1 to 4	✓	1 to 10	<1	5	1 to 10	
Disorientation	-	-	-	-	✓	-	-	-	✓	
Dizziness	1 to 10	9	2 to 11	3 to 20	1 to 10	1 to 10	<1	2 to 10	-	
Drowsiness	-	-	2	-	-	-	-	-	>10	

Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendro-flumethiazide	
Emotional lability	-	-	✓	<1	-	-	-	-	-	
Fatigue	1 to 10	8	3 to 17	5 to 20	1 to 10	1 to 10	6 to 8	1 to 10	✓	
Hallucinations	-	<1	✓	-	✓	<1	<1	✓	<1	
Headache	1 to 10	-	1 to 9	2 to 12	-	1 to 10	<1	✓	<1	
Hyper/hypoesthesia	-	-	-	-	-	-	1 to 2	-	-	
Insomnia	<1	10	3 to 8	2 to 4	✓	1 to 10	2 to 3	✓	>10	
Lethargy	<1	-	4	-	-	1 to 10	-	-	-	
Lightheadedness	-	-	✓	12	-	-	-	-	-	
Malaise	-	-	-	-	-	-	<1	-	-	
Memory loss	-	-	-	-	✓	-	<1	✓	-	
Mental impairment	-	-	-	-	-	1 to 10	-	-	-	
Nervousness	-	7	2	-	✓	-	<1	✓	<1	
Nightmares/vivid dreams	<1	5	✓	-	✓	1 to 10	-	✓	✓	
Paresthesia	-	-	-	-	-	<1	<1	✓	-	
Psychosis	-	-	✓	-	-	<1	-	-	-	
Sleep disturbance	-	-	-	1 to 8	-	-	<1	✓	-	
Somnolence	-	-	✓	-	✓	-	<1	✓	✓	
Vertigo	-	-	✓	<1	-	-	<1	✓	-	
<b>Dermatologic</b>										
Acne	-	-	-	-	-	-	<1	-	-	
Alopecia	-	-	✓	<1	1 to 10	<1	<1	<1	-	
Cutaneous ulcers	-	-	✓	-	-	-	-	-	-	
Dermatitis	-	-	✓	-	-	-	-	-	-	
Eczematous eruptions	-	-	✓	-	-	-	-	-	-	
Erythema multiforme	-	-	✓	-	-	-	<1	<1	✓	
Exfoliative dermatitis	-	-	✓	-	-	-	<1	<1	✓	
Hyperkeratosis	-	-	✓	-	-	-	-	-	-	
Nail changes	-	-	✓	-	-	-	-	-	-	
Oculomucocutaneous reactions	-	-	✓	-	-	-	-	-	-	
Photosensitivity	-	-	-	<1	-	1 to 10	1 to 10	1 to 10	✓	
Pruritus	-	1	✓	<1	-	-	<1	5	-	
Pseudo pemphigoid	-	-	-	-	✓	-	-	-	-	
Psoriasisiform rash	-	-	✓	-	✓	<1	<1	-	-	
Psoriasis (exacerbated)	-	-	-	-	1 to 10	-	<1	✓	✓	
Purpura	-	-	-	-	-	<1	<1	-	✓	
Rash	-	-	0 to 2	2 to 5	1 to 10	<1	<1	5	-	
Red crusted skin	-	-	-	<1	-	-	-	-	-	
Skin necrosis after extravasation	-	-	-	<1	-	-	-	-	-	
Stevens-Johnson syndrome	-	-	✓	-	-	-	<1	<1	-	
Sweating, excessive	-	≤2	2	<1	-	-	-	✓	-	
Toxic epidermal necrolysis	-	-	✓	-	-	-	<1	<1	✓	
Ulcers	-	-	✓	-	-	-	-	-	-	
Urticaria	-	-	✓	5	1 to 10	<1	-	✓	-	
<b>Endocrine and Metabolic</b>										

Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendro-flumethiazide	
Diabetes (exacerbated)	-	-	-	-	-	-	-	✓	✓	
Glycosuria	-	-	-	-	-	<1	-	-	-	
Gout	-	-	-	-	-	<1	<1	-	-	
Hypoglycemia masked	-	-	-	-	1 to 10	-	-	-	-	
Libido decreased	-	-	-	-	✓	-	-	✓	-	
<b>Gastrointestinal</b>										
Abdominal pain	-	-	1	1 to 4	-		<1	✓	-	
Anorexia	-	-	✓	-	✓	1 to 10	1 to 10	1 to 10	✓	
Constipation	-	-	0 to 2	-	1 to 10	1 to 10	<1	1	1 to 10	
Cramping	-	-	✓	-	-				✓	
Diarrhea	1 to 10	≤2	2 to 7	2 to 7	1 to 10	1 to 10	3 to 4	5	1 to 10	
Dry mouth	-	-	-	-	✓				-	
Dyspepsia	1 to 10	-	1 to 7	2 to 3	1 to 10		<1		✓	
Epigastric distress	-	-	-	-	-	1 to 10	1 to 10	1 to 10	-	
Flatulence	-	-	4	1 to 2	-			1	-	
Gastritis/gastric irritation	-	-	-	-	-	-	<1		-	
Heartburn	-	-	-	-	-	-	-	1	-	
Ischemic colitis	<1	-	✓	-	-				-	
Melena	-	-	-	-	-				✓	
Nausea	1 to 10	5	1 to 6	4 to 10	1 to 10	1 to 10	2	1	1 to 10	
Pancreatitis	-	-	-	-	-	<1		<1	-	
Peptic ulcer	-	-	-	-	-	-	<1			
Periodontitis	-	-	-	-	-				✓	
Retroperitoneal fibrosis	-	-	-	-	✓			✓	-	
Stomach discomfort	-	-	✓	3 to 6	-				1 to 10	
Taste disorder	-	-	-	-	-		<1	✓	✓	
Vomiting	-	≤2	✓	4 to 10	-	<1	1 to 2	✓	1 to 10	
Weight gain	-	≤2	-	-	-		<1	✓	-	
Xerostomia	-	-	-	-	1 to 10	-	<1	-	-	
<b>Genitourinary</b>										
Cystitis	-	-	-	-	-	-	<1	-	-	
Impotence	-	≤2	1	2	1 to 10	1 to 10	<1	✓	-	
Interstitial nephritis	-	-	✓	-	-	-	-	-	-	
Micturition (frequency)	-	-	1	-	-	-	-	-	-	
Oliguria	-	-	✓	-	-	-	-	-	-	
Polyuria	-	≤2	-	-	-	<1	<1	-	-	
Proteinuria	-	-	✓	-	-	-	-	-	-	
Sexual ability decreased	-	-	-	3	-	-	<1	-	>10	
<b>Hematologic</b>										
Agranulocytosis	-	-	✓	-	-	<1	-	<1	-	
Anemia (aplastic/hemolytic)	-	-	-	-	-	<1	-	<1	✓	
Bleeding	-	-	-	2	-	-	-	-	-	
Claudication	-	-	-	-	✓	-	-	✓	-	
Eosinophilia	-	-	-	<1	-	-	-	-	-	

Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendro-flumethiazide	
Leukopenia	-	-	-	<1	-	<1	<1	<1	<1	
Prothrombin decreased	-	-	-	-	-	-	-	-	✓	
Purpura	<1	-	✓	-	-	<1	-	-	-	
Thrombocytopenia	<1	-	✓	<1	-	<1	<1	<1	<1	
<b>Hepatic</b>										
Cholestatic jaundice	-	-	-	-	-	-	-	-	✓	
Hepatic impairment	-	-	-	-	-	<1	-	<1	-	
Hepatitis	-	-	-	-	-	-	-	✓	-	
Increase liver enzymes	-	7	-	-	-	<1	-	-	-	
Transaminases increase	-	-	✓	<1	-	-	<1	✓	-	
<b>Laboratory Test Abnormalities</b>										
Alkaline phosphatase increased	-	<1	✓	-	-	-	-	✓	-	
Electrolyte imbalance	-	-	-	-	-	-	-	-	✓	
Hypercalcemia	-	-	-	-	-	<1	-	<1	-	
Hypercholesterolemia	-	-	-	-	-	-	-	-	-	
Hyperglycemia	-	-	✓	-	-	<1	-	-	-	
Hyperkalemia	-	-	✓	-	-	-	<1	-	-	
Hyperlipidemia	-	-	✓	<1	-	-	-	-	-	
Hypernatremia	-	-	-	-	-	<1	-	-	-	
Hyperphosphatemia	-	-	-	-	-	-	-	-	✓	
Hypertriglyceridemia	-	-	-	-	-	-	<1	-	✓	
Hyperuricemia	-	<1	-	-	-	<1	<1	-	-	
Hypoglycemia	<1	-	✓	-	-	-	<1	-	-	
Hypokalemia	-	-	-	-	-	1 to 10	-	1 to 10	-	
Hyponatremia	-	-	-	-	-	<1	-	-	-	
Lactate dehydrogenase increased	-	<1	-	-	-	-	-	✓	-	
<b>Musculoskeletal</b>										
Arthralgia	1 to 10	7	1	-	-	-	1 to 10	✓	-	
Arthritis	-	-	-	-	-	-	-	✓	-	
Arthropathy	-	-	✓	-	-	-	-	-	-	
Asthenia	-	-	-	-	-	-	≤2	-	-	
Back pain	-	-	-	3	-	-	<1	-	-	
Carpal Tunnel syndrome	-	-	✓	-	-	-	-	-	-	
Extremity pain	-	-	-	7	-	-	-	-	-	
Joint pain	-	-	-	-	-	-	<1	-	-	
Muscle cramps	-	3	-	-	-	<1	<1	-	✓	
Muscle pain	-	10	-	-	-	-	<1	✓	-	
Myalgia	-	-	1	<1	-	<1	-	-	-	
Myasthenia gravis exacerbated	-	-	-	-	✓	-	-	-	-	
Myotonus	-	-	✓	-	-	-	-	-	-	
Neuralgia	-	-	-	-	-	-	-	-	✓	
Paralysis	-	-	-	<1	-	-	-	-	-	
Paresthesia	-	3	✓	4	✓	-	-	-	-	
Polyarthritis	-	-	✓	-	-	-	-	-	-	

Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendro-flumethiazide	
Restlessness	-	-	-	-	-	<1	<1	-	-	
Tremor	-	-	-	-	-	-	<1	-	✓	
Twitching	-	-	-	-	-	-	<1	-	-	
Weakness	-	4	1	4 to 13	-	<1	-	-	✓	
<b>Renal</b>										
Blood urea nitrogen increase	-	-	✓	-	-	-	<1	-	-	
Creatinine increase	-	-	-	-	-	-	<1	-	-	
Hematuria	-	-	-	-	-	-	1 to 10	-	-	
Interstitial nephritis	-	-	-	-	-	-	-	<1	-	
Renal colic	-	-	-	-	-	-	<1	-	-	
Renal failure	-	-	-	-	-	-	-	<1	-	
<b>Respiratory</b>										
Asthma	-	-	-	1 to 2	-	-	<1	-	-	
Bronchitis	-	-	-	-	-	-	<1	-	-	
Bronchospasm	<1	-	✓	-	✓	-	<1	1	1 to 10	
Cough	<1	-	1	-	✓	-	<1	-	-	
Dyspnea	-	5	1 to 6	5 to 21	1 to 10	<1	1 to 2	1 to 3	<1	
Eosinophilic pneumonitis	-	-	-	-	-	-	-	<1	-	
Laryngospasm	-	-	✓	-	-	-	-	-	-	
Nasal congestion	-	-	-	-	✓	-	-	-	-	
Nasopharyngitis	-	-	-	-	-	-	-	-	✓	
Pharyngitis	-	-	✓	-	-	-	<1	-	-	
Pulmonary edema	-	-	✓	<1	✓	-	-	-	-	
Respiratory failure	-	-	✓	-	✓	-	-	<1	-	
Rhinitis	-	-	1	-	-	-	3 to 4	✓	-	
Sinusitis	-	-	-	-	-	-	2	-	-	
Upper respiratory infection	-	-	5	5 to 8	-	-	5	-	-	
Wheezing	-	≤2	✓	-	-	<1	-	1	✓	
<b>Special Senses</b>										
Abnormal/blurred vision	-	-	3	-	-	-	-	✓	-	
Burning	-	≤2	-	-	-	-	-	-	-	
Corneal sensitivity decrease	-	-	-	-	✓	-	-	-	-	
Cystoid macular edema	-	-	-	-	✓	-	-	-	-	
Diplopia	-	-	-	-	✓	-	-	-	-	
Dry eyes	-	-	-	-	✓	-	-	✓	-	
Eye discomfort/pain	-	≤2	-	-	-	-	<1	-	-	
Hearing decreased	-	-	-	-	-	-	<1	-	-	
Hyperemia of conjunctiva	-	-	✓	-	-	-	-	-	-	
Keratitis	-	-	-	-	1 to 10	-	-	-	-	
Lacrimation abnormal	-	-	-	-	-	-	<1	-	✓	
Mydriasis	-	-	✓	-	-	-	-	-	-	
Ocular discharge	-	-	-	-	✓	-	-	-	-	
Ocular pain	-	-	-	-	✓	-	-	-	-	
Ptosis	-	-	-	-	✓	-	-	-	-	



Adverse Events	Single Entity Agents (O-Z)					Combination Products				
	Penbutolol	Pindolol	Propranolol	Sotalol	Timolol	Atenolol and Chlor-thalidone	Bisoprolol and HCTZ	Metoprolol and HCTZ	Nadolol and Bendro-flumethiazide	
Refractive changes	-	-	-	-	✓	-	-	-	-	
Tinnitus	-	-	-	-	1 to 10	-	<1	-	-	
Visual disturbances	-	≤2	✓	1 to 5	✓	-	<1	✓	-	
Xerophthalmia	-	-	✓	-	-	-	-	-	-	
<b>Other</b>										
Allergy	-	-	-	-	✓	-	-	<1	-	
Anaphylactoid reaction	-	-	✓	-	-	-	-	-	-	
Angioedema	-	-	-	-	1 to 10	-	-	-	✓	
Cholecystitis	-	-	-	-	-	<1	-	-	-	
Cutaneous vasculitis	-	-	-	-	-	<1	<1	-	-	
Gangrene	-	-	-	-	-	-	-	✓	-	
Hypervolemia	-	-	-	-	-	-	-	-	✓	
Lupus syndrome	-	-	✓	-	1 to 10	<1	-	-	✓	
Mesenteric arterial thrombosis	<1	-	-	-	-	-	-	-	-	
Necrotizing angitis	-	-	-	-	-	<1	-	-	-	
Peyronie's disease	-	-	✓	-	1 to 10	<1	<1	<1	-	
Positive antinuclear antibody test	-	-	-	-	-	<1	<1	-	-	
Tinnitus	-	-	-	-	-	-	-	✓	-	

✓ Percent not specified

- Event not reported

**Table 10. Boxed Warning for Atenolol<sup>1</sup>**

<b>WARNING</b>
Advise patients with coronary artery disease who are being treated with atenolol against abrupt discontinuation of therapy. Severe exacerbation of angina and the occurrence of myocardial infarction and ventricular arrhythmias have been reported in patients with angina following the abrupt discontinuation of therapy with $\beta$ -blockers. The last two complications may occur with or without preceding exacerbation of the angina pectoris. As with other $\beta$ -blockers, when discontinuation of atenolol is planned, observe the patient carefully and advise the patient to limit physical activity to a minimum. If the angina worsens or acute coronary insufficiency develops, it is recommended that atenolol be promptly reinstated, at least temporarily. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue atenolol therapy abruptly, even in patients treated only for hypertension.

**Table 11. Boxed Warning for Metoprolol<sup>1</sup>**

<b>WARNING</b>
Ischemic heart disease: Following abrupt cessation of therapy with certain $\beta$ -blocking agents, exacerbations of angina pectoris and, in some cases, myocardial infarction have occurred. When discontinuing chronically administered metoprolol, particularly in patients with ischemic heart disease, gradually reduce the dosage over a period of one to two weeks and carefully monitor the patient. If angina markedly worsens or acute coronary insufficiency develops, reinstate metoprolol administration promptly, at least temporarily, and take other measures appropriate for the management of unstable angina. Warn patients against interruption or discontinuation of therapy without their health care provider's advice. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue metoprolol tartrate therapy abruptly, even in patients treated only for hypertension.

**Table 12. Boxed Warning for Nadolol<sup>1</sup>**

<b>WARNING</b>
Exacerbation of ischemic heart disease following abrupt withdrawal: Hypersensitivity to catecholamines has been observed in patients withdrawn from $\beta$ -blocker therapy; exacerbation of angina and, in some cases, myocardial infarction have occurred after abrupt discontinuation of such therapy. When discontinuing chronically administered nadolol, particularly in patients with ischemic heart disease, gradually reduce the dosage over a period of one to two weeks and carefully monitor the patient. If angina markedly worsens or acute coronary insufficiency develops, reinstitute nadolol administration promptly, at least temporarily, and take other measures appropriate for the management of unstable angina. Warn patients against interruption or discontinuation of therapy without the physician's advice. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue nadolol therapy abruptly, even in patients treated only for hypertension.

**Table 13. Boxed Warning for Propranolol<sup>1</sup>**

<b>WARNING</b>
Angina pectoris: There have been reports of exacerbation of angina and, in some cases, myocardial infarction, following abrupt discontinuance of propranolol therapy. Therefore, when discontinuance of propranolol is planned, the dosage should be gradually reduced over at least a few weeks, and the patient should be cautioned against interruption or cessation of therapy without a health care provider's advice. If propranolol therapy is interrupted and exacerbation of angina occurs, it is usually advisable to reinstitute propranolol therapy and take other measures appropriate for the management of angina pectoris. Because coronary artery disease may be unrecognized, it may be prudent to follow the above advice in patients considered at risk of having occult atherosclerotic heart disease who are given propranolol for other indications.

**Table 14. Boxed Warning for Sotalol<sup>1</sup>**

<b>WARNING</b>
To minimize the risk of induced arrhythmia, place patients initiated or reinitiated on sotalol AF or sotalol for a minimum of three days (on their maintenance dose) in a facility that can provide cardiac resuscitation, continuous electrocardiographic monitoring, and calculations of creatinine clearance. Calculate creatinine clearance prior to dosing. Do not substitute sotalol for sotalol AF because of significant differences in labeling (i.e., patient package insert, dosing administration, safety information).

**Table 15. Boxed Warning for Timolol<sup>1</sup>**

<b>WARNING</b>
Exacerbation of ischemic heart disease following abrupt withdrawal: Hypersensitivity to catecholamines has been observed in patients withdrawn from $\beta$ -blocker therapy; exacerbation of angina and, in some cases, myocardial infarction have occurred after abrupt discontinuation of such therapy. When discontinuing chronically administered timolol, particularly in patients with ischemic heart disease, gradually reduce the dosage over a period of one to two weeks and carefully monitor the patient. If angina markedly worsens or acute coronary insufficiency develops, reinstitute timolol administration promptly, at least temporarily, and take other measures appropriate for the management of unstable angina. Warn patients against interruption of discontinuation of therapy without the physician's advice. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue timolol therapy abruptly, even in patients treated only for hypertension.

## VII. Dosing and Administration

The usual dosing regimens for the  $\beta$ -adrenergic blocking agents are listed in Table 16.

**Table 16. Usual Dosing Regimens for the Beta-Adrenergic Blocking Agents<sup>3-22</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Acebutolol	<p><u>Hypertension:</u> Capsule: initial, 400 mg/day, twice daily dosing may be required for adequate control; maintenance, 200 to 1,200 mg/day in two divided doses; maximum, 1,200 mg/day</p> <p><u>Ventricular arrhythmias:</u> Capsule: initial: 200 mg twice daily; maintenance, gradual increase until optimal response, usually 600 to 1,200 mg/day; maximum, 1,200 mg/day</p>	Safety and efficacy in children have not been established.	Capsule: 200 mg 400 mg
Atenolol	<p><u>Angina pectoris:</u> Tablet: initial, 50 mg once daily; maintenance, if optimal response not achieved after one week, increase to 100 mg daily; maximum, 200 mg/daily</p> <p><u>Hypertension:</u> Tablet: initial: 50 mg once daily; maintenance, if optimal response not achieved, increase dose to 100 mg once daily; maximum, 100 mg/day</p> <p><u>Myocardial infarction:</u></p>	Safety and efficacy in children have not been established.	Tablet: 25 mg 50 mg 100 mg

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	Tablet: 50 mg twice daily, or 100 mg once daily for 6 to 9 days or until hospital discharge		
Betaxolol	<u>Hypertension:</u> Tablet: initial, 10 mg once daily; maintenance, if optimal response not seen after seven to 14 days, may increase the dose to 20 mg/day; maximum, 40 mg/day	Safety and efficacy in children have not been established.	Tablet: 10 mg 20 mg
Bisoprolol	<u>Hypertension:</u> Tablet: initial, 2.5 to 5 mg once daily; maintenance, if optimal control is not achieved, dose may be increased to 10 mg daily and again to 20 mg/day if needed; maximum, 20 mg/day	Safety and efficacy in children have not been established.	Tablet: 5 mg 10 mg
Carvedilol	<p><u>Heart failure:</u> Extended-release capsule: initial, 10 mg once daily; maintenance, if tolerated, double the dose at intervals of &gt;14 days as needed; maximum, 80 mg once daily</p> <p>Tablet: initial, 3.125 mg twice daily; maintenance, if tolerated, double the dose at intervals of &gt;14 days as needed up to 50 mg twice daily; maximum, 25 mg twice daily (patients ≤85 kg) or 50 mg twice daily (patients &gt;85 kg)</p> <p><u>Hypertension:</u> Extended-release capsule: initial, 20 mg once daily; maintenance, if tolerated, double the dose every seven to 14 days as needed; maximum, 80 mg once daily</p> <p>Tablet: initial, 6.25 mg twice daily; maintenance, if tolerated, double the dose every seven to 14 days as needed; maximum, 25 mg twice daily</p> <p><u>Myocardial Infarction:</u> Capsule ER: initial, 10 to 20 mg once daily; maintenance: if tolerated, double the dose every 3 to 10 days as needed up to a maximum of 80 mg once daily</p> <p>Tablet IR: initial, 6.25 mg twice daily; maintenance: if tolerated, double the dose every 3 to 10 days as needed up to a maximum of 25 mg twice daily</p>	Safety and efficacy in children have not been established.	<p>Extended-release capsule: 10 mg 20 mg 40 mg 80 mg</p> <p>Tablet: 3.125 mg 6.25 mg 12.5 mg 25 mg</p>
Labetalol	<u>Hypertension:</u> Injection, tablet: initial: 100 mg twice daily; maintenance, titrate by increments of 100 mg twice daily every two to three days, usual dose is 200 to 400 mg twice daily; larger doses may be administered three times daily to improve tolerability;	Safety and efficacy in children have not been established.	<p>Injection: 5 mg/mL</p> <p>Tablet: 100 mg 200 mg 300 mg</p>

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	maximum, doses of 1,200 to 2,400 mg/day have been used		
Metoprolol	<p><u>Angina pectoris:</u> Extended-release capsule, tablet: initial, 100 mg once daily; maintenance, gradually increase dose in weekly intervals; maximum, 400 mg/day</p> <p>Injection, tablet: initial, 100 mg/day in two divided doses; maintenance, gradually increase dose in weekly intervals, usual dose is 100 to 400 mg/day; maximum, 400 mg/day</p> <p><u>Heart failure:</u> Extended-release capsule, tablet: initial, 25 mg/day; maintenance, double the dose every two weeks up to 200 mg/day or highest dose tolerated</p> <p><u>Hypertension:</u> Extended-release capsule, tablet: initial, 25 to 100 mg once daily; maintenance, gradually increase dose in weekly intervals up to 400 mg/day</p> <p>Injection, tablet: initial, 50 to 100 mg/day in single or divided doses; maintenance, gradually increase dose in weekly intervals, usual dose is 100 to 450 mg/day; maximum, 450 mg/day</p> <p><u>Myocardial infarction:</u> Injection, tablet: initial, 100 mg twice daily; maintenance, 100 mg twice daily for at least three months</p>	<p><u>Hypertension in children <math>\geq 6</math> years of age:</u> Extended-release capsule, tablet: initial: 1 mg/kg once daily (maximum: 50 mg once daily); maintenance, adjust dose to optimal response up to 2 mg/kg or 200 mg/day; maximum, 2 mg/kg/day or 200 mg/day</p> <p>Safety and efficacy in children &lt;6 years of age have not been established.</p>	<p>Extended-release capsule (succinate): 25 mg 50 mg 100 mg 200 mg</p> <p>Extended-release tablet (succinate): 25 mg 50 mg 100 mg 200 mg</p> <p>Injection (tartrate): 5 mg/5 mL</p> <p>Tablet (tartrate): 25 mg 37.5 mg 50 mg 75 mg 100 mg</p>
Nadolol	<p><u>Angina pectoris:</u> Tablet: initial, 40 mg once daily; maintenance, increase dose by 40 to 80 mg every three to seven days until optimal response; maximum, 240 mg/day</p> <p><u>Hypertension:</u> Tablet: initial, 40 mg once daily; maintenance, increase dose gradually by 40 to 80 mg increments every seven to 21 days until optimal response; maximum, 320 mg/day</p>	Safety and efficacy in children have not been established.	Tablet: 20 mg 40 mg 80 mg
Nebivolol	<u>Hypertension:</u> Tablet: initial: 5 mg once daily; maintenance, increase in two week intervals until optimal response; maximum, 40 mg/day	Safety and efficacy in children have not been established.	Tablet: 2.5 mg 5 mg 10 mg 20 mg
Penbutolol	<u>Hypertension:</u> Tablet: initial, 20 mg once daily;	Safety and efficacy in children have not been	Tablet: 20 mg

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	maintenance, 20 mg once daily, usual dose 10 to 40 mg once daily; maximum, 80 mg/day	established.	
Pindolol	<u>Hypertension:</u> Tablet: initial, 5 mg twice daily; maintenance, after three to four weeks, may be increase by 10 mg/day increments as needed; maximum, 60 mg/day	Safety and efficacy in children have not been established.	Tablet: 5 mg 10 mg
Propranolol	<p><u>Angina pectoris:</u> Extended-release capsule (Inderal LA<sup>®</sup>): initial, 80 mg once daily; maintenance, may gradually increase dose in three to seven day increments up to 160 mg once daily or higher, usual dose is 160 mg daily; maximum, 320 mg/day</p> <p>Solution, tablet: maintenance, 80 to 320 mg/day administered in two, three or four divided doses; maximum, 320 mg/day</p> <p><u>Cardiac arrhythmias:</u> Injection (ventricular arrhythmias): usual dose, 1 to 3 mg</p> <p>Solution, tablet (atrial fibrillation): maintenance, 10 to 30 mg in three to four divided doses before meals and at bedtime</p> <p><u>Essential tremor:</u> Solution, tablet: initial, 40 mg twice daily; maintenance, usual dose is 120 mg/day; maximum, 320 mg/day</p> <p><u>Hypertension:</u> Extended-release capsule (Inderal LA<sup>®</sup>): initial, 80 mg once daily; maintenance, may titrate dose up to 120 mg/day or higher, usual dose is 120 to 160 mg/day; maximum, 640 mg/day</p> <p>Extended-release capsule (InnoPran XL<sup>®</sup>): initial, 80 mg once daily at bedtime (around 10 pm); maintenance, may titrate dose up to 120 mg/day; maximum, 120 mg/day</p> <p>Solution, tablet: initial, 40 mg twice daily; maintenance, gradually increase the dose up to 640 mg/day divided into two to three doses, usual dose is 120 to 240 mg/day divided into two to three doses; maximum, 640 mg/day</p> <p><u>Hypertrophic subaortic stenosis:</u> Extended-release capsule (Inderal LA<sup>®</sup>):</p>	<p>Infantile hemangioma: Solution (Hemangeol<sup>®</sup>): Initiate treatment at 5 weeks to 5 months; initial, 0.15 mL/kg (0.6 mg/kg) twice daily at least 9 hours apart; after one week increase to 0.3 mL/kg (1.1 mg/kg) twice daily; after another week increase the dose to 0.4 mL/kg (1.7 mg/kg) twice daily and maintain for six months, readjusting for weight changes</p> <p>Safety and effectiveness for infantile hemangioma have not been established in pediatric patients greater than one year of age</p>	<p>Extended-release capsule: 60 mg 80 mg 120 mg 160 mg</p> <p>Injection: 1 mg/mL</p> <p>Solution: 4.28 mg/mL 20 mg/5 mL 40 mg/5 mL</p> <p>Tablet: 10 mg 20 mg 40 mg 60 mg 80 mg</p>

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<p>maintenance, 80 to 160 mg once daily</p> <p>Solution, tablet: 20 to 40 mg three to four times daily before meals and at bedtime</p> <p><u>Migraine:</u> Extended-release capsule (Inderal LA<sup>®</sup>): initial, 80 mg once daily; maintenance, may increase dose gradually up to 160 to 240 mg once daily, usual dose is 160 to 240 mg once daily; maximum, 240 mg/day</p> <p>Solution, tablet: initial, 80 mg daily in divided doses; maintenance, increase dose gradually up to 160 to 240 mg/day; maximum, 240 mg/day</p> <p><u>Myocardial Infarction:</u> Solution, tablet: initial, 40 mg three times daily; maintenance, after one month, titrate up to 60 to 80 mg three times daily as tolerated, usual dose is 180 to 240 mg in divided doses; maximum, 240 mg/day</p> <p><u>Pheochromocytoma:</u> Solution, tablet (operable tumors): 60 mg/day in divided doses for three days preoperatively as adjunct to <math>\alpha</math>-adrenergic blockade</p> <p>Solution, tablet (inoperable tumors): 30 mg/day in divided doses as adjunct to <math>\alpha</math>-adrenergic blockade</p>		
Sotalol	<p><u>Cardiac arrhythmias:</u> Solution, tablet (Betapace AF<sup>®</sup>, Sotylize<sup>®</sup>; maintenance of normal sinus rhythm): initial, 80 mg twice daily; maintenance, increase dose gradually with three days between increments up to 120 mg twice daily; maximum, 160 mg twice daily</p> <p>Solution, tablet (Betapace<sup>®</sup>, Sotylize<sup>®</sup>; ventricular arrhythmias): initial, 80 mg twice daily; maintenance, increase dose gradually with three days between increments up to 120 to 160 mg twice daily; maximum, 480 to 640 mg/day</p>	<p><u>Cardiac arrhythmias in children &gt;2 years of age:</u> Solution, tablet (Betapace AF<sup>®</sup>, Sotylize<sup>®</sup>; maintenance of normal sinus rhythm): initial, 30 mg/m<sup>2</sup> three times daily; maintenance, increase dose gradually with three days between increments up to 60 mg/m<sup>2</sup> three times daily; maximum, 60 mg/m<sup>2</sup> three times daily; neonate dosing available for Sotylize<sup>®</sup></p> <p>Solution, tablet (Betapace<sup>®</sup>, Sotylize<sup>®</sup>; ventricular Arrhythmias): initial,</p>	<p>Solution: 5 mg/mL</p> <p>Tablet: 80 mg, 120 mg, 160 mg, 240 mg</p>

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
		30 mg/m <sup>2</sup> three times daily; maintenance, increase dose gradually with three days between increments up to 60 mg/m <sup>2</sup> three times daily; maximum, 60 mg/m <sup>2</sup> three times daily; neonate dosing available for Sotylize®	
Timolol	<p><u>Hypertension:</u> Tablet: initial, 10 mg twice daily; maintenance, increase dose gradually in seven day increments up to 60 mg/day, usual dose is 20 to 40 mg/day; maximum, 60 mg/day divided into two doses</p> <p><u>Migraine:</u> Tablet: initial, 10 mg twice daily; maintenance, may increase dose up to 30 mg/day; maximum, 30 mg/day divided into two doses</p> <p><u>Myocardial infarction:</u> Tablet: 10 mg twice daily</p>	Safety and efficacy in children have not been established.	Tablet: 5 mg 10 mg 20 mg
<b>Combination Products</b>			
Atenolol and chlorthalidone	<p><u>Hypertension:</u> Tablet: initial: 50-25 mg once daily; maintenance, if optimum response is not achieved after one to two weeks, may increase to 100-25 mg once daily</p>	Safety and efficacy in children have not been established.	Tablet: 50-25 mg 100-25 mg
Bisoprolol and HCTZ	<p><u>Hypertension:</u> Tablet: initial, 2.5-6.25 mg once daily; maintenance, may titrate dose every seven to 14 days; maximum, 20-12.5 mg once daily</p>	Safety and efficacy in children have not been established.	Tablet: 2.5-6.25 mg 5-6.25 mg 10-6.25 mg
Metoprolol and HCTZ	<p><u>Hypertension:</u> Extended-release tablet: dosing must be individualized, the usual dose of metoprolol is 25 to 100 mg/day, and the usual dose of HCTZ is 12.5 to 50 mg/day</p> <p>Tablet: initial, 100-25 mg/day in single or divided doses; maintenance, may titrate dose gradually until desired effect is achieved, usual dose of metoprolol is 100 to 450 mg/day, and usual dose of HCTZ is 12.5 to 50 mg/day, may be administered in single or divided doses</p>	Safety and efficacy in children have not been established.	Tablet: 50-25 mg 100-25 mg 100-50 mg
Nadolol and bendroflumethiazide	<p><u>Hypertension:</u> Tablet: initial, 40-5 mg once daily; maintenance, if desired effect is not achieved, may increase dose to 80-5 mg once daily</p>	Safety and efficacy in children have not been established.	Tablet: 80-5 mg

HCTZ=hydrochlorothiazide, NYHA=New York Heart Association



## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the  $\beta$ -adrenergic blocking agents are summarized in Table 17.

**Table 17. Comparative Clinical Trials with the Beta-Adrenergic Blocking Agents**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Angina</b>				
<p>Pandhi et al.<sup>57</sup> (1985)</p> <p>Acebutolol 100 to 400 mg TID</p> <p>vs</p> <p>propranolol 40 to 160 mg TID</p> <p>vs</p> <p>placebo</p>	<p>DB, XO</p> <p>Patients with classical anginal symptoms of effort with <math>\geq 7</math> attacks per week and angina being stable for <math>\geq 8</math> to 12 weeks</p>	<p>N=24</p> <p>18 weeks</p>	<p>Primary: Incidence of anginal attack, number of nitroglycerin tablets used, exercise tolerance, side effects</p> <p>Secondary: Not reported</p>	<p>Primary: Both acebutolol and propranolol significantly reduced the incidence of anginal attacks per week compared to placebo (<math>P &lt; 0.001</math> for both groups), but the difference between the two groups was not significant (<math>P &gt; 0.05</math>).</p> <p>Both acebutolol and propranolol significantly reduced the number of nitroglycerin tablets used per week compared to placebo (<math>P &lt; 0.001</math> for both groups), but the difference between the two groups was not significant (<math>P &gt; 0.05</math>).</p> <p>Both acebutolol and propranolol significantly improved exercise tolerance compared to placebo (<math>P &lt; 0.001</math>), but the difference between the two groups was not significant (<math>P &gt; 0.05</math>).</p> <p>Side effects reported (i.e., insomnia, sweating, bitter taste, heart burn, muscle weakness) were similar between the two treatment groups. Clinical significance of the side effects was not reported.</p> <p>Secondary: Not reported</p>
<p>Jackson et al.<sup>58</sup> (1980)</p> <p>Atenolol 25, 50, 100, and 200 mg/day, each dose administered for a 2 week period</p> <p>vs</p>	<p>XO</p> <p>Adult patients with clinically stable exercise-induced angina for <math>\geq 3</math> months</p>	<p>N=10</p> <p>12 weeks</p>	<p>Primary: Anginal attack rate, nitroglycerin consumption, exercise data</p> <p>Secondary: Not reported</p>	<p>Primary: Compared to placebo, atenolol reduced the angina attack rate during all periods (<math>P &lt; 0.001</math>). A dose response was present with a decreasing number of attacks with increasing dosage. Doses of 100 and 200 mg were significantly more effective to 25 mg (<math>P &lt; 0.001</math> for both), but there was no significant difference between the 50 and 100 mg, or 100 and 200 mg (<math>P</math> values not reported).</p> <p>Nitroglycerin consumption declined in a parallel, dose-related fashion. Compared to placebo, all doses of atenolol decreased nitroglycerin consumption significantly (<math>P &lt; 0.001</math>), with no significant difference</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>placebo</p> <p>All patients received SB placebo for the first 4 weeks of the trial.</p>				<p>between 50 vs 100 and 200 mg, or 100 vs 200 mg (P values not reported).</p> <p>All doses of atenolol significantly reduced resting and exercise heart rate at three hours (P&lt;0.001) and 24 hours (P&lt;0.001) after ingestion. Atenolol was significantly more effective at 100 and 200 mg, with no significant difference between the two doses (P value not reported). The maximal exercise double product (heart rate times SBP) at the occurrence of chest pain was significantly reduced at peak and trough testing with all atenolol doses (P&lt;0.001 for all), but 100 and 200 mg were significantly more effective than 25 and 50 mg (P&lt;0.001 for both). The amount of exercise necessary to produce angina three hours after drug ingestion was increased by all atenolol doses; however, only 50 (P&lt;0.001), 100 (P&lt;0.005) and 200 mg (P&lt;0.001) showed significant improvement compared to placebo.</p> <p>Secondary: Not reported</p>
<p>Oh et al.<sup>59</sup> (2016)</p> <p>Atenolol 25 mg BID</p> <p>vs</p> <p>carvedilol 12.5 mg BID</p> <p>After a week of treatment, the initial dose of the study medication could be doubled</p>	<p>OL, PG, PRO, RCT</p> <p>Patients 20 to 80 years of age with stable angina pectoris who had a positive exercise treadmill test according to American College of Cardiology Foundation and American Heart Association guidelines</p>	<p>N=89</p> <p>6 months</p>	<p>Primary: BP, heart rate, treadmill exercise test, Seattle Angina Questionnaire, metabolic parameters</p> <p>Secondary: Not reported</p>	<p>Primary: Office SBP and DBP at baseline was similar between the two groups and had not changed at the end of the study. Both carvedilol and atenolol significantly reduced heart rate from baseline (76 ± 11 to 66 ± 9 beat/min, P&lt;0.001; 74 ± 9 to 64 ± 9 beat/min, P&lt;0.001, respectively). Improvement of time to ST-segment depression during the treadmill exercise and the Seattle Angina Questionnaire scores for angina stability and frequency after six months of treatment were similar between groups. There was no significant change from baseline in the level of fasting glucose, insulin, or glycated hemoglobin in either group. However, TC and LDL-C levels significantly reduced to a greater extent with carvedilol than with atenolol (-23 vs -10 and -38 vs -24%, respectively, P&lt;0.05 for both), although the rate of statin use was comparable. No changes were seen in HDL cholesterol and triglyceride levels after six months of treatment in both groups compared with baseline.</p> <p>Secondary: Not reported</p>
<p>Kardas et al.<sup>60</sup> (2007)</p>	<p>OL, PG, RCT</p> <p>Patients 40 to 75</p>	<p>N=112</p> <p>8 weeks</p>	<p>Primary: Overall compliance</p>	<p>Primary: The overall compliance significantly higher in the betaxolol group compared to the metoprolol group (86.5±21.3 vs 76.1±26.3%,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Betaxolol 20 mg QD  vs  metoprolol 50 mg BID	years with ischemic heart disease NYHA class I to II, no prior $\beta$ -blocker treatment, and whose mental state enabled conscious participation in the study		Secondary: Drug effectiveness, health-related QOL	respectively; P=0.002).  Secondary: There was not a significant difference in chest pain episodes observed between the betaxolol and metoprolol groups compared from baseline (0.42/week and 0.46/week change in episodes, respectively; P>0.05).  Overall, QOL dimensions were similar among both treatment groups, with the exception of physical function in which a significantly greater improvement was observed in the betaxolol group compared to the metoprolol group (42.9 vs 15.2 patients improved, respectively; P<0.01).
van der Does et al. <sup>61</sup> (1999)  Carvedilol 25 to 50 mg BID  vs  metoprolol 50 to 100 mg BID	DB, MC, RCT  Patients $\leq$ 80 years of age with CHD and chronic stable angina for $\geq$ 2 months, exertional angina with symptoms improving after taking short acting nitrates or after a period of rest, and 1 exercise test performed that was limited by moderate anginal pain	N=368  3 months	Primary: Moderate anginal pain and time to ST- 1-mm segment depression  Secondary: Not reported	Primary: Compared to baseline, both carvedilol and metoprolol significantly decreased time to anginal pain during exercise test (+77s [+20 to +140] and +76 [+25 to +155], respectively; P<0.001 for both).  Compared to baseline, both carvedilol and metoprolol significantly decreased time to ST- 1-mm segment depression during exercise test (+75.5 s [+47 to +154 s] and +60 [0 to +146 s], respectively; P<0.001 for both).  Carvedilol significantly improved the time to 1-mm ST-segment depression compared to metoprolol (RR, 1.386; 95% CI, 1.045 to 1.839; P<0.05)  Secondary: Not reported
Weiss et al. <sup>62</sup> (1998)  Carvedilol 12.5 to 50 mg BID  vs  placebo	DB, MC, XO  Patients with 2 stress tests which evoked ischemic signs and symptoms	N=122  12 weeks	Primary: Efficacy, safety  Secondary: Not reported	Primary: The carvedilol 25 and 50 mg groups significantly reduced the time to angina compared to placebo (25 mg: 337 s, P=0.0039; 50 mg: 345 s; P<0.001 vs 316 s).  The carvedilol 25 and 50 mg groups significantly reduced the time to 1-mm ST-segment depression compared to placebo (25 mg: 313 s; 50 mg: 323 s vs 301 s; P<0.0001 for both).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The percentage of patients reporting any adverse experience was slightly less in those receiving placebo (placebo: 28.4%; 12.5 mg: 33.1%; 25 mg: 34.5%; 50 mg: 31.9%). Adverse events included dizziness, fatigue, headache, dyspepsia, and any hypotensive event. The clinical significance of the adverse events was not reported.</p> <p>Secondary: Not reported</p>
<p>Hauf-Zachariou et al.<sup>63</sup> (1997)</p> <p>Carvedilol 25 mg BID</p> <p>vs</p> <p>verapamil 120 mg TID</p>	<p>DB, MC, PG, RCT</p> <p>Patients 18 to 75 years with a confirmed diagnosis of CAD, exertional chest pain relieved by rest or glyceryl trinitrate for <math>\geq 2</math> months and 2 exercise tests with signs and symptoms of ischemia</p>	<p>N=313</p> <p>12 weeks</p>	<p>Primary: Total exercise time, time to onset of angina, and time to 1 mm ST-segment depression, blood pressure, heart rate, rate pressure product</p> <p>Secondary: Not reported</p>	<p>Primary: There was not a significant difference in total exercise time observed between the carvedilol (increased from 378 s to 436 s) and verapamil (increased from 386 s to 438 s) groups (RR, 1.14; 90% CI, 0.85±1.52).</p> <p>There was not a significant difference observed between the carvedilol and verapamil groups in time to onset of angina (increase from 296 s to 325 s vs 285 s to 326 s) and in time to 1 mm ST-segment depression (increase from 267 s to 298 s vs 286 s to 302 s).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced SBP (from 175 to 166 mm Hg) compared to verapamil (from 173 to 173 mm Hg).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced heart rate (from 123 to 112 mm Hg) compared to verapamil (from 124 to 120 mm Hg).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced rate pressure product (from 21564 to 18802 mm Hg) compared to verapamil (from 21488 to 20992 mm Hg).</p> <p>Secondary: Not reported</p>
<p>Savanitto et al.<sup>64</sup> (1996)</p> <p>Weeks 1 to 6: Metoprolol ER</p>	<p>DB, MC, RCT</p> <p>Patients with typical anginal symptoms that had been stable</p>	<p>N=280</p> <p>6 weeks</p>	<p>Primary: Angina frequency, exercise tolerance, safety</p>	<p>Primary: At week six, both metoprolol (mean change, -1.95; 95 % CI, -1.25 to -2.64) and nifedipine (mean change, -1.57; 95 % CI, -0.69 to -2.45) significantly reduced the frequency of angina compared to baseline, but there was not a statistical difference between groups. At the end of 10</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>200 mg QD</p> <p>vs</p> <p>nifedipine 20 mg BID</p> <p>Weeks 7 to 10: Metoprolol ER 200 mg QD plus placebo</p> <p>vs</p> <p>metoprolol ER 200 mg QD and nifedipine 20 mg BID</p> <p>vs</p> <p>nifedipine 20 mg BID plus placebo</p>	<p>for approximately 6 months, who showed a positive response to exercise stress testing with 23 min of exercise tolerance and were in sinus rhythm and had an analyzable ST segment on ECG</p>		<p>Secondary: Not reported</p>	<p>weeks, there was not a statistical difference observed between the groups.</p> <p>At week six, both metoprolol and nifedipine significantly increased the mean exercise time to 1-mm ST-segment depression compared to baseline (both <math>P&lt;0.01</math>); but metoprolol was significantly more effective than nifedipine (<math>P&lt;0.05</math>).</p> <p>At week 10, the groups randomized to combination therapy had a further increase in time to 1-mm ST-segment depression (<math>P&lt;0.05</math> vs placebo).</p> <p>There were 14 cardiovascular events including one sudden death, three acute myocardial infarctions, eight cases of unstable angina, one of syncope and one of stroke and the incidence of these events did not differ among the treatment groups.</p> <p>Secondary: Not reported</p>
<p>Turner et al.<sup>65</sup> (1978)</p> <p>Propranolol 40 to 240 mg/day, administered in 4 divided doses</p> <p>vs</p> <p>nadolol 40 to 240 mg/day, administered in 2 divided doses</p>	<p>DB, PC, RCT, XO</p> <p>Men with ischemic heart disease with presence of stable angina pectoris and absence of acute MI during the preceding 4 months, ECG evidence of myocardial ischemia during treadmill exercise testing and/or</p>	<p>N=14</p> <p>Up to 18 weeks</p>	<p>Primary: Glyceryl trinitrate consumption, exercise tolerance, heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Mean glyceryl trinitrate consumption decreased significantly from placebo with both propranolol and nadolol (<math>P&lt;0.05</math> for all). There was no significant difference between propranolol and nadolol, with nadolol 240 mg/day producing a significant decrease in consumption of glyceryl trinitrate compared to 160 mg/day (<math>P&lt;0.05</math>).</p> <p>Both treatments resulted in similar improvements in exercise tolerance (30%; <math>P&lt;0.01</math>) and external work performed (48%; <math>P&lt;0.01</math>).</p> <p>A slightly greater suppression of heart rate during exercise was observed with nadolol compared to propranolol (<math>P&lt;0.05</math>).</p> <p>Both treatments resulted in significant decreases in resting heart rate;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo	arteriographic evidence of >60% obstruction of the lumen of $\geq 2$ major coronary arteries, the absence of CHF, a resting DBP <90 mm Hg, absence of contra- indications to $\beta$ -blocker therapy and the absence of other cardiac or severe systemic disease			<p>however, the rate corrected systolic time intervals changed very little from control.</p> <p>The effects of the two treatments could not be differentiated by echocardiography or phonocardiography.</p> <p>Secondary: Not reported</p>
<b>Arrhythmias</b>				
Lui et al. <sup>66</sup> (1983)  Acebutolol 200 or 400 mg/day  vs placebo	DB, PC, RCT, XO  Adult patients with $\geq 30$ ventricular ectopic beats per hour on 3 control ambulatory monitoring	N=25  Not reported	<p>Primary: Resting heart rate, ventricular arrhythmias, paired ventricular ectopic beats, ventricular tachycardia, electro-physiologic effects, adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: Both doses of acebutolol produced a significant decrease in heart rate (P&lt;0.01 for both), with no significant differences between 200 and 400 mg (P value not reported).</p> <p>Mean ventricular ectopic beat reduction from the control period was 34.9% during the two placebo periods. Following acebutolol, mean ectopic beat suppression was greater, although not significantly different when compared to placebo, at 44.9 and 49.5% using 200 and 400 mg, respectively (P values not reported).</p> <p>Nineteen of the 25 patients achieved episodes of paired ventricular ectopic beats (couplets) on control ambulatory monitoring. The mean reduction of paired beats was significantly higher than placebo (48.8%) with 70.5 (P&lt;0.05) and 74.5% (P&lt;0.01) with acebutolol 200 and 400 mg, respectively.</p> <p>Five patients who had ventricular tachycardia during both control and placebo periods had complete suppression during acebutolol treatment.</p> <p>Mean QRS and QTc intervals revealed no significant difference as compared to the control period.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>There were no significant adverse effects related to acebutolol administration. Patients did not develop any bronchospasm, significant bradycardia, heart block, CHF or any central nervous system adverse effect.</p> <p>Secondary: Not reported</p>
<p>Lee et al.<sup>67</sup> (2008)</p> <p>Amiodarone</p> <p>vs</p> <p>sotalol</p> <p>vs</p> <p>β-blockers (agents not specified)</p> <p>Doses of the agents were not specified</p>	<p>RETRO</p> <p>Patients with AF and/or CHF (NYHA class ≥III) and an implantable cardioverter defibrillator</p>	<p>N=55</p> <p>2.6 years</p>	<p>Primary: Cumulative rates of inappropriate shocks</p> <p>Secondary: Not reported</p>	<p>Primary: Amiodarone demonstrated a significantly lower rate of inappropriate shock was compared β-blocker group (27.3 vs 70.6% at four years; P=0.003). This demonstrated an 83% reduction compared to the β-blockers (HR, 0.17; 95% CI, 0.05 to 0.64; P=0.008).</p> <p>There was not a significant difference in rates of inappropriate shocks observed between the amiodarone and sotalol groups (27.3 vs 54.3% at four years; P=0.29).</p> <p>There was not a significant difference in rates of inappropriate shocks observed between the sotalol and β-blocker groups (54.3 vs 70.6% at four years; P=0.16).</p> <p>Secondary: Not reported</p>
<p>Connolly et al.<sup>68</sup> (2006)</p> <p>OPTIC</p> <p>β-blocker (bisoprolol, carvedilol or metoprolol)</p> <p>vs</p> <p>sotalol 240 mg/day</p>	<p>DB, MC, RCT</p> <p>Patients who received an implantable cardioverter defibrillator within 21 days of randomization, had sustained ventricular tachycardia,</p>	<p>N=412</p> <p>12 months</p>	<p>Primary: Implantable cardioverter defibrillator shock for any reason</p> <p>Secondary: Not reported</p>	<p>Primary: Shocks occurred in 41 patients (38.5%) in the β-blocker group, 26 (24.3%) in the sotalol group, and 12 (10.3%) in the amiodarone plus β-blocker group.</p> <p>A reduction in the risk of shock was observed with use of amiodarone plus β-blocker or sotalol vs β-blocker alone (HR, 0.44; 95% CI, 0.28 to 0.68; P&lt;0.001).</p> <p>The amiodarone plus β-blocker group significantly reduced the risk of shock compared to β-blocker alone (HR, 0.27; 95% CI, 0.14 to 0.52; P&lt;0.001) and sotalol (HR, 0.43; 95% CI, 0.22 to 0.85; P=0.02).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>in two to three divided doses</p> <p>vs</p> <p>amiodarone 200 mg QD plus <math>\beta</math>-blocker (bisoprolol, carvedilol or metoprolol)</p> <p>Amiodarone was loaded at 400 mg BID for 2 weeks, followed by 400 mg/day for 4 weeks, and then 200 mg/day until then end of the study</p>	<p>ventricular fibrillation or cardiac arrest, LVEF <math>\leq</math>40%, inducible ventricular tachycardia or ventricular fibrillation by programmed ventricular stimulation with LVEF <math>\leq</math>40% or unexplained syncope with ventricular tachycardia or ventricular fibrillation, inducible by programmed stimulation</p>			<p>Sotalol did not significantly reduce the risk of shock compared to the <math>\beta</math>-blocker alone group (HR, 0.61; 95% CI, 0.37 to 1.01; P=0.055).</p> <p>Secondary: Not reported</p>
<p>Balcetyte-Harris et al.<sup>69</sup> (2002)</p> <p>Esmolol 0.5 mg/kg over 5 minutes then 0.05 mg/kg/min titrated to heart rate of 55 to 65 bpm and SBP &gt;100 mm Hg for up to 24 hours</p> <p>vs</p>	<p>OL, RCT</p> <p>Patients referred for elective CABG without concomitant valve replacement who were in sinus rhythm</p>	<p>N=50</p> <p>72 hours</p>	<p>Primary: Development of AF lasting &gt;30 mins</p> <p>Secondary: Development of adverse events, hypotension (SBP &lt;90 mm Hg), symptomatic bradycardia or CHF (left ventricular failure)</p>	<p>Primary: There was not a significant difference in development of AF after CABG between the esmolol and <math>\beta</math>-blocker group (seven [26%] vs six [26%] patients, respectively).</p> <p>Secondary: Significantly more patients in the esmolol group experienced significant adverse events compared to the patients in the <math>\beta</math>-blocker group (11 [41%] vs one [4%] patient(s), respectively; P=0.006).</p> <p>Significantly more patients in the esmolol group experienced hypotension compared to the patients in the <math>\beta</math>-blocker group (eight vs one patient(s), respectively; P=0.03).</p> <p>There was not a statistically significant difference between the esmolol</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
oral $\beta$ -blocker (metoprolol $\geq$ 50 mg/day was the preferred agent)				and the $\beta$ -blocker group in the development bradycardia requiring pacing (two vs zero patients, respectively) and in left ventricular failure (one vs zero patient(s), respectively).
Kettering et al. <sup>70</sup> (2002)  Metoprolol 25 to 200 mg/day  vs  sotalol 40 to 480 mg/day	PRO, RCT  Symptomatic patients between 18 and 80 years with sustained ventricular tachycardia and/or ventricular fibrillation requiring an implantable cardioverter defibrillator	N=100  2 years	Primary: Ventricular tachycardia or ventricular fibrillation recurrence requiring implantable cardioverter defibrillator intervention  Secondary: Total mortality	Primary: There was not a significant difference in ventricular tachycardia/ventricular fibrillation recurrence rates observed between the metoprolol group (33 patients) and the sotalol group (30 patients; P=0.68).  After one year of treatment, 46.3% of patients in the metoprolol group and 54.7% of patients in the sotalol group were free of a recurrence of ventricular tachycardia or ventricular fibrillation (P=0.68). After two years, rates were 31.5 and 36.6%, respectively.  Secondary: There was not a significant difference in mortality rates observed between the metoprolol group (eight deaths) and the sotalol group (six patients; P=0.43).
Seidl et al. <sup>71</sup> (1998)  Metoprolol 50 mg/day  vs  sotalol 80 mg/day  The doses of the study medications were titrated to the maximum titrates dose.	OL, RCT  Patients >18 years of age requiring treatment if life-threatening ventricular tachycardia/ventricular fibrillation who required an implantable cardioverter defibrillator due to non-inducible or drug refractory ( $\geq$ 1 unsuccessful antiarrhythmic trial) arrhythmias	N=70  26 $\pm$ 16 months	Primary: Recurrence of ventricular tachycardia requiring antitachycardia pacing, fast ventricular tachycardia or ventricular fibrillation requiring implantable cardioverter defibrillator, discharges, total mortality  Secondary:	Primary: Actuarial rates for absence of ventricular tachycardia recurrence were significantly higher in the metoprolol group vs the sotalol group at one and two years (83 and 80 vs 57 and 51%, respectively; P=0.016).  Actuarial rates for absence of recurrence of a fast ventricular tachycardia or ventricular fibrillation were significantly higher in the metoprolol group vs the sotalol group one and two years (88 and 80 vs 54 and 46%, respectively; P=0.002)  Actuarial survival rates at one and two years were not significantly different between the metoprolol and sotalol groups (94 and 91 vs 86 and 83%, respectively; P=0.287)  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Not reported	
Steeds et al. <sup>72</sup> (1999)  Sotalol 80 mg BID  vs  atenolol 50 mg QD	OL, PRO, RCT, XO  Symptomatic patients >50 years of age with paroxysmal AF documented on ECG	N=47  2 months	Primary: Frequency of paroxysmal AF  Secondary: Average and total duration of paroxysmal AF, total ectopic count, symptom assessments	Primary: There was not a significant difference in frequency of episodes of paroxysmal AF observed between the sotalol and atenolol groups (median difference, 0 min; 95% CI, 0 to 1; P=0.47).  Secondary: There was not a significant difference in average duration of episodes of paroxysmal AF observed between the sotalol and atenolol groups (median difference, 0 min; 95% CI, 0 to 1 min; P=0.31) or in total duration of episodes of paroxysmal AF (median difference, 0 min; 95% CI, -1 to 2 min; P=0.51).  There was not a significant difference in total ectopic count observed between the sotalol and atenolol groups (median difference, -123; 95% CI, -362 to 135; P=0.14) during either treatment period.  There was not a significant difference in tolerance and symptom scores observed between the sotalol and atenolol groups (median difference, -5; 95% CI, -20 to 5; P=0.26)
<b><i>Essential Tremor</i></b>				
Calzetti et al. <sup>73</sup> (1981)  Metoprolol 150 mg/dose  vs  propranolol 120 mg/dose  vs  placebo	DB, PC, RCT  Patients 19 to 72 years with essential tremor and symptomatic for ≥1 year prior to the study	N=23  3 weeks	Primary: Tremor magnitude, heart rate, blood pressure  Secondary: Not reported	Primary: Both metoprolol (47±9.7%) and propranolol (55±5.0%) significantly decreased tremor magnitude from baseline compared to placebo (22±7.3%; P<0.01 for both treatments compared to placebo), but there was not a significant difference observed between the metoprolol and propranolol groups.  Both propranolol (0.073) and metoprolol (0.01) significantly diminished the normal increase in pulse rate on standing (P<0.01) and placebo had no effect on such pulse rate. There was not a significant difference observed between the metoprolol and propranolol groups.  Both metoprolol and propranolol significantly reduced the SBP from baseline compared to placebo, in the supine and standing positions (P<0.05).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Yetimalar et al. <sup>74</sup> (2005)  Propranolol 120 mg/day  vs  olanzapine 20 mg/day	DB, RCT, XO  Patients with essential tremor and previous therapy with $\geq 1$ medications for essential tremor without significant benefit, which was withdrawn $\geq 1$ month before study drug was given	N=38  74 days	Primary: Tremor, global QOL  Secondary: Not reported	Primary: After 30 days, both propranolol and olanzapine significantly reduced the all tremor evaluation measures (i.e., speaking, eating, dressing, writing working) compared to baseline (P=0.000), but at the end of the study, olanzapine significantly improved all tremor evaluation measures (P<0.05) except hygiene (P =0.08) as compared to propranolol.  Both propranolol (63%) and olanzapine (87%) significantly improved global QOL from baseline, but olanzapine significantly improved the global QOL score compared to propranolol (4.5 $\pm$ 0.7 vs 3.6 $\pm$ 0.9; P=0.000).  Secondary: Not reported
Gironell et al. <sup>75</sup> (1999)  Propranolol 40 mg TID  vs  gabapentin 400 mg TID  vs  placebo	DB, PC, XO  Patients with moderate to severe essential tremor that was chronic ( $\geq 5$ years), persistent, and bilateral postural tremor with or without kinetic tremor involving hands or forearms, with no other neurological abnormalities or explanation for tremor	N=16  66 days	Primary: Tremor Clinical Rating Scale, accelerometric recordings, self-reported disability scale  Secondary: Not reported	Primary: Both gabapentin and propranolol significantly reduced the clinical examination and motor task performance components of the Tremor Clinical Rating Scale compared to placebo (-3.10 $\pm$ 1.10; P=0.01 and -4.50 $\pm$ 1.10; P=0.001, respectively), and significant differences were not observed between the gabapentin and propranolol groups (1.40 $\pm$ 1.16; P=0.23).  Both gabapentin and propranolol significantly reduced the activities of daily living component of the Tremor Clinical Rating Scale compared to placebo (-3.03 $\pm$ 1.46; P<0.05 and -4.95 $\pm$ 1.46; P=0.002, respectively), and significant differences were not observed between the gabapentin and propranolol groups (1.92 $\pm$ 1.46; P=0.20).  Both gabapentin and propranolol significantly reduced the patient's subjective assessment of the Tremor Clinical Rating Scale compared to placebo (1.37 $\pm$ 0.46; P=0.006 and 1.44 $\pm$ 0.46; P=0.004, respectively). Significant differences were not observed between the gabapentin and the propranolol groups (-0.07 $\pm$ 0.46; P=0.89).  Both gabapentin and propranolol significantly reduced the absolute power of the dominant frequency peak of accelerometry compared to placebo

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>(-2352.0±1153.3; P=0.05 and -2282.14±1116.58; P=0.05, respectively), but significant differences were not observed between the gabapentin and the propranolol groups (-70.39±1165.22; P=0.95).</p> <p>Gabapentin significantly reduced the self-reported disability scale score more than placebo (-6.04±2.75; P=0.04) and propranolol did not (-4.48±2.75; P=0.11), but there were no significant differences between the gabapentin and propranolol groups (-1.55±2.75; P=0.58).</p> <p>Secondary: Not reported</p>
<b>Heart Failure</b>				
<p>CIBIS Investigators and Committees<sup>76</sup> (1994) CIBIS</p> <p>Bisoprolol 1.25 to 5 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patient received standard therapy (diuretic and vasodilator)</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients 18 to 75 years with NYHA functional class III or IV due to idiopathic dilated cardiomyopathy, ischemia, HTN or valvular heart disease, a LVEF of &lt;40%, and background therapy with a diuretic and a vasodilator</p>	<p>N=641</p> <p>1.9 years</p>	<p>Primary: Total mortality</p> <p>Secondary: Tolerability, analysis critical events</p>	<p>Primary: There was no statistical significance between bisoprolol and placebo in total mortality (53 vs 67; RR, 0.80; 95% CI, 0.56 to 1.15; P=0.22).</p> <p>Secondary: Bisoprolol was well tolerated with no between group difference in premature treatment withdrawals (82 on placebo, 75 on bisoprolol; not significant).</p> <p>Significantly fewer patients in the bisoprolol group required hospitalization for cardiac decompensation (90 in placebo versus 61 in bisoprolol; P&lt;0.01), and more patients improved by at least one NYHA functional class (48 on placebo versus 68 on bisoprolol; P=0.04) by the end of follow-up period.</p>
<p>CIBIS-II Investigators and Committees<sup>77</sup> (1999) CIBIS-II</p> <p>Bisoprolol 1.25 to 10 mg QD added</p>	<p>DB, MC, PC, RCT</p> <p>Symptomatic patients 18 to 80 years in NYHA class III or IV, with LVEF of 35% or less receiving</p>	<p>N=2,647</p> <p>1.3 years</p>	<p>Primary: All-cause mortality</p> <p>Secondary: All-cause hospital admissions, cardiovascular mortality,</p>	<p>Primary: CIBIS-II was stopped early, after the second interim analysis, because bisoprolol showed a significant mortality benefit. All-cause mortality was significantly lower with bisoprolol than on placebo (156 [11.8%] vs 228 [17.3%] deaths, respectively; HR, 0.66; 95% CI, 0.54 to 0.81; P&lt;0.0001).</p> <p>Significantly fewer sudden deaths among patients on bisoprolol than in those on placebo (48 [3.6%] vs 83 [6.3%] deaths, respectively; HR, 0.56;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>to usual therapy (diuretic and vasodilator)</p> <p>vs</p> <p>placebo</p>	<p>standard therapy with diuretics and ACE inhibitor or other vasodilator</p>		<p>cardiovascular mortality and cardiovascular hospital admissions (composite endpoint), permanent premature treatment withdrawals</p>	<p>95% CI, 0.39 to 0.80; P=0.0011).</p> <p>Secondary: All-cause hospital admissions was significantly lower with bisoprolol than on placebo (440 [33%] vs 513 [39%] patients, respectively; HR, 0.80; 95% CI, 0.71 to 0.91; P=0.0006).</p> <p>All-cardiovascular deaths was significantly lower with bisoprolol than on placebo (119 [9%] vs 161 [12%] patients, respectively; HR, 0.71; 95% CI, 0.56 to 0.90; P=0.0049).</p> <p>Occurrence of composite endpoints of all cardiovascular deaths and cardiovascular admissions was significantly lower with bisoprolol than on placebo (388 [29%] vs 463 [35%] patients, respectively; HR, 0.79; 95% CI, 0.69 to 0.90; P=0.0004).</p> <p>Occurrence of treatment withdrawals was not statistically different between bisoprolol and the placebo group (194 [15%] vs 192 [15%] patients, respectively; HR, 1.00; 95% CI, 0.82 to 1.22; P=0.98).</p>
<p>Contini et al.<sup>78</sup> (2013) CARNEBI</p> <p>Bisoprolol</p> <p>vs</p> <p>carvedilol</p> <p>vs</p> <p>nebivolol</p> <p>each at maximal clinically tolerated dose</p>	<p>RCT, XO</p> <p>Patients aged 18 to 80 years with diagnosis of either idiopathic or ischemic dilated cardiomyopathy, previous evidence of LVEF ≤ 40%, NYHA class I to III with stable clinical conditions and optimized drug regimen</p>	<p>N=61</p> <p>Each patient performed a 2-month therapy with each β-blocker</p>	<p>Primary: Clinical conditions, quality of life, laboratory data, echocardiographic evaluation, spirometry, alveolar capillary membrane diffusion, chemoreceptor response, cardiopulmonary exercise test, and response to hypoxia during constant workload exercise</p>	<p>Primary: Clinical conditions, NYHA class, Minnesota questionnaire, renal function, hemoglobin concentration, brain natriuretic peptide, Echocardiographic data, and Doppler data were unaffected by the different β-blockers studied.</p> <p>Carbon monoxide diffusing capacity was lower on Carvedilol (18.3 ± 4.8* mL/min/mm Hg) compared to Nebivolol (19.9 ± 5.1) and Bisoprolol (20.0 ± 5.0) due to membrane diffusion 20% reduction (*= P&lt; 0.0001). Constant workload exercise showed in hypoxia a faster VO<sub>2</sub> (oxygen uptake) kinetic and a lower ventilation with Carvedilol. Peripheral and central sensitivity to CO<sub>2</sub> was lower in Carvedilol while response to hypoxia was higher in Bisoprolol.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Willenheimer et al.<sup>79</sup> (2005) CIBIS-III</p> <p>Bisoprolol 1.25 to 10 mg QD</p> <p>vs</p> <p>enalapril 2.5 to 10 mg BID</p>	<p>BE, MC, OL, PG, RCT</p> <p>Patients ≥65 years with stable mild to moderate CHF (NYHA class II to III), LVEF of ≤35% ≥3 months prior to randomization, not on an ACE inhibitor, β-blocker or ARB therapy and no clinically relevant fluid retention of diuretic adjustment within the 7 days prior to randomization</p>	<p>N=1,010</p> <p>1.22±0.42 years</p>	<p>Secondary: Not reported</p> <p>Primary: Combined all-cause mortality or hospitalization</p> <p>Secondary: Combined end point at the end of the monotherapy phase and the individual components of the primary end point, cardiovascular death and cardiovascular hospitalization, permanent treatment cessation and the need for early introduction of the second drug as indicators of drug tolerability</p>	<p>Primary: There were 178 patients (35.2%) with a primary end point of combined all-cause mortality or all-cause hospitalization in the bisoprolol-first group, compared to 186 (36.8%) patients in the enalapril-first group (absolute difference, -1.6%; 95% CI, -7.6 to 4.4; HR, 0.94; 95% CI, 0.77 to 1.16; non-inferiority for bisoprolol-first vs enalapril-first treatment; P=0.019).</p> <p>Secondary: The combined endpoint at the end of the monotherapy phase occurred in 109 patients in the bisoprolol-first group compared to 108 patients in the enalapril-first group (HR, 1.02; 95% CI, 0.78 to 1.33; between-group difference P=0.90); 23 vs 32 patients died, respectively (HR, 0.72; 95% CI, 0.42 to 1.24; between-group difference P=0.24); and 99 vs 92 patients had been a hospitalization, respectively (HR, 1.08; 95% CI, 0.81 to 1.43; between-group difference P=0.59).</p> <p>There were 65 deaths in the bisoprolol-first group, as compared to 73 in the enalapril-first group (HR, 0.88; 95% CI, 0.63 to 1.22; between-group difference P=0.44).</p> <p>In the bisoprolol-first group, 151 patients were hospitalized, compared to 157 patients in the enalapril-first group (HR, 0.95; 95% CI, 0.76 to 1.19; between-group difference P=0.66).</p> <p>There was not a significant difference in cardiovascular death rate observed between the bisoprolol-first (55) and enalapril-first (56) treatment groups (HR, 0.97; 95% CI, 0.67 to 1.40; between-group difference P=0.86).</p> <p>During the monotherapy phase, 35 (6.9%) patients in the bisoprolol-first group permanently discontinued therapy, compared to 49 (9.7%) patients in the enalapril-first group. During the combined-therapy phase, 19 patients (4.2%) in the bisoprolol-first group permanently discontinued</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>bisoprolol therapy and 47 (10.4%) discontinued enalapril therapy. In the enalapril-first group, 24 patients (5.5%) permanently discontinued bisoprolol and 16 (3.7%) discontinued enalapril.</p> <p>There was not a statistical significant difference observed in the early introduction of the second drug between the bisoprolol-first group (39 [7.7%] patients) compared to the enalapril-first group (37 [7.3%] patients; P=0.81).</p>
<p>Packer et al.<sup>80</sup> (2001) COPERNICUS</p> <p>Carvedilol 3.125 to 25 mg BID</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients with severe chronic heart failure as a result of ischemic or nonischemic cardiomyopathy, dyspnea or fatigue at rest or on minimal exertion for <math>\geq 2</math> months and a LVEF <math>&lt; 25\%</math> despite appropriate conventional therapy with diuretics, and an ACE inhibitor, or ARB</p>	<p>N=2,280</p> <p>10.4 months</p>	<p>Primary: Total mortality</p> <p>Secondary: Combined risk of death or hospitalization for any reason, withdrawal rates</p>	<p>Primary: The study was stopped early due to statistical significance.</p> <p>The annual mortality in the placebo group was 19.7% (190) versus 12.8% (130 deaths) in the carvedilol group, a 35% reduction in mortality (95% CI, 19 to 48%; P&lt;0.00013).</p> <p>Secondary: Carvedilol reduced the combined risk of death or hospitalization for any reason by 24% compared to placebo (425 vs 507 patients; 95% CI, 13 to 33%; P&lt;0.001)</p> <p>Withdrawal rates were significantly higher in the placebo group compared to the carvedilol group (18.5 vs 14.8; P=0.02).</p>
<p>Packer et al.<sup>81</sup> (2002) COPERNICUS</p> <p>Carvedilol 3.125 mg BID, titrated up to 25 mg BID</p> <p>vs</p>	<p>DB, PC, RCT</p> <p>Patients with dyspnea or fatigue at rest or on minimal exertion for <math>\geq 2</math> months and a LVEF <math>&lt; 25\%</math> as a result of an</p>	<p>N=2,289</p> <p>10.4 months</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Combined risk of death or hospitalization for any reason, combined risk of</p>	<p>Primary: The annual mortality rate with placebo was 19.7% per patient year of follow up, which was reduced to 12.8% by treatment with carvedilol, corresponding to a 35% reduction in the risk of death (P=0.00013).</p> <p>Secondary: Carvedilol reduced the risk of death or any hospitalization by 24% (P=0.00004).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo	ischemic or nonischemic cardiomyopathy, being treated with a diuretic and either an ACE inhibitor or ARB		death or hospitalization for any cardiovascular reason, combined risk of death or hospitalization for heart failure, patient global assessment	<p>Carvedilol reduced the combined risk of death or hospitalization for cardiovascular reason by 27% (P=0.0002) and the combined risk of death or hospitalization for heart failure by 31% (P=0.000004).</p> <p>Patients receiving carvedilol spent 27% fewer days in the hospital for any reason (P=0.005) and 40% fewer days in the hospital for heart failure (P&lt;0.0001).</p> <p>More patients receiving carvedilol felt improved and fewer patients felt worse compared to patients receiving placebo after six months of maintenance therapy (P=0.0009).</p> <p>Patients receiving carvedilol were less likely to experience a serious adverse event (P=0.002), especially worsening heart failure, sudden death, cardiogenic shock or ventricular tachycardia.</p>
Packer et al. <sup>82</sup> (1996)  Carvedilol 3.125 mg BID, titrated up to 50 mg BID  vs  placebo	DB, PC, RCT  Patients with symptoms of heart failure for ≥3 months and an ejection fraction ≤35%, despite ≥2 months of treatment with diuretics and an ACE inhibitor (if tolerated)	N=1,094  6 to 12 months	Primary: All-cause mortality, cardiovascular morbidity  Secondary: Not reported	<p>Primary: Thirty one (7.8%) patients receiving placebo died compared to 22 (3.2%) deaths in patients receiving carvedilol; this difference represents a 65% decrease in the risk of death (95% CI, 39 to 80; P&lt;0.001). Treatment with carvedilol was associated with a large decrease in the risk of dying of progressive heart failure and in the risk of sudden death.</p> <p>Ninety eight (14.1%) patients receiving carvedilol and 78 patients (19.6%) receiving placebo had at least one hospitalization for cardiovascular causes; this difference represents a 27% reduction in the risk of hospitalization (95% CI, 3 to 45; P=0.036).</p> <p>Secondary: Not reported</p>
Dargie et al. <sup>83</sup> (2001) CAPRICORN  Carvedilol 6.25 to 25 mg BID mg  vs	DB, MC, PC, RCT  Patients 18 years and older with a stable MI occurring 3 to 21 days prior to randomization, LVEF ≤40% and	N=1,959  1.3 years	Primary: All-cause mortality, all-cause mortality or cardiovascular hospital admissions	<p>Primary: There was not a significant difference observed between the carvedilol and placebo groups in the combined endpoint of all-cause mortality and hospital admissions due to cardiovascular events (340 [35%] vs 367 [37%], respectively; HR, 0.92; 95% CI, 0.80 to 1.07; P=0.296).</p> <p>All-cause mortality alone was statistically better in the carvedilol group than the placebo group (116 [12%] vs 151 [15%], respectively; HR, 0.77;</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo	ACE inhibitor therapy for $\geq 48$ hours		Secondary: Sudden death, hospital admission for heart failure, recurrent nonfatal MI, all-cause mortality or recurrent nonfatal MI	95% CI, 0.60 to 0.98; P=0.031).  Secondary: There was not a significant difference observed between the carvedilol and placebo groups in sudden death (51 [5%] vs 69 [7%], respectively; HR, 0.74; 95% CI, 0.51 to 1.06; P=0.098) or in hospital admissions for heart failure (118 [12%] vs 138 [14%], respectively; HR, 0.86; 95% CI, 0.67 to 1.09; P=0.215).  The carvedilol group, compared to placebo, experienced significantly lower rates of nonfatal MIs (34 [3%] vs 57 [6%], respectively; HR, 0.59; 95% CI, 0.39 to 0.90; P=0.014) and all-cause mortality or recurrent nonfatal MI (139 [14%] vs 192 [20%], respectively; HR, 0.71; 95% CI, 0.57 to 0.89; P=0.002).
Krum et al. <sup>84</sup> (abstract) (1995)  Carvedilol 25 mg BID  vs  placebo	DB, PC, RCT  Patients with severe chronic HF receiving digitalis, diuretics and an ACE inhibitor (if tolerated)	N=56  14 weeks	Primary: Cardiac performance; symptom score; combined risk of death, worsening heart failure, and life-threatening ventricular tachycardia  Secondary: Not reported	Primary: Compared to placebo, carvedilol improved cardiac performance, as reflected by an increase of LVEF (P=0.005) and stroke volume index (P=0.010), and a decrease in pulmonary wedge pressure (P=0.003), mean right atrial pressure (P=0.002) and systemic vascular resistance (P=0.017).  Compared to placebo, carvedilol improved symptom scores (P=0.002), functional class (P=0.013) and submaximal exercise tolerance (P=0.006).  The combined risk of death, worsening heart failure and life-threatening ventricular tachyarrhythmia was lower with carvedilol compared to placebo (P=0.028).  Carvedilol was associated with more dizziness and advanced heart block.  Secondary: Not reported
Bristow et al. <sup>85</sup> (1996)  Carvedilol 6.25 mg BID	DB, MC, PC, RCT  Symptomatic ( $\geq 3$ months) patients, 18 to 85 years with stable heart failure	N=345  6 months	Primary: Submaximal exercise improvement  Secondary:	Primary: There were no differences on submaximal exercise with any dose compared to placebo. Walk distances between in each group ranged between 300 to 400 m in both the 6-minute and 9-minute walk tests; P=0.50 and P=0.27, respectively).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs carvedilol 12.5 mg BID vs carvedilol 25 mg BID vs placebo</p> <p>All patients remained on their standard medications.</p>	<p>from ischemic or nonischemic dilated cardiomyopathy, an LVEF of <math>\leq 35\%</math>, a 6-minute walk test between 150 to 425 m and on stable doses of diuretics and ACE inhibitors for 2 weeks before baseline testing</p>		<p>Minnesota questionnaire, changes in NYHA functional class, changes in LVEF, hospitalization, changes in signs and symptoms of heart failure, occurrence of adverse clinical experiences, survival</p>	<p>Secondary: There were no significant changes in the overall Minnesota Questionnaire scores incorporating both physical and emotional dimensions (changes from baseline in the placebo and low-, medium-, and high-dose carvedilol groups of -7.3, -7.9, -7.3, and -6.6, respectively; <math>P=0.512</math> in difference from placebo).</p> <p>There were no significant improvements in NYHA functional classes in the carvedilol groups compared to placebo (actual values not reported; <math>P=0.64</math>).</p> <p>Carvedilol treatment resulted in a dose-related significant improvement in LVEF; carvedilol 6.25 mg (~5 ejection fraction units; <math>P&lt;0.005</math>), 12.5 mg (~6 ejection fraction units; <math>P&lt;0.005</math>) and 25 mg (~7.5 ejection fraction units; <math>P&lt;0.0001</math>) compared to placebo (2 ejection fraction unit improvement).</p> <p>The mean number of hospitalizations per patient were significantly reduced in each of the carvedilol groups (~0.1 hospitalizations) compared to placebo (~0.35; <math>P&lt;0.01</math>).</p> <p>Bradycardia was significantly higher in the carvedilol 12.5 mg group (10 [11%]) and the 25 mg group (10 [11%]) compared to placebo (1 [1%]; <math>P&lt;0.05</math>). Also, dizziness was significantly higher in the carvedilol 25 mg group (34 [38%]) compared to the placebo group (19 [23%]; <math>P&lt;0.05</math>). The clinical significance of these advents was not mentioned.</p> <p>There was a dose-related, statistically significant reduction in mortality in the carvedilol-treated groups, with respective mortality rates of 6.0% for the carvedilol 6.25 mg group (RR, 0.356; 95% CI, 0.127 to 0.998; <math>P&lt;0.05</math>), 6.7% for the 12.5 mg group (HR, 0.416; 95% CI, 0.158 to 1.097; <math>P=0.07</math>), and 1.1% in the 25 mg group (HR, 0.067; 95% CI, 0.009 to 0.512; <math>P&lt;0.001</math>) compared to 15.5% mortality in the placebo group.</p> <p>Combining all three carvedilol arms of the study compared to the placebo arm showed statistical significance in all-cause mortality, risk reduced by 73% (<math>P&lt;0.001</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Fröhlich et al. <sup>86</sup> (2015)  Carvedilol with a median dose of 38 mg/day (75% of target dose)  vs  metoprolol succinate with a median dose of 103 mg/day (53% of target dose)	Cohort, PRO  Patients with stable systolic chronic heart failure who were using either carvedilol or metoprolol succinate	N=4,016  Mean follow-up of 52.8±33.6 months	Primary: Mortality  Secondary: Not reported	Primary: In the complete sample, 304 (27.2%) patients died in the carvedilol group and 1,066 (36.8%) in the metoprolol group. In a univariable analysis of the general sample, metoprolol therapy was associated with higher mortality compared with carvedilol therapy (HR, 1.49; 95% CI, 1.31 to 1.69; P<0.001). This difference was not seen after multivariable adjustment (HR, 0.93; 95% CI, 0.57 to 1.50; P=0.75) and adjustment for propensity score and dose equivalents (HR, 1.06; 95% CI, 0.94 to 1.20; P=0.36) or in the propensity and dose equivalent-matched sample (HR, 1.00; 95% CI, 0.82 to 1.23; P=0.99). These results were essentially unchanged for all prespecified subgroups.  Secondary: Not reported
Poole-Wilson et al. <sup>87</sup> (2003) COMET  Carvedilol 25 mg BID  vs  metoprolol 50 mg BID	DB, MC, PG, RCT  Patients with NYHA class II to IV heart failure, admission for a cardiovascular reason in the previous 2 years, an LVEF of <35%, and were stable and optimized with diuretics for ≥2 weeks and ACE inhibitor for ≥4 weeks unless not tolerated	N=3,029  58 months	Primary: All-cause mortality, composite endpoint of mortality or all-cause admission  Secondary: Not reported	Primary: All-cause mortality was significantly lower in the carvedilol group compared to the metoprolol group (512 [34%] vs 600 [40%], respectively; HR, 0.83; 95% CI, 0.74 to 0.93; P=0.0017).  Cardiovascular deaths were significantly lower in the carvedilol group compared to the metoprolol group (438 [29%] vs 534 [35%], respectively; HR, 0.80; 95% CI, 0.70 to 0.90; P=0.0004).  There was not a significant difference in the composite endpoints of all-cause mortality or all-cause admission observed between the carvedilol and metoprolol groups (1,116 [74%] vs 1,160 [76%], respectively; HR, 0.94; 95% CI, 0.86 to 1.02; P=0.122).  Secondary: Not reported
Packer et al. <sup>88</sup> (2001)  Carvedilol 50 to 100 mg/day	MA (19 trials)  Patients with NYHA class II or III and LVEF	N=2,779  8.3 months	Primary: Change in LVEF  Secondary: Not reported	Primary: In the six placebo-controlled trials, metoprolol significantly increased the mean LVEF by 0.063±0.002 compared to the increase with placebo of 0.025±0.001 (difference of 0.038±0.005; P<0.0001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  metoprolol 50 to 150 mg/day or metoprolol ER 150 to 200 mg/day  or  placebo	dysfunction			In the nine placebo-controlled trials, carvedilol significantly increased the mean LVEF by $0.079 \pm 0.001$ compared to the increase with placebo of $0.012 \pm 0.001$ (difference of $0.065 \pm 0.005$ ; $P < 0.0001$ ). Comparing the two agents, carvedilol increased the LVEF significantly greater than metoprolol (difference of $0.026 \pm 0.007$ ; $P = 0.0002$ ).  In the four direct comparator trials, carvedilol significantly increased the mean LVEF by $0.089 \pm 0.002$ compared to the increase with metoprolol of $0.055 \pm 0.002$ (difference of $0.029 \pm 0.011$ ; $P = 0.009$ ).  Secondary: Not reported
Arumanayagam et al. <sup>89</sup> (2001)  Carvedilol 25 mg BID  vs  metoprolol 50 mg BID	DB, RCT  Symptomatic Chinese patients with CHF and LVEF of <45%	N=24  12 weeks	Primary: Plasma total antioxidant status, erythrocyte superoxide dismutase and glutathione peroxidase  Secondary: Not reported	Primary: Neither carvedilol nor metoprolol significantly reduced total antioxidant status activities after 12 weeks of therapy ( $1.65 \pm 0.06$ to $1.68 \pm 0.09$ and $1.44 \pm 0.05$ to $1.51 \pm 0.06$ mmol/L, respectively).  Carvedilol significantly reduced erythrocyte superoxide dismutase activity after 12 weeks of therapy, ( $986 \pm 46$ to $871 \pm 22$ U/g Hb; $P < 0.001$ ), but metoprolol did not ( $790 \pm 43$ to $836 \pm 46$ U/g Hb).  Carvedilol significantly reduced glutathione peroxidase activity after 12 weeks of therapy, ( $145 \pm 7$ to $132 \pm 9$ U/g Hb; $P < 0.05$ ), but metoprolol did not ( $143 \pm 8$ to $138 \pm 9$ U/g Hb).  Secondary: Not reported
Sanderson et al. <sup>90</sup> (1999)  Carvedilol 25 mg BID  vs  metoprolol 50 mg BID	DB, PG, RCT  Symptomatic patients with CHF, LVEF of <45%, and on standard therapy (diuretics, digoxin and ACE inhibitor)	N=51  12 weeks	Primary: Symptom score (QOL questionnaire and NYHA class), exercise tolerance time, LVEF  Secondary: Not reported	Primary: A significant improvement in symptom scores from baseline were experienced in both the carvedilol ( $17.2 \pm 3$ to $8.1 \pm 2$ ; $P < 0.001$ ) and metoprolol ( $13.1 \pm 1.8$ to $4.8 \pm 1.4$ ; $P < 0.001$ ) groups, but there was not a significant difference between the agents.  A significant improvement in NYHA class from baseline were experienced in both the carvedilol ( $2.6 \pm 0.11$ to $2.2 \pm 0.12$ ; $P < 0.001$ ) and metoprolol ( $2.7 \pm 0.09$ to $2.1 \pm 0.09$ ; $P < 0.001$ ) groups, but there was not a significant difference between the agents.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
All patients continued on their standard therapy.				<p>A significant improvement in exercise tolerance time from baseline were experienced in both the carvedilol (1122±51 to 1194±63; P&lt;0.05) and metoprolol (1164±46 to 1263±52; P&lt;0.01) groups, but there was not a significant difference between the agents.</p> <p>A significant improvement in LVEF from baseline were experienced in both the carvedilol (26±1.8 to 35±2.6; P&lt;0.001) and metoprolol (25±1.8 to 31±2.5; P&lt;0.001) groups, but there was not a significant difference between the agents.</p> <p>Secondary: Not reported</p>
<p>Lechat et al.<sup>91</sup> (1998)</p> <p>β-blockers (bisoprolol, bucindolol, carvedilol, metoprolol, and nebivolol)</p> <p>vs</p> <p>placebo</p>	<p>MA (18 trials)</p> <p>Patients with NYHA class I to IV chronic heart failure</p>	<p>N=3,023</p> <p>1.5 to 15 months</p>	<p>Primary: All-cause mortality, hospitalizations due to heart failure, combination of all-cause mortality and hospitalizations for worsened heart failure, changes in functional status, changes in LVEF</p> <p>Secondary: Not reported</p>	<p>Primary: All endpoints showed a significant effect for β-blockers (P&lt;0.05).</p> <p>β-blockers demonstrated a 32% reduction in risk of death compared to placebo (130 vs 156 deaths; 95% CI, 12% to 47%; P=0.003).</p> <p>β-blockers demonstrated a 41% reduction in hospitalizations due to heart failure compared to placebo (166 vs 223 hospitalizations; 95% CI, 26% to 52%; P&lt;0.001).</p> <p>β-blockers demonstrated a 37% reduction in the combination of mortality and morbidity compared to placebo (239 vs 293; 95% CI, 24% to 49%; P&lt;0.001).</p> <p>β-blockers demonstrated a 32% increase in the likelihood of improvement in NYHA class (95% CI, 1% to 74%; P=0.04) and a 30% decrease in the likelihood of worsening NYHA (95% CI, 4% to 50%; P=0.03) compared to placebo</p> <p>β-blockers demonstrated a 29% increase in ejection fraction compared to placebo (0.23±0.04 vs 0.31±0.04; P&lt;10<sup>-9</sup>).</p> <p>β-adrenergic agents did not differ in respect to any outcome measure except that reduction in mortality risk. Beta-selective agents were less</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				robust than the nonselective agents (P=0.049).  Secondary: Not reported
Brophy et al. <sup>92</sup> (2001)  β-blockers (bisoprolol, bucindolol, carvedilol, metoprolol and nebivolol)  vs  placebo	MA (22 trials)  Patients with CHF of various etiologies	N=10,135  3 to 23 months	Primary: Overall mortality, hospitalizations for CHF  Secondary: Not reported	Primary: β-blockers significantly reduced mortality compared to placebo (444 vs 624; OR, 0.65; 95% CI, 0.53 to 0.80).  β-blockers significantly reduced hospitalizations due to CHF compared to placebo (540 vs 754; RR, 0.64; 95% CI, 0.53 to 0.79).  The probability that β-blocker therapy reduced total mortality and hospitalizations for congestive heart failure was almost 100%. The best estimates of these advantages are 3.8 lives saved and four fewer hospitalizations per 100 patients treated in the first year after therapy. The probability that these benefits are clinically significant (>2 lives saved or >2 fewer hospitalizations per 100 patients treated) is 99%.
Whorlow et al. <sup>93</sup> (2000)  β-blockers (bisoprolol, bucindolol, carvedilol metoprolol, nebivolol)  vs  placebo	MA (18 trials)  Patients with NYHA class IV heart failure currently taking background therapy (ACE inhibitors and diuretics with or without digoxin)	N=8,119  3 to 21 months	Primary: Mortality in NYHA class IV patients  Secondary: Not reported	Primary: β-blockers demonstrated a 29% reduction in mortality compared to placebo in patients with NYHA class IV (RR, 0.71; 95% CI, 0.52 to 0.96).  The 29% risk reduction is similar to risk reduction seen with β-adrenergic blockers in other NYHA classes.  β-blockers demonstrated a 32% reduction in mortality compared to placebo in patients with NYHA class I to IV (HR, 0.68; 95% CI, 0.61 to 0.77).  Secondary: Not reported
Bouzamondo et al. <sup>94</sup> (2003)  β-blockers (bisoprolol, bucindolol,	MA  Randomized controlled evaluating patients with heart failure depending on	N=not specified  Duration varied	Primary: Overall mortality, hospitalized for worsening heart failure  Secondary:	Primary: β-blockers reduced overall mortality by 22% compared to placebo (95% CI, 16% to 28%).  β-blockers reduced hospitalizations due to worsening heart failure by 24% compared to placebo (95% CI, 20% to 29%).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
carvedilol, and metoprolol)  vs  placebo	NYHA class		Not reported	Benefits were similar for bisoprolol, metoprolol, and carvedilol regardless of NYHA class.  Secondary: Not reported
Jabbour et al. <sup>95</sup> (2010)  β-blockers (bisoprolol, carvedilol, metoprolol)	OL, XO  Patients with NYHA class I to III heart failure with a subgroup of patients with coexisting COPD	N=51  16 weeks	Primary: Post-bronchodilator FEV <sub>1</sub>  Secondary: Not reported	Primary: FEV <sub>1</sub> was significantly higher in patients receiving bisoprolol vs carvedilol, both in those with coexisting COPD (P<0.01) and without (P=0.02).  There was a significant difference between all patients receiving carvedilol versus those receiving metoprolol (P=0.04), however, when compared for coexisting COPD, there was no difference in FEV <sub>1</sub> .  There was no significant difference for all patients, those with COPD, or those with CHF only when metoprolol and bisoprolol were compared.
MERIT-HF Study Group <sup>96</sup> (1999) MERIT-HF  Metoprolol CR/XL 12.5 mg up to 200 mg QD  vs  placebo	DB, MC, PC, RCT  Symptomatic patients 40 to 80 years in NYHA class II to IV, with LVEF of 40% or less stabilized on standard therapy (diuretic and vasodilator)	N=3,991  1 year	Primary: All-cause mortality, all-cause mortality in combination with all-cause admission to hospital (time to first event)  Secondary: Not reported	Primary: Study was stopped early on the recommendation of the independent safety committee. All-cause mortality was significantly lower in the metoprolol CR/XL group than in the placebo group (145 [7.2%] vs 217 [11.0 %] deaths, RR, 0.66; 95% CI, 0.53 to 0.81; P=0.00009).  There were significantly fewer sudden deaths in the metoprolol CR/XL group than in the placebo group (79 vs 132; RR, 0.59; 95% CI, 0.45 to 0.78; P=0.0002) and deaths from worsening heart failure (30 vs 58; RR, 0.51; 95% CI, 0.33 to 0.79; P=0.0023).  Study drug was permanently stopped early in 13.9% of the patients in the metoprolol CR/XL group and in 15.3% of patients in the placebo group (RR, 0.90; 95% CI, 0.77 to 1.06).  Secondary: Not reported
Goldstein et al. <sup>97</sup> (2001) MERIT-HF	Sub group analysis of MERIT-HF	N=795  1 year	Primary: All-cause mortality,	Primary: There were 45 deaths (11.7% per patient year of follow-up) with metoprolol and 72 deaths (19.1%) with placebo. Metoprolol decreased

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Metoprolol CR/XL 12.5 mg, titrated up to 200 mg QD</p> <p>vs</p> <p>placebo</p>	<p>Patients with NYHA Class III to IV heart failure with LVEF &lt;25%</p>		<p>composite of all-cause mortality and all-cause admission to hospital (time to first event)</p> <p>Secondary: Not reported</p>	<p>total mortality by 39%, sudden death by 45% and death due to worsening heart failure by 55%.</p> <p>Metoprolol also decreased the combined end points of all-cause mortality or all-cause hospitalization by 29%, all-cause mortality or hospitalization for worsening heart failure by 44% and cardiac death or nonfatal MI by 46%.</p> <p>Metoprolol reduced the total number of hospitalizations (all-cause) by 27% (0.709 vs 0.965 per patient year of follow up; P=0.0037).</p> <p>During the up titration phase of the trial, the cumulative numbers of patients hospitalized (all-cause) were: 17 vs 21 after two weeks, 28 vs 30 after four weeks, 39 vs 40 after six weeks, 46 vs 56 after eight weeks and 76 vs 102 after three months. The total number of hospitalizations for cardiovascular causes was reduced by 34% (0.475 vs 0.715 per patient year of follow up; P=0.0005) and for worsening heart failure by 45% (0.273 vs 0.497; P&lt;0.0001).</p> <p>Improvement in NYHA functional class was recorded in 46.2 vs 36.7% of patients receiving metoprolol and placebo (P=0.0031).</p> <p>Secondary: Not reported</p>
<p>Waagstein et al.<sup>98</sup> (1993) MDC</p> <p>Metoprolol 5 mg BID, titrated up to 100 to 150 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients 16 to 75 years of age with symptomatic dilated cardiomyopathy, an ejection fraction &lt;40% and being treated with diuretics, ACE inhibitors and nitrates</p>	<p>N=383</p> <p>18 months</p>	<p>Primary: Combined all-cause mortality and clinical deterioration to a point at which cardiac transplantation would normally be offered as a treatment option</p> <p>Secondary:</p>	<p>Primary: Thirty eight patients receiving placebo reached the primary endpoint compared to 25 patients receiving metoprolol, which corresponded to a risk reduction of 34% (95% CI, -6 to 62; P=0.058).</p> <p>With regard to the individual endpoints, 21 patients met the non-fatal endpoint of need for heart transplantation; two and 19 patients receiving metoprolol and placebo (P=0.0001). During the 12 or 18 months of follow up, all-cause mortality were 23 and 21 patients receiving metoprolol and placebo (P value not reported).</p> <p>Secondary: There was a significantly greater increase in ejection fraction with</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Cardiac function, exercise capacity, QOL, hospital admission or emergency visits for HF treatment	<p>metoprolol compared to placebo by six and 12 months (P value not reported).</p> <p>QOL improved significantly more with metoprolol compared to placebo (P=0.01).</p> <p>With metoprolol, exercise capacity was significantly greater at six and 12 months compared to baseline (P=0.0006 and P=0.0007). With placebo there was a significant improvement from baseline at six months (P=0.007), but not at 12 months (P=0.46). The difference between the two treatments was significant only at 12 months (P=0.046).</p> <p>There was no difference between the treatments in the number of patients readmitted to the hospital (28 vs 20%; P=0.12), but the number of readmissions for all patients in the group was significantly lower with metoprolol (83 vs 51) as was the mean number of readmissions per patient (0.47 vs 0.28; P&lt;0.04).</p>
<p>Di Lenarda et al.<sup>99</sup> (1999)</p> <p>Metoprolol 142±44 mg QD</p> <p>vs</p> <p>carvedilol 12.5 mg to 50 mg BID</p>	<p>OL, PG, RCT</p> <p>Symptomatic (&gt;12 months) patients with stable dilated cardiomyopathy, LVEF of ≤40% and who poorly responded to chronic treatment with metoprolol plus conventional therapy (metoprolol plus ACE inhibitor, digitalis, diuretics), persistent moderate-to-severe left ventricular dysfunction and reduced exercise</p>	<p>N=30</p> <p>12 months</p>	<p>Primary: Improvement in left ventricular function and remodeling</p> <p>Secondary: Effects on symptoms, QOL, exercise tolerance, ventricular arrhythmias</p>	<p>Primary: LVEF significantly improved in the carvedilol group (7±3%) compared to the metoprolol group (-1±2%; P=0.045).</p> <p>LV end-systolic volume was significantly improved in the carvedilol group (-7±5) compared to the metoprolol group (6±4 mL/m<sup>2</sup>; P=0.047). There was not a significant difference in LV end-diastolic volume observed between the carvedilol (-8±7) and the metoprolol group (7±6 mL/m<sup>2</sup>; P=0.053).</p> <p>Secondary: There was not a significant difference observed in the NYHA class, the Heart Failure Score, the Minnesota “Living With Heart Failure” Questionnaire and submaximal exercise tolerance did not significantly change between the carvedilol and metoprolol groups.</p> <p>Carvedilol, compared to metoprolol, demonstrated a positive effect on ventricular ectopic beats (-12±9 vs 62±50 n/h; P=0.05) and couplets (-0.5±0.4 vs 1.5±0.6 n/h; P=0.048), but not a significant effect on episodes of nonsustained ventricular tachycardia (-0.02±0.03 vs 0.03±0.01).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Maack et al.<sup>100</sup> (2001)</p> <p>Metoprolol 12.5 to 100 mg BID</p> <p>vs</p> <p>carvedilol 3.125 to 25 mg BID</p>	<p>tolerance</p> <p>OL, XO</p> <p>Patients with stable NYHA class I to III heart failure due to ischemic or idiopathic dilated cardiomyopathy and an LVEF of &lt;35%</p>	<p>N=80</p> <p>6 months</p>	<p>Primary: Change in LVEF and change in baseline hemodynamic properties (left ventricular end diastolic, end systolic volume, NYHA class)</p> <p>Secondary: Not reported</p>	<p>Primary: After six months of treatment, LVEF improved in the carvedilol group (32±3 to 36±4%; P&lt;0.05 vs baseline) and in the metoprolol group (27±4 to 30±5%; P&lt;0.05 vs baseline). There was not a statistical difference between the agents.</p> <p>There were no differences between the groups in left ventricular end diastolic, end systolic volume, NYHA functional class or any other hemodynamic parameters at rest.</p> <p>Secondary: Not reported</p>
<p>Metra et al.<sup>101</sup> (2000)</p> <p>Metoprolol 5 to 100 mg BID</p> <p>vs</p> <p>carvedilol 3.125 to 50 mg BID</p> <p>All patients continued on their usual treatment for heart failure.</p>	<p>DB, PRO, RCT</p> <p>Symptomatic (≥6 months) patients with CHF caused by ischemic or nonischemic cardiomyopathy, NYHA class II to IV, LVEF ≤35% and a peak oxygen uptake ≤25 mL/kg-1/min-1 and on constant background therapy (furosemide and ACE inhibitor or ARB) for 1 week prior to the study</p>	<p>N=150</p> <p>15 months</p>	<p>Primary: Change in LVEF</p> <p>Secondary: Hemodynamic variables at rest and peak exercise, maximal and submaximal exercise tolerance, QOL, NYHA functional class, frequency of death and urgent transplantation</p>	<p>Primary: Both agents significantly increased LVEF from baseline (P&lt;0.001 for both), but carvedilol increased LVEF significantly greater than metoprolol (10.9±11 vs 7.2±7.7%; P=0.038).</p> <p>Secondary: At the end of the study, both agents carvedilol and metoprolol increased stroke volume and stroke work indexes and decreased mean pulmonary artery pressure, pulmonary wedge pressure, and heart rate from baseline (all P&lt;0.05 from baseline). However, the increase in stroke volume and stroke work indexes during exercise and the decreases in mean pulmonary artery pressure and pulmonary wedge pressure at both rest and exercise were greater with carvedilol than with metoprolol (all P&lt;0.05).</p> <p>Carvedilol increased rest and exercise cardiac index from baseline (both P&lt;0.05).</p> <p>Heart rate declined with both drugs at rest and exercise, but the decrease in exercise heart rate with carvedilol was greater than with metoprolol (P&lt;0.05 for the difference between the groups).</p> <p>Both metoprolol and carvedilol significantly improved NYHA class, 6-minute walk distance, and QOL scores from baseline (all P&lt;0.05), and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>there were no differences between the two treatments.</p> <p>Overall, 21 patients in the metoprolol group and 17 patients in the carvedilol group died or underwent urgent transplantation.</p>
<b>Hypertension</b>				
<p>Reim et al.<sup>102</sup> (1985)</p> <p>Acebutolol 400 mg QD</p> <p>vs</p> <p>propranolol 160 mg QD</p>	<p>DB, MC, XO</p> <p>Patients 18 to 70 years with essential HTN and blood pressure of &gt;150/90 mm Hg</p>	<p>N=18</p> <p>14 weeks</p>	<p>Primary: Blood pressure and heart rate during ergometer exercise test</p> <p>Secondary: Not reported</p>	<p>Primary: There was not a significant difference observed between the acebutolol and propranolol groups in decreases in blood pressure (systolic and diastolic) and heart rate at rest (P=0.123, P=0.230 and P=0.210, respectively).</p> <p>At the ergometer 25 watt load, heart rate and DBP were not significantly different between acebutolol and propranolol (P=0.087 and P=0.068, respectively), but SBP was significantly lower in the acebutolol group (P=0.042)</p> <p>At the higher ergometer loads of 50 and 75 watts, acebutolol had a significantly lower increase in SBP and heart rate compared to propranolol during exercise (50 watts: P=0.004 and P=0.012, respectively; 75 watts: P=0.005 and P=0.001, respectively), but there was not a significant difference observed between the groups in DBP in the 50 and 75 watt loads (P=0.057 and P=0.058, respectively).</p> <p>At the highest ergometer load of 100 watts, acebutolol significantly reduced systolic and DBPs and heart rate compared to propranolol (P=0.003, P=0.001, and P=0.001, respectively).</p> <p>Secondary: Not reported</p>
<p>Fogari et al.<sup>103</sup> (1984)</p> <p><u>Weeks 1 to 4:</u> Atenolol 50 mg QD</p> <p>vs</p>	<p>RCT, SB</p> <p>Patients 61 to 80 years inadequately controlled (SBP &gt;170 mm Hg and/or DBP &gt;100 mm Hg) on antihypertensive</p>	<p>N=38</p> <p>6 months</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: After the first four weeks, atenolol (from 177.5 to 161.1 mm Hg) significantly reduced blood pressure compared to baseline, but chlorthalidone did not (from 176.6 to 179.1 mm Hg).</p> <p>The combination atenolol-chlorthalidone therapy significantly reduced mean standing SBP and DBP, supine SBP and DBP, supine and standing heart rate, compared to previous therapies (P&lt;0.001 for all comparisons).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>chlorthalidone 12.5 mg QD</p> <p><u>Weeks 5 to study end:</u> atenolol and chlorthalidone 50-12.5 mg QD (fixed-dose combination product)</p>	<p>medications</p>			<p>The combination atenolol-chlorthalidone therapy significantly reduced mean standing SBP and DBP, supine SBP and DBP, supine and standing heart rate, compared to atenolol and chlorthalidone monotherapy (P&lt;0.001 or P&lt;0.01 for all comparisons).</p> <p>Mean blood pressure reduction obtained by the atenolol and chlorthalidone combination product was 30/15 mm Hg in the standing position (P&lt;0.001).</p> <p>Serum potassium increased with atenolol-chlorthalidone (4.45 mEq/L) compared to chlorthalidone alone (4.01 mEq/L; P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Leonetti et al.<sup>104</sup> (1986)</p> <p>Atenolol 50 mg QD</p> <p>vs</p> <p>atenolol 100 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 mg QD</p> <p>vs</p> <p>atenolol and chlorthalidone 50-12.5 mg QD (fixed-dose</p>	<p>DB, RCT</p> <p>Patients 24 to 68 years with mild to moderate HTN (WHO stage I or II), with supine DBP ≥95 mm Hg at the end of the 4-week washout period</p>	<p>N=28</p> <p>16 weeks</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: Mean supine blood pressure was significantly reduced in all treatment groups compared to placebo: 153±18/93±9 mm Hg for atenolol 50 mg patients, 155±22/91±8 mm Hg for atenolol 100 mg patients, 148±17/93±11 mm Hg for chlorthalidone 12.5 mg patients, and 144±16/89±6 mm Hg for the atenolol-chlorthalidone combination patients. All of the changes in blood pressure were significant (P&lt;0.01) versus placebo.</p> <p>Supine SBP was lower with atenolol-chlorthalidone than with the atenolol 100 mg alone (P&lt;0.05).</p> <p>Upright SBP was lower with atenolol-chlorthalidone than with atenolol 50 mg alone (P&lt;0.05) and atenolol 100 mg alone (P&lt;0.05).</p> <p>Mean supine heart rate was 77±7 bpm after placebo which decreased to 69±10 bpm (P&lt;0.01) after atenolol 50 mg, to 67±6 bpm (P&lt;0.01) after atenolol 100 mg, to 77±10 bpm (P=not significant, was not reported) after chlorthalidone alone.</p> <p>Chlorthalidone alone demonstrated a significant reduction in serum potassium levels compared to placebo (3.88 vs 4.09 mEq/L; P&lt;0.05) and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
combination product)				<p>no change when the atenolol-chlorthalidone combination was compared to placebo (3.98 vs 4.09; P=not significant, value was not reported).</p> <p>Chlorthalidone alone and atenolol-chlorthalidone demonstrated a significant increase in serum uric acid levels compared to placebo (4.90±1.52 mg/dL, 5.07±1.33 mg/dL, respectively, vs 4.24±1.12 for placebo; P&lt;0.05 for both).</p> <p>All treatments were well tolerated. Some adverse events reported included dyspnea, precordial discomfort and cold extremities. Incidence, severity and P values were not reported.</p>
<p>Nissinen et al.<sup>105</sup> (1980)</p> <p>Atenolol 100 mg QD plus chlorthalidone 25 mg in the morning</p> <p>vs</p> <p>atenolol and chlorthalidone 100-25 mg in the morning (fixed-dose combination product)</p> <p>vs</p> <p>placebo</p>	<p>DB, RCT</p> <p>Patients with newly diagnosed mild to moderate HTN (supine DBP 100 mm Hg on ≥3 occasions)</p>	<p>N=23</p> <p>16 weeks</p>	<p>Primary: Changes in blood pressure and heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Each of the active drug combinations lowered standing, supine, and post-exercise blood pressure significantly compared to placebo at two and four weeks (P&lt;0.001, P&lt;0.01 and P&lt;0.05). There was not a statistical difference between the active treatment regimens (P value not significant).</p> <p>Each of the active drug combinations lowered standing, supine, and post-exercise heart rate significantly compared to placebo at two and four weeks (P&lt;0.001, P&lt;0.01 and P&lt;0.05). There was not a statistical difference between the active treatment regimens (P value not significant).</p> <p>Side effects did not differ between treatment groups and placebo in terms of frequency or severity. Reported side effects included dizziness, headache and tiredness.</p> <p>Secondary: Not reported</p>
<p>Johnson et al.<sup>106</sup> (2009)</p> <p>Atenolol 50 to 100 mg QD for 9 weeks, followed</p>	<p>RCT</p> <p>Patients 17 to 65 years of age mild to moderate essential HTN</p>	<p>N=368</p> <p>15 to 18 weeks</p>	<p>Primary: Blood pressure lowering effect of drug initiation order: the addition of a β-blocker to a</p>	<p>Primary: When analyzed by order of initiation of the two drugs, the response to HCTZ and atenolol was greater overall than that seen for atenolol and HCTZ (P=0.0007 and P&lt;0.0001).</p> <p>This study suggests that initiation of HCTZ followed by atenolol results in</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>by atenolol 50 to 100 mg QD and HCTZ 12.5 to 25 mg QD for 9 weeks</p> <p>vs</p> <p>HCTZ 12.5 to 25 mg QD for 9 weeks, followed by HCTZ 12.5 to 25 mg QD and atenolol 50 to 100 mg QD for 9 weeks</p>			<p>thiazide versus the addition of a thiazide to a <math>\beta</math>-blocker</p> <p>Secondary: Not reported</p>	<p>greater blood pressure lowering as compared with initiation in the reverse order, with differences that are potentially clinically important.</p> <p>Secondary: Not reported</p>
<p>Dhakam et al.<sup>107</sup> (2008)</p> <p>Atenolol 50 mg QD</p> <p>vs</p> <p>nebivolol 5 mg QD</p> <p>vs</p> <p>placebo QD</p>	<p>DB, RCT, XO</p> <p>Never-treated subjects with isolated systolic HTN</p>	<p>N=16</p> <p>17 weeks</p>	<p>Primary: Change in central blood pressure</p> <p>Secondary: Change in peripheral blood pressure, AIx, aPWV and N-terminal proBNP.</p>	<p>Primary: There was not a statistically significant difference observed in the change in aortic SBP between the nebivolol and atenolol groups (<math>125\pm 3</math> vs <math>127\pm 3</math> mm Hg; <math>P=0.4</math>), but both agents were significantly better than placebo (<math>131\pm 2</math> mm Hg).</p> <p>There was not a statistically significant difference observed in the change in aortic DBP between the nebivolol and atenolol groups (<math>75\pm 2</math> vs <math>73\pm 2</math> mm Hg; <math>P=0.3</math>), but both agents were better than placebo (<math>82\pm 2</math> mm Hg).</p> <p>Secondary: There was not a statistically significant difference observed in the change in brachial SBP between the nebivolol and atenolol groups (<math>136\pm 3</math> vs <math>137\pm 3</math> mm Hg; <math>P=0.4</math>), but both agents were significantly better than placebo (<math>149\pm 3</math> mm Hg).</p> <p>There was not a statistically significant difference observed in the change in brachial DBP between the nebivolol and atenolol groups (<math>75\pm 2</math> vs <math>73\pm 2</math> mm Hg; <math>P=0.5</math>), but both agents were better than placebo (<math>82\pm 2</math> mm Hg).</p> <p>There was a statistically significant reduction in AIx in the atenolol group</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>compared to the nebivolol group (32±2 vs 28±2%; P=0.4), but both agents were significantly better than placebo (22±2%).</p> <p>There was not a statistically significant difference observed in the reduction of aPWV in the atenolol group compared to the nebivolol group (8.9±0.3 vs 9.1±0.3 m/s; P=0.2), but both agents were significantly better than placebo (10.0±0.4 m/s; P was not reported).</p> <p>There was not a statistically significant difference observed in the rise in N-terminal pro-BNP in the atenolol group compared to the nebivolol group (157 vs 138 pg/mL; P=0.6), but both agents were significantly better than placebo (75 mg/mL).</p>
<p>Fogari et al.<sup>108</sup> (1997)</p> <p>Atenolol 50 mg QD</p> <p>vs</p> <p>nebivolol 5 mg QD</p>	<p>DB, PG, RCT</p> <p>Patients 18 to 70 years of age with stable type 2 diabetes (HbA<sub>1c</sub> ≤8% during previous 6 months with diet and/or oral therapy stable for ≥6 months), and mild to moderate HTN (DBP ≥95 and &lt;116 mm Hg) at the end of the 4-week run-in period with placebo</p>	<p>N=30</p> <p>6 months</p>	<p>Primary: Changes in blood pressure, heart rate, 24-hour urinary C-peptide excretion, HbA<sub>1c</sub>, plasma glucose, lipid levels</p> <p>Secondary: Euglycemic hyperinsulinemic clamp test (body glucose utilization)</p>	<p>Primary: Both atenolol and nebivolol significantly reduced blood pressure and heart rate from baseline (P&lt;0.001 for all measures), but there was not a significant difference between the treatment groups at weeks 0, 2, and 24 (P&gt;0.05 for all measures).</p> <p>There no significant changes from baseline in mean 24-hour urinary C-peptide excretion, HbA<sub>1c</sub>, plasma glucose, and lipid levels (P&gt;0.05). There were also no significant differences observed between treatment groups in any of these measures (P&gt;0.05).</p> <p>Secondary: There was not a significant decrease from baseline in mean values for whole body glucose utilization observed in neither the atenolol group nor the nebivolol group (mean decrease of 0.9 vs 2.6%, respectively; P&gt;0.05) and the groups were significant from each other (P&gt;0.05).</p>
<p>Dietz et al.<sup>109</sup> (2008)</p> <p>Atenolol 50 to 100 mg QD</p> <p>vs</p>	<p>DB, MC, RCT</p> <p>Patients ≥18 years of age with HTN (mean sitting DBP ≥95 and &lt;110 mm Hg)</p>	<p>N=694</p> <p>12 weeks</p>	<p>Primary: Changes in mean sitting SBP and mean sitting DBP, rates of blood pressure control (&lt;140/90 mm Hg), pulse pressure and</p>	<p>Primary: Treatment with aliskiren and atenolol combination therapy led to a significantly greater reduction in mean sitting SBP by 17.3 mm Hg compared to aliskiren monotherapy (difference, -2.9 mm Hg; P=0.039) or atenolol monotherapy (difference, -3.0 mm Hg; P=0.034). There was no difference between mean sitting SBP reductions with aliskiren and atenolol monotherapy (difference, -0.1 mm Hg; P=0.954).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and atenolol 50 to 100 mg QD</p>			<p>pulse rate, plasma renin concentration, plasma renin activity</p> <p>Secondary: Not reported</p>	<p>Treatment with aliskiren and atenolol combination therapy led to a significantly greater reduction in mean sitting DBP by 14.1 mm Hg compared to aliskiren monotherapy (difference, -2.9 mm Hg; P&lt;0.001), but not atenolol monotherapy (difference, -0.5 mm Hg; P=0.545). Reductions in mean sitting DBP with atenolol were larger compared to those observed with aliskiren (difference, 2.4 mm Hg; P=0.003).</p> <p>Rates of blood pressure control were higher with aliskiren and atenolol combination therapy (51.3%) compared to aliskiren monotherapy (36.1%, P&lt;0.001) or atenolol monotherapy (42.2%, P=0.009). There was no significant difference in blood pressure control rates between aliskiren and atenolol monotherapy (P=0.388).</p> <p>Mean pulse pressure was reduced by 3.0 mm Hg with aliskiren and atenolol combination therapy and aliskiren monotherapy. Atenolol monotherapy did not affect pulse pressure. Aliskiren monotherapy did not affect pulse rate. Significant mean reductions in pulse rate of &gt;10 bpm were observed with atenolol monotherapy and the aliskiren and atenolol combination (P&lt;0.001 vs aliskiren monotherapy for both).</p> <p>Aliskiren monotherapy increased plasma renin concentration by 241% and aliskiren/atenolol increased plasma renin concentration by 85% (P=0.010 vs aliskiren). Atenolol monotherapy decreased plasma renin concentration by 24% (P&lt;0.001 vs aliskiren and aliskiren/atenolol). Aliskiren, atenolol and aliskiren/atenolol reduced plasma renin activity by 65, 52, and 61%, respectively.</p> <p>Secondary: Not reported</p>
<p>Wald et al.<sup>110</sup> (2008)</p> <p>Atenolol 25 mg QD</p> <p>vs</p>	<p>DB, DD, RCT, XO</p> <p>Patients ≥ 40 years enrolled in a HTN or anticoagulation clinic</p>	<p>N=47</p> <p>16 weeks</p>	<p>Primary: Reduction in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: The mean reductions in SBP in the atenolol alone, lisinopril alone and atenolol plus lisinopril groups were 16.1, 12.5 and 22.9 mm Hg, respectively. The mean reductions in DBP in the atenolol alone, lisinopril alone and atenolol plus lisinopril groups were 9.8, 6.8 and 13.9 mm Hg, respectively. The reductions with lisinopril plus atenolol group were significantly higher than either agent as monotherapy (P&lt;0.001).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
lisinopril 5mg QD  vs  lisinopril 5 mg and atenolol 25 mg QD  vs  placebo				Secondary: Not reported
Pareek et al. <sup>111</sup> (2010)  Atenolol 25 to 50 mg QD  vs  amlodipine 2.5 to 5 mg and atenolol 25 to 50 mg QD	AC, MC, OL, RCT  Adults with either untreated or pretreated essential HTN	N=190  12 weeks	Primary: Change in SBP and DBP  Secondary: Not reported	Primary: At the end of four weeks, the mean change in SBP (-30.0±10.4 vs -25.08±9.05; P=0.008) and DBP (-18.10±7.45 vs -14.78±7.48; P=0.021) was significantly greater in the low-dose combination therapy as compared to the low-dose monotherapy.  At the end of 12 weeks, the mean SBP (127.82±8.90 vs 138.0±14.4; P=0.001) and mean DBP (81.73±8.78 vs 87.35±5.50; P=0.011) were significantly lower in the high-dose combination group as compared to the high-dose monotherapy group.  Secondary: Not reported
Chapman et al. <sup>112</sup> (2007) ASCOT-BPLA  Atenolol 50 to 100 mg titrated to target blood pressure <140/90 mm Hg (or <130/90 mm Hg in diabetic patients); bendroflumethiazide* plus potassium	Subanalysis of ASCOT-BPLA evaluating effects of spironolactone on treatment-resistant HTN  Patients 40 to 79 years of age with HTN and ≥3 cardiovascular risk factors, with SBP ≥160 mm Hg and/or DBP ≥100 mm Hg	N=1,411  1.3 years	Primary: Change in DBP and SBP, adverse effects  Secondary: Not reported	Primary: Spironolactone-treated patients lead to a significant 21.9 mm Hg reduction in SBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 20.8 to 23.0 mm Hg; P<0.001).  Spironolactone-treated patients lead to a significant 9.5 mm Hg reduction in DBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 9.0 to 10.1; P<0.001).  Spironolactone-treated patients exhibited small but significant decreases in sodium, LDL-C and TC as well as increases in potassium, glucose, creatinine and HDL-C (P<0.05).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>1.25 to 2.5 mg plus doxazosin were added for additional blood pressure control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p> <p>vs</p> <p>amlodipine 5 to 10 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); perindopril 4 to 8 mg and doxazosin were added for additional control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p>	<p>(not on antihypertensive therapy) or SBP <math>\geq</math>140 mm Hg and/or DBP <math>\geq</math>90 mm Hg (on antihypertensive therapy)</p>			<p>The most common adverse effect reported in the trial was gynecomastia in men (P value not reported).</p> <p>Secondary: Not reported</p>
<p>Pepine et al.<sup>113</sup> (2006) INVEST</p>	<p>Post hoc analysis of INVEST</p> <p>Patients with</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Risk for adverse outcome associated with baseline</p>	<p>Primary: Previous heart failure (adjusted HR, 1.96), as well as diabetes (HR, 1.77), increased age (HR, 1.63), United States residency (HR, 1.61), renal impairment (HR, 1.50), stroke/TIA (HR, 1.43), smoking (HR, 1.41), MI</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Atenolol (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p> <p>vs</p> <p>verapamil SR (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p>	<p>essential HTN</p>		<p>factors, follow-up blood pressure and drug treatments</p> <p>Secondary: Not reported</p>	<p>(HR, 1.34), PVD (HR, 1.27), and revascularization (HR, 1.15) predicted increased risk.</p> <p>Follow-up SBP &lt;140 mm Hg (HR, 0.82) or DBP &lt;90 mm Hg (HR, 0.70) and trandolapril with verapamil SR (HR, 0.78 and 0.79) were associated with reduced risk.</p> <p>Secondary: Not reported</p>
<p>Denardo et al.<sup>114</sup> (2015) INVEST</p> <p>Atenolol (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p>	<p>Subgroup analysis of INVEST</p> <p>INVEST patients (patients with clinically stable hypertension and CAD) who underwent 24-hour ambulatory monitoring prior to randomization (“baseline”) and after one year of</p>	<p>N=117</p> <p>One year</p>	<p>Primary: BP, HR, pulse pressure</p> <p>Secondary: Not reported</p>	<p>Primary: Hourly SBP and DBP decreased after one year for both verapamil SR- and atenolol-based treatment strategies compared with baseline (P&lt;0.0001). Atenolol also decreased hourly HR (P&lt;0.0001). Both treatment strategies decreased SBP variability (weighted standard deviation: P=0.012 and 0.021, respectively). Compared with verapamil SR, atenolol also increased the prevalence of BP and HR nighttime dipping among prior non-dippers (BP: OR,3.37; 95% CI, 1.26 to 8.97; P=0.015; HR: OR, 4.06; 95% CI, 1.35 to 12.17; P=0.012) and blunted HR morning surge (2.8 vs 4.5 beats/min/hr; P=0.019).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>verapamil SR (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p>	<p>treatment</p>			
<p>Hilleman et al.<sup>115</sup> (1999)</p> <p>Monotherapy (atenolol, HCTZ, captopril, enalapril, lisinopril, amlodipine, diltiazem, nifedipine, verapamil)</p> <p>vs</p> <p>amlodipine and benazepril (fixed-dose combination)</p>	<p>MA (82 trials)</p> <p>Patients with mild-to-moderate essential HTN</p>	<p>N=not reported</p> <p>≥4 weeks</p>	<p>Primary: Absolute change in supine DBP from baseline</p> <p>Secondary: Percent of patients who achieved blood pressure control, safety</p>	<p>Primary: The mean absolute decrease in supine DBP ranged from 9.7 to 13.3 mm Hg with verapamil showing the greatest effect and captopril the least. When studies were weighted by sample size, amlodipine and benazepril, atenolol, lisinopril, and verapamil showed the greatest blood pressure effect.</p> <p>Secondary: The average percentage of patients defined as controlled after treatment varied from 53.5 to 79.0%, with amlodipine and benazepril (74.3%) and lisinopril (79.0%) showing the highest percentage control (P=0.096).</p> <p>The incidence of adverse events ranged from 12.1 to 41.8%, with lisinopril and verapamil showing the lowest incidences (12.1% and 14.1%, respectively) and nifedipine the highest incidence. Lisinopril demonstrated significantly less overall side effects compared to nifedipine (P=0.030).</p> <p>Nifedipine demonstrated a higher withdrawal rate due to side effects compared to atenolol, HCTZ, enalapril, amlodipine, and diltiazem (P=0.002). Although amlodipine and benazepril had the lowest rate of withdrawals due to adverse events, lack of significant change was due to the low number of cohorts available for analysis.</p>
<p>Davidov et al.<sup>116</sup> (1988)</p>	<p>DB, MC, RCT</p>	<p>N=141</p>	<p>Primary: Change in blood</p>	<p>Primary: Both betaxolol and propranolol significantly reduced SBP from baseline</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Betaxolol 10 to 40 mg QD  vs  propranolol 40 to 160 mg BID	Patients 21 to 73 years with mild to moderate HTN (supine DBP of 95 to 115 mm Hg)	24 weeks	pressure and heart rate  Secondary: Not reported	(7±2.5 and 7±2.0 mm Hg; P<0.01 for both).  Both betaxolol and propranolol significantly reduced DBP from baseline (11±0.9 and 9±1.2 mm Hg; P<0.01 for both).  Both betaxolol and propranolol significantly heart rate from baseline (6±1.3 and 7±1.1 bpm; P<0.01 for both).  At the end of the study, there was not a significant difference in response between groups.  Secondary: Not reported
Czuriga et al. <sup>117</sup> (2003) NEBIS  Bisoprolol 5 mg QD  vs  nebivolol 5 mg QD	MC, PG, RCT, SB  Patients 30 to 65 years with mild to moderate HTN, a DBP 95 to 110 mm Hg and a SBP ≤180 mm Hg at the end of the placebo run-in period who were either newly diagnosed or previously treated hypertensives and required a change of therapy in consequence of side-effects or poor compliance	N=273  16 weeks	Primary: Percentage of responders achieving DBP normalization (≤90 mm Hg) or a DBP reduction of at least 10 mm Hg and heart sitting rate  Secondary: Adverse events, symptom questionnaire	Primary: There was not a significant difference between percentage of responders between the nebivolol group (92%) and the bisoprolol group (89.6%).  There was not a significant difference in the mean change in blood pressure observed between the nebivolol and bisoprolol (SBP: -20.5±12.9 vs -20.0±12.0 mm Hg, respectively; P=0.7434) and DBP (-15.7±6.4 vs -16.0 ± 6.8 mm Hg, respectively; P=0.8230).  There was not a significant difference in mean heart rate observed between the nebivolol (68.7±8.5 per minute) and the bisoprolol group (68.1±7.5 per minute).  Secondary: There was not significant difference in rates of adverse events reported between the nebivolol (eight patients [5.8%]) and the bisoprolol group (12 patients [8.9%]; P>0.05). All adverse events were either mild (55%) or moderate (45%) in intensity.  Both treatments demonstrated a significant reduction in the basal score index at visit 5 (nebivolol, -0.7 vs bisoprolol, -0.5; P<0.02), but there was no significant difference between treatment groups (P>0.05).
Stoschitzky et al. <sup>118</sup>	DB, PC, RCT, XO	N=16	Primary: Heart rate and	Primary: Compared to baseline, heart rate at exercise was decreased at three hours

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2006)</p> <p>Bisoprolol 10 mg on day 1, then 5 mg QD</p> <p>vs</p> <p>carvedilol 50 mg on day 1, then 25 mg BID</p> <p>vs</p> <p>nebivolol 10 mg on day 1, then 5 mg QD</p>	<p>Male patients between 22 and 34 years with a height between 177 and 189 cm, and body weight between 66 and 86 k</p>	<p>1 week</p>	<p>blood pressure at rest and exercise</p> <p>Secondary: Effects on nocturnal melatonin release, QOL</p>	<p>after the first dose by bisoprolol (-24%), carvedilol (-17%) and nebivolol (-15%); (P&lt;0.05 for each group). Bisoprolol was significantly better than nebivolol (P&lt;0.05).</p> <p>Compared to baseline, heart rate at exercise was decreased at 24 hours after the first dose by bisoprolol (-18%), carvedilol (12 hours; -15%) and nebivolol (-13%); (P&lt;0.05 for each group). There was not a statistical significance observed between the groups.</p> <p>Compared to baseline, heart rate at exercise was decreased at 24 hours after the respective last dose at the end of one week of chronic administration by bisoprolol (-14%), carvedilol (12 hours; -15%) and nebivolol (-13%); (P&lt;0.05 in all cases). There was not a statistical significance observed between the groups.</p> <p>All of the agents significantly decreased SBP both at rest and exercise at three and 24 hrs after the first dose as well at 24 hr after the last dose after seven days of chronic administration (P&lt;0.05 in all cases). None of the agents had a significant effect on DBP at rest or at exercise.</p> <p>Secondary: Compared to placebo, nocturnal melatonin release was decreased by bisoprolol (-44%, P&lt;0.05) whereas nebivolol (-16%) and carvedilol (-19%) had no effect.</p> <p>Total QOL with carvedilol (8.0±0.8) was slightly but significantly lower than that with placebo (8.6±0.4), nebivolol (8.5±0.6) and bisoprolol (8.4±0.5); (P&lt;0.05 in all cases).</p>
<p>Lewin et al.<sup>119</sup> (1993)</p> <p>Bisoprolol and HCTZ 5-6.25 mg QD (fixed-dose combination product)</p>	<p>MC, PC</p> <p>Adult patients with stable mild to moderate (sitting DBP 95 to 114 mm Hg) essential HTN</p>	<p>N=36</p> <p>4 weeks</p>	<p>Primary: Changes in 24-hr ambulatory daytime and nighttime blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: There were statistically significant reductions in blood pressure and pulse (P&lt;0.01) at weeks two and four of treatment.</p> <p>There were statistically significant reductions (P&lt;0.01) in 24 hr SBP and DBP, daytime and nighttime blood pressure, compared to the end of the placebo phase. There was a reduction in systolic and diastolic load also (P&lt;0.01).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo				The combination was well tolerated. The scores from the overall QOL questionnaire indicated an improvement with the combination (P=0.02).
Benetos et al. <sup>120</sup> (2000)  Bisoprolol and HCTZ 2.5-6.25 mg QD (fixed-dose combination product)  vs  amlodipine 5 mg QD	DB, MC, PG, RCT  Patients over 60 years with supine SBP 160 to 210 mm Hg and DBP <90 mm Hg	N=164  12 weeks	Primary: Changes in blood pressure, heart rate, adverse events, QOL scores  Secondary: Not reported	Primary: Both bisoprolol and HCTZ and amlodipine significantly reduced SBP (-20.0±13.7 and -19.6±14.2 mm Hg, respectively; P<0.001) and DBP (-4.5±7.4 and -2.4±8.4 mm Hg, respectively from baseline to week 12, but there was not a significant difference between the agents (SBP; P=0.85 and DBP; P=0.09).  Bisoprolol and HCTZ significantly reduced heart rate from baseline, but amlodipine did not (-7.6±8.4 [P<0.001] and -0.2±11.4 bpm, respectively).  Bisoprolol and HCTZ significantly reduced heart rate when compared to amlodipine (P=0.0001).  Overall adverse events were not significantly different between the amlodipine and the bisoprolol and HCTZ group (39 and 40%, respectively). Adverse events reported included headache, leg edema, fatigue and bradycardia but severity of events was not reported.  Overall QOL scores were not significantly different between the amlodipine and the bisoprolol and HCTZ group.  Secondary: Not reported
Prisant et al. <sup>121</sup> (1995)  Bisoprolol and HCTZ 2.5-6.25, 5-6.25, or 10-6.25 mg/day (fixed-dose combination product)  vs	DB, MC, PG, RCT  Patients ≥21 years with mild to moderate essential HTN, (average sitting DBP 95 to 114 mm Hg) each treatment was once daily and titrated to effect	N=218  17 weeks	Primary: Mean change from baseline in SBP and DBP, lab measurements, adverse events, QOL questionnaire  Secondary: Not reported	Primary: Mean decreases in SBP and DBP from baseline were 13.4/10.7 mm Hg for bisoprolol and HCTZ patients, 12.8/10.2 mm Hg for amlodipine patients, and 7.3/6.6 mm Hg for enalapril patients. The hypotensive effects were significant for all three groups (P<0.001).  SBP and DBP mean changes from baseline for the bisoprolol and HCTZ group and the amlodipine group were greater than the change from baseline for the enalapril group (P<0.01).  Response rates (DBP ≤90 mm Hg or ≥10 mm Hg decrease from baseline)

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>enalapril 5, 10, or 20 mg</p> <p>vs</p> <p>amlodipine 2.5, 5, or 10 mg</p>				<p>were 71% for the bisoprolol and HCTZ group, 69% for the amlodipine group, and 45% for the enalapril group. The response rates for the bisoprolol and HCTZ and the amlodipine groups differed significantly from the enalapril group (P&lt;0.01).</p> <p>Twenty nine percent of bisoprolol patients had adverse experiences compared to 42% of amlodipine patients (P=0.12). Nearly 47% of enalapril patients had adverse experience compared to bisoprolol (P=0.04). Adverse events reported included headache, fatigue, peripheral edema, and dizziness.</p> <p>Drug related adverse events were 16% for the bisoprolol and HCTZ patients, 21% for the amlodipine patients, and 23% for the enalapril patients. There was no significant difference between the groups.</p> <p>Enalapril demonstrated a mean decrease from baseline of 7.9 mg/dL for TC (P=0.02 vs amlodipine) and 6.6 mg/dL for LDL-C (P=0.04 vs amlodipine) which were not significantly different from the increase from the bisoprolol and HCTZ group of 1.7 mg/dL (P=0.07 vs enalapril) for TC and +0.6 mg/dL in LDL-C. However, the increase in TGs was highest for bisoprolol and HCTZ-treated patients compared to amlodipine- and enalapril-treated patients (P=0.08, for bisoprolol and HCTZ vs enalapril).</p> <p>There was not a significant difference from baseline or between treatment groups in QOL scores: 0.9 for the bisoprolol and HCTZ group, 0.5 for the amlodipine group, and 2.3 for the enalapril group.</p>
<p>Frishman et al.<sup>122</sup> (1994)</p> <p>Bisoprolol 2, 5, 10, or 40 mg QD</p> <p>vs</p> <p>HCTZ 6.25 or 25 mg QD</p>	<p>DB, MC, PC, RCT</p> <p>Patients 21 years and older with mild to moderate essential HTN whose weight was 35% of the ideal for height and frame and mean sitting DBP was stable and</p>	<p>N=512</p> <p>12 weeks</p>	<p>Primary: Changes in DBP and SBP</p> <p>Secondary: Not reported</p>	<p>Primary: All treatment groups (all doses) of bisoprolol, HCTZ and the combination of bisoprolol and HCTZ significantly reduced sitting DBP from baseline (P&lt;0.01).</p> <p>The reduction in blood pressure was significantly greater as the doses of the bisoprolol, HCTZ and the combination of bisoprolol-HCTZ were increased (P&lt;0.05).</p> <p>The combination bisoprolol and HCTZ significantly reduced sitting DBP compared to the separate agents as monotherapy (P&lt;0.01).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  bisoprolol plus HCTZ, all possible combinations	between 95 to 115 mm Hg			<p>With higher doses of HCTZ, there was a significantly higher incidence of hypokalemia, defined as potassium &lt;3.5 mmol/L (P&lt;0.01). Incidence of hyperuricemia also significantly increased with the increase in HCTZ dose (P&lt;0.01). Adverse events associated with hypokalemia and hyperuricemia were not reported.</p> <p>As the dose of bisoprolol was increased, the frequency and severity of adverse events reported significantly increased (P&lt;0.05). Adverse events reported included asthenia, diarrhea, dyspepsia and somnolence, but severity of effects was not reported.</p> <p>Secondary: Not reported</p>
Frishman et al. <sup>123</sup> (1995)  Bisoprolol 5 mg QD  vs  HCTZ 25 mg QD  vs  bisoprolol and HCTZ 5-6.25 mg QD (fixed-dose combination product)  vs  placebo	DB, MC, PC, PG, RCT  Patients ≥21 years with mild to moderate (stage II or II) systemic HTN whose body weight was not >10% below or 35% above the ideal weight for height and frame, and were off all antihypertensive medications before study entry and sitting DBP was 95 to 115 mm Hg on 3 consecutive weekly visits	N=547  10 weeks	Primary: Changes in blood pressure and adverse events  Secondary: Not reported	Primary: All active treatment groups significantly reduced sitting DBP and SBP from baseline compared to placebo (P<0.01).  Addition of HCTZ 6.25 mg contributed significantly to the blood pressure lowering effects of bisoprolol 5 mg.  The combination bisoprolol and HCTZ 5-6.25 mg produced a significantly greater reduction in mean sitting DBP from baseline (-12.6±0.5 mm Hg) compared to bisoprolol 5 mg alone (-10.5±0.5 mm Hg; P=0.02) and HCTZ 25 mg alone (-8.5±0.5 mm Hg; P<0.01). Bisoprolol 5 mg monotherapy was significantly better a reducing DBP compared to HCTZ 25 mg alone (P=0.03).  The combination bisoprolol and HCTZ 5-6.25 mg produced a significantly greater reduction in mean sitting SBP from baseline (-15.8 mm Hg) compared to bisoprolol 5 mg alone (-10 mm Hg; P<0.01) and HCTZ 25 mg alone (-15.8 mm Hg; P<0.01). There was not a significant difference in mean reduction between bisoprolol 5 mg alone and HCTZ 25 mg alone.  Bisoprolol and HCTZ 5-6.25 mg in combination had a 73% response rate compared to 61% for the bisoprolol group and 47% for the HCTZ group.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Bisoprolol and HCTZ 5-6.25 mg in combination was found to be significantly more effective compared to bisoprolol 5 mg or HCTZ 25 mg in all subgroups of patients regardless of age, race, gender, or smoking history (<math>P&gt;0.05</math> for all comparisons).</p> <p>Bisoprolol and HCTZ 5-6.25 mg in combination did not have an increase in frequency or severity of adverse events. The adverse events were comparable to that in the placebo group and frequency among groups was not significant. The most common adverse events reported were headache, dizziness, fatigue, and cough.</p> <p>Significantly greater number patients in the HCTZ 25 mg group (6.5%) experienced hypokalemia (potassium <math>&lt;3.4</math> mEq/L) compared to the bisoprolol 5 mg group (0.7%; <math>P&lt;0.01</math>), the bisoprolol and HCTZ combination group (0.7%; <math>P&lt;0.01</math>), and placebo (0%; <math>P&lt;0.01</math>).</p> <p>Hyperglycemia occurred in 7.4% of patients in the HCTZ 25 mg group, which was significantly higher than in the placebo group (5.2%; <math>P=0.03</math>). Also, the incidence of hyperuricemia (uric acid <math>&gt;7.5</math> mg/dL) was significantly higher in the HCTZ 25 mg group (24.4%) compared to placebo (2.7%; <math>P&lt;0.01</math>).</p> <p>Secondary: Not reported</p>
<p>Williams et al.<sup>124</sup> (2015) PATHWAY-2</p> <p>Twelve weeks of once daily treatment with each of spironolactone (25 to 50 mg), bisoprolol (5 to 10 mg), doxazosin modified release (4</p>	<p>DB, PC, XO</p> <p>Patients 18 to 79 years of age with seated clinic SBP <math>\geq 140</math> mmHg (or <math>\geq 135</math> mmHg for patients with diabetes) and home SBP (18 readings over four days) <math>\geq 130</math> mmHg, despite treatment for at least three</p>	<p>N=335</p> <p>12 months</p>	<p>Primary: Average home SBP, recorded in the morning and the evening in triplicate, on four consecutive days before study visits</p> <p>Secondary: Clinic SBP, BP control rates, adverse events</p>	<p>Primary: The average reduction in home SBP by spironolactone was significantly greater compared to placebo (<math>-8.70</math> mmHg; 95% CI, <math>-9.72</math> to <math>-7.69</math>; <math>P&lt;0.0001</math>), compared to the mean of the other two active treatments (doxazosin and bisoprolol; <math>-4.26</math>; 95% CI, <math>-5.13</math> to <math>-3.38</math>; <math>P&lt;0.0001</math>), and compared to the individual treatments; versus doxazosin (<math>-4.03</math>; 95% CI, <math>-5.04</math> to <math>-3.02</math>; <math>P&lt;0.0001</math>) and versus bisoprolol (<math>-4.48</math>; 95% CI, <math>-5.50</math> to <math>-3.46</math>; <math>P&lt;0.0001</math>).</p> <p>Secondary: The results for seated clinic SBP largely mirror those seen with home SBP except that there was a large placebo effect on clinic BP that was not seen with home BP measurement.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
to 8 mg), and placebo, in addition to their baseline blood pressure drugs	months with maximally tolerated doses of three drugs (an ACE or ARB, a CCB, and a diuretic)			<p>Overall 219 (68.9%; 95% CI, 63.6 to 73.8) of 314 patients achieved target home SBP of &lt;135 mmHg. 58% of patients had their BP controlled with spironolactone, which was significantly greater than rates for other treatments (P&lt;0.001 when compared to doxazosin, bisoprolol, and placebo). Most patients who were controlled by doxazosin or bisoprolol had a still greater fall in blood pressure on spironolactone, which was consequently the most effective treatment in almost 60% of patients. This was at least three times the proportion in whom doxazosin or bisoprolol were the most effective.</p> <p>All active treatments were well tolerated with similar low rates of adverse events and withdrawals due to adverse events.</p>
<p>Hamaad et al.<sup>125</sup> (2007)</p> <p>Carvedilol 3.125 to 25 mg BID</p> <p>vs</p> <p>bisoprolol 1.25 to 10 mg QD</p>	<p>RCT</p> <p>Patients with stable LVEF of &lt;40% and treated with diuretic and ACE inhibitor or ARB</p>	<p>N=31</p> <p>12 weeks</p>	<p>Primary: Blood pressure, heart rate responses and both time and frequency domain heart rate variability</p> <p>Secondary: Not reported</p>	<p>Primary: Carvedilol significantly reduced DBP from baseline to week 12 of therapy (stage 6), but bisoprolol did not: 10±16 mm Hg (P=0.045) and 7±16 mm Hg, respectively (P=0.159), but there was not a significant difference between groups.</p> <p>Both carvedilol and bisoprolol significantly reduced SBP from baseline to week 12 of therapy (stage 6): 18±28 mm Hg (P=0.045) and 12±16 mm Hg, respectively (P&lt;0.003) but there was not a significant difference between groups.</p> <p>Both carvedilol and bisoprolol significantly decreased mean heart rate from baseline to week 12 of therapy (stage 6): 25±20 bpm and 23±10 bpm, respectively (P&lt;0.01 for both agents vs baseline) but there was not a significant difference between groups (P=0.708).</p> <p>Neither carvedilol nor bisoprolol significantly increased four of the five heart rate variability indices measured including SDNN, RMSSD, low frequency power or high frequency power. But both carvedilol and bisoprolol significantly increased triangular index from baseline to week 12 of therapy (stage 6): 7±6 (P&lt;0.01) and 5±6 (P=0.01), respectively, but there was not a significant difference between groups.</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
<p>Erdogan et al.<sup>126</sup> (2011)</p> <p>Carvedilol 25 mg QD for 1 month</p> <p>vs</p> <p>nebivolol 5 mg QD for 1 month</p> <p>All patients went through a 10 day placebo run in period.</p>	<p>DB, PC, PRO, RCT, XO</p> <p>Patients with mild to moderate HTN</p>	<p>N=20</p> <p>2 months</p>	<p>Primary: Blood pressure, heart rate</p> <p>Secondary: Safety</p>	<p>Primary: Treatment with carvedilol (133.8±9/86.6±8.6 mmHg) and nebivolol (134±8.7/85.6±7.4 mmHg) significantly decreased SBP and DBP compared to placebo (143.9±8.9/94.4±9.2 mmHg; P&lt;0.05). There was no difference between carvedilol and nebivolol (P&gt;0.05).</p> <p>Mean heart rate was significantly decreased after initiating treatment with carvedilol (70.2±5.2 bpm) and nebivolol (64.9±3.9 bpm) compared to placebo (78.8±5.2; P&lt;0.05).</p> <p>Secondary: No adverse events were reported with either treatment.</p>
<p>Saunders et al.<sup>127</sup> (1987)</p> <p>Labetalol 100 to 800 mg BID</p> <p>vs</p> <p>propranolol 40 to 320 mg</p>	<p>DB, PG</p> <p>Patients with mild to moderate HTN</p>	<p>N=153</p> <p>Duration not specified</p>	<p>Primary: Blood pressure, heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Labetalol was significantly better than propranolol at the end of monotherapy at lowering DBP (P&lt;0.05) but there was no difference in lowering SBP.</p> <p>Propranolol was significantly better at lowering heart rate compared to labetalol (P&lt;0.01).</p> <p>No difference in the decrease in blood pressure after a diuretic was added.</p> <p>Secondary: Not reported</p>
<p>McAreavey et al.<sup>128</sup> (1984)</p> <p>Labetalol 200 mg QD up to 1,600 mg BID</p> <p>vs</p>	<p>DB, PG, RCT</p> <p>Patients with inadequately controlled HTN while receiving atenolol 100 mg/day and bendrofluazide* 5 mg/day</p>	<p>N=238</p> <p>6 months</p>	<p>Primary: Comparative safety and efficacy, target blood pressure &lt;140/95 mm Hg</p> <p>Secondary: Not reported</p>	<p>Primary: Target blood pressure was reached in 25% of patients receiving hydralazine, 23% of patients receiving minoxidil, 19% of patients receiving prazosin, 17% of patients receiving methyldopa and zero percent of patients receiving placebo (P values not reported).</p> <p>Labetalol had the highest withdrawal rate compared to the other treatments with 78% (P&lt;0.05). Minoxidil had the second highest withdrawal rate with 57% (P&lt;0.05), due to fluid retention. There were no significant differences</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>prazosin 0.5 mg QD up to 10 mg BID</p> <p>vs</p> <p>hydralazine 12.5 mg QD up to 100 mg BID</p> <p>vs</p> <p>methyldopa 125 mg QD up to 1,000 mg BID</p> <p>vs</p> <p>placebo</p> <p>Minoxidil as add on therapy was given to men only.</p> <p>Doses were titrated upward at 2 week intervals until target blood pressure or maximum dose was reached.</p>				<p>in withdrawal rates among the other treatments.</p> <p>Secondary: Not reported</p>
<p>Wright et al.<sup>129</sup> (2002) AASK Metoprolol 50 to</p>	<p>DB, MC, RCT</p> <p>Patients were self-identified African Americans aged 18</p>	<p>N=1,094</p> <p>3-6.4 years</p>	<p>Primary: Rate of change in GFR (grouped by usual blood pressure [MAP</p>	<p>Primary: No significant difference in primary outcome was reported between the usual blood pressure group compared to the lower blood pressure group (P=0.24).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>200 mg/day</p> <p>vs</p> <p>ramipril 2.5 to 10 mg/day</p> <p>vs</p> <p>amlodipine 5 to 10 mg/day</p>	<p>to 70 years with HTN and a GFR between 20 and 65 mL/min/ 1.73 m<sup>2</sup> and no other identified cause of renal insufficiency</p>		<p>goal 102 to 107 mm Hg] vs lower blood pressure [<math>\leq</math>92 mm Hg])</p> <p>Secondary: Clinical composite outcome (reduction in GFR by 50% or more, ESRD, or death)</p>	<p>None of the drug group comparisons showed consistently significant differences in the GFR slope.</p> <p>Secondary: The lower blood pressure goal did not significantly reduce the rate of the clinical composite outcome (risk reduction for lower blood pressure group, 2%; 95% CI, -22 to 21; P=0.85).</p> <p>Ramipril resulted in significant risk reductions in the clinical composite outcomes compared to amlodipine (38%; 95% CI, 14 to 56; P=0.004) and metoprolol (22%; 95% CI, 1 to 38; P=0.04).</p> <p>There was no significant difference in the clinical composite outcome between the amlodipine and metoprolol groups.</p>
<p>Dafgard et al.<sup>130</sup> (1981)</p> <p>Metoprolol and HCTZ 200-25 mg QD in the morning (fixed-dose combination product)</p> <p>vs</p> <p>HCTZ 50 mg QD in the morning</p> <p>vs</p> <p>HCTZ 25 mg QD in the morning</p>	<p>DB, MC, RCT</p> <p>Patients with essential HTN (WHO stages I or II) not adequately controlled (<math>\geq</math>160/95 mm Hg) on HCTZ 25 mg/day</p>	<p>N=31</p> <p>32 weeks</p>	<p>Primary: Blood pressure, heart rate, adverse events, laboratory values</p> <p>Secondary: Not reported</p>	<p>Primary: After the eight week run-in period with HCTZ 25 mg alone, the mean supine blood pressure was significantly reduced from 183/110 to 172/103 mm Hg (P&lt;0.01/P&lt;0.01). The increased dose of HCTZ 50 mg following the run-in period did not further significantly reduce the mean blood pressure (165/104 mm Hg).</p> <p>A small but statistically significant reduction in supine heart rate was seen when the HCTZ dose was increased from 25 to 50 mg (82 down to 78 bpm; P&lt;0.05).</p> <p>After the 12 week double-blind period, the mean supine blood pressure was 153/98 mm Hg in the HCTZ 50 mg group. After the 12 week follow-up period, there was not any additional decrease in blood pressure (153/97 mm Hg).</p> <p>Fixed-dose combination product of metoprolol and HCTZ produced a significant reduction in supine blood pressure after 12 weeks of therapy from 172/105 mm Hg on HCTZ 25 mg alone to 154/97 mm Hg on the combination therapy (P&lt;0.001/P&lt;0.01). Similar results were found with the standing blood pressure reductions, from 165/108 to 147/97 mm Hg (P&lt;0.001/P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>After the eight week run-in period, the supine heart rate was 80 bpm which decreased to 64 bpm with the metoprolol and HCTZ fixed-dose combination (P&lt;0.001). The values for standing heart rate demonstrated similar significant reductions (85 to 66 bpm; P&lt;0.001).</p> <p>After the additional 12 week follow-up, the patients in the metoprolol and HCTZ fixed-dose combination group did not demonstrate a significant further reduction in heart rate or blood pressure in any position.</p> <p>Both agents were tolerated and the most common adverse events reported included insomnia, headache, tiredness, and shortness of breath. The majority of events were mild, few were moderate, and none were severe. The only significant changes in laboratory values occurred with the HCTZ 25 and 50 mg groups, where an increase in serum uric acid was observed from 0.30 to 0.34 and 0.35 mmol/L, respectively (P&lt;0.01 and P&lt;0.05; respectively).</p> <p>Secondary: Not reported</p>
<p>Smilde et al.<sup>131</sup> (1983)</p> <p>Metoprolol 400 mg QD in the morning for 5 weeks, followed by metoprolol and HCTZ 200-25 mg QD in the morning (fixed-dose combination product) (group 1)</p> <p>vs</p> <p>metoprolol and HCTZ 200-25 mg</p>	<p>DB, PG, RCT, XO</p> <p>Patients &lt;65 years with essential HTN (supine DBP ≥95 mm Hg) not controlled on metoprolol 200 mg alone</p>	<p>N=37</p> <p>15 weeks</p>	<p>Primary: Changes in DBP, SBP, and heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Both group 1 and 2 significantly reduced DBP (P&lt;0.01) from baseline and the two groups were not significantly different from each other.</p> <p>The combination products significantly reduced SBP from baseline (P&lt;0.05, P&lt;0.01 depending on comparison)</p> <p>Group 2 significantly reduced heart rate at the end of the study compared to baseline (P&lt;0.05).</p> <p>Clinically relevant changes in laboratory parameters or mean body weight were not observed between the groups.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QAM for 5 weeks (fixed-dose combination product), followed by metoprolol 400 mg QD in the morning for 5 weeks (group 2)</p>				
<p>Liedholm et al.<sup>132</sup> (1981)</p> <p>Metoprolol and HCTZ 100-12.5 mg BID (fixed-dose combination product) (group A)</p> <p>vs</p> <p>metoprolol and HCTZ 100-25 mg BID (fixed-dose combination product) (group B)</p> <p><u>Extended Study:</u> Metoprolol and HCTZ 100-12.5 mg, 2 tablets QD in the morning (fixed-dose combination product)</p>	<p>RCT</p> <p>Patients 18 to 72 years with mild to moderate essential HTN (WHO I or II)</p> <p><u>Extended Study:</u> OL</p> <p>Those patients who participated in the initial trial, had poor blood pressure control on existing antihypertensive therapy, and were being treated with a <math>\beta</math>-blocker and additional diuretic therapy</p>	<p>N=55</p> <p>12 weeks</p> <p><u>Extended Study:</u> N=49</p> <p>6 months</p>	<p>Primary: Change in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>In group A, there was a significant decrease in supine blood pressure from 189/112 to 172/105 mm Hg with metoprolol monotherapy and further reduction to 148/92 mm Hg with the metoprolol and HCTZ 100-12.5 mg (P&lt;0.001/P&lt;0.001).</p> <p>In group B, there was a significant decrease in supine blood pressure from 184/111 to 170/104 mm Hg with metoprolol monotherapy and further reduced to 152/96 mm Hg with metoprolol and HCTZ 100-25 mg (P&lt;0.01/P&lt;0.05) after 12 weeks.</p> <p>Supine heart rate fell in group A from 78 to 68 bpm with metoprolol monotherapy (P&lt;0.001). No further heart rate reduction was noted with the metoprolol and HCTZ 100-12.5 mg. In group B, supine heart rate fell from 76 to 69 bpm (P&lt;0.05). No further heart rate reduction was seen with metoprolol and HCTZ 100-25 mg.</p> <p>In group A, serum sodium fell from 143 to 140 mmol/L (P&lt;0.01). In group B, serum potassium fell with from 4.4 to 4.0 mmol/L (P&lt;0.001).</p> <p><u>Extended Study:</u> After six months of extended the therapy, there was no further significant reductions in supine or standing blood pressure, but there was a reduction in standing DBP from 97 to 95 mm Hg (P&lt;0.05).</p>
<p>Materson et al.<sup>133</sup> (1990)</p>	<p>DB, MC, RCT</p> <p>Men <math>\geq</math>60 years with</p>	<p>N=690</p> <p>12 months</p>	<p>Primary: The average reduction in SBP</p>	<p>Primary:</p> <p>Across all four treatments, there was an additional average reduction in BP of 13.1/10.6 mm Hg. The average reduction in SBP from baseline to</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Metoprolol 50, 100 or 200 mg BID</p> <p>vs</p> <p>hydralazine 25, 50 or 100 mg BID</p> <p>vs</p> <p>methyldopa 250, 500 or 1,000 mg BID</p> <p>vs</p> <p>reserpine 0.05, 0.10 or 0.25 mg QD</p> <p>All patients received HCTZ 25 to 100 mg QD.</p>	<p>HTN not currently receiving antihypertensive therapy and DBP 90 to 114 mm Hg and SBP &lt;240 mm Hg or a DBP &lt;100 mm Hg and a SBP &lt;240 mm Hg if currently taking antihypertensive therapy and the blood pressure criteria was met after ≥2 weeks without medication</p>	<p>N=811</p>	<p>and DBP, the number of patients achieving the goal blood pressure, the average change in heart rate</p> <p>Secondary: The rates of drug intolerances, adverse effects</p>	<p>endpoint for hydralazine, methyldopa, metoprolol and reserpine were -11.5±10.1 (P&lt;0.001), -15.0±13.7 (P&lt;0.001), -13.0±15.4 (P&lt;0.001) and -12.7±11.5 (P&lt;0.001), respectively. There was no significant difference in SBP reductions among the different treatments (P=0.43). The average reduction in DBP from baseline to endpoint for hydralazine, methyldopa, metoprolol and reserpine were -11.3±5.9 (P&lt;0.001), -10.6±6.3 (P&lt;0.001), -10.6±6.7 (P&lt;0.001) and -9.8±6.3 (P&lt;0.001), respectively. There was no significant difference in DBP reductions among the different treatments (P=0.59).</p> <p>The average change in heart rate from baseline to endpoint for hydralazine, methyldopa, metoprolol and reserpine were 1.4±10.5 (P value not significant), -1.6±9.3 (P value not significant), 15.9±11.9 (P&lt;0.05) and -7.9±10.7 (P&lt;0.05), respectively. There was a significant difference in change in heart rate among the different treatments (P&lt;0.001).</p> <p>The percentage of patients achieving the goal blood pressure at endpoint with hydralazine, methyldopa, metoprolol and reserpine were 85.3, 81.7, 76.9 and 72.3%, respectively (P=0.28).</p> <p>Secondary: Drug intolerance, defined as adverse effects prompting dose reduction or discontinuation, was present in 23.3% of patients not achieving goal blood pressure compared to 2.8% of those who did (P&lt;0.001). This was significant with hydralazine, methyldopa and metoprolol, but not with reserpine.</p> <p>There were 27 (10%) treatment discontinuations due to adverse effects (hydralazine [n=3], methyldopa [n=8], metoprolol [n=9] and reserpine [n=7]). There were two treatment discontinuations with methyldopa and one with reserpine due to depression.</p> <p>The overall incidence of volunteered moderate or severe adverse effects, not prompting treatment discontinuation, was significantly greater (P&lt;0.01) with methyldopa (31%) and hydralazine (25%) compared to reserpine (15%) or metoprolol (9%).</p>
<p>Greathouse.<sup>134</sup></p>	<p>DB, PC, PG, RCT</p>	<p>N=811</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2010)</p> <p>Nebivolol 5, 10 or 20 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patients entered a 4 to 6 week washout, SB, placebo run in period.</p>	<p>Patients <math>\geq 18</math> years of age with stage I to II HTN (average sitting DBP <math>\geq 95</math> and <math>\leq 109</math> mm Hg)</p>	<p>12 weeks</p>	<p>Change in mean sitting DBP at trough drug concentration (24<math>\pm</math>2 hours after the previous morning's dose)</p> <p>Secondary: Mean changes in trough sitting SBP, responder rate (mean trough SBP <math>&lt; 90</math> mm Hg or a decrease of <math>\geq 10</math> mm Hg from baseline), safety and tolerability</p>	<p>Least squares mean reductions in trough sitting DBP at week 12 were significantly greater with all doses of nebivolol compared to placebo (P=0.002 for 5 mg and P&lt;0.001 for 10 and 20 mg).</p> <p>All doses of nebivolol reduced peak sitting DBP in a dose-dependent manner. The least squares mean reductions in peak sitting DBP following treatment with 5, 10, and 20 mg of nebivolol were -10.5, -11.6, and -12.2 mm Hg (P&lt;0.001 vs placebo for all).</p> <p>Secondary: All doses of nebivolol resulted in least squares mean reductions in trough sitting SBP from baseline, with only the 20 mg dose reaching significance compared to patients receiving placebo (P&lt;0.001). All doses of nebivolol reduced peak sitting SBP in a dose-dependent manner. The least squares mean reductions with nebivolol in peak sitting SBP were -7.7, -10.7 and -4.7 mm Hg (P=0.004 vs placebo for 10 mg and P&lt;0.001 vs placebo for 20 mg).</p> <p>Significantly more patients receiving nebivolol were treatment responders compared to placebo (66.0 [P=0.009 vs placebo], 66.8 [P=0.005 vs placebo] and 68.9% [P=0.002 vs placebo] vs 49.3%).</p> <p>A total of 27 (36.0%) and 311 (42.5%) patients receiving placebo and nebivolol experienced an adverse event. The most commonly reported adverse events for the combined nebivolol group (all doses) compared to the placebo group were headache (7.5 vs 5.3%), fatigue (3.8 vs 1.3%) and nasopharyngitis (3.7 vs 4.0%).</p>
<p>Neutel et al.<sup>135</sup> (2010)</p> <p>Nebivolol 5, 10 or 20 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, PG, RCT</p> <p>Patients <math>\geq 18</math> years of age with stage I to II HTN who were inadequately controlled by antihypertensive medication (SBP <math>\geq 90</math> and <math>\leq 109</math> mm</p>	<p>N=669</p> <p>12 weeks</p>	<p>Primary: Change in mean clinic sitting DBP at trough (24<math>\pm</math>3 hours after previous morning's dose)</p> <p>Secondary: Change in mean</p>	<p>Primary: Addition of nebivolol to background antihypertensive therapy led to significant additional blood pressure reductions compared to placebo. Nebivolol 5, 10, and 20 mg significantly lowered trough sitting DBP by -3.3, -3.5, and -4.6 mm Hg, respectively (P&lt;0.001 for all doses).</p> <p>Secondary: Nebivolol 5, 10 and 20 mg significantly lowered trough sitting SBP by -5.7, -3.7, and -6.2 mm Hg, respectively (P&lt;0.001 for 5 and 20 mg and P=0.015 for 10 mg).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	Hg) and stable on a regimen of antihypertensive medications consisting of $\geq 1$ and $\leq 2$ of an ACE inhibitor, ARB or diuretic		trough sitting SBP and mean sitting DBP, change in mean sitting SBP at peak (two to three hours after dosing), mean peak and trough supine and standing DBP and SBP, mean 24 hour DBP and SBP as measured by ambulatory blood pressure monitoring, responder rate (sitting SBP <90 mm Hg or an absolute reduction $\geq 10$ mm Hg)	<p>Reductions in trough blood pressure in the standing and supine positions were comparable to sitting blood pressure reductions for all nebivolol doses.</p> <p>All doses of nebivolol also significantly reduced peak sitting DBP (-3.2, -4.0, and -4.3 mm Hg) and sitting SBP (-5.7, -5.6, and -5.9 mm Hg) at week 12 compared to placebo (P&lt;0.001 for both).</p> <p>Reductions from baseline to week 12 in peak blood pressure with nebivolol in both supine and standing positions were consistent with those for sitting DBP and sitting SBP (data not reported).</p> <p>After 12 weeks, the proportion of patients responding to treatment was significantly higher with nebivolol 5 mg (53.0%; P=0.028), 10 mg (60.1%; P=0.001) and 20 mg (65.1%; P&lt;0.001) compared to placebo (41.3%). In addition, a significantly higher percentage of patients receiving nebivolol achieved blood pressure control (&lt;140/90 mm Hg) (43.0, 41.3 and 52.7 vs 29.3%; P<math>\leq</math>0.029).</p>
Weiss et al. <sup>136</sup> (2011)  Nebivolol 1.25 to 30 or 40 mg/day  vs  placebo	Pooled analysis of 3 PC, RCT, SB  Patients with stage I-II HTN	N=2,016  $\geq 12$ weeks	Primary: Mean change from baseline in sitting DBP, sitting SBP, and heart rate at 12 weeks  Secondary: Safety	Primary: Compared to placebo, reductions in DBP, SBP, and heart rate were significantly greater with nebivolol at the recommended dosages of 5-30/40 mg/day (P<0.001 for all).  Secondary: The most commonly reported adverse events were headache (7.1 vs 5.9%), fatigue (3.6 vs 1.5%), and nasopharyngitis (3.1 vs 4.4%).
Rosei et al. <sup>137</sup> (2003)  Nebivolol 5 mg QD  vs	DB, MC, PG, RCT  Patients between 24 and 65 years with mild to moderate uncomplicated essential HTN that was newly	N=65  12 weeks	Primary: Response rates, changes in sitting blood pressure  Secondary: Standing blood pressure, sitting	Primary: There was not a significant difference in response rates observed between the two treatment groups.  Both treatment groups significantly reduced sitting SBP (P<0.0001) and DBP (P<0.0001) throughout the study compared to baseline but there were no significant differences observed between the treatment groups at most visits, but at week eight, DBP was significantly lower in the nebivolol

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
lisinopril 20 mg QD	diagnosed, or previous antihypertensive therapy was withdrawn at >1 month before active treatment, and had a sitting DBP of >95 and <114 mm Hg		and standing heart rate	<p>group compared to the lisinopril group (P&lt;0.05).</p> <p>Secondary: There was not a significant difference observed between treatment groups in standing blood pressure measurements.</p> <p>Both treatment groups significantly reduced sitting heart rate (P&lt;0.01) throughout the study compared to baseline but there were no significant differences observed between the treatment groups at most visits, but at week eight, heart rate were significantly lower in the nebivolol group compared to the lisinopril group (P&lt;0.05).</p>
<p>Mazza et al.<sup>138</sup> (2002)</p> <p>Nebivolol 2.5 to 5 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>DB, MC, PG, RCT</p> <p>Patients between 65 to 89 years of age with mild to moderate essential HTN and DBP ranging from 95 to 114 mm Hg</p>	<p>N=168</p> <p>16 weeks</p>	<p>Primary: Change in sitting blood pressure, response rates</p> <p>Secondary: Standing blood pressure changes, standing and sitting heart rate changes</p>	<p>Primary: There was not a significant difference observed between the amlodipine and nebivolol treatments groups in changes in sitting DBP (blood pressure values and P values not reported). At weeks four and eight, a slightly lower sitting SBP was observed in per-protocol patients in the amlodipine groups vs those in the nebivolol group (blood pressure values not reported, P&lt;0.005).</p> <p>Response rates were not significantly difference between the amlodipine group and the nebivolol group (86 vs 88%, respectively). The percentage of patients who reached normalization (blood pressure &lt;140/90 mm Hg) was no significant between the amlodipine and the nebivolol groups (47 vs 50%).</p> <p>Secondary: There were significant differences in standing blood pressure observed between the groups.</p> <p>Heart rate was significantly lower in the nebivolol group compared to the amlodipine group at all treatment visits (P&lt;0.001).</p> <p>Patients in the amlodipine group experienced a significantly greater rate of headache (seven vs five patients) and ankle edema (12 vs zero patients) compared to the patients in the nebivolol group (P&lt;0.05 for both).</p>
Van Bortel et al. <sup>139</sup> (2005)	DB, MC, PG, RCT	N=314	Primary: Effects on blood	<p>Primary: At the end of 12 weeks, both nebivolol and losartan significantly reduced</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Nebivolol 5 mg QD</p> <p>vs</p> <p>losartan 50 mg QD</p> <p>If after 6 weeks, DBP was not normalized, then HCTZ 12.5 mg QD was added to therapy</p>	<p>Patients &lt;70 years of age with DBP at randomization between 95 and 114 mm Hg</p>	<p>12 weeks</p>	<p>pressure, overall QOL</p> <p>Secondary: Comparison of different aspects of QOL</p>	<p>SBP compared to baseline (P&lt;0.0001 for both), but the agents were not significantly different from each other.</p> <p>Both agents also significantly decreased DBP compared to baseline (P&lt;0.0001), but nebivolol significantly reduced DBP compared to losartan (P&lt;0.02).</p> <p>At the end of 12 weeks, both nebivolol and losartan significantly improved QOL scores compared to baseline (P&lt;0.007), but the agents were not significantly different from each other.</p> <p>Secondary: At week 12 there was not a significant difference observed in the individual questions of the QOL questionnaire between the groups. Questions inquired about headaches, lightheadedness, sleepiness, flushing, and sexual function.</p>
<p>Van Bortel et al.<sup>140</sup> (2008)</p> <p>Nebivolol</p> <p>vs</p> <p>ACE inhibitor, ARB, β-blocker, calcium channel blocker, or placebo</p>	<p>MA</p> <p>12 RCTs involving &gt;25 patients with essential HTN where nebivolol 5 mg QD was compared to placebo or other active drugs for &gt;1 month</p>	<p>N=2,653</p> <p>Duration varied</p>	<p>Primary: Antihypertensive effect and tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: Overall, higher response rates were observed with nebivolol than all other antihypertensive agents combined (OR, 1.41; 95% CI, 1.15 to 1.73; P=0.001) and compared to the ACE inhibitors (OR, 1.92; 1.30 to 2.85; P=0.001), but response rates to nebivolol were similar to β-blockers (OR, 1.29; 95% CI, 0.81 to 2.04; P=0.283), calcium channel blockers (OR, 1.19; 95% CI, 0.83 to 1.70; P=0.350) and losartan (OR, 1.35; 95% CI, 0.84 to 2.15; P=0.212).</p> <p>Overall, a higher percentage of patients obtained normalized blood pressure with nebivolol compared to the other antihypertensive agents combined (OR, 1.35; 95% CI, 1.07 to 1.72; P=0.012). A higher percentage of patient receiving nebivolol obtained normalized blood pressure compared to losartan (OR, 1.98; 95% CI, 1.24 to 3.15; P=0.004) and calcium channel blockers (OR, 1.96; 95% CI, 1.05 to 1.96; P=0.024), but not when compared to other β-blockers (OR, 1.29; 95% CI, 0.81 to 1.65; P=0.473).</p> <p>Overall, the percentage of adverse events was significantly lower with nebivolol compared to the other antihypertensive agents combined (OR, 0.59; 95% CI, 0.48 to 0.72; P&lt;0.001) and similar to placebo (OR, 1.16;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>95% CI, 0.76 to 1.67; P=0.482). In comparing nebivolol to the individual treatments, nebivolol had a lower percentage of adverse events compared to losartan (OR, 0.52; 95% CI, 0.30 to 0.89; P=0.016), the other <math>\beta</math>-blockers (OR, 0.56; 95% CI, 0.36 to 0.85; P=0.007) and calcium channel blockers (OR, 0.49; 95% CI 0.33 to 0.72; P&lt;0.001), but was similar to ACE inhibitors (OR, 0.75; 95% CI 0.52 to 1.08).</p> <p>Secondary: Not reported</p>
<p>Veterans Administration Cooperative Study Group on Antihypertensive Agents<sup>141</sup> (1983)</p> <p>Nadolol 80 to 240 mg QD in the morning</p> <p>vs</p> <p>bendroflumethiazide 5 to 10 mg* QD in the morning</p> <p>vs</p> <p>nadolol and bendroflumethiazide*</p>	<p>DB, RCT</p> <p>Men 20 to 69 years with pretreatment DBP of 95 to 114 mm Hg</p>	<p>N=365</p> <p>12 weeks</p>	<p>Primary: Changes in blood pressure, change in blood pressure among races, heart rate, adverse events, laboratory values</p> <p>Secondary: Not reported</p>	<p>Primary: DBP of &lt;90 mm Hg was achieved in 49% of the nadolol patients, 46% of the bendroflumethiazide patients, and 85% of the combination patients. There was a significantly higher percentage of patients who achieved the DBP goal compared to the nadolol alone group and bendroflumethiazide group alone (P&lt;0.01 for both).</p> <p>The reduction in SBP was significantly greater in the combination group compared to the nadolol alone and bendroflumethiazide group (-25.3±1.4, -10.5±1.6, and -17.4±1.7 mm Hg, respectively; P&lt;0.001 for both) and bendroflumethiazide produced a significantly greater reduction compared to nadolol alone (P&lt;0.01).</p> <p>The reduction of DBP in white patients was significantly greater than the decrease in African American (decrease of 15.6 vs 9.6 mm Hg, respectively; P&lt;0.001). In addition, 77% of white patients achieved DBP of &lt;90 mm Hg compared to only 31% of African American patients (P&lt;0.001).</p> <p>Adverse events were infrequent. The most common were impotence, lethargy, weakness, and postural dizziness, which occurred more often with bendroflumethiazide than nadolol.</p> <p>Significant reductions in average heart rate from baseline were observed with nadolol alone (decrease by 16.1 bpm; P&lt;0.001) and with the combination product (decrease by 15.8 bpm; P&lt;0.001).</p> <p>Serum potassium levels significantly decreased from baseline in the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>bendroflumethiazide group by <math>-0.57 \pm 0.06</math> mEq/L (<math>P &lt; 0.001</math>) and in the combination group by <math>-0.44 \pm 0.05</math> mEq/L (<math>P &lt; 0.001</math>).</p> <p>Serum uric acid levels significantly increased from baseline in the bendroflumethiazide group by <math>1.7 \pm 0.2</math> mg/dL (<math>P &lt; 0.001</math>), in the nadolol group by <math>0.4 \pm 0.1</math> mg/dL (<math>P &lt; 0.01</math>) and in the combination group by <math>-1.9 \pm 0.1</math> mg/dL (<math>P &lt; 0.001</math>).</p> <p>Fasting glucose levels significantly increased from baseline in the bendroflumethiazide group by <math>6.1 \pm 2.1</math> mg/dL (<math>P &lt; 0.001</math>) and in the combination group by <math>7.4 \pm 1.1</math> mg/dL (<math>P &lt; 0.001</math>).</p> <p>Cholesterol significantly increased from baseline in the bendroflumethiazide group by <math>11.5 \pm 4.3</math> mg/dL (<math>P &lt; 0.001</math>).</p> <p>TGs significantly increased from baseline in the bendroflumethiazide group by <math>34.6 \pm 14.8</math> mg/dL (<math>P &lt; 0.01</math>), in the nadolol group by <math>38.7 \pm 13.2</math> mg/dL (<math>P &lt; 0.01</math>) and in the combination group by <math>67.8 \pm 11.9</math> mg/dL (<math>P &lt; 0.001</math>).</p> <p>Secondary: Not reported</p>
<p>Frick et al.<sup>142</sup> (1978)</p> <p>Penbutolol 40 mg BID</p> <p>vs</p> <p>propranolol 160 mg BID</p>	<p>DB, XO</p> <p>Patients 29 to 64 years of age with HTN</p>	<p>N=20</p> <p>13 weeks</p>	<p>Primary: Blood pressure, heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Penbutolol significantly reduced supine and standing blood pressures (both SBP and DBP) from baseline (<math>P &lt; 0.05</math>). Propranolol also significantly reduced blood pressures from baseline (SBP: <math>P &lt; 0.02</math> and diastolic: <math>P &lt; 0.01</math>), but there was not significant difference between agents.</p> <p>Penbutolol significantly reduced supine and standing heart rates from baseline (from <math>76 \pm 10</math> to <math>61 \pm 9</math>; <math>P &lt; 0.001</math> and from <math>85 \pm 13</math> to <math>67 \pm 8</math>; <math>P &lt; 0.001</math>, respectively). Propranolol also significantly reduced heart rates from baseline (to <math>59 \pm 8</math>; <math>P &lt; 0.001</math> and to <math>63 \pm 7</math>; <math>P &lt; 0.001</math>, respectively), but there was not significant difference between agents.</p> <p>Secondary: Not reported</p>
<p>Finnerty et al.<sup>143</sup></p>	<p>SB</p>	<p>N=59</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1979)</p> <p>Propranolol 80 mg to 320 mg QD</p> <p>vs</p> <p>reserpine 0.125 mg to 0.25 mg QD</p> <p>vs</p> <p>methyldopa 500 mg to 2,000 mg QD</p> <p>All patients received hydroflumethiazide* 50 or 100 mg QD.</p>	<p>Patients with HTN unresponsive to hydroflumethiazide alone</p>	<p>9 weeks</p>	<p>Percentage of patients achieving a DBP below 90 mm Hg</p> <p>Secondary: Not reported</p>	<p>At study endpoint, the DBP below 90 mm Hg was achieved in all 20 patients (100%) treated with hydroflumethiazide plus reserpine, 13 of the 19 patients (68.4%) treated with hydroflumethiazide plus methyldopa, and in 16 of the 20 patients (80%) treated with hydroflumethiazide plus propranolol.</p> <p>Secondary: Not reported</p>
<p>VA Cooperative Study<sup>144</sup> (1977)</p> <p>Propranolol 40 to 160 mg TID (P), propranolol 40- to 160 mg TID plus HCTZ 35 mg (P+T), propranolol 40 to 160 mg TID plus hydralazine 35 mg (P+H), or propranolol 40 to 160 mg TID plus HCTZ 35 mg plus hydralazine 35 mg</p>	<p>DB, RCT</p> <p>Men 18 to 59 years with DBP of 90 to 114 mm Hg</p>	<p>N=450</p> <p>18 months</p>	<p>Primary: Percent of patients who achieved a DBP &lt;90 mm Hg at 6 months, heart rate, withdrawal rate</p> <p>Secondary: Not reported</p>	<p>Primary: At six months, significantly more patients in the R+T arm (88%) attained a DBP &lt;90 mm Hg and <math>\geq 5</math> mm Hg less than the initial blood pressure compared to the P arm (52%; <math>P &lt; 0.01</math>) and the P+H arm (72%; <math>P &lt; 0.05</math>). The other arms: P+T (81%) and P+T+H (92%) were not significantly different than the R+T arm.</p> <p>The 12 and 18 month results do not have the statistical validity of the six months results due to the reduced sample size. The following percentage of patients attained DBP &lt;90 mm Hg and <math>\geq 5</math> mm Hg less than the initial pressure: R+T=89.1 and 82.6%, P=59.5 and 58.1%, P+T=86.0 and 86.4%, P+H=67.4 and 76.1%, and P+T+H=89.4 and 91.8%.</p> <p>There was not a significance difference in heart rate reductions at six and 18 months between the groups (R+T=5.0<math>\pm</math>1.3 and 5.0<math>\pm</math>1.3 mean change in heart rate, P=9.1<math>\pm</math>1.3 and 9.2<math>\pm</math>1.8, P+T=8.8<math>\pm</math>1.2 and 6.3<math>\pm</math>1.5, P+H=8.9<math>\pm</math>1.3 and 7.8<math>\pm</math>1.5, and P+T+H=5.9<math>\pm</math>1.1 and 7.7<math>\pm</math>1.5).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(P+T+H)</p> <p>vs</p> <p>reserpine 35 mg plus HCTZ 35 mg (R+T)</p>				<p>Withdrawals for any reason were similar between the treatment arms and were not statistically significant (R+T=14 patients, P=11, P+T=12, P+H=14, and P+T+H=16).</p>
<p>Stevens et al.<sup>145</sup> (1982)</p> <p><u>Dose-finding phase:</u> Propranolol 80, 160, 240, or 320 mg/day in 2 divided doses</p> <p>vs</p> <p>propranolol and HCTZ 80-50, 160-50, 240-50, 320-50 mg/day in 2 divided doses (fixed-dose combination product)</p> <p><u>Double-blind phase:</u> Propranolol and HCTZ (fixed-dose combination product)</p> <p>vs</p>	<p>DB, PG, RCT</p> <p>Patients with mild to moderate essential HTN (DBP 100 to 125 mm Hg)</p>	<p>N=158</p> <p>25 weeks</p>	<p>Primary: Mean changes of SBP and DB, heart rate, lab values</p> <p>Secondary: Not reported</p>	<p>Primary: After the 12 week dose finding-phase, 94% of patients had a decrease <math>\geq 10</math> mm Hg in DBP. The mean SBP and DBP reduced from 158.0 (<math>\pm 17.3</math>)/105.6 (<math>\pm 6.0</math>) mm Hg to 131.5 (<math>\pm 14.4</math>)/86.4 (<math>\pm 6.7</math>) mm Hg (P&lt;0.001).</p> <p>After the 10 week portion of the study, there were significantly greater increases (P&lt;0.05) in mean SBP or DBP with propranolol and HCTZ alone vs the combination product of propranolol and HCTZ from the end of the dose-finding to the last four biweekly visits to the mean of those visits, and to the last visit. The mean increases of SBP and DBP at the endpoint were: propranolol, 10.2/6.3 mm Hg; HCTZ 13.1/9.3 mm Hg; propranolol-HCTZ combination product 3/1.5 mm Hg.</p> <p>There was a significant decrease in heart rate as the dose of propranolol was increased though the trial (P&gt;0.30).</p> <p>The only lab value that showed a statistically significant change was serum chloride. The percent of patients that fell outside of the normal range were as follows: propranolol 6/36 (17%), HCTZ 14/37 (38%), and combination 4/28 (14%); P&lt;0.05.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>propranolol</p> <p>vs</p> <p>HCTZ</p>				
<p>de Leeuw et al.<sup>146</sup> (1997)</p> <p>Verapamil SR and trandolapril 180-2 mg/day, atenolol and chlorthalidone 100-25 mg/day, or lisinopril and HCTZ 20-12.5 mg/day (fixed-dose combination products)</p> <p>vs</p> <p>placebo</p> <p>All patients entered a SB, placebo 4 week run in period.</p>	<p>DB, MC, PC, RCT</p> <p>Patients 18 to 70 years of age with essential HTN (WHO I or II) newly or unsuccessfully treated, with supine DBP 101 to 114 mm Hg in week 4 of the run in period</p>	<p>N=205</p> <p>12 weeks</p>	<p>Primary: Changes in supine blood pressure, standing blood pressure response rates, normalization rates</p> <p>Secondary: Not reported</p>	<p>Primary: Each of the three treatments was significantly more effective than placebo in reducing seated DBP. Changes in DBP were as follows: verapamil SR and trandolapril, -13 (95% CI, -16 to -9); atenolol and chlorthalidone, -13 (95% CI, -16 to -9); lisinopril and HCTZ, -12 (95% CI, -15 to -9) and placebo, -3 (95% CI, -7 to 0) (P=0.0001 for all vs placebo), but there was not a significance among the treatments (P values not reported).</p> <p>Each of the three treatments was significantly more effective than placebo in reducing seated SBP. Changes in SBP were as follows: verapamil SR and trandolapril, -27 (95% CI, -33 to -21); atenolol and chlorthalidone, -28 (95% CI, -34 to -22); lisinopril and HCTZ, -23 (95% CI, -29 to -17) and placebo, -3 (95% CI, -9 to 3) (P=0.0001 for all vs placebo), but there was not a significance among the treatments (P values not reported).</p> <p>Effects on standing blood pressure demonstrated similar results as the effects on sitting blood pressure (P values not reported).</p> <p>Normalization of DBP (&lt;90 mm Hg), corrected for placebo, were significantly higher with all treatments compared to placebo (verapamil SR and trandolapril, 33% [95% CI, 16 to 50; P&lt;0.0005]; atenolol and chlorthalidone, 31% [95% CI, 14 to 48; P&lt;0.002] and lisinopril and HCTZ, 25% [95% CI, 9 to 42; P&lt;0.005]).</p> <p>Response rates (normalization of DBP or a reduction in DBP &gt;10 mm Hg), corrected for placebo, were significantly higher with all treatments compared to placebo (verapamil SR and trandolapril, 40% [95% CI, 22 to 58; P&lt;0.0001], atenolol and chlorthalidone, 44% [95% CI, 27 to 61; P&lt;0.0001] and lisinopril and HCTZ, 37% [95% CI, 19 to 55; P&lt;0.0002]).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Casas et al.<sup>147</sup> (2005)</p> <p>ACE inhibitor or ARBs compared to other antihypertensive drugs (<math>\beta</math>-adrenergic blocking agents, <math>\alpha</math>-adrenergic blocking agents, calcium-channel blocking agents, or combinations)</p> <p>vs</p> <p>ACE inhibitor or ARBs compared to placebo</p> <p>Specific agents and doses were not specified.</p>	<p>MA (127 trials)</p> <p>Studies in adults that examined the effect of any drug treatment with a blood pressure lowering action on progression of renal disease</p>	<p>N=not reported</p> <p>4.2 years (mean)</p>	<p>Primary: Doubling of serum creatinine, and ESRD</p> <p>Secondary: Serum creatinine, urine albumin excretion and GFR</p>	<p>Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>Secondary: Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).</p> <p>Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).</p> <p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>
<p>Baguet et al.<sup>148</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan,</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the <math>\beta</math>-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)</p> <p>Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>reported).</p> <p>The weighted average reduction of SBP and DBP for each drug class were as follows:            Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively.            β-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively.            Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively.            ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.            ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively.            Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary: Not reported</p>
<b>Post Myocardial Infarction and Other Cardiovascular Outcomes Trials</b>				
<p>Gottlieb et al.<sup>149</sup> (2001)</p> <p>Atenolol vs metoprolol</p>	<p>RETRO</p> <p>Patients discharged from the hospital with the diagnosis of an acute MI and on a β-blocker</p>	<p>N=69,338</p> <p>2 years</p>	<p>Primary: Mortality rates at 1 and 2 year(s)</p> <p>Secondary: Not reported</p>	<p>Primary: β-blockers demonstrated a 40% overall reduction in mortality compared to those patient who did not receive β-blocker therapy.</p> <p>One year mortality rates in the three groups were metoprolol 8.32% (CI, 8.07 to 8.58, atenolol 8.16% (CI, 7.76 to 8.58), propranolol 9.55% (CI, 9.69 to 10.48), and other 9.19% (CI, 8.16 to 10.33).</p> <p>Two year mortality rates in the three groups were metoprolol 13.52% (CI,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs propranolol  vs other (not specified)				13.21 to 13.84), atenolol 13.41% (CI, 12.91 to 13.93), propranolol 15.91% (CI, 14.83 to 17.05), and other 15.17% (CI, 13.88 to 16.56). There were no differences between atenolol and metoprolol at the end of the two years, both of which were statistically better than propranolol.  Compared to metoprolol, patients discharged on propranolol had 15% increased mortality at one year and 18% increased mortality at two years, which were significantly higher than metoprolol.  Secondary: Not reported
Testa et al. <sup>150</sup> (2014)  Patients taking atenolol  vs  Patients not taking atenolol	Observational  Patients aged ≥65 years with isolated HTN	N=972  12 years	Primary: Mortality  Secondary: Not reported	Primary: Univariate analysis shows that elderly participants taking atenolol show greater mortality than those not taking atenolol (52.4 vs 66.7%; P=0.047).  Cox regression analysis on 12-year mortality showed that age, number of diseases, number of drugs, basic activity of daily living ≥1%, and social support score were predictive; whereas female sex and Mini-Mental State Examination score were protective of long-term mortality. Additionally, pulse arterial pressure (HR, 1.02; 95% CI, 1.01 to 1.03; P=0.035) and atenolol use (HR, 1.89; 95% CI, 1.03 to 4.25; P<0.05) were predictive of long-term mortality.  Secondary: Not reported
Black et al. <sup>151</sup> (2003) CONVINCE  Atenolol 50 mg QD  vs  verapamil ER 180 mg QD	AC, DB, MC, RCT  Patients 55 years of age and older with HTN and ≥1 risk factor for cardiovascular disease	N=16,476  3 years	Primary: Composite first occurrence of acute MI, stroke or cardiovascular disease-related death  Secondary: Cardiovascular endpoints expanded, all-	Primary: There was no significant difference between the verapamil treatment group and the atenolol or HCTZ treatment groups in the composite primary endpoint (HR, 1.02; 95% CI, 0.88 to 1.18; P=0.77).  Secondary: There was no significant difference between the verapamil treatment group and the atenolol or HCTZ treatment group in rates of cardiovascular-related hospitalization (P=0.31), death (all-cause mortality) (P=0.32) and cancer rates (P=0.46).  Patients treated with verapamil experienced a significantly higher rate of

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs HCTZ 12.5 mg QD			cause mortality, cancer, hospitalization for bleeding, incidence of primary endpoints between 6AM and noon, adverse events	death or bleeding unrelated to stroke (HR, 1.54; 95% CI, 1.15 to 2.04; P=0.003).  Primary endpoints did not differ significantly based on time of day (P=0.43).  Patients treated with verapamil were more likely to withdraw for adverse events or symptoms than those treated with atenolol or HCTZ (P=0.02).
Dahlöf et al. <sup>152</sup> (2002) LIFE Atenolol 50 to 100 mg QD vs losartan 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	DB, DD, PG, RCT  Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200 to 95 to 115 mm Hg) and left ventricular hypertrophy	N=9,193  ≥4 years	Primary: Composite of cardiovascular death, MI and stroke  Secondary: All-cause mortality, hospitalization for angina or heart failure, revascularization procedures, resuscitated cardiac arrest, new-onset diabetes	Primary: SBP fell by 30.2 and 29.1 mm Hg in the losartan and atenolol groups, respectively (treatment difference, P=0.017) and DBP fell by 16.6 and 16.8 mm Hg, respectively (treatment difference, P=0.37). MAP was 102.2 and 102.4 mm Hg, respectively (P value not significant). Heart rate decreased more in patients assigned to atenolol than losartan (-7.7 vs -1.8 beats/minute, respectively; P<0.0001).  Compared to atenolol, the primary composite occurred in 13.0% fewer patients receiving losartan (RR, 0.87; 95% CI, 0.77 to 0.98; P=0.021).  While there was no difference in the incidence cardiovascular mortality (P=0.206) and MI (P=0.491), losartan treatment resulted in a 24.9% relative risk reduction in stroke compared to atenolol (P=0.001).  Secondary: A 25% lower incidence of new-onset diabetes was reported with losartan compared to atenolol (P=0.001). There was no significant difference among the other secondary end points between the two treatment groups.  Note: At end point or end of follow-up, 18 and 26% of patients on losartan were receiving HCTZ alone or with other drugs, respectively. In the atenolol group, 16 and 22% of patients were receiving HCTZ alone or with other drugs, respectively.
Julius et al. <sup>153</sup> (2004) LIFE Black Subset Atenolol 50 to 100	Post hoc analysis  Patients 55 to 80 years old with essential HTN	N=523  ≥4 years	Primary: Composite of cardiovascular death, MI and stroke	Primary: Compared to atenolol (11.2%), losartan in the United States African American population resulted in a greater incidence of the composite end point (17.4%; P=0.033).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg QD vs losartan 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	(sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy		Secondary: Not reported	HRs favored atenolol across all parameters (P=0.246 for cardiovascular mortality, P=0.140 for MI, and P=0.030 for stroke).  In African American patients, blood pressure reduction was similar in both groups, and regression of electrocardiographic-left ventricular hypertrophy was greater with losartan.  Secondary: Not reported
Lindholm et al. <sup>154</sup> (2002) LIFE Diabetic Subset  Atenolol 50 to 100 mg QD vs losartan 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	Post hoc analysis  Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy	N=1,195  ≥4 years	Primary: Composite of cardiovascular death, MI and stroke  Secondary: All-cause mortality	Primary: Compared to atenolol, losartan resulted in a 24% decrease in the primary composite end point (P=0.031).  Losartan treatment resulted in a 37% risk reduction in cardiovascular deaths vs atenolol (P=0.028).  Losartan treatment resulted in a 39% risk reduction in all-cause mortality vs atenolol (P=0.002).  Mean blood pressure fell to 146/79 mm Hg in losartan patients and 148/79 mm Hg in atenolol patients.  Secondary: Mortality from all causes was 63 and 104 in the losartan and atenolol groups, respectively (RR, 0.61; P=0.002).
Kjeldsen et al. <sup>155</sup> (2002) LIFE Isolated Systolic Hypertension Subset  Atenolol 50 to 100 mg QD	Post hoc analysis  Patients 55 to 80 years old with isolated systolic HTN (SBP of 160 to 200 mm Hg and DBP <90 mm Hg) and left ventricular	N=1,326  ≥4 years	Primary: Composite of cardiovascular death, MI, or stroke  Secondary: All-cause mortality	Primary: Compared to atenolol, losartan resulted in a trend towards a 25% reduction in the primary end point (P=0.06).  Losartan treatment resulted in a 46% risk reduction in cardiovascular mortality (P=0.01) and 40% risk reduction in stroke compared to atenolol (P=0.02). There was no difference in the incidence of MI.  Blood pressure was reduced by 28/9 and 28/9 mm Hg in the losartan and

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>losartan 50 to 100 mg QD</p> <p>HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.</p>	<p>hypertrophy</p>			<p>atenolol arms.</p> <p>Secondary: Patients receiving losartan also had reductions in all-cause mortality (28%; <math>P &lt; 0.046</math>).</p>
<p>Fossum et al.<sup>156</sup> (2006) ICARUS, a LIFE substudy</p> <p>Atenolol 50 to 100 mg QD</p> <p>vs</p> <p>losartan 50 to 100 mg QD</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy</p>	<p>N=81</p> <p>3 years</p>	<p>Primary: Amount and density of atherosclerotic lesions in the common carotid arteries and carotid bulb</p> <p>Secondary: Not reported</p>	<p>Primary: The amount of plaque decreased in the losartan group and increased in the atenolol group, though the difference between groups was not statistically significant (<math>P=0.471</math>).</p> <p>Patients in the atenolol group had a greater increase in plaque index compared to the losartan group, though the difference between groups was not statistically significant (<math>P=0.742</math>)</p> <p>Secondary: Not reported</p>
<p>Kizer et al.<sup>157</sup> (2005) (LIFE substudy)</p> <p>Atenolol 50 to 100 mg QD</p> <p>vs</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular</p>	<p>N=9,193</p> <p><math>\geq 4</math> years</p>	<p>Primary: Reduction in the risk of different stroke subtypes and neurological deficits</p> <p>Secondary: Not reported</p>	<p>Primary: The risk of fatal stroke was significantly decreased in the losartan group compared to the atenolol group (<math>P=0.032</math>).</p> <p>The risk of atherothrombotic stroke was significantly decreased in the losartan group compared to the atenolol group (<math>P=0.001</math>).</p> <p>Comparable risk reductions were observed for hemorrhagic and embolic stroke but did not reach statistical significance.</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>losartan 50 to 100 mg QD</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>hypertrophy</p>			<p>The risk of recurrent stroke was significantly reduced in the losartan arm compared to the atenolol arm (P=0.017).</p> <p>The number of neurological deficits per stroke was similar (P=0.68), but there were fewer strokes in the losartan group for nearly every level of stroke severity.</p> <p>Secondary: Not reported</p>
<p>Wachtell et al.<sup>158</sup> (2005) (LIFE substudy)</p> <p>Atenolol 50 to 100 mg QD</p> <p>vs</p> <p>losartan 50 to 100 mg QD</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy</p>	<p>N=8,851 (patients in LIFE with no baseline history of AF but at risk for AF)</p> <p>≥4 years</p>	<p>Primary: Incidence of new-onset AF and outcome</p> <p>Secondary: Not reported</p>	<p>Primary: Significantly fewer patients in the losartan group experienced new-onset AF compared to the atenolol group (P&lt;0.001).</p> <p>Randomization to losartan treatment was associated with a 33% lower rate of new onset AF independent of other risk factors (P&lt;0.001).</p> <p>Patients in the losartan group had a 40% lower rate of composite events consisting of cardiovascular death, fatal or non-fatal stroke, and fatal or non-fatal MI (P=0.03).</p> <p>Significantly fewer strokes occurred in the losartan group compared to the atenolol group (P=0.01), and there was a trend toward fewer MIs in the losartan group (P=0.16).</p> <p>There was no significant difference in cardiovascular mortality between groups.</p> <p>In contrast, the atenolol group experienced significantly fewer hospitalizations for heart failure (P=0.004) and a trend toward fewer sudden cardiac deaths (P=0.07).</p> <p>Secondary: Not reported</p>
<p>Wachtell et al.<sup>159</sup> (2005) (LIFE substudy)</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80</p>	<p>N=342 (LIFE patients with AF at the</p>	<p>Primary: Cardiovascular morbidity and</p>	<p>Primary: Patients with a history of AF had significantly higher rates of cardiovascular and all-cause mortality, fatal and non-fatal stroke, heart</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Atenolol 50 to 100 mg QD vs losartan 50 to 100 mg QD</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy</p>	<p>start of the LIFE study)  ≥4 years</p>	<p>mortality  Secondary: Not reported</p>	<p>failure, revascularization and sudden cardiac death compared to patients without AF (P&lt;0.001).</p> <p>Patients with a history of AF had similar rates of MI and hospitalization for angina pectoris (P≥0.209).</p> <p>The primary composite endpoint of cardiovascular mortality, stroke and MI occurred in significantly fewer patients in the losartan group compared to the atenolol group (P=0.009).</p> <p>The difference in MI between groups was not significant.</p> <p>Treatment with losartan trended toward lower all-cause mortality (P=0.09) and fewer pacemaker implantations (P=0.065).</p> <p>Secondary: Not reported</p>
<p>Dahlöf et al.<sup>160</sup> (1991) Hypertension (STOP)</p> <p>Atenolol 50 mg QD, HCTZ 25 mg QD plus amiloride 2.5 mg QD, metoprolol 100 mg QD, or pindolol 5 mg QD vs placebo</p>	<p>DB, MC, RCT</p> <p>Swedish men and women 70 to 84 years old with treated or untreated essential HTN defined as SBP ≥180 mm Hg with a DBP of ≥90 mm Hg, or DBP &gt;105 mm Hg irrespective of the SBP measured on 3 separate occasions during a 1-month placebo run-in phase in previously untreated patients</p>	<p>N=1,627  25 months</p>	<p>Primary: Frequency of stroke, MI, and other cardiovascular death  Secondary: Not reported</p>	<p>Primary: The active treatments significantly reduced the number of all primary endpoints (94 vs 58; RR, 0.60; 95% CI, 0.43 to 0.85; P=0.0031), frequency of stroke (53 vs 29; RR, 0.53; 95% CI, 0.33 to 0.86; P=0.0081) and frequency of other cardiovascular deaths (13 vs 4; RR, 0.30; 95% CI, 0.07 to 0.97) compared to placebo.</p> <p>There was not a statistically significant decrease observed in the rate of MI between the active treatments and placebo (28 vs 25; RR, 0.87; 95% CI, 0.49 to 1.56).</p> <p>Secondary: Not reported</p>
<p>Hansson et al.<sup>161</sup></p>	<p>BE, MC, OL, RCT</p>	<p>N=6,614</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1999) HYPERTENSION -2 (STOP)</p> <p><u>Conventional drug group</u> Atenolol 50 mg QD, HCTZ 25 mg QD plus amiloride 2.5 mg QD, metoprolol 100 mg QD, or pindolol 5 mg QD</p> <p>vs</p> <p><u>Newer drug group</u> ACE inhibitors (enalapril 10 mg QD or lisinopril 10 mg QD) or calcium channel blockers (felodipine 2.5 mg QD, or isradipine 2 to 5 mg QD)</p>	<p>Swedish men and women between 70 to 84 years old with treated or untreated essential with HTN on 3 separate occasions defined by SBP <math>\geq</math>180 mm Hg, DBP &gt;105 mm Hg, or both</p>	<p>60 months</p>	<p>Combined fatal stroke, MI, and other fatal cardiovascular disease; combined fatal and nonfatal stroke, MI, and other cardiovascular Mortality</p> <p>Secondary: Not reported</p>	<p>The combined fatal mortality endpoints occurred in 221 of the 2,213 patients in the conventional drugs group and in 438 of 4,401 in the newer drugs group (RR, 0.99; 95% CI, 0.84 to 1.16; P=0.89).</p> <p>The combined fatal and nonfatal mortality endpoints occurred in 460 patients taking conventional drugs and in 887 taking newer drugs (RR, 0.96; 95% CI, 0.86 to 1.08; P=0.49).</p> <p>Secondary: Not reported</p>
<p>Dalhof et al.<sup>162</sup> (2005) ASCOT-BPLA</p> <p>Atenolol 50 to 100 mg/day adding bendroflumethiazide* 1.25 to 2.5 mg/day and potassium as needed</p>	<p>MC, OL, RCT</p> <p>Patients 40 to 79 years of age with HTN and <math>\geq</math>3 other cardiovascular risk factors (left ventricular hypertrophy, other specified abnormalities on</p>	<p>N=19,257</p> <p>5.5 years</p>	<p>Primary: Nonfatal MI (including silent MI) and fatal CHD</p> <p>Secondary: All-cause mortality, total stroke, primary end points minus silent MI, all coronary</p>	<p>Primary: No statistically significant difference in nonfatal MI and fatal CHD was reported between the amlodipine plus perindopril group compared to the atenolol plus bendroflumethiazide groups (HR, 0.90; 95% CI, 0.79 to 1.2; P=0.1052).</p> <p>Secondary: Significantly greater reductions in the following secondary end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: all- cause mortality (P=0.0247), total stroke (P=0.0003), primary end points minus silent MI (P=0.0458), all coronary</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>amlodipine 5 to 10 mg/day adding perindopril 4 to 8 mg/day as needed</p> <p>If blood pressure was still not achieved, doxazosin 4 to 8 mg/day was added to the regimen.</p>	<p>ECG, type 2 diabetes, PAD, history of stroke or TIA, male, age <math>\geq 55</math> years, microalbuminuria or proteinuria, smoking, TC:HDL-C ratio <math>\geq 6</math>, or family history of CHD)</p>		<p>events, total cardiovascular events and procedures, cardiovascular mortality, nonfatal and fatal heart failure, effects on primary end point and on total cardiovascular events and procedures among prespecified subgroups</p> <p>Tertiary: Silent MI, unstable angina, chronic stable angina, PAD, life-threatening arrhythmias, development of diabetes, development of renal impairment</p>	<p>events (P=0.0070), total cardiovascular events and procedures (P&lt;0.0001), and cardiovascular mortality (P=0.0010).</p> <p>There were no significant differences in nonfatal and fatal heart failure between the two treatment groups (P=0.1257).</p> <p>The study was terminated early due to higher mortality and worse outcomes on several secondary end points observed in the atenolol study group.</p> <p>Tertiary: Significantly greater reductions in the following end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: unstable angina (P=0.0115), PAD (P=0.0001), development of diabetes (P&lt;0.0001), and development of renal impairment (P=0.0187).</p> <p>There were no significant differences in the incidence of silent MI (P=0.3089), chronic stable angina (P=0.8323) or life-threatening arrhythmias (P=0.8009) between the two treatment groups.</p> <p>There was no significant difference in the percent of patients who stopped therapy because of an adverse event between the two treatment groups (overall 25%). There was, however, a significant difference in favor of amlodipine plus perindopril in the proportion of patients who stopped trial therapy because of a serious adverse events (2 vs 3%; P&lt;0.0001).</p>
<p>Pepine et al.<sup>163</sup> (2003) INVEST</p> <p>Atenolol 50 mg/day (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3),</p>	<p>MC, OL, RCT</p> <p>Patients with essential HTN</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: First occurrence of death (all cause), nonfatal MI or stroke</p> <p>Secondary: Cardiovascular death, angina, cardiovascular</p>	<p>Primary: At 24 months, in the calcium antagonist strategy subgroup, 81.5% of patients were taking verapamil SR, 62.9% trandolapril, and 43.7% HCTZ. In the non-calcium antagonist strategy, 77.5% of patients were taking atenolol, 60.3% HCTZ, and 52.4% trandolapril.</p> <p>After a follow-up of 61,835 patient-years (mean, 2.7 years per patient), 2,269 patients had a primary outcome event with no statistically significant difference between treatment strategies (9.93% in calcium antagonist strategy vs 10.17% in non-calcium antagonist strategy; RR,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>then add trandolapril (step 4) (non-calcium antagonist strategy)</p> <p>vs</p> <p>verapamil SR 240 mg/day (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p> <p>Trandolapril was recommended for all patients with heart failure, diabetes, or renal insufficiency.</p>			<p>hospitalization, angina, blood pressure control (SBP/DBP &lt;140/90 mm Hg or &lt;130/85 mm Hg if diabetic or renal impairment), safety</p>	<p>0.98; 95% CI, 0.90 to 16; P=0.57).</p> <p>Secondary: There was no significant difference in the rate of cardiovascular death (P=0.94) or cardiovascular hospitalization (P=0.59) between the two treatment groups.</p> <p>At 24 months, angina episodes decreased in both groups, but the mean frequency was lower in the calcium antagonist strategy group (0.77 episodes/week) compared to the non-calcium antagonist strategy group (0.88 episodes/week; P=0.02).</p> <p>Two-year blood pressure control was similar between groups. The blood pressure goals were achieved by 65.0% (systolic) and 88.5% (diastolic) of calcium antagonist strategy patients and 64.0% (systolic) and 88.1% (diastolic) of non-calcium antagonist strategy patients. A total of 71.7% of calcium antagonist strategy patients and 70.7% of non-calcium antagonist strategy patients achieved an SBP &lt;140 mm Hg and DBP &lt;90 mm Hg.</p> <p>Both regimens were generally well tolerated. Patients in the calcium antagonist strategy group reported constipation and cough more frequently than patients in the non-calcium antagonist strategy group, while non-calcium antagonist strategy patients experienced more dyspnea, lightheadedness, symptomatic bradycardia and wheezing (all were statistically significant with P≤0.05).</p>
<p>Mancia et al.<sup>164</sup> (2007) INVEST</p> <p>Atenolol 25 to 200 mg QD</p> <p>vs</p> <p>verapamil SR 120 to 480 mg QD</p>	<p>MC, open blinded endpoint, PRO, RCT</p> <p>Patients with HTN, requiring drug therapy (BP&gt;140/90 or &gt;130/80 mm Hg if diabetic or with renal impairment), and CAD</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Occurrence of death, nonfatal MI and nonfatal stroke</p> <p>Secondary: Blood pressure control rates</p>	<p>Primary: Rates (death, nonfatal MI and nonfatal stroke) were similar for both treatment groups (P value not reported).</p> <p>Secondary: Rates of death, MI and stroke declined as the number of office visits for which blood pressure was controlled increased (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Bangalore et al.<sup>165</sup> (2008) INVEST</p> <p>Verapamil SR 120 to 480 mg QD</p> <p>vs</p> <p>atenolol 25 to 200 mg QD</p> <p>Trandolapril and/or HCTZ were added to control blood pressure.</p>	<p>INVEST substudy</p> <p>Patients 50 years of age and older with hypertension requiring drug therapy (blood pressure &gt;140/90 or &gt;130/80 mm Hg if diabetic or with renal impairment), and documented coronary artery disease</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: First occurrence of death, nonfatal MI, nonfatal stroke</p> <p>Secondary: Death, total MI, total stroke</p>	<p>Primary: No significant difference was observed between groups in the primary endpoint (P=0.30).</p> <p>Among patients with the primary outcome, no significant difference was observed between groups in the risk of death (P=0.94).</p> <p>There was no significant difference between groups in the risk of nonfatal MI (P=0.41).</p> <p>There was a trend toward a 29% reduction in the risk of nonfatal stroke in the verapamil group compared to the atenolol group (P=0.06).</p> <p>Secondary: The risks of fatal and nonfatal MI were similar between groups.</p> <p>No significant differences were observed between groups in fatal and nonfatal stroke (P=0.18).o</p>
<p>Iliuta et al.<sup>166</sup> (2009)</p> <p>Betaxolol 20 mg/day</p> <p>vs</p> <p>metoprolol 100 mg BID</p>	<p>OL, MC</p> <p>Patients who were admitted for CABG surgery</p>	<p>N=1352</p> <p>30 days</p>	<p>Primary: Mortality, in-hospital occurrence of AF, total hospital stay and immobilization (days)</p> <p>Secondary: Not reported</p>	<p>Primary: Betaxolol significantly decreased 30 day mortality (P=0.001) and in-hospital AF (P=0.0001) compared to metoprolol.</p> <p>Patients taking betaxolol were less likely to be hospitalized for &gt;15 days (9.94 vs 13.27, P=0.01) or immobilized for &gt;3 days (5.19 vs 8.26, p=0.002) compared to metoprolol.</p> <p>Secondary: Not reported</p>
<p>Jonsson et al.<sup>167</sup> (2005)</p> <p>Carvedilol 6.25 to 25 mg BID</p> <p>vs</p> <p>atenolol 12.5 to 50</p>	<p>OL, RCT</p> <p>Patients between 18 to 80 years of age with chest pain consistent with an acute MI, admitted to the hospital 24 hours after onset</p>	<p>N=232</p> <p>1.5±1.3 years</p>	<p>Primary: Change in global or regional LVEF after 12 months, cardiovascular endpoints, adverse events</p> <p>Secondary:</p>	<p>Primary: At baseline, mean global LVEF was 54.8% in the carvedilol and 53.0% in the atenolol group and increased after 12 months to 57.1% in the carvedilol and 56.0% in the atenolol group. There was not a significant difference between treatment groups for change in global or regional LVEF (values were not reported).</p> <p>There was not a significant difference in the rates of occurrence of the first serious cardiovascular events observed between the carvedilol and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg BID	and a confirmed diagnosis with significant increase in cardiac enzymes		Not reported	atenolol groups after adjustment for diuretic use (0.247 vs 0.299; RR, 0.83; 95% CI, 0.56 to 1.23; P=0.39).  Of the nonserious adverse events reported, a greater incidence of colds hand and feet were reported in the atenolol group (38 [33.3%]) compared to the carvedilol group (24 [20%]; P=0.025).  Secondary: Not reported
Pasternak et al. <sup>168</sup> (2014)  Carvedilol  vs  metoprolol succinate	RETRO  Danish patients aged 50 to 84 years with HF and LVEF ≤40% who received carvedilol or metoprolol succinate treatment	N=11,664  Up to 3 years (Median 2.4)	Primary: All-cause mortality  Secondary: Cardiovascular mortality	Primary: The cumulative incidence of all-cause mortality was 18.3 and 18.8% in the carvedilol and metoprolol groups, respectively. After adjustment for propensity score, the risk of mortality did not differ significantly between carvedilol and metoprolol users (aHR, 0.99; 95% CI, 0.88 to 1.11).  Secondary: The risk of cardiovascular mortality was not significantly different between carvedilol and metoprolol users (aHR, 1.05; 95% CI, 0.88 to 1.26).
Seo et al. <sup>169</sup> (2015)  Carvedilol  vs  β1-selective β blockers (bisoprolol, metoprolol, and nebivolol)	PRO, propensity-score matched cohort  Patients ≥18 years of age with acute MI undergoing percutaneous coronary intervention	N=7,863  mean follow-up of 243 ± 144 days	Primary: All-cause death or MI during follow-up  Secondary: All-cause death, cardiac death, and MI	Primary: Of the 7,863 patients examined, 6,231 (79.2%) were treated with carvedilol and 1,632 (20.7%) were treated with β1-selective β-blockers. During the follow-up period, the primary end point of all-cause death or MI occurred in 94 patients (1.5%) in the carvedilol group and 31 patients (1.9%) in the β1-selective β-blockers group. In the multivariate Cox regression model, no differences in the risk of all-cause death or MI were observed between the carvedilol and β1-selective β-blocker groups.  Secondary: The risks of all-cause death, cardiac death, and MI were also similar between the carvedilol and β1-selective β-blocker groups during the follow-up period.
Olsson et al. <sup>170</sup> (1992)  Metoprolol 100 mg BID	MA (5 trials)  Patients with a past history of MI	N=5,474  3 months to 3 years	Primary: All-cause mortality, sudden deaths	Primary: Metoprolol significantly reduced all-cause mortality compared to placebo (188 vs 223 deaths; P=0.036).  Metoprolol significantly reduced sudden deaths compared to placebo (62

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			Secondary: Not reported	vs 104 deaths; P=0.002).  Secondary: Not reported
Piccini et al. <sup>171</sup> (2014)  Amiodarone  vs  sotalol  vs  no antiarrhythmic drug (AAD)	RETRO  Patients with CAD and AF	N=2,838  Median follow-up 4.2 years	Primary: All-cause mortality  Secondary: Not reported	Primary: In unadjusted and adjusted settings, mortality rates were lower in patients treated with sotalol compared with amiodarone or no AAD. After adjustment for baseline characteristics only, the 1-year mortality rate was 10% in those treated with sotalol, 20% in those treated with amiodarone, and 14% in those treated with no AAD (no P-value reported).  Landmark analysis at 60 days and one year was also performed. After adjustment and weighting, sotalol was associated with improved survival from 0 to 60 days compared with amiodarone (HR, 0.14; 95% CI, 0.06 to 0.32) but not at later time points ( $\geq 60$ days or $\geq 1$ year). Similarly, compared with no AAD therapy, sotalol was not associated with improved survival beyond 60 days. Cumulative survival after one year in patients treated with sotalol vs no AAD was also not improved (P=0.64).  Secondary: Not reported
No authors listed <sup>172</sup> (abstract) (1983)  Timolol  vs  placebo	DB, MC, PC, RCT  Patients <75 years of age surviving an acute MI	N=1,884  12 to 33 months	Primary: All-cause mortality  Secondary: Not reported	Primary: Long term treatment with timolol improved prognosis. A significant difference in life table mortality of 39.3% between treatments was observed (13.3 vs 21.9%; P=0.0003). The difference was due to a lower rate of sudden cardiac death with timolol compared to placebo (7.7 vs 13.9%; P=0.0001).  Secondary: Not reported
Patel et al. <sup>173</sup> (2014)  $\beta$ -blocker therapy (carvedilol, metoprolol)	RETRO  Medicare patients in the OPTIMIZE-HF registry (having a primary discharge)	N=2,198 (1099 propensity-matched pairs)  Up to 6 years	Primary: composite endpoint of all-cause mortality or HF rehospitalization	Primary: Discharge prescriptions for $\beta$ -blockers to older HF with preserved ejection fraction patients who were not receiving these drugs prior to admission had no association with the primary composite endpoint during a median of 2.2 years of follow-up (HR, 1.03; 95% CI, 0.94 to 1.13; P=0.569). This association was homogeneous across various clinically relevant



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
succinate, and bisoprolol at their respective guideline-recommended target doses of 50, 200, and 10 mg/day)  vs  no $\beta$ -blocker therapy	diagnosis of HF), aged $\geq 65$ years with EF $\geq 40\%$ who were eligible for new discharge prescriptions of $\beta$ -blockers	(Median 2.2)	Secondary: All-cause mortality, HF rehospitalization, and all-cause rehospitalization	subgroups.  Secondary: HRs for all-cause mortality and HF rehospitalization associated with a prescription for initiation of $\beta$ -blocker therapy were 0.99 (95% CI, 0.90 to 1.10; P=0.897) and 1.17 (95% CI, 1.03 to 1.34; P=0.014), respectively. The latter association lost significance when higher EF cutoffs of $\geq 45\%$ , $\geq 50\%$ and $\geq 55\%$ were used.
Hansson et al. <sup>174</sup> (2000) NORDIL  Conventional therapy (diuretic, $\beta$ -blocker or both)  vs  diltiazem 180 to 360 mg QD	BE, MC, OL, PRO, RCT  Patients 50 to 74 years of age with DBP $\geq 100$ mm Hg and previously untreated	N=10,881  4.5 years	Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death  Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI	Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/ $\beta$ -blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).  Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/ $\beta$ -blocker patients (P=0.04).  Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/ $\beta$ -blocker patients (P=0.17).  Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57) and congestive heart failure (P=0.42).
Messerli et al. <sup>175</sup> (1998)  $\beta$ -blockers (atenolol, metoprolol or pindolol)  vs	MA  10 RCTs lasting $\geq 1$ year, which used as first line agents diuretics and/or $\beta$ -blockers and reported morbidity and	N=16,164  1 year	Primary: Cardiovascular morbidity and mortality, all-cause morbidity  Secondary: Not reported	Primary: Diuretic treatment significantly reduced the odds for cardiovascular mortality by 25% (OR, 0.75; 95% CI, 0.64 to 0.87), while $\beta$ -blockers did not reduce cardiovascular mortality (OR, 0.98; 95% CI, 0.78 to 1.23; P values not reported).  Diuretic treatment significantly reduced the odds for all-cause mortality by 14% (OR, 0.86; 95% CI, 0.77 to 0.96), while $\beta$ -blockers did not reduce all-cause mortality (OR, 1.05; 95% CI, 0.88 to 1.25; P values not

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
diuretics (amiloride, chlorthalidone, HCTZ, HCTZ and triamterene [fixed-dose combination product], or thiazide)	mortality outcomes in patients $\geq 60$ years of age with HTN			reported).  Secondary: Not reported
Wysong et al. <sup>176</sup> (2007)  $\beta$ -blockers (atenolol, metoprolol, oxprenolol*, or propranolol)  vs  other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)	MA  13 RCTs evaluating patients $\geq 18$ years of age with HTN	N=91,561  Duration varied	Primary: All-cause mortality  Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions	Primary: There was not a significant difference observed in all-cause mortality between $\beta$ -blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with $\beta$ -blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).  Secondary: There was a significant decrease in stroke observed with $\beta$ -blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with $\beta$ -blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).  CHD risk was not significantly different between $\beta$ -blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07)), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).  The risk of total cardiovascular disease was lower with $\beta$ -blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of $\beta$ -blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with <math>\beta</math>-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>
<p>Lindholm et al.<sup>177</sup> (2005)</p> <p><math>\beta</math>-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol, or propranolol)</p> <p>vs</p> <p>other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a <math>\beta</math>-blocker as first-line treatment (in <math>\geq 50\%</math> of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>The RR of stroke was 16% higher with <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non <math>\beta</math>-blockers (RR, 1.26%; 95% CI, 1.15 to 1.38; P&lt;0.0001).</p> <p>The relative risk of MI was 2% higher for <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).</p> <p>The RR of all-cause mortality was 3% higher for <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
or placebo				
Freemantle et al. <sup>178</sup> (1999)  β-blockers (acebutolol, alprenolol, atenolol, betaxolol, carvedilol, labetalol, oxprenolol*, pindolol, practolol*, propranolol, sotalol, timolol and xamoterol*)  vs  control (agents were not specified)	MA (82 trials)  Patients with acute or past MI	N=54,234  6 to 48 months	Primary: All-cause mortality  Secondary: Nonfatal reinfarction and withdrawal from treatment	Primary: The pooled random effects in short term trials demonstrated a mortality rate of 10.5% (3,062 out of 29,260 patients) which is a 4% reduction compared to the controlled groups (OR, 0.96; 95% CI, 0.85 to 1.08).  The pooled random effects in long term trials demonstrated a mortality rate of 9.7% (2415 out of 24974 patients) which is 23% reduction when compared to the controlled groups (OR, 0.77; 95% CI, 0.69 to 0.85).  Individually, only four drugs achieved a statistically significant reduction in the death: propranolol (OR, 0.71; CI, 0.59 to 0.85]), timolol (OR, 0.59; CI, 0.46 to 0.77), metoprolol (OR, 0.80; CI, 0.66 to 0.96; and acebutolol (OR, 0.49; CI, 0.25 to 0.93).  Secondary: A reduction in nonfatal re-infarctions of 0.9 events in every 100 (0.3 to 1.6) annually is suggested by this analysis; therefore about 107 patients would require treatment for one year to avoid one nonfatal reinfarction.  Overall, 5,151 of 21,954 patients (23.5%) withdrew from treatment. with withdrawal occurring more often in the β-blocker groups. When comparing to placebo, the difference in annualized rate of withdrawal was 1.16 in 100 patients treated (1.16; 95% CI, 0.56 to 1.76).
<b>Miscellaneous</b>				
Schellenburg et al. <sup>179</sup> (2008)  Metoprolol 47.5 to 142.5 mg/day  vs  nebivolol 5	DB, PRO, RCT  Patients 18 to 65 years of age with the diagnosis of migraine with/ without aura, ≥1 year history, onset prior to 50 years of age, written record	N=38  30 weeks	Primary: Number of migraine attacks  Secondary: Onset of action, duration of attacks, responder rate, severity, use of pain medication,	Primary: There was not a significant difference in the frequency of migraine attacks observed between metoprolol and nebivolol (1.3±1.0 vs 1.6±1.5, respectively; P value not reported).  Secondary: There was not a significant difference in any of the secondary endpoints observed between metoprolol and nebivolol (P values not reported).  Use of acute pain medication decreased with both treatments, as well as

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day	of attacks for the previous 3 months and $\geq 2$ attack/month during screening		migraine disability assessment, QOL score	accompanying symptoms. Both patient and physician evaluations of disability and disease status were similarly favorable to the two treatments (P values not reported).
Silberstein et al. <sup>180</sup> (2011)  Propranolol ER 240 mg/day  vs  placebo	DB, MC, PC, RCT  Patients with chronic migraine inadequately controlled ( $\geq 10$ headaches/month) with topiramate (50 to 100 mg/day)	N=191  6 months	Primary: 28 day moderate to severe headache rate reduction at six months (weeks 16 to 24) compared to baseline (weeks -4 to 0)  Secondary: Not reported	Primary: The six month reduction in moderate to severe 28 day headache rate and total 28 day headache rate for combination therapy vs topiramate was not significantly different (4.0 vs 4.5 days; P=0.57 and 6.2 vs 6.1; P=0.91).  Secondary: Not reported
Tfelt-Hansen et al. <sup>181</sup> (1984)  Timolol 10 mg BID  vs  propranolol 80 mg BID  vs  placebo  All patients entered a 4 week pretreatment period.	DB, PC, RCT, XO  Patients 18 to 65 years of age with a history of 2 to 6 common migraine attacks per month	N=96  40 weeks	Primary: Frequency, duration and severity of attacks; number of responders ( $\geq 50\%$ reduction in the frequency of attacks compared to baseline)  Secondary: Frequency of attacks with associated symptoms, frequency of attacks requiring relief medication	Primary: Both timolol and propranolol decreased the frequency of attacks from baseline (P<0.01 for both).  For severity of headache attacks, a small but significant reduction was observed with timolol (P<0.05 vs baseline).  There was no effect on duration of attacks with either timolol or propranolol.  The number of responders was significantly higher with timolol (n=44) and propranolol (n=48) compared to placebo (n=24; P<0.01 for both).  Secondary: Both timolol and propranolol decreased the frequency of attacks associated with nausea or frequency of attacks associated with symptomatic therapy (P<0.01 for both vs baseline).
Linde et al. <sup>182</sup> (2004)	MA	N=5,072	Primary: Headache and	Primary: Compared to placebo, propranolol showed a significant advantage in

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Propranolol 60 to 320 mg/day</p> <p>vs</p> <p>placebo or another agent (calcium channel blockers, other <math>\beta</math>-blockers or other agent)</p>	<p>26 randomized and quasi-randomized clinical trials of <math>\geq 4</math> weeks duration comparing clinical effects of propranolol with placebo or another drug in adult patients with migraine</p>	<p>4 to 30 weeks</p>	<p>migraine frequency</p> <p>Secondary: Not reported</p>	<p>response to treatment with overall RR of response (“responder ratio”) of 1.94 (95% CI, 1.61 to 2.35).</p> <p>Compared to placebo, propranolol showed a significant advantage for the reduction of frequency of migraines with overall mean difference of -0.40 (95% CI, -0.56 to -0.24).</p> <p>Propranolol did not demonstrate a significantly greater response to treatment compared to calcium channel blockers with an overall responder ratio of 1.00 (95% CI, 0.92 to 1.09).</p> <p>Propranolol did not demonstrate a significantly greater reduction in migraine frequency compared to calcium channel blockers with an overall mean difference of -0.02 (95% CI, 0.12 to 0.08).</p> <p>In the three trials comparing propranolol and nadolol, the overall responder ratio favored nadolol (responder ratio, 0.60; 95% CI, 0.37 to 0.97), but the results of the three trials were contradictory.</p> <p>In the three trials comparing propranolol and metoprolol, there was not a significant difference observed in the overall responder ratio between the two treatments (responder ratio, 0.78; 95% CI, 0.56 to 1.09).</p> <p>Propranolol did not demonstrate a significantly greater reduction in migraine frequency compared to other <math>\beta</math>-blockers with an overall mean difference of -0.01 (95% CI, 0.24 to 0.22).</p> <p>A quantitative MA was not performed on trials comparing propranolol to other drugs due to the great variety of comparator drugs used. One trial was significantly in favor of propranolol (vs amitriptyline), five with a trend in favor of propranolol, 11 showing no difference, two with a trend in favor of the comparator drug and one not interpretable; one of the two comparisons of propranolol alone and propranolol in combination with amitriptyline was classified as no difference, and the other as showing a trend in favor of the combination (P values not reported).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
Léauté-Labrèze et al. <sup>183</sup> (2015)  Propranolol (1 or 3 mg/kg/day, divided into two daily doses)  vs  placebo BID	DB, PC, RCT  Patients 35 to 150 days of age with a proliferating infantile hemangioma requiring systemic therapy	N=460  24 to 96 weeks	Primary: Success (complete or nearly complete resolution of the target hemangioma) or failure of trial treatment at week 24 versus baseline according to centralized evaluation  Secondary: Success or failure of trial treatment according to on-site assessments by the investigator at week 48 versus baseline	Primary: At the time of the interim analysis (188 patients completing 24 weeks of therapy), 2 of 25 patients (8%) receiving placebo had successful treatment at week 24, as compared with 4 of 41 patients (10%) receiving 1 mg/kg/day of propranolol for 3 months, 3 of 39 patients (8%) receiving 3 mg/kg/day for 3 months, 15 of 40 patients (38%) receiving 1 mg/kg/day for 6 months (P=0.004 for the comparison with placebo), and 27 of 43 patients (63%) receiving 3 mg/kg/day for 6 months (P<0.001 for the comparison with placebo).  Overall, 61 of 101 patients (60%) assigned to the selected propranolol regimen and 2 of 55 patients (4%) assigned to placebo had successful treatment at week 24 (P<0.001).  Improvement between baseline and week 5 (according to centralized assessment) occurred in 88% of patients assigned to the selected regimen and 5% of patients assigned to placebo (P<0.001).  Secondary: Not reported

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, CR=controlled-release, ER=extended-release, QD=once daily, SR=sustained-release, TID=three times daily, XL=extended-release

Study design abbreviations: AC=active comparator, BE=blinded endpoint, DB=double blind, DD=double dummy, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, RETRO=retrospective, SB=single blind, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitor, AF=atrial fibrillation, Aix=augmentation index, aPWV=aortic pulse wave velocity, ARB=angiotensin II receptor blocker, CABG=coronary artery bypass graft, CAD=coronary artery disease, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, COPD=chronic obstructive pulmonary disease, DBP=diastolic blood pressure, ECG=electrocardiogram, ESRD=end stage renal disease, FEV<sub>1</sub>=forced expiratory volume in one second, GFR=glomerular filtration rate, HDL-C=high-density lipoprotein cholesterol, HR=hazard ratio, HTN=hypertension, LDL-C=low-density lipoprotein cholesterol, LVEF=left ventricular ejection fraction, MAP=mean arterial pressure, MI=myocardial infarction, NYHA=New York Heart Association, OR=odds ratio, PAD=peripheral arterial disease, pro-BNP= pro-B-type natriuretic peptide, PVD=peripheral vascular disease, QOL=quality of life, RMSSD=root mean square of successive RR intervals, RR=relative risk, SBP=systolic blood pressure, SDNN=standard deviation of the normal RR intervals, TC=total cholesterol, TG=triglyceride, TIA=transient ischemic attack, WHO=World Health Organization

**Additional Evidence**

Dose Simplification

Nissinen et al. evaluated newly diagnosed hypertensive patients who received atenolol 100 mg and chlorthalidone 25 mg given as single entity products or as a fixed-dose combination. Each of the active drug combinations significantly lowered standing, supine and postexercise blood pressure. There was no significant difference among the treatment regimens.<sup>114</sup>

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 18. Relative Cost of the Beta-Adrenergic Blocking Agents**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Acebutolol	capsule	N/A	N/A	\$\$\$
Atenolol	tablet	Tenormin <sup>®*</sup>	\$\$\$\$\$	\$
Betaxolol	tablet	N/A	N/A	\$\$
Bisoprolol	tablet	N/A	N/A	\$
Carvedilol	extended-release capsule, tablet	Coreg <sup>®*</sup> , Coreg CR <sup>®*</sup>	\$\$\$\$\$	\$
Labetalol	injection, tablet	N/A	N/A	\$
Metoprolol	extended-release capsule, extended-release tablet, injection, tablet	Kapsargo Sprinkle <sup>®</sup> , Lopressor <sup>®*</sup> , Toprol-XL <sup>®*</sup>	\$\$\$ to \$\$\$\$	\$
Nadolol	tablet	Corgard <sup>®*</sup>	\$\$\$\$\$	\$
Nebivolol	tablet	Bystolic <sup>®*</sup>	\$\$\$\$	\$\$\$
Penbutolol	tablet	Levatol <sup>®</sup>	N/A <sup>†</sup>	N/A
Pindolol	tablet	N/A	N/A	\$\$\$



Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Propranolol	extended-release capsule, injection, solution, tablet	Hemangeol <sup>®</sup> , Inderal LA <sup>®*</sup> , Inderal XL <sup>®</sup> , InnoPran XL <sup>®</sup>	\$\$\$\$\$	\$
Sotalol	tablet, solution	Betapace <sup>®*</sup> , Betapace AF <sup>®*</sup> , Sotylize <sup>®</sup>	\$\$\$\$\$	\$
Timolol	tablet	N/A	N/A	\$\$\$\$
<b>Combination Products</b>				
Atenolol and chlorthalidone	tablet	Tenoretic <sup>®*</sup>	\$\$\$\$	\$
Bisoprolol and HCTZ	tablet	Ziac <sup>®*</sup>	\$\$\$\$\$	\$
Metoprolol and HCTZ	tablet	N/A	N/A	\$\$\$
Nadolol and bendroflumethiazide	tablet	N/A	N/A	\$\$\$

\*Generic is available in at least one dosage form or strength.

†Product was discontinued by manufacturer, but remains active in pharmacy system. Pricing information not available.

HCTZ=hydrochlorothiazide, N/A=not available

## X. Conclusions

All of the beta-adrenergic blocking agents ( $\beta$ -blockers) are approved for the treatment of hypertension, with the exception of sotalol.<sup>1-22</sup> Some of the products are also approved for the treatment of angina, arrhythmias, essential tremor, heart failure, hypertension, hypertrophic aortic stenosis, infantile hemangiomas, migraine prophylaxis, myocardial infarction, and pheochromocytoma.<sup>1-22</sup> These agents differ with regards to their adrenergic-receptor blockade, membrane stabilizing and intrinsic sympathomimetic activities, as well as lipophilicity.<sup>1,2,25</sup> All of the agents are available in a generic formulation, with the exception of penbutolol.

Several national and international guidelines address the use of  $\beta$ -blockers.<sup>26-56</sup> Due to improvements in cardiovascular morbidity and mortality, treatment guidelines recommend the use of a  $\beta$ -blocker in patients with the following conditions: acute coronary syndromes, angina, arrhythmias, coronary artery disease, heart failure, left ventricular dysfunction, post-myocardial infarction, and infantile hemangiomas.<sup>26-56</sup> There are several published guidelines on the treatment of hypertension. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>40-48</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>42</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>40-48</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>40-47</sup>  $\beta$ -blockers are recommended as one of several initial options for the prevention of migraine headaches (metoprolol and propranolol), as well as for the treatment of essential tremor (propranolol).<sup>50-52,54,55</sup> The Clinical Practice Guideline for the Management of Infantile Hemangiomas was published in January 2019 and recommends the use of oral propranolol as the first-line agent for the treatment.<sup>56</sup> The oral solution formulation of propranolol, under the brand name Hemangeol<sup>®</sup>, is the first agent to gain FDA-approval for this indication.<sup>12</sup>

Numerous clinical trials have shown that the  $\beta$ -blockers can effectively lower blood pressure when administered alone or in combination with other antihypertensive agents. Comparative studies have demonstrated similar efficacy among the  $\beta$ -blockers.<sup>102-148</sup> Most patients will require more than one antihypertensive agent to achieve blood pressure goals.<sup>40-47</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>43,44</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations.<sup>105</sup>

In patients with chronic stable angina,  $\beta$ -blockers improve exercise tolerance and reduce the frequency of attacks. Head-to-head trials have demonstrated similar efficacy among several of the  $\beta$ -blockers.<sup>57-65</sup> In patients with heart

failure,  $\beta$ -blockers (bisoprolol, carvedilol, and metoprolol succinate) have been shown to reduce mortality, sudden death, cardiovascular deaths, and death due to heart failure. Clinical trials supporting the use of carvedilol in patients with mild-to-severe heart failure were conducted with the immediate-release formulation.<sup>76-83,86,87,91-94,96-98</sup> Data to support the use of the extended-release capsules for the treatment of heart failure is based on pharmacokinetic and pharmacodynamic parameters that demonstrated bioequivalence with the immediate-release formulation.<sup>6</sup>

In general, adverse events are similar among the  $\beta$ -blockers. Common adverse effects include fatigue, cold hands, dizziness, and weakness.<sup>1-22</sup>  $\beta$ -blockers that are more selective for the  $\beta_1$ -receptors (atenolol and metoprolol) may be safer to use in those with reactive airway disease as they are less likely to cause bronchospasm.<sup>23,25</sup>

There is insufficient evidence to support that one brand beta-adrenergic blocking agent is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand beta-adrenergic blocking agents within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand beta-adrenergic blocking agent is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. Sectral [package insert]. Bridgewater (NJ): Promius Pharma, LLC; 2012 Jan.
4. Tenormin® [package insert]. Morristown (NJ): Almatica Pharma LLC; 2021 Jan.
5. Coreg® [package insert]. Research Triangle Park (NC): GlaxoSmithKline; 2017 Sep.
6. Coreg CR® [package insert]. Research Triangle Park (NC): GlaxoSmithKline; 2020 Nov.
7. Lopressor® [package insert]. East Hanover (NJ): Validus Pharmaceuticals LLC; 2019 Mar.
8. Toprol-XL® [package insert]. Wilmington (DE): AstraZeneca LP; 2018 Dec.
9. Corgard® [package insert]. Louisville (KY): USWM, LLC; 2021 Aug.
10. Bystolic® [package insert]. Madison (NJ): Allergan USA, Inc.; 2019 Jun.
11. Levatol® [package insert]. Smyrna (GA): Schwarz Pharma; 2011 May.
12. Hemangeol® [package insert]. Parsippany (NJ): Pierre Fabre Pharmaceuticals, Inc.; 2021 Jun.
13. Inderal LA® [package insert]. Cranford (NJ): Akrimax Pharmaceuticals, LLC; 2019 Aug.
14. InnoPran XL® [package insert]. Cranford (NJ): Akrimax Pharmaceuticals, LLC; 2021 Aug.
15. Betapace® [package insert]. Zug (Switzerland): Covis Pharma; 2021 Jun.
16. Betapace AF® [package insert]. Wayne (NJ): Covis Pharma; 2021 Jun.
17. Tenoretic® [package insert]. Morristown (NJ): Almatica Pharma LLC; 2021 Jan.
18. Ziac® [package insert]. Parsippany (NJ): Teva Pharmaceuticals USA, Inc; 2021 Mar.
19. Dutoprol® [package insert]. Cary (NC): Concordia Pharmaceuticals, Inc; 2020 Aug.
20. Lopressor HCT® [package insert]. Suffern (NY): Novartis Pharmaceuticals Corporation; 2019 Mar.
21. Corzide® [package insert]. New York (NY): Pfizer, Inc; 2019 Mar.
22. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
23. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed Jan 2022.
24. Metry DW. Infantile hemangiomas: Management. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2022 Jan]. Available from: <http://www.utdol.com/utd/index.do>.
25. Podrid PJ. Major side effects of beta blockers. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2022 Jan]. Available from: <http://www.utdol.com/utd/index.do>.
26. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425.
27. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline From the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
28. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
29. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>.
30. O’Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.

31. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
32. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].
33. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
34. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
35. January CT, Wann L, Calkins H, et al. 2019 Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2019;140:e125-e151. doi:10.1161/CIR.0000000000000665.
36. National Institute for Health and Care Excellence. Atrial fibrillation: diagnosis and management. NG196. 2021. <https://www.nice.org.uk/guidance/ng196>.
37. Frendl G, Sodickson AC, Chung MK, et al. 2014 AATS Guidelines for the Prevention and Management of Peri-Operative Atrial Fibrillation and Flutter (POAF) for Thoracic Surgical Procedures. *The Journal of thoracic and cardiovascular surgery*. 2014;148(3):e153-e193. doi:10.1016/j.jtcvs.2014.06.036.
38. Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, et al. 2017 ACC/AHA/ESC guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines and the Heart Rhythm Society. *Circulation*. 2018;138:e272-e391. Doi:10.1161/CIR.0000000000000549.
39. The Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy. *European Heart Journal* 2014;35:2733–2779. doi:10.1093/eurheartj/ehu284.
40. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
41. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
42. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
43. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104
44. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
45. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
46. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003
47. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of

- the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Hypertension. 2018 Jun; 71(6): 1269-1324.
48. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. Diabetes Care 2022;45(Suppl. 1).
  49. Mayans L, Walling A.. Acute Migraine Headache: Treatment Strategies. Am Fam Physician. 2018 Feb 15;97(4):243-251.
  50. Ha H, Gonzalez, A. Migraine Headache Prophylaxis. Am Fam Physician. 2019 Jan 199(1):17-24.
  51. Silberstein SD, Holland S, Freitag F, Dodick DW, Argoff C, Ashman E. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults. Neurology 2012;78(17):1337-1345.
  52. Evers S, Afra J, Frese A, et al.; European Federation of Neurological Societies. EFNS guideline on the drug treatment of migraine--revised report of an EFNS task force. Eur J Neurol 2009;16:968-81.
  53. National Cancer Institute. Pheochromocytoma and paraganglioma treatment (PDQ®)-Health Professional Version [Internet]. Bethesda (MD): National Institutes of Health (US), National Cancer Institute; 2021 Sep 30 [cited 2022 Jan]. Available from: <https://www.cancer.gov/types/pheochromocytoma/hp/pheochromocytoma-treatment-pdq>.
  54. Zesiewicz TA, Elble R, Louis ED, Hauser RA, Sullivan KL, Dewey RB Jr, et al. Practice Parameter: Therapies for essential tremor: Report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology. 2005 Jun 28;64(12):2008-20.
  55. Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: Treatment of essential tremor. Neurology 2011;77(19):1752-1755.
  56. Krowchuck DP, Frieden IJ, Mancini AJ, et al. Clinical Practice Guideline for the Management of Infantile Hemangiomas. American Academy of Pediatrics. 2019 Jan 143(1):e20183475; DOI: 10.1542/peds.2018-3475
  57. Pandhi P, Sethi V, Sharma BK, Wahi PL, Sharma PL. Double blind cross-over clinical trial of acebutolol and propranolol in angina pectoris. Int J Clin Pharmacol Ther Toxicol. 1985 Nov;23(11):598-600.
  58. Jackson G, Schwartz J, Kates RE, Winchester M, Harrison DC. Atenolol: once-daily cardioselective beta blockade for angina pectoris. Circulation. 1980;61(3):555-60.
  59. Oh PC, Kang WC, Moon J, Park YM, Kim S, Kim MG, et al. Anti-Anginal and Metabolic Effects of Carvedilol and Atenolol in Patients with Stable Angina Pectoris: A Prospective, Randomized, Parallel, Open-Label Study. Am J Cardiovasc Drugs. 2016 Jun;16(3):221-8.
  60. Kardas P. Compliance, clinical outcome, and quality of life of patients with stable angina pectoris receiving once-daily betaxolol versus twice daily metoprolol: a randomized controlled trial. Vasc Health Risk Manag. 2007 Jun;3(2):235-42.
  61. van der Does R, Hauf-Zachariou U, Pfarr E, Holtbrügge W, König S, Griffiths M, Lahiri A. Comparison of safety and efficacy of carvedilol and metoprolol in stable angina pectoris. Am J Cardiol. 1999 Mar 1;83(5):643-9.
  62. Weiss R, Ferry D, Pickering E, Smith LK, Dennish G 3rd, Krug-Gourley S, et al; Effectiveness of three different doses of carvedilol for exertional angina. Carvedilol-Angina Study Group. Am J Cardiol. 1998 Oct 15; 82(8):927-31.
  63. Hauf-Zachariou U, Blackwood RA, Gunawardena KA, O'Donnell JG, Garnham S, Pfarr E. Carvedilol versus verapamil in chronic stable angina: a multicentre trial. Eur J Clin Pharmacol. 1997;52(2):95-100.
  64. Savonitto S, Ardissio D, Egstrup K, Rasmussen K, Bae EA, Omland T, et al. Combination therapy with metoprolol and nifedipine versus monotherapy in patients with stable angina pectoris. Results of the International Multicenter Angina Exercise (IMAGE) Study. J Am Coll Cardiol. 1996 Feb;27(2):311-6.
  65. Turner GG, Nelson RR, Nordstrom LA, Diefenthal HC, Gobel FL. Comparative effect of nadolol and propranolol on exercise tolerance in patients with angina pectoris. Br Heart J. 1978;40:1361-70.
  66. Lui HK, Lee G, Dhurandhar R, Hungate EJ, Laddu A, Dietrich P, et al. Reduction of ventricular ectopic beats with oral acebutolol: a double-blind, randomized crossover study. Am Heart J. 1983;105:722-6.
  67. Lee CH, Nam GB, Park HG, Kim HY, Park KM, Kim J, et al. Effects of antiarrhythmic drugs on inappropriate shocks in patients with implantable cardioverter defibrillators. Circ J. 2008 Jan;72(1):102-5.
  68. Connolly SJ, Dorian P, Roberts RS, et al. Comparison of  $\beta$ -blockers, amiodarone plus  $\beta$ -blockers, or sotalol for prevention of shocks from implantable cardioverter defibrillators: The OPTIC study: A randomized trial. JAMA. 2006 Jan 11;295:165-71.
  69. Balcetyte-Harris N, Tamis JE, Homel P, Menchavez E, Steinberg JS. Randomized study of early intravenous esmolol versus oral beta-blockers in preventing post-CABG atrial fibrillation in high risk patients identified by signal-averaged ECG: Results of a pilot study. A N E. 2002 Apr;7(2):86-91.

70. Kettering K, Mewis C, Dörnberger V, Vonthein R, Bosch RF, Kühlkamp V. Efficacy of metoprolol and sotalol in the prevention of recurrences of sustained ventricular tachyarrhythmias in patients with an implantable cardioverter defibrillator. *Pacing Clin Electrophysiol.* 2002 Nov;25(11):1571-6.
71. Seidl K, Hauer B, Schwick NG, Zahn R, Senges J. Comparison of metoprolol and sotalol in preventing ventricular tachyarrhythmias after the implantation of a cardioverter/defibrillator. *Am J Cardiol.* 1998 Sep 15;82(6):744-8.
72. Steeds RP, Birchall AS, Smith M, Channer KS. An open label, randomized, crossover study comparing sotalol and atenolol in the treatment of symptomatic paroxysmal atrial fibrillation. *Heart.* 1999 Aug;82(2):170-5.
73. Calzetti S, Findley LJ, Gresty MA, Perucca E. Metoprolol and propranolol in essential tremor: a double-blind, controlled study. *J Neurol Neurosurg Psychiatry.* 1981 Jul;44(9):814-9.
74. Yetimalar Y, Gulumser I, Kurt T, Basoglu M. Olanzapine versus propranolol in essential tremor. *Clin Neurol Neurosurg.* 2005 Dec;108(1):32-5.
75. Gironell A, Kulisevsky J, Barbanoj M, López-Villegas D, Hernández G, Pascual-Sedano B. A randomized placebo-controlled comparative trial of gabapentin and propranolol in essential tremor. *Arch Neurol.* 1999 Apr;56(4):475-80.
76. CIBIS Investigators and Committees. A randomized trial of beta-blockade in heart failure. The Cardiac Insufficiency Bisoprolol Study (CIBIS). *Circulation.* 1994 Oct;90(4):1765-73.
77. CIBIS-II Investigators and Committees. The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomized trial. *Lancet.* 1999 Jan 2;353(9146):9-13.
78. Contini M, Apostolo A, Cattadori G, et al. Multiparametric comparison of CARvedilol, vs. NEbivolol, vs. Bisoprolol in moderate heart failure: the CARNEBI trial. *Int J Cardiol.* 2013 Oct 3;168(3):2134-2140.
79. Willenheimer R, van Veldhuisen DJ, Silke B, Erdmann E, Follath F, Krum H, et al; CIBIS III Investigators. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite sequence: results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation.* 2005 Oct 18;112(16):2426-35.
80. Packer M, Coats AJ, Fowler MB, Katus HA, Krum H, Mohacsi P, et al; Carvedilol Prospective Randomized Cumulative Survival Study Group. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med.* 2001 May 31;344(22):1651-8.
81. Packer M, Fowler MB, Roecker EB, Coats AJS, Katus HA, Krum H, et al. Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the carvedilol prospective randomized cumulative survival (COPERNICUS) study. *Circulation.* 2002;106:2194-99.
82. Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM, et al. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. *N Engl J Med.* 1996 May 23; 334(21):1349-55.
83. Dargie HJ; CAPRICORN Investigators. Effect of carvedilol on outcome after myocardial infarction in patients with left-ventricular dysfunction: the CAPRICORN randomised trial. *Lancet.* 2001 May 5;357(9266):1385-90.
84. Krum H, Sackner-Berstein JD, Goldsmith RL, Kukin ML, Schwartz B, Penn J, et al. Double-blind, placebo-controlled study of the long-term efficacy of carvedilol in patients with severe chronic heart failure (abstract). *Circulation.* 1995;92:1499-506.
85. Bristow MR, Gilbert EM, Abraham WT, Adams KF, Fowler MB, Hershberger RE, et al. Carvedilol produces dose-related improvements in left ventricular function and survival in subjects with chronic heart failure. MOCHA Investigators. *Circulation.* 1996 Dec 1;94(11):2807-16.
86. Fröhlich H, Zhao J, Täger T, Cebola R, Schellberg D, Katus HA, et al. Carvedilol Compared With Metoprolol Succinate in the Treatment and Prognosis of Patients With Stable Chronic Heart Failure: Carvedilol or Metoprolol Evaluation Study. *Circ Heart Fail.* 2015 Sep;8(5):887-96.
87. Poole-Wilson PA, Swedberg K, Cleland JG, Di Lenarda A, Hanrath P, et al; Carvedilol Or Metoprolol European Trial Investigators. Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the Carvedilol Or Metoprolol European Trial (COMET): randomised controlled trial. *Lancet.* 2003 Jul 5;362(9377):7-13.
88. Packer M, Antonopoulos GV, Berlin JA, Chittams J, Konstam MA, Udelson JE. Comparative effects of carvedilol and metoprolol on left ventricular ejection fraction in heart failure: results of a meta-analysis. *Am Heart J.* 2001 Jun;141(6):899-907.
89. Arumanayagam M, Chan S, Tong S, Sanderson JE. Antioxidant properties of carvedilol in heart failure: a double-blind randomized controlled trial. *J Cardiovas Pharmacol.* 2001 Jan;37(1):48-54.
90. Sanderson JE, Chan SK, Yip G, Yeung LY, Chan KW, Raymond K, Woo KS. Beta-blockade in heart failure: a comparison of carvedilol with metoprolol. *J Am Coll Cardiol.* 1999 Nov 1;34(5):1522-8.

91. Lechat P, Packer M, Chalon S, Cucherat M, Arab T, Boissel JP. Clinical effects of beta-adrenergic blockade in chronic heart failure: a meta-analysis of double-blind, placebo-controlled, randomized trials. *Circulation*. 1998 Sep 22;98(12):1184-91.
92. Brophy JM, Joseph L, Rouleau JL. Beta-blockers in congestive heart failure. A Bayesian meta-analysis. *Ann Intern Med*. 2001 Apr 3;134(7):550-60.
93. Whorlow SL, Krum H. Meta-analysis of effect of beta-blocker therapy on mortality in patients with New York Heart Association class IV chronic congestive heart failure. *Am J Cardiol*. 2000 Oct 15;86(8):886-9.
94. Bouzamondo A, Hulot JS, Sanchez P, Lechat P. Beta-blocker benefit according to severity of heart failure. *Eur J Heart Fail*. 2003 Jun;5(3):281-9.
95. Jabbour A, Macdonald, PS, Keogh AM, Kotlyar E, Mellemkjaer S, et al. Differences between beta-blockers in patients with chronic heart failure and chronic obstructive pulmonary disease. *J Am Coll Cardiol* 2010;55:1780-7.
96. MERIT-HF Study Group. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomized Intervention Trial in Congestive Heart Failure (MERIT-HF) *Lancet*. 1999 Jun 12;353(9169):2001-7.
97. Goldstein S, Fagerberg B, Hjalmarson A, Kjekshus J, Waagstein F, Wedel H, et al. Metoprolol controlled release/extended release in patients with severe heart failure. *J Am Coll Cardiol*. 2001;38:932-8.
98. Waagstein F, Bristow MR, Swedberg K, Camerini F, Fowler MB, Silver MA, et al. Beneficial effects of metoprolol in idiopathic dilated cardiomyopathy. Metoprolol in Dilated Cardiomyopathy (MDC) Trial Study Group. *Lancet*. 1993 Dec 11;342(8885):1441-6.
99. Di Lenarda A, Sabbadini G, Salvatore L, Sinagra G, Mestroni L, Pinamonti B, et al. Long-term effects of carvedilol in idiopathic dilated cardiomyopathy with persistent left ventricular dysfunction despite chronic metoprolol. The Heart-Muscle Disease Study Group. *J Am Coll Cardiol*. 1999 Jun;33(7):1926-34.
100. Maack C, Elter T, Nickenig G, LaRosee K, Crivaro M, Stäblein A, et al. Prospective crossover comparison of carvedilol and metoprolol in patients with chronic heart failure. *J Am Coll Cardiol*. 2001 Oct;38(4):939-46.
101. Metra M, Giubbini R, Nodari S, Boldi E, Modena MG, Dei Cas L. Differential effects of beta-blockers in patients with heart failure: A prospective, randomized, double-blind comparison of the long-term effects of metoprolol versus carvedilol. *Circulation*. 2000 Aug 1;102(5):546-51.
102. Reim HG, Wagner W. Hemodynamic effects of acebutolol and propranolol in hypertensive patients during exercise. *Int J Clin Pharmacol Ther Toxicol*. 1985 May;23(5):238-43.
103. Fogari R, Zoppi A. Half-strength atenolol-chlorthalidone combination (Tenoretic mite) in the treatment of elderly hypertensive patients. *Int J Clin Pharmacol Ther Toxicol*. 1984 Jul;22(7):386-93.
104. Leonetti G, Pasotti C, Capra A. Low-dose atenolol-chlorthalidone combination for treatment of mild hypertension. *Int J Clin Pharmacol Ther Toxicol*. 1986 Jan;24(1):43-7.
105. Nissinen A, Tuomilehto J. Evaluation of the antihypertensive effect of atenolol in fixed or free combination with chlorthalidone. *Pharmatherapeutica*. 1980;2(7):462-8.
106. Johnson JA, Gong Y, Bailey KR, et al. Hydrochlorothiazide and atenolol combination antihypertensive therapy: Effects of drug initiation order. *Clin Pharmacol Ther*. 2009;86:533-9.
107. Dhakam Z, Yasmin, McEniery CM, Burton T, Brown MJ, Wilkinson IB, et al. A comparison of atenolol and nebivolol in isolated systolic hypertension. *J Hypertens*. 2008 Feb;26(2):351-6.
108. Fogari R, Zoppi A, Lazzari P, Mugellini A, Lusardi P, Preti P, Van Nueten L, Vertommen C. Comparative effects of nebivolol and atenolol on blood pressure and insulin sensitivity in hypertensive subjects with type II diabetes. *J Hum Hypertens*. 1997 Nov;11(11):753-7.
109. Dietz R, Dechend R, Yu C, et al. Effects of the direct renin inhibitor aliskiren and atenolol alone or in combination in patients with hypertension. *J Renin Angiotensin Aldosterone Syst* 2008;9:163-175.
110. Wald DS, Law M, Mills S, Bestwick JP, Morris JK, Wald NJ. A 16-week, randomized, double-blind, placebo-controlled, crossover trial to quantify the combined effect of an angiotensin-converting enzyme inhibitor and a beta-blocker on blood pressure reduction. *Clin Ther*. 2008 Nov;30(11):2030-9.
111. Pareek A, Salkar H, Mulay P, Desai S, Chandurkar N, Redkar N. A randomized, comparative, multicenter, evaluation of atenolol/amlodipine combination with atenolol alone in essential hypertension patients. *Am J Ther*. 2010;17:46-52.
112. Chapman N, Dobson J, Wilson S, Dahlöf B, Sever PS, Wedel H, Poulter NR; Anglo-Scandinavian Cardiac Outcomes Trial Investigators. Effect of spironolactone on blood pressure in subjects with resistant hypertension. *Hypertension*. 2007 Apr;49(4):839-45.
113. Pepine CJ, Kowey PR, Kupfer S, et al. Predictors of adverse outcome among patients with hypertension and coronary artery disease. *J Am Coll Cardiol*. 2006;47(3):547-51.

114. Denardo SJ, Gong Y, Cooper-DeHoff RM, Farsang C, Keltai M, Szirmai L, et al. Effects of verapamil SR and atenolol on 24-hour blood pressure and heart rate in hypertension patients with coronary artery disease: an international verapamil SR-trandolapril ambulatory monitoring substudy. *PLoS One*. 2015 Apr 2;10(4):e0122726.
115. Hilleman DE, Ryschon KL, Mohiuddin SM, Wurdeman RL. Fixed-dose combination vs monotherapy in hypertension: a meta-analysis evaluation. *J Hum Hypertens*. 1999;13:477-83.
116. Davidov ME, Glazer N, Wollam G, Zager PG, Cangiano J. Comparison of betaxolol, a new beta 1-adrenergic antagonist, to propranolol in the treatment of mild to moderate hypertension. *Am J Hypertens*. 1988 Jul;1(3 Pt 3):206S-210S.
117. Czuriga I, Rieicansky I, Bodnar J, Fulop T, Kruzsicz V, Kristof E, Edes I, For The NEBIS Investigators ; NEBIS Investigators Group. Comparison of the new cardioselective beta-blocker nebivolol with bisoprolol in hypertension: the Nebivolol, Bisoprolol Multicenter Study (NEBIS). *Cardiovasc Drugs Ther*. 2003 May;17(3):257-63.
118. Stoschitzky K, Stoschitzky G, Brussee H, Bonelli C, Dobnig H. Comparing beta-blocking effects of bisoprolol, carvedilol and nebivolol. *Cardiology*. 2006;106(4):199-206.
119. Lewin AJ, Lueg MC, Targum S, et al. A clinical trial evaluating the 24-hour effects of bisoprolol/hydrochlorothiazide 5 mg/6.25 mg combination in patients with mild to moderate hypertension. *Clin Cardiol*. 1993 Oct;16(10):732-6.
120. Benetos A, Consoli S, Safavian A, Dubanchet A, Safar M. Efficacy, safety, and effects on quality of life of bisoprolol/hydrochlorothiazide versus amlodipine in elderly patients with systolic hypertension. *Am Heart J*. 2000 Oct;140(4):E11.
121. Prisant LM, Weir MR, Papademetriou V, et al. Low-dose combination therapy: an alternative first-line approach to hypertension treatment. *Am Heart J*. 1995 Aug;130(2):359-66.
122. Frishman WH, Bryzinski BS, Coulson LR, et al. A multifactorial trial design to assess combination therapy in hypertension. Treatment with bisoprolol and hydrochlorothiazide. *Arch Intern Med*. 1994 Jul 11;154(13):1461-8.
123. Frishman WH, Burris JF, Mroczek WJ, et al. First-line therapy option with low-dose bisoprolol fumarate and low-dose hydrochlorothiazide in patients with stage I and stage II systemic hypertension. *J Clin Pharmacol*. 1995 Feb;35(2):182-8.
124. Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G, et al. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial.
125. Hamaad A, Lip GYH, Nicholls D, MacFadyen RJ. Comparative dose titration response to the introduction of bisoprolol or carvedilol in stable chronic systolic heart failure. *Cardiovasc Drugs Ther*. 2007(Sep)26;21:437-44.
126. Erdogan O, Ertem B, Altun A. Comparison of antihypertensive efficacy of carvedilol and nebivolol in mild-to-moderate primary hypertension: a randomized trial. *Anadolu Kardiyol Derg*. 2011;11:310-3.
127. Saunders E, Curry C, Hinds J, Kong BW, Medakovic M, Poland M, Roper K. Labetalol compared with propranolol in the treatment of black hypertensive patients. *J Clin Hypertens*. 1987 Sep;3(3):294-302.
128. McAreavey D, Ramsey LE, Latham L, McLaren AD, Lorimer AR, Reid JL, et al. Third drug trial: comparative study of antihypertensive agents added to treatment when blood pressure remains uncontrolled by a beta blocker plus thiazide diuretic. *Br Med J (Clin Res Ed)*. 1984 Jan 14;288(6411):106-11.
129. Wright JT Jr, Bakris G, Greene T, et al. Effects of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA*. 2002 Nov 20;288(19):2421-31.
130. Dafgard T, Forsen B, Lindahl T. Comparative study of hydrochlorothiazide and a fixed combination of metoprolol and hydrochlorothiazide essential hypertension. *Ann Clin Res*. 1981;13 Suppl 30:37-44.
131. Smilde JG. Comparison of the antihypertensive effect of a double dose of metoprolol versus the addition of hydrochlorothiazide to metoprolol. *Eur J Clin Pharmacol*. 1983;25(5):581-3.
132. Liedholm H, Ursing D. Antihypertensive effect and tolerability of two fixed combination of metoprolol and hydrochlorothiazide followed by a long-term tolerance study with one combination. *Ann Clin Res*. 1981;13 Suppl 30:45-53.
133. Materson BJ, Cushman WC, Goldstein G, et al. Treatment of hypertension in the elderly: I. Blood pressure and clinical changes: results of a Department of Veterans Affairs cooperative study. *Hypertension*. 1990 Apr;15(4):348-60.
134. Greathouse M. Nebivolol efficacy and safety in patients with stage I-II hypertension. *Clin Cardiol*. 2010;33(4):E20-E7.



135. Neutel JM, Smith DHG, Gradman AH. Adding nebivolol to ongoing antihypertensive therapy improves blood pressure and response rates in patients with uncontrolled stage I-II hypertension. *J Hum Hyperten.* 2010;24:64-73.
136. Weiss RJ, Saunders E, Greathouse M. Efficacy and tolerability of nebivolol in stage I-II hypertension: a pooled analysis of data from three randomized, placebo-controlled monotherapy trials. *Clin Ther.* 2011;33:1150-61.
137. Rosei EA, Rizzoni D, Comini S, Boari G; Nebivolol-Lisinopril Study Group. Evaluation of the efficacy and tolerability of nebivolol versus lisinopril in the treatment of essential arterial hypertension: a randomized, multicentre, double-blind study. *Blood Press Suppl.* 2003 May;1:30-5.
138. Mazza A, Gil-Extremera B, Maldonado A, Toutouzas T, Pessina AC. Nebivolol vs amlodipine as first-line treatment of essential arterial hypertension in the elderly. *Blood Press.* 2002;11(3):182-8.
139. Van Bortel LM, Bulpitt CJ, Fici F. Quality of life and antihypertensive effect with nebivolol and losartan. *Am J Hypertens.* 2005 Aug;18(8):1060-6.
140. Van Bortel LM, Fici F, Mascagni F. Efficacy and tolerability of nebivolol compared with other antihypertensive drugs: a meta-analysis. *Am J Cardiovasc Drugs.* 2008;8(1):35-44.
141. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Efficacy of nadolol alone and combined with bendroflumethiazide and hydralazine for systemic hypertension. *Am J Cardiol.* 1983 Dec 1;52(10):1230-37.
142. Frick MH, Hartikainen M, Pörsti P. Penbutolol, a new non-selective beta-adrenergic blocking compound in the treatment of hypertension. A comparison with propranolol. *Ann Clin Res.* 1978 Apr;10(2):105-6.
143. Finnerty FA Jr, Gyftopoulos A, Berry C, McKenney A. Step 2 regimens in hypertension: an assessment. *JAMA.* 1979 Feb 9;241(6):579-81.
144. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Propranolol in the treatment of essential hypertension. *JAMA.* 1977 May 23;237(21):2303-10.
145. Stevens JD, Mullane JF. Propranolol-hydrochlorothiazide combination in essential hypertension. *Clin Ther.* 1982;4(6):497-509.
146. de Leeuw PW, Notter T, Zilles P. Comparison of different fixed antihypertensive combination drugs: a double-blind, placebo-controlled parallel group study. *J Hypertens.* 1997 Jan;15(1):87-91.
147. Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet.* 2005 Dec 10;366:2026-33.
148. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig.* 2007;27(11):735-53.
149. Gottlieb SS, McCarter RJ. Comparative effects of three beta blockers (atenolol, metoprolol, and propranolol) on survival after acute myocardial infarction. *Am J Cardiol.* 2001 Apr 1;87(7):823-6.
150. Testa G, Cacciatore F, Della-Morte D, et al. Atenolol use is associated with long-term mortality in community-dwelling older adults with hypertension. *Geriatr Gerontol Int.* 2014 Jan;14(1):153-158.
151. Black HR, Elliott WJ, Grandits G, Grambsch P, Lucente T, White WB, et al; CONVINCe Research Group. Principal results of the Controlled Onset Verapamil Investigation of Cardiovascular End Points (CONVINCE) Trial. *JAMA.* 2003 Apr 23-30;289(16):2073-82.
152. Dahlöf B, Devereux RB, Kjeldsen SE, Julius S, Beevers G, de Faire U, et al; LIFE Study Group. Cardiovascular morbidity and mortality in the Losartan Intervention For End point reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet.* 2002 Mar 23;359(9311):995-1003.
153. Julius S, Alderman MH, Beevers G, et al. Cardiovascular risk reduction in hypertensive black patients with left ventricular hypertrophy: the LIFE study. *J Am Coll Cardiol.* 2004;43:1047-55.
154. Lindholm LH, Ibsen H, Dahlöf B, Devereux RB, Beevers G, de Faire U, et al; LIFE Study Group. Cardiovascular morbidity and mortality in patients with diabetes in the Losartan Intervention For End point reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet.* 2002;359:1004-10.
155. Kjeldsen SE, Dahlöf B, Devereux RB, Julius S, Aurup P, Edelman J, et al, Snapinn S, Wedel H; LIFE (Losartan Intervention for End point Reduction) Study Group. Effects of losartan on cardiovascular morbidity and mortality in patients with isolated systolic hypertension and left ventricular hypertrophy: a Losartan Intervention for End point Reduction (LIFE) substudy. *JAMA.* 2002 Sep 25;288(12):1491-8.
156. Fossum E, Olsen M, Hoiegggen A, Wachtell K, Reims H, Kjeldsen S et al. Long-term effects of a losartan-compared with and atenolol-based treatment regimen on carotid artery plaque development in hypertensive patients with left ventricular hypertrophy: ICARUS, a LIFE substudy. *J Clin Hypertens.* 2006;8:169-73.

157. Kizer J, Dahlof B, Kjeldsen S, Julius S, Beevers G, de Faire U et al. Stroke reduction in hypertensive adults with cardiac hypertrophy randomized to losartan versus atenolol: the losartan intervention for endpoint reduction in hypertension study. *Hypertension*. 2005;45:46-52.
158. Wachtell K, Lehto M, Gerds E, Olsen M, Hornestam B, Dahlof B et al. Angiotensin II receptor blockade reduces new-onset atrial fibrillation and subsequent stroke compared to atenolol: the losartan intervention for endpoint reduction in hypertension (LIFE) study. *J Am Coll Cardiol*. 2005;45:712-9.
159. Wachtell K, Hornestam B, Lehto M, Slotwiner D, Gerds E, Olsen M et al. Cardiovascular morbidity and mortality in hypertensive patients with a history of atrial fibrillation: the losartan intervention for endpoint reduction in hypertension (LIFE) study. *J Am Coll Cardiol*. 2005;45:705-11.
160. Dahlöf B, Lindholm LH, Hansson L, Scherstén B, Ekblom T, Wester PO. Morbidity and mortality in the Swedish Trial in Old Patients with Hypertension (STOP-Hypertension). *Lancet*. 1991 Nov 23;338(8778):1281-5.
161. Hansson L, Lindholm LH, Ekblom T, Dahlöf B, Lanke J, Scherstén B, et al. Randomized trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 (STOP) study. *Lancet*. 1999 Nov 20;354(9192):1751-6.
162. Dahlof B, Sever PS, Poulter NR, et al; for the ASCOT Investigators. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomized controlled trial. *Lancet*. 2005 Sep 10;366(9489):895-906.
163. Pepine CJ, Handberg EM, Cooper-DeHoff RM, et al. A calcium antagonist vs a non-calcium antagonist hypertension treatment strategy for patients with coronary artery disease: the international verapamil-trandolapril study (INVEST): a randomized controlled trial. *JAMA*. 2003 Dec 3;290(21):2805-16.
164. Mancia G, Messerli F, Bakris G et al. Blood Pressure Control and Improved Cardiovascular Outcomes in the International Verapamil SR-Trandolapril Study. *Hypertension*. 2007;50:299-305.
165. Bangalore S, Messerli F, Cohen J, Bacher P, Sleight P, Mancia G, et al. Verapamil-sustained release-based treatment strategy at reducing cardiovascular events in patients with prior myocardial infarction: an International Verapamil SR-Trandolapril (INVEST) substudy. *Am Heart J*. 2008;156:241-7.
166. Iliuta L, Christodorescu R, Filipescu D, Moldovan H, Radulescu B, Vasile R. Prevention of perioperative atrial fibrillation with beta blockers in coronary surgery: betaxolol versus metoprolol. *Interact Cardiovasc Thorac Surg*. 2009;9:89-93.
167. Jonsson G, Abdelnoor M, Müller C, Kjeldsen SE, Os I, Westheim A. A comparison of the two beta-blockers carvedilol and atenolol on left ventricular ejection fraction and clinical endpoints after myocardial infarction. a single-centre, randomized study of 232 patients. *Cardiology*. 2005 Mar 21;103(3):148-55.
168. Pasternak B, Svanström H, Melbye M, Hviid A. Association of treatment with carvedilol vs metoprolol succinate and mortality in patients with heart failure. *JAMA Intern Med*. 2014 Oct;174(10):1597-1604.
169. Seo GW, Kim DK, Kim KH, Seol SH, Jin HY, Yang TH, et al. Impact of Carvedilol versus  $\beta$ 1-selective  $\beta$  blockers (bisoprolol, metoprolol, and nebivolol) in patients with acute myocardial infarction undergoing percutaneous coronary intervention. *Am J Cardiol*. 2015 Nov 15;116(10):1502-8.
170. Olsson G, Wikstrand J, Warnold I, Manger Cats V, McBoyle D, Herlitz J, et al. Metoprolol-induced reduction in postinfarction mortality: pooled results from five double-blind randomized trials. *Eur Heart J*. 1992 Jan;13(1):28-32.
171. Piccini JP, Al-Khatib SM, Wojdyla DM, et al. Comparison of safety of sotalol versus amiodarone in patients with atrial fibrillation and coronary artery disease. *Am J Cardiol*. 2014;114(5):716-722.
172. No authors listed. A multicenter study on timolol in secondary prevention after myocardial infarction (abstract). *Acta Med Scand Suppl*. 1983;674:1-129.
173. Patel K, Fonarow GC, Ekundayo OJ, et al. Beta-blockers in older patients with heart failure and preserved ejection fraction: class, dosage, and outcomes. *Int J Cardiol*. 2014 May 15;173(3):393-401.
174. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, et al. Randomized trial of effects of calcium antagonists compared with diuretics and  $\beta$ -blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet*. 2000 Jul 29;356(9227):359-65.
175. Messerli FH, Grossman E, Goldbourt U. Are beta-blockers efficacious as first-line therapy for hypertension in the elderly? A systematic review. *JAMA*. 1998 Jun 17;279(23):1903-7.
176. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.

177. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*. 2005 Oct 29-Nov 4;366(9496):1545-53.
178. Freemantle N, Cleland J, Young P, Mason J, Harrison J. Beta blockade after myocardial infarction: systematic review and meta regression analysis. *BMJ*. 1999 Jun 26;318(7200):1730-7.
179. Schellenberg R, Lichtenthal A, Wohling H, Graf C, Brixius K. Nebivolol and metoprolol for treating migraine: An advance in  $\beta$ -blocker treatment? *Headache*. 2008 Jan;48(1):118-25.
180. Silberstein SD, Dodick DW, Lindblad AS, Holroyd K, Harrington M, Matthew NT, et al. Randomized, placebo-controlled trial of propranolol added to topiramate in chronic migraine. *Neurology*. 2012;78:976-84.
181. Tfelt-Hansen P, Standnes B, Kangasneimi P, Hakkarainen H, Olesen J. Timolol vs propranolol vs placebo in common migraine prophylaxis a double-blind multicenter trial. *Acta Neurol Scand*. 1984;69:1-8.
182. Linde K, Rossnagel K. Propranolol for migraine prophylaxis. *Cochrane Database Syst Rev*. 2004;(2):CD003225. doi: 10.1002/14651858.CD003225.pub2.
183. Léauté-Labrèze C, Hoeger P, Mazereeuw-Hautier J, et al. A randomized, controlled trial of oral propranolol in infantile hemangioma. *N Engl J Med*. 2015 Feb 19;372(8):735-46.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Dihydropyridines  
AHFS Class 242808  
May 4, 2022**

**I. Overview**

The movement of calcium ions is essential for the function of all types of muscle, including cardiac and vascular smooth muscle. When this flow is reduced, the result is a weakening of muscle contraction and relaxation of muscle tissue.<sup>1-3</sup> Relaxation of coronary vascular smooth muscle increases the flow of oxygenated blood into the myocardium, while relaxation of arteriolar smooth muscle decreases peripheral vascular resistance.<sup>3-5</sup> Both coronary and systemic vasodilation serve to reduce cardiac workload. The calcium-channel blocking agents include dihydropyridines and nondihydropyridines. Although they have different binding sites on the L-type calcium channel, both block the transmembrane influx of calcium ions into cardiac and vascular smooth muscle. The nondihydropyridines also block the T-type calcium channel in the atrioventricular node.<sup>1-5</sup>

The dihydropyridines are approved for the treatment of angina and hypertension. Amlodipine is also indicated to reduce the risk of hospitalization due to angina and to reduce the risk of a coronary revascularization procedure in patients with recently documented coronary artery disease.<sup>1,2,6-17</sup> They are potent vasodilators and have little effect on cardiac muscle contractility or conduction. The dihydropyridines are available in a variety of single entity formulations. Amlodipine is also available in combination with benazepril, olmesartan, valsartan, or valsartan-hydrochlorothiazide. Angiotensin converting enzyme (ACE) inhibitors block the conversion of angiotensin I to angiotensin II, and also inhibit the breakdown of bradykinin, a potent vasodilator. Angiotensin II receptor antagonists block the angiotensin II receptor subtype AT<sub>1</sub>, preventing the negative effects of angiotensin II, regardless of its origin. Hydrochlorothiazide inhibits the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride.<sup>1,2</sup>

The dihydropyridines that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the products with the exception of clevidipine is available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Dihydropyridines Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Amlodipine	suspension, tablet	Katerzia <sup>®</sup> , Norvasc <sup>®*</sup>	amlodipine
Clevidipine	injection <sup>^</sup>	Cleviprex <sup>®</sup>	none
Felodipine	extended-release tablet	N/A	felodipine
Isradipine	capsule	N/A	isradipine
Nicardipine	capsule*, injection*	Cardene-NACL <sup>®^*</sup>	nicardipine
Nifedipine	capsule, extended-release tablet	Adalat CC <sup>®*</sup> , Procardia XL <sup>®*</sup>	nifedipine
Nimodipine	capsule*, solution	Nymalize <sup>®</sup>	nimodipine
Nisoldipine	extended-release tablet*	Sular ER <sup>®*</sup>	nisoldipine
<b>Combination Products</b>			
Amlodipine and benazepril	capsule	Lotrel <sup>®*</sup>	amlodipine and benazepril
Amlodipine and olmesartan	tablet	Azor <sup>®*</sup>	amlodipine and olmesartan
Amlodipine and valsartan	tablet	Exforge <sup>®*</sup>	amlodipine and valsartan
Amlodipine, valsartan, and hydrochlorothiazide	tablet	Exforge HCT <sup>®*</sup>	amlodipine, valsartan, and hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.

<sup>^</sup>Product is primarily administered in an institution.

PDL=Preferred Drug List

N/A=Not available

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the dihydropyridines are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Dihydropyridines**

Clinical Guideline	Recommendations
<p>American College of Cardiology/American Heart Association: <b>2007 Chronic Angina Focused Update of the 2002 Guidelines for the Management of Patients With Chronic Stable Angina (2007)</b><sup>18</sup></p>	<ul style="list-style-type: none"> <li>• Aspirin should be started at 75 to 162 mg/day and continued indefinitely in all patients, unless contraindicated.</li> <li>• Use of warfarin in conjunction with aspirin and/or clopidogrel is associated with an increased risk of bleeding and should be monitored closely.</li> <li>• Patients with hypertension and established coronary artery disease (CAD) should be treated with blood pressure medication(s) as tolerated, including angiotensin converting enzyme (ACE) inhibitors and/or <math>\beta</math>-adrenergic antagonists (<math>\beta</math>-blockers) with the addition of other medications as needed to achieve blood pressure goals of &lt;140/90 or &lt;130/80 mm Hg for patients with chronic kidney disease or diabetes.</li> <li>• Long-acting calcium channel blocking agents or long-acting nitrates may be used if <math>\beta</math>-blockers are contraindicated. Immediate-release and short-acting dihydropyridine calcium channel blockers can increase adverse cardiac events and should not be used.</li> <li>• Long-acting calcium channel blockers or long-acting nitrates may be used with <math>\beta</math>-blockers if initial treatment is not successful.</li> <li>• ACE inhibitors should be used indefinitely in patients with a left ventricular ejection fraction (LVEF) <math>\leq</math>40% and in those with hypertension, diabetes or chronic kidney disease, unless contraindicated.</li> <li>• ACE inhibitors should also be used indefinitely in patients at lower risk (mildly reduced or normal LVEF in whom cardiovascular risk factors remain well controlled and revascularization has been performed), unless contraindicated.</li> <li>• Angiotensin II receptor blockers (ARBs) are recommended in patients with hypertension, those who have an indication for an ACE inhibitor and are intolerant to them, who have heart failure, or who have had a myocardial infarction (MI) and have a LVEF <math>\leq</math>40%.</li> <li>• ARBs may be considered in combination with an ACE inhibitor for heart failure due to left ventricular systolic dysfunction.</li> <li>• Aldosterone blockade is recommended in patients post-MI without significant renal dysfunction or hyperkalemia who are already receiving therapeutic doses of an ACE inhibitor and a <math>\beta</math>-blocker, have a LVEF <math>\leq</math>40% and have either diabetes or heart failure.</li> <li>• It is beneficial to start and continue <math>\beta</math>-blocker therapy indefinitely in all patients who have had a MI, acute coronary syndrome or left ventricular dysfunction with or without heart failure symptoms, unless contraindicated.</li> <li>• Annual influenza vaccination is recommended in patients with cardiovascular disease.</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)</b><sup>19</sup></p>	<p><u>Pharmacological management of stable coronary artery disease (CAD) patients</u></p> <ul style="list-style-type: none"> <li>• The two aims of the pharmacological management of stable CAD patients are to obtain relief of symptoms and to prevent CV events.</li> <li>• Optimal medical treatment indicates at least one drug for angina/ischaemia relief plus drugs for event prevention.</li> <li>• It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>• It is indicated to review the patient's response soon after starting therapy.</li> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin,</li> </ul>

Clinical Guideline	Recommendations
	<p>consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</p> <ul style="list-style-type: none"> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief: <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a <math>\beta</math>-blocker and/or a non-DHP-calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</li> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</li> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischaemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and beta-blockers avoided.</li> </ul> </li> <li>• Event prevention: <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><u>Treatment in patients with microvascular angina</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>• <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>• Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>• ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>• Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul> <p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>• Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy (DAPT).</li> <li>• Aspirin is recommended for elective stenting.</li> <li>• Clopidogrel is recommended for elective stenting.</li> <li>• Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>• GP IIb/IIIa antagonists should be considered for bailout situation only.</li> <li>• Platelet function testing or genetic testing may be considered in specific or high risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>• Prasugrel or ticagrelor may be considered in specific high-risk situations of</li> </ul>

Clinical Guideline	Recommendations
	<p>elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</p> <ul style="list-style-type: none"> <li>• Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>• Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>• Prasugrel or ticagrelor is not recommended in low-risk elective stenting.</li> <li>• After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low</li> <li>• Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months</li> </ul> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all revascularized patients receive a secondary prevention and be scheduled for follow-up visit.</li> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after second generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/ American College of Cardiology Foundation/ American Heart Association/ American Association for Thoracic Surgery/ Preventive Cardiovascular Nurses Association/ Society of Thoracic Surgeons: <b>Management of Stable Ischemic Heart Disease (2012)</b><sup>20</sup></p>	<p><u>Medical therapy to prevent MI and death in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</li> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</li> <li>• Patients should receive an annual influenza vaccine.</li> </ul>



Clinical Guideline	Recommendations
	<p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> <li>• Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes (2014)</b><sup>21</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation &lt;90%, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications <ul style="list-style-type: none"> <li>○ Nitrates <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>○ Beta-adrenergic blockers <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval &gt;0.24 second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</li> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> </li> <li>○ Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval &gt;0.24 seconds, or second or third degree</li> </ul> </li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>atrioventricular block without a cardiac pacemaker.</p> <ul style="list-style-type: none"> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> <ul style="list-style-type: none"> <li>○ Other anti-ischemic interventions <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> </li> </ul> <ul style="list-style-type: none"> <li>• Inhibitors of renin-angiotensin-aldosterone system <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF &lt;0.40 and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF &lt;0.40 who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine &gt;2.5 mg/dL in men or &gt;2.0 mg/dL in women) or hyperkalemia (K &gt;5.0 mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF &lt;0.40, diabetes mellitus, or heart failure.</li> </ul> </li> <li>• Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include: <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatid or tirofiban.</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet agents               <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatide, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> <li>• Anticoagulant therapy               <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>• Timing of CABG in relation to use of antiplatelet agents               <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> </li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>• Medications at discharge               <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</p> <ul style="list-style-type: none"> <li>○ All patients who are post–NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post–NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post–NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> <ul style="list-style-type: none"> <li>● Late hospital and post-hospital oral antiplatelet therapy <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation</b> (2020)<sup>22</sup></p>	<p><u>Pharmacological treatment of ischemia</u></p> <ul style="list-style-type: none"> <li>● Sublingual or intravenous nitrates and early initiation of <math>\beta</math>-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>● Continuation of chronic <math>\beta</math>-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>● Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>● In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>● Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds.               <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul>

Clinical Guideline	Recommendations
	<p><b>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</b></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li>• If at low bleeding risk (HAS-BLED <math>\leq 2</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED <math>\geq 3</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED <math>\geq 3</math> and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p><b>Recommendations for post-interventional and maintenance treatment</b></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>

Clinical Guideline	Recommendations
<p>American College of Cardiology/American Heart Association: <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>23</sup></p>	<p><u>Routine medical therapies: calcium channel blockers</u></p> <ul style="list-style-type: none"> <li>Evidence demonstrates that beneficial effect on infarct size or the rate of reinfarction when calcium channel blocker therapy was initiated during either the acute or convalescent phase of ST-segment elevation myocardial infarction (STEMI). However, calcium channel blockers may be useful to relieve ischemia, lower blood pressure, or control the ventricular response rate to atrial fibrillation in patients who are intolerant to <math>\beta</math>-blockers.</li> <li>Use of immediate-release nifedipine is contraindicated in patients with STEMI due to hypotension and reflex sympathetic activation with tachycardia.</li> </ul> <p><u>Routine medical therapies: <math>\beta</math>-blockers</u></p> <ul style="list-style-type: none"> <li>Oral <math>\beta</math>-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral <math>\beta</math>-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li><math>\beta</math>-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>Patients with initial contraindications to the use of <math>\beta</math>-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>It is reasonable to administer intravenous <math>\beta</math>-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia.</li> </ul> <p><u>Routine medical therapies: Renin-Angiotensin-Aldosterone System Inhibitors</u></p> <ul style="list-style-type: none"> <li>An angiotensin-converting enzyme (ACE) inhibitor should be administered within the first 24 hours to all patients with STEMI with anterior location, HF, or ejection fraction (EF) <math>\leq 40\%</math>, unless contraindicated.</li> <li>An angiotensin receptor blocker (ARB) should be given to patients with STEMI who have indications for but are intolerant of ACE inhibitors.</li> <li>An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq 40\%</math> and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>24</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of ST-elevation myocardial infarction (STEMI)</u></p> <ul style="list-style-type: none"> <li>Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>In patients with an indication for oral anticoagulation, oral anticoagulants are indicated in addition to antiplatelet therapy.</li> <li>In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the estimated risk of recurrent coronary events and bleeding).</li> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age <math>\geq 50</math> years, and at least one of the following risk factors: age <math>\geq 65</math> years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR <math>&lt; 60</math> mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</li> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with <math>\beta</math>-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with <math>\beta</math>-blockers is indicated in patients with heart failure or left ventricular dysfunction, LVEF <math>\leq 40\%</math> unless contraindicated.</li> <li>• Intravenous <math>\beta</math>-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> <li>• Intravenous <math>\beta</math>-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of <math>&lt; 1.8</math> mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C <math>&gt; 1.8</math> mmol/L (<math>&gt; 70</math> mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction <math>\leq 40\%</math> and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
American College of Cardiology/ American Heart Association:	<p><u>Top 10 messages for the primary prevention of cardiovascular disease</u></p> <ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> </ul>

Clinical Guideline	Recommendations
<p><b>Guideline on the Primary Prevention of Cardiovascular Disease (2019)</b><sup>25</sup></p>	<ul style="list-style-type: none"> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</li> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</li> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</li> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be <math>&lt; 130/80</math> mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul> <p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a</li> </ul>



Clinical Guideline	Recommendations
	<p>decision is made for statin therapy, a moderate-intensity statin should be recommended.</p> <ul style="list-style-type: none"> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) patients, LDL-C levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</li> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> </li> <li>• In patients at borderline risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul> <p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>• In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include: <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>• In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>• In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>• In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>• In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than 130/80 mm Hg.</li> <li>• In adults with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering</li> </ul>

Clinical Guideline	Recommendations
	<p>medication are recommended.</p> <ul style="list-style-type: none"> <li>In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul> <p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> <li>To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)<sup>26</sup></p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have</li> </ul>

Clinical Guideline	Recommendations
	<p>reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</p> <ul style="list-style-type: none"> <li>• In patients with LVEF <math>\leq 40\%</math>, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF <math>\leq 50\%</math>, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF <math>\leq 50\%</math>, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal</li> </ul>

Clinical Guideline	Recommendations
	<p>therapy. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq 35\%</math>) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq 70</math> beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and</li> </ul>

Clinical Guideline	Recommendations
	<p>cardiovascular mortality. (LoE: B)</p> <ul style="list-style-type: none"> <li>In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><b>Treatment of Stage D (advanced/refractory) HF</b></p> <ul style="list-style-type: none"> <li>For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>27</sup></p>	<p><b>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</b></p> <ul style="list-style-type: none"> <li>An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HF<sub>r</sub>EF to reduce the risk of HF hospitalization and death.</li> <li>A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HF<sub>r</sub>EF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>Dapagliflozin or empagliflozin are recommended for patients with HF<sub>r</sub>EF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HF<sub>r</sub>EF regardless of diabetes status.</li> <li>Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HF<sub>r</sub>EF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq</math>35%, in sinus rhythm and a resting heart rate <math>\geq</math>70 bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq</math>35%, in sinus rhythm and a resting heart rate <math>\geq</math>70 bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>An ARB may be considered to reduce the risk of HF hospitalization and death in</li> </ul>

Clinical Guideline	Recommendations
	<p>patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</p> <ul style="list-style-type: none"> <li>• Vericiguat may be considered in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul>

Clinical Guideline	Recommendations
	<p><b>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</b></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>28</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: Global Hypertension Practice Guidelines (2020)<sup>29</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>(HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</p> <ul style="list-style-type: none"> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediated drug treatment in all patients.</li> <li>● Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>30</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>● Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt;</math>60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul>

Clinical Guideline	Recommendations
	<p><b>Guidelines for patients with hypertension who have had a recent myocardial infarction</b></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><b>Treatment of hypertension in association with heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction &lt;40%), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq</math>30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><b>Treatment of hypertension in association with stroke</b></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><b>Treatment of hypertension in association with LVH</b></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or</li> </ul>

Clinical Guideline	Recommendations
	<p><b>minoxidil should not be used.</b></p> <p><b>Treatment of hypertension in association with nondiabetic chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><b>Treatment of hypertension in association with renovascular disease</b></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><b>Treatment of hypertension in association with diabetes mellitus</b></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><b>Resistant hypertension</b></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin,</li> </ul>

Clinical Guideline	Recommendations
	<p>amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</p> <ul style="list-style-type: none"> <li>Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>31</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>Some agents should be considered as the preferential choice in specific conditions</li> </ul>

Clinical Guideline	Recommendations
	<p>because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</p> <ul style="list-style-type: none"> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should</li> </ul>

Clinical Guideline	Recommendations
	<p>be avoided.</p> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered</li> </ul>

Clinical Guideline	Recommendations
	<p>according to its tolerability and impact on renal function and electrolytes.</p> <ul style="list-style-type: none"> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq</math>220 mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq</math>140/90 mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq</math>130 mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>



Clinical Guideline	Recommendations
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>32</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person’s medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>33</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p><b>Kidney Disease</b> <b>Improving Clinical</b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP)</li> </ul>

Clinical Guideline	Recommendations
<p>Outcomes Group: <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>34</sup></p>	<p>measurement in preference to routine office BP measurements for the management of high BP in adults.</p> <ul style="list-style-type: none"> <li>The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>35</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq 140/90</math>) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq 180/110</math> and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of <math>&lt;130/80</math> mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;15\%</math>), treat to blood pressure target of <math>&lt;140/90</math>.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;120/80</math>, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with type 2 diabetes and CKD, consider use of sodium–glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium–glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin–to–creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin–to–creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin–to–creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American College of Cardiology/American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>36</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>• Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>• Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>• Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>• For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different</li> </ul>

Clinical Guideline	Recommendations
	<p>classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</p> <ul style="list-style-type: none"> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>Adults with SIHD and hypertension (BP <math>\geq</math>130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq</math>300 mg/d, or <math>\geq</math>300 mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq</math>220/120 mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq</math>140/90 mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension</li> </ul>

Clinical Guideline	Recommendations
	<p>with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> <li>• For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP <math>\geq 180</math> mmHg or DBP <math>\geq 110</math> mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> </ul>



<b>Clinical Guideline</b>	<b>Recommendations</b>
	<ul style="list-style-type: none"><li data-bbox="488 207 1377 264">• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li><li data-bbox="488 268 1406 325">• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li></ul>

\*Agent not available in the United States.

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the dihydropyridines are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Dihydropyridines<sup>1,6-17</sup>**

Indication(s)	Single Entity Agents							Combination Products			
	Amlodipine	Felodipine	Isradipine	Nicardipine	Nifedipine	Nimodipine	Nisoldipine	Amlodipine and Benazepril	Amlodipine and Olmesartan	Amlodipine and Valsartan	Amlodipine and Valsartan and HCTZ
<b>Angina Pectoris</b>											
Treatment of chronic stable angina	✓ *			✓ (IR)†							
Treatment of chronic stable angina without evidence of vasospasm in patients who remain symptomatic despite adequate doses of β-blockers and/or organic nitrates or who cannot tolerate those agents					✓ (capsule)						
Treatment of vasospastic angina	✓ ‡				✓ (capsule, ER tablet)§						
<b>Coronary Artery Disease (CAD)</b>											
Reduce the risk of hospitalization due to angina and to reduce the risk of a coronary revascularization procedure in patients with recently documented CAD by angiography and without heart failure or an ejection fraction <40%	✓										
<b>Hypertension</b>											
Treatment of hypertension	✓	✓	✓ ¶	✓   #	✓ (ER)		✓	✓ **	✓	✓ ††	✓ †††
<b>Miscellaneous</b>											
Improvement of neurological outcome by reducing the incidence and severity of ischemic deficits in patients with subarachnoid hemorrhage from ruptured intracranial berry aneurysms regardless of their post-ictus neurological condition (i.e., Hunt and Hess Grades I-V)						✓					

\*Alone or in combination with other antianginal agents.

†Alone or in combination with β-blockers.

‡Confirmed or suspected vasospastic angina. Alone or may be used in combination with other antianginal agents.

§Vasospastic angina confirmed by any of the following criteria: 1) classical pattern of angina at rest accompanied by ST segment elevation, 2) angina or coronary artery spasm provoked by ergonovine, or 3) angiographically demonstrated coronary artery spasm.

|| Alone or in combination with other antihypertensive agents.

¶ Alone or in combination with thiazide-type diuretics.

#Cardene IV<sup>®</sup> is indicated for the short term treatment of hypertension when oral therapy is not feasible or not desirable. For prolonged control of blood pressure, transfer patients to oral medication as soon as their clinical condition permits.

\*\*Not adequately controlled on monotherapy with either agent

††Not adequately controlled on monotherapy or as initial therapy in patients likely to need multiple drugs to achieve their blood pressure goals

‡‡This fixed combination drug is not indicated for the initial therapy of hypertension.

ER=extended-release, IR=immediate-release

#### IV. Pharmacokinetics

The pharmacokinetic parameters of the dihydropyridines are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Dihydropyridines<sup>2</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Amlodipine	64 to 90	93	Liver, extensive	Renal (70)	30 to 60
Felodipine	13 to 20	>99	Liver, extensive	Renal (70) Feces (10)	26.7 to 33.2
Isradipine	14 to 24 IR: 90 to 95 CR: 15 to 24	95	Liver, complete	Renal (60 to 65) Feces (25 to 30)	8
Nicardipine	35	>95	Liver, nearly 100%	Oral: Renal (60) Feces (35) IV with oral dose: Renal (49) Feces (43 within 96 hours)	IV: 14.4 Oral: 8.6
Nifedipine	IR: rapid and complete ER: complete	92 to 98	Liver, extensive	Renal (80) Feces (20)	2
Nimodipine	13	>95	Liver, extensive	Renal (<1) Feces (% not reported)	1 to 2
Nisoldipine	5	>99	Liver, extensive	Renal (60 to 80) Feces (6 to 12)	13.7 ± 4.3
<b>Combination Products</b>					
Amlodipine and benazepril	64 to 90/≥37	93/93	Liver, extensive (% not reported)/ Liver, extensive (% not reported)	Renal (60% inactive, 10% unchanged)/ Renal (20)	48/ 10 to 11
Amlodipine and olmesartan	64 to 90/26	93/99	Liver, extensive (90)/ Intestinal wall (100)	Renal (10)/ Renal (35 to 50) Feces (50 to 65)	45/7
Amlodipine and valsartan	64 to 90/25	93/95	Liver, extensive (90)/ Not reported	Renal (60)/ Renal (13) Feces (83)	30 to 50/ 6
Amlodipine and valsartan and HCTZ	64 to 90/ 25/ 60 to 80	93/ 95/ 40	Liver, extensive (90)/ Liver, minimal (20)/ Not metabolized	Renal (60)/ Renal (7 to 13) Feces (83)/ Renal (>61)	45/ 6 to 9/ 5.8 to 18.9

CR=controlled-release, ER=extended-release, HCTZ=hydrochlorothiazide, IR=immediate-release

#### V. Drug Interactions

Major drug interactions with the dihydropyridines are listed in Table 5.

**Table 5. Major Drug Interactions with the Dihydropyridines<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
-----------------	-------------	-----------

Generic Name(s)	Interaction	Mechanism
ARBs (olmesartan, valsartan)	Potassium-sparing diuretics	The risk of hyperkalemia may be increased when potassium-sparing diuretics are co-administered with ACE inhibitors or ARBs.
Amlodipine	Simvastatin	Simvastatin plasma concentrations may be elevated, increasing the risk of toxicity (e.g., myositis, rhabdomyolysis).
ACE inhibitors (benazepril)	Aldosterone blockers	Serious hyperkalemia, possibly with cardiac arrhythmias or arrest, may occur with the combination of aldosterone blockers and benazepril.
ACE inhibitors (benazepril)	Immunosuppressants (Azathioprine and Mercaptopurine)	Concurrent use of immunosuppressants and benazepril may result in myelosuppression.
ACE inhibitors (benazepril)	Sacubitril	Concurrent use of sacubitril and ACE inhibitors may result in increased risk of angioedema.
ACE inhibitors (benazepril)	Potassium-sparing diuretics	The risk of hyperkalemia may be increased when potassium-sparing diuretics are co-administered with ACE inhibitors or ARBs.
Dihydropyridines (amlodipine, felodipine, isradipine, nifedipine, nimodipine, nisoldipine)	Macrolide antibiotics	Inhibition of nifedipine metabolism (CYP3A4) by macrolide and related antibiotics may lead to elevated dihydropyridine plasma concentrations, increasing the pharmacologic effects and risk of adverse reactions (e.g., severe hypotension).
Dihydropyridines (amlodipine)	Rifampin	Concurrent use of amlodipine and rifampin may result in reduced amlodipine efficacy.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics may induce hypokalemia which may increase the risk of torsades de pointes.
Thiazide diuretics (HCTZ)	Lithium	Thiazide diuretics may promote enhanced proximal tubular reabsorption of lithium leading to elevated serum concentrations. Thiazide diuretics may increase the therapeutic and toxic effects of lithium.
Dihydropyridines (felodipine, isradipine, nifedipine, nimodipine, nisoldipine)	Azole antifungals	Dihydropyridine serum levels may increase resulting from a decrease metabolism due to CYP3A4 inhibition by azole antifungal agents.
Dihydropyridines (amlodipine, nimodipine)	HCV protease inhibitors	Dihydropyridine serum levels may increase resulting from a decrease metabolism due to CYP3A4 inhibition by protease inhibitors.
Dihydropyridines (amlodipine, nifedipine, nimodipine)	HIV protease inhibitors	Dihydropyridine serum levels may increase resulting from a decrease metabolism due to CYP3A4 inhibition by protease inhibitors.
Dihydropyridines (nimodipine)	Barbiturates	Metabolism of dihydropyridines may be increased due to induction of mixed function oxidases by barbiturates, causing an increase in first-pass metabolism and decreased bioavailability, reducing the effects of dihydropyridines.
ARBs (olmesartan, valsartan)	Aliskiren	Concurrent use of aliskiren may result in an increased risk of hyperkalemia, renal impairment, and hypotension.
ARBs (olmesartan, valsartan)	ACE Inhibitors	Coadministration of ARBs and ACE inhibitors may be associated with an increased risk of renal dysfunction and/or hyperkalemia.
ARBs (olmesartan, valsartan)	Lithium	Elevations in plasma lithium levels may occur.
ARBs (olmesartan,	Potassium	Hyperkalemia, possibly with cardiac arrhythmias or cardiac

Generic Name(s)	Interaction	Mechanism
valsartan)	preparations	arrest, may occur with the combination of olmesartan and potassium preparations.
Dihydropyridines (amlodipine, felodipine, isradipine, nicardipine, nifedipine, nisoldipine)	Clopidogrel	Concurrent use may result in decreased antiplatelet effect and increased risk of thrombotic events.
Dihydropyridines (isradipine)	Mefloquine	Concurrent use of isradipine and mefloquine may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Dihydropyridines (nimodipine)	Hydantoins	Dihydropyridine serum levels may decrease due to increased first-pass metabolism caused by hydantoins.
ACE inhibitors (benazepril)	Aliskiren	The risk of hyperkalemia, renal impairment, and hypotension may be increased when aliskiren is coadministered with benazepril.
ACE inhibitors (benazepril)	mTOR inhibitors (everolimus and sirolimus)	The risk of angioedema may be increased with concurrent administration of mTOR inhibitors and benazepril.
ACE inhibitors (benazepril)	Alteplase	Concurrent use of alteplase and ACE inhibitors may result in an increased risk of orolingual angioedema.
ACE inhibitors (benazepril)	HIV protease inhibitors	Pharmacologic effects of benazepril may be increased by HIV protease inhibitors.
ACE inhibitors (benazepril)	Imidazoles	Imidazoles may increase the plasma concentrations and pharmacologic effects of benazepril.
ACE inhibitors (benazepril)	NSAIDs	The antihypertensive effects of benazepril may be decreased by NSAIDs. Nephrotoxicity associated with benazepril or NSAIDs may be increased by this drug combination.
ACE inhibitors (benazepril)	Lithium	Pharmacologic effects of lithium may be increased by benazepril. Elevated lithium serum concentrations with toxicity may occur.
ACE inhibitors (benazepril)	Potassium preparations	Hyperkalemia, possibly with cardiac arrhythmias or cardiac arrest, may occur with the combination of benazepril and potassium preparations.
ACE inhibitors (benazepril)	Trimethoprim	Hyperkalemia, possibly with cardiac arrhythmias or cardiac arrest, may occur with the combination of trimethoprim and benazepril.
ACE inhibitors (benazepril, felodipine, nimodipine)	Vasopressin receptor antagonists	Plasma concentrations of benazepril may be increased by co-administration of vasopressin receptor antagonists.
Dihydropyridines (isradipine)	Inhaled anesthetics	Concurrent use may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Dihydropyridines (isradipine)	Phenothiazines	Concurrent use of isradipine and phenothiazines may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Dihydropyridines (isradipine)	Tricyclic antidepressants	Concurrent use of isradipine and tricyclic antidepressants may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Dihydropyridines (amlodipine, felodipine, nicardipine)	Cyclosporine	Cyclosporine serum levels may increase due to inhibited metabolism by dihydropyridines.
Dihydropyridines	Carbamazepine	Carbamazepine may decrease plasma concentrations and

Generic Name(s)	Interaction	Mechanism
(felodipine, nifedipine, nimodipine)		effects of nifedipine.
Dihydropyridines (nicardipine, nifedipine)	Fentanyl	Concurrent use of fentanyl and nicardipine may result in severe hypotension.
Dihydropyridines (amlodipine, nicardipine, nifedipine)	Tacrolimus	Tacrolimus serum levels may be elevated due to inhibition of metabolism by dihydropyridines.
Thiazide diuretics (HCTZ)	Diazoxide	The combination of diazoxide with a thiazide diuretic may lead to hyperglycemia through an unknown mechanism; therefore, the combination should be avoided.
Thiazide diuretics (HCTZ)	Digitalis glycosides	Thiazide diuretics may induce electrolyte disturbances which may predispose patients to digitalis-induced arrhythmias.
Dihydropyridines (clevidipine, isradipine, nicardipine, nimodipine, nisoldipine)	Digitalis glycosides	Dihydropyridines may induce electrolyte disturbances which may predispose patients to digitalis-induced arrhythmias.
Calcium channel blockers (amlodipine)	Digoxin	Concurrent use of digoxin and calcium channel blockers may result in increased risk of complete heart block.
Nimodipine	Rifapentine	Rifapentine can reduce plasma concentrations of nimodipine and reduce efficacy levels of nimodipine
Nimodipine	Dexamethasone	Dexamethasone can reduce plasma concentrations of nimodipine and reduce efficacy levels of nimodipine
Nimodipine	rifampin	Rifampin can reduce plasma concentrations of nimodipine and reduce efficacy levels of nimodipine
Nimodipine	Phenobarbital	Phenobarbital can reduce plasma concentrations of nimodipine and reduce efficacy levels of nimodipine
Nimodipine	Phenytoin	Phenytoin can reduce plasma concentrations of nimodipine and reduce efficacy levels of nimodipine
Nimodipine	Nefazodone	Nefazodone can increase serum nimodipine levels
Nimodipine	Delavirdine	Delavirdine can increase serum nimodipine levels.
Nicardipine	Vecuronium	The combination of vecuronium and nicardipine can result in enhanced neuromuscular blockade
Isradipine	Arsenic trioxide	The combination of arsenic trioxide and isradipine can result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest)
Isradipine	Zolmitriptan	The combination of zolmitriptan and isradipine can result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest)
Isradipine	Fluoxetine	The combination of fluoxetine and isradipine can result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest)
Isradipine	Octreotide	The combination of octreotide and isradipine can result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest)
Isradipine	Pentamidine	The combination of isradipine and pentamidine can result in increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest)
Isradipine	Dolasetron, ondansetron	The combination of isradipine and dolasetron can increase the risk of cardiotoxicity (QT prolongation, torsades de pointes,

Generic Name(s)	Interaction	Mechanism
		cardiac arrest)
Isradipine	Gemifloxacin	The combination of isradipine and Gemifloxacin can increase the risk of cardiotoxicity (QT prolongation, torsades de pointes cardiac arrest)
Dihydropyridines (felodipine, isradipine, nicardipine, nimodipine)	St. John's Wort	St. John's wort may reduce serum levels of dihydropyridines creating subtherapeutic levels of dihydropyridines for antihypertensive activity.
Calcium Channel Blockers (amlodipine, felodipine, isradipine, nicardipine)	Droperidol	Combination of droperidol and calcium channel blockers can increase risk of cardiotoxicity including QT prolongation, torsades de pointes, or cardiac arrest.
Felodipine	Nilotinib	The combination of nilotinib and felodipine can increase exposure of either felodipine or nilotinib.
Dihydropyridines (clevidipine, isradipine, nicardipine, nifedipine, nimodipine, nisoldipine)	Lacosamide	The combination of dihydropyridines and lacosamide can result in prolonged PR interval, AV block, brady cardia, and ventricular tachyarrhythmia
Calcium Channel Blockers (clevidipine, felodipine, isradipine, nicardipine, nifedipine, nimodipine, nisoldipine)	Dantrolene	The combination of dantrolene and calcium channel blockers may result in hyperkalemia with cardiovascular collapse.

ACE inhibitors=angiotensin converting enzyme inhibitors, ARB=angiotensin II receptor blocker, CYP=cytochrome P450 isoenzyme, HCTZ=hydrochlorothiazide, HIV=human immunodeficiency virus, NSAID=nonsteroidal anti-inflammatory drug,

## VI. Adverse Drug Events

The most common adverse drug events reported with the dihydropyridines are listed in Tables 6 and 7. The boxed warnings for the dihydropyridines are listed in Tables 8 through 12.

**Table 6. Adverse Drug Events (%) Reported with the Dihydropyridines (Amlodipine-containing Products)<sup>1,6-17</sup>**

Adverse Events	Single Entity Agents	Combination Products			
	Amlodipine	Amlodipine and Benazepril	Amlodipine and Olmesartan	Amlodipine and Valsartan	Amlodipine and Valsartan and HCTZ
<b>Cardiovascular</b>					
Arrhythmia	1	-	-	✓	-
Atrial fibrillation	1	-	-	1	-
Bradycardia	1	-	-	1	-
Cardiac murmur	-	-	-	✓	-
Chest pain	1	-	-	1	-
Edema	2 to 11	2	2 to 15	✓	7
Hypotension	-	-	✓	✓	-
Orthostatic hypotension	-	-	✓	<1	-
Palpitations	1 to 5	✓	1 to 5	✓	0.5
Peripheral ischemia	1	-	-	1	-
Peripheral edema	18 to 26	✓	✓	5	-
Pitting edema	-	-	-	✓	-
Postural dizziness	1	-	-	-	-



Adverse Events	Single Entity Agents	Combination Products			
	Amlodipine	Amlodipine and Benazepril	Amlodipine and Olmesartan	Amlodipine and Valsartan	Amlodipine and Valsartan and HCTZ
Postural hypotension	-	-	-	1	-
Pulse irregularity	-	-	-	✓	-
Syncope	1	-	-	-	-
Tachycardia	1	-	-	✓	✓
Vasculitis	1	-	-	1	-
Ventricular tachycardia	1	-	-	1	-
<b>Central Nervous System</b>					
Abnormal dreams	1	-	-	1	-
Agitation	1	-	-	✓	-
Amnesia	1	-	-	✓	-
Anxiety	-	✓	-	3	-
Apathy	1	-	-	✓	-
Asthenia	1	✓	✓	✓	-
Ataxia	-	-	-	✓	-
Carpal tunnel syndrome	-	-	-	✓	-
Cervicobrachial syndrome	-	-	-	✓	-
Depersonalization	1	-	-	1	-
Depression	1 to 2	-	-	✓	-
Dizziness	1 to 3	1	1 to 3	2	8
Headache	7	2	-	11	5
Hypoesthesia	1	-	-	✓	-
Insomnia	1	✓	-	✓	-
Migraine	-	-	-	✓	-
Nervousness	1	✓	-	1	-
Paresthesia	1	-	-	✓	-
Peripheral neuropathy	1	-	-	1	-
Postural dizziness	-	✓	-	1	<1
Pyrexia	-	-	-	✓	-
Sciatica	-	-	-	✓	-
Sinus headache	-	-	-	✓	-
Somnolence	<2	✓	<2	3	✓
Syncope	-	✓	-	1	✓
Tremor	1	✓	-	1	✓
Vertigo	1	-	-	✓	-
<b>Dermatologic</b>					
Alopecia	-	-	✓	✓	-
Cold and clammy skin	1	-	-	✓	-
Dermatitis	-	✓	-	✓	-
Eczema	-	-	-	✓	-
Erythema	-	-	-	✓	-
Erythema multiforme	1	-	-	1	-
Exanthema	-	-	-	✓	-
Flushing	1 to 3	✓	1 to 3	✓	-
Hyperhidrosis	1	-	-	✓	-
Pruritus	1	-	✓	✓	✓
Rash	1 to 2	✓	✓	✓	✓
Rash, erythematous	1 to 2	-	-	1	-
Rash, maculopapular	✓	-	-	1	-
Skin discoloration	1	-	-	✓	-
Skin dryness	1	-	-	✓	-
Urticaria	1	-	✓	✓	-
<b>Endocrine and Metabolic</b>					
Decreased libido	-	✓	-	-	-
Gout	-	-	-	✓	-
Gynecomastia	✓	-	✓	-	-
Diabetes mellitus	-	-	-	✓	-
Thirst	1	-	-	1	-
<b>Gastrointestinal</b>					
Abdominal discomfort	-	-	-	✓	-
Abdominal distension	-	-	-	✓	-
Abdominal pain	2	✓	-	3	-
Anorexia	1	-	-	1	-
Colitis	-	-	-	✓	-

Adverse Events	Single Entity Agents	Combination Products			
	Amlodipine	Amlodipine and Benazepril	Amlodipine and Olmesartan	Amlodipine and Valsartan	Amlodipine and Valsartan and HCTZ
Constipation	1	✓	-	✓	✓
Diarrhea	✓	✓	✓	3	✓
Dry mouth	✓	✓	-	✓	✓
Dyspepsia	1	-	-	✓	2
Dysphagia	1 to 2	-	-	1	-
Esophagitis	-	✓	-	-	-
Flatulence	1	-	-	✓	-
Gastritis	-	-	-	✓	✓
Gastroenteritis	-	-	-	✓	-
Hemorrhoids	-	-	-	✓	✓
Hepatitis	-	-	-	✓	-
Increased appetite	-	-	-	✓	-
Jaundice	✓	-	✓	✓	✓
Loose stools	1	-	-	✓	-
Nausea	3	✓	-	3	2
Pancreatitis	1	-	-	1	-
Sprue-like enteropathy	-	-	✓	-	-
Vomiting	1	-	✓	✓	-
<b>Genitourinary</b>					
Cystitis	-	-	-	✓	-
Dysuria	-	-	-	✓	-
Erectile dysfunction	-	-	-	✓	✓
Hematuria	-	-	-	✓	-
Impotence	-	✓	-	✓	-
Micturition disorder	1	-	-	1	-
Nephrolithiasis	-	-	-	✓	-
Nocturia	1	-	✓	1	-
Pollakiuria	-	-	-	✓	-
Polyuria	-	✓	-	✓	-
Sexual dysfunction	1 to 2	-	-	1	-
Urinary frequency	1	-	✓	✓	-
Urinary tract infection	-	-	-	✓	-
<b>Hematological</b>					
Leukopenia	1	-	-	1	-
Neutropenia	-	✓	-	✓	✓
Purpura	1	-	-	1	-
Thrombocytopenia	1	-	-	✓	-
Blood urea nitrogen increased	-	-	-	✓	✓
Creatinine increases	-	-	✓	✓	✓
Hepatic enzyme elevations	✓	-	✓	✓	✓
Hypercholesterolemia	-	-	-	✓	-
Hyperglycemia	1	-	-	1	-
Hyperkalemia	-	✓	✓	3 to 10	-
Hypokalemia	-	✓	-	-	-
<b>Musculoskeletal</b>					
Arthralgia	1	-	-	✓	✓
Arthrosis	1	-	-	1	-
Back pain	1 to 2	✓	-	✓	2
Hypertonia	-	-	-	✓	-
Joint sprain	-	-	-	✓	-
Joint swelling	-	-	-	✓	✓
Limb injury	-	-	-	✓	-
Malaise	1	-	-	1	-
Muscle cramps	1	✓	-	1	-
Muscle spasms	-	-	-	✓	2
Muscle weakness	-	-	-	✓	✓
Musculoskeletal chest pain	-	-	-	✓	-
Myalgia	1 to 2	✓	-	✓	-
Osteoarthritis	-	-	-	✓	✓
Pain	1	-	-	✓	-
Rhabdomyolysis	✓	-	✓	✓	-
Twitching	1	-	-	✓	-
<b>Respiratory</b>					

Adverse Events	Single Entity Agents	Combination Products			
	Amlodipine	Amlodipine and Benazepril	Amlodipine and Olmesartan	Amlodipine and Valsartan	Amlodipine and Valsartan and HCTZ
Bronchitis	-	-	-	✓	-
Cough	-	3	-	2	✓
Dysphonia	-	-	-	✓	-
Dyspnea	1	-	-	1	-
Epistaxis	1 to 2	-	-	✓	-
Influenza	-	-	-	2	-
Nasal congestion	-	-	-	✓	✓
Nasopharyngitis	-	-	-	4	2
Pharyngitis	-	✓	-	✓	-
Pharyngolaryngeal pain	-	-	-	✓	✓
Pharyngotonsillitis	-	-	-	✓	-
Pneumonia	-	-	-	✓	-
Rhinitis	-	-	-	✓	-
Seasonal allergies	-	-	-	✓	-
Sinus congestion	-	-	-	✓	-
Sinusitis	-	-	-	✓	-
Upper respiratory tract infection	-	-	-	3	-
<b>Special Senses</b>					
Abnormal visual accommodation	1	-	-	✓	-
Conjunctivitis	1	-	-	1	-
Diplopia	1	-	-	1	-
Eye pain	1	-	-	1	-
Ear pain	-	-	-	✓	-
Parosmia	1	-	-	✓	-
Taste perversion	-	-	-	✓	-
Tinnitus	1	-	-	✓	-
Visual disturbance	-	-	-	✓	-
Xerophthalmia	1	-	-	✓	-
<b>Other</b>					
Acute renal failure	-	-	✓	✓	-
Allergic reaction	1	-	✓	1	-
Angioedema	1	-	✓	✓	-
Contusion	-	-	-	✓	-
Epicondylitis	-	-	-	✓	-
Fatigue	4.5	✓	-	✓	2
Gingival hyperplasia	1	-	-	1	-
Hot flush	1	✓	-	✓	-
Hypersensitivity	-	-	-	✓	-
Lymphadenopathy	-	-	-	✓	-
Renal insufficiency	-	-	-	-	-
Rigors	1	-	-	1	-
Tooth abscess	-	-	-	✓	-
Toothache	-	-	-	✓	-
Tonsillitis	-	-	-	✓	-
Viral infection	-	-	-	✓	-
Weight gain	1	-	-	1	-
Weight loss	1	-	-	1	-

✓ Percent not specified  
- Event not reported

**Table 7. Adverse Drug Events (%) Reported with the Dihydropyridines (Drugs B - Z)<sup>1,6-17</sup>**

Adverse Events	Felodipine	Isradipine	Nicardipine	Nifedipine	Nimodipine	Nisoldipine
<b>Cardiovascular</b>						
Angina (increased)	-	-	6	-	-	2
Arrhythmia	1 to 2	-	-	-	-	-
Atrial fibrillation	-	≤1	<1	-	-	-
Bradycardia	-	-	-	-	≤1	-
Cardiac failure	-	≤1	-	-	-	-
Cerebrovascular accident	-	-	-	-	-	1
Chest pain	1 to 2	-	-	-	-	-

Adverse Events	Felodipine	Isradipine	Nicardipine	Nifedipine	Nimodipine	Nisoldipine
Edema	-	4 to 36	≤1	-	≤1	-
Electrocardiogram abnormalities	-	-	≤1	-	≤1	-
Epistaxis	-	≤1	-	-	-	-
Erythromelalgia	-	-	-	1	-	-
Hypotension	1 to 2	≤1	-5	5	1 to 50	-
Myocardial infarction	1 to 2	≤1	≤1	✓ ≤4	-	✓
Orthostatic hypotension	-	-	-	-	-	1
Palpitations	<3	1 to 5	3 to 4	<7	-	3
Pedal edema	-	-	6 to 8	-	-	-
Peripheral edema	2 to 17	4 to 40	6	7 to 29	-	22
Pericarditis	-	-	1	-	-	-
Peripheral ischemia	-	-	✓	-	-	-
Postural hypotension	-	-	≤1	-	-	-
Pulse irregularity	1 to 2	-	-	-	-	-
Rebound vasospasm	-	-	-	-	1	-
Tachycardia	1 to 2	1 to 3	1 to 4	-	≤1	-
Vasodilatation/vasodilation	-	-	1 to 5	-	-	4
Ventricular fibrillation	-	≤1	-	1	-	-
Ventricular tachycardia	-	-	≤1	-	-	-
<b>Central Nervous System</b>						
Anxiety	1 to 2	-	✓	-	-	-
Asthenia	2 to 4	1 to 6	-	<3	-	-
Ataxia	-	-	-	1	-	-
Balance difficulties	-	-	-	<2	-	-
Chills	-	-	-	<2	-	1
Confusion	-	-	✓	-	-	-
Depression	1 to 2	≤1	✓	1	≤1	-
Dizziness	3 to 4	3 to 8	1 to 7	4 to 27	-	5
Drowsiness	-	≤1	-	-	-	-
Fatigue	-	3 to 9	-	6	-	-
Headache	11 to 15	10 to 22	6 to 15	10 to 23	≤4	22
Insomnia	1 to 2	≤1	≤1	<3	-	-
Irritability	1 to 2	-	-	-	-	-
Migraine	-	-	-	1	-	1
Nervousness	1 to 2	≤1	≤1	<7	-	-
Numbness	-	≤1	-	-	-	-
Paresthesia	1 to 2	≤1	≤1	<3	-	-
Sleep disturbance	-	-	-	<2	-	-
Somnolence	1 to 2	-	≤1	<3	-	-
Stroke	-	≤1	-	-	-	-
Syncope	1 to 2	≤1	≤1	-	-	-
Transient ischemic attack	-	≤1	-	-	-	-
Tremor	-	-	≤1	<8	-	-
Vertigo	-	-	✓	1	-	-
<b>Dermatologic</b>						
Acne	-	-	-	-	≤1	-
Alopecia	-	-	-	<1	-	-
Dermatitis	-	-	-	1 to 2	-	-
Erythema	1 to 2	-	-	-	-	-
Flushing	4 to 7	1 to 5	6 to 10	3 to 25	-	✓
Hematoma	-	-	-	-	1	-
Hyperhidrosis	-	≤1	11	<2	-	-
Pruritus	-	≤1	-	<2	1	-
Rash	<2	≤3	≤1	<3	1 to 2	2
Urticaria	1 to 2	≤1	-	<2	-	-
<b>Endocrine and Metabolic</b>						
Breast pain	-	-	-	1	-	-
Decreased libido	1 to 2	-	-	-	-	-
Gout	-	-	-	1	-	-
Gynecomastia	1 to 2	-	-	-	-	-
<b>Gastrointestinal</b>						
Abdominal discomfort	-	≤5	-	<2	-	-
Abdominal pain	1 to 2	<1	-	<3	-	-
Acid regurgitation	1 to 2	-	-	-	-	-
Anorexia	-	-	-	-	-	1
Colitis	-	-	-	-	-	1
Constipation	<2	1 to 4	≤1	3	-	-

Adverse Events	Felodipine	Isradipine	Nicardipine	Nifedipine	Nimodipine	Nisoldipine
Diarrhea	1 to 2	≤3	-	<2	2 to 4	-
Dry mouth	1 to 2	≤1	≤1	<3	-	-
Dyspepsia	1 to 4	-	1 to 2	3 to 11	-	-
Dysphagia	-	-	-	-	-	1
Flatulence	1 to 2	-	-	<2	-	1
Gastritis	-	-	-	-	-	1
Gastrointestinal hemorrhage	-	-	-	-	1	1
Gastrointestinal symptoms	-	-	-	-	≤2	-
Hepatitis	-	-	-	-	1	1
Increased appetite	-	≤1	-	-	-	1
Jaundice	-	-	-	-	1	-
Nausea	1 to 2	1 to 5	2 to 5	3 to 11	≤1	2
Vomiting	1 to 2	≤1	≤5	-	-	-
<b>Genitourinary</b>						
Decreased libido	-	≤1	-	-	-	-
Dysuria	1 to 2	≤1	-	1	-	-
Hematuria	-	-	-	1	-	-
Impotence	1 to 2	≤1	✓	<3	-	-
Nocturia	-	≤1	-	1	-	-
Pollakiuria	-	1 to 3	-	-	-	-
Polyuria	1 to 2	-	-	1 to 3	-	-
Sexual dysfunction	-	-	-	<2	-	-
Urinary frequency/urgency	1 to 2	-	-	-	-	-
<b>Hematological</b>						
Anemia	1 to 2	-	-	<1	1	-
Leukopenia	-	≤1	-	-	-	-
Thrombocytopenia	-	-	✓	-	1	-
<b>Laboratory Test Abnormalities</b>						
Hepatic enzyme elevations	1 to 2	≤1	✓	-	≤1	-
Hyponatremia	-	-	-	-	1	-
<b>Musculoskeletal</b>						
Arthralgia	1 to 2	-	✓	<3	-	-
Back pain	1 to 2	≤1	-	1	-	-
Hypertonia	-	-	✓	1	-	-
Inflammation	-	-	-	<2	-	-
Joint sprain	1 to 2	≤1	-	-	-	-
Malaise	-	-	≤1	1	-	1
Muscle cramps	1 to 2	≤1	-	3 to 8	≤1	-
Muscle weakness	-	≤1	-	10 to 12	-	-
Musculoskeletal chest pain	1 to 2	2 to 3	-	<3	-	-
Myalgia	1 to 2	-	1	1	-	-
Neck pain	-	≤1	✓	-	-	-
Pain	-	-	≤1	<3	-	-
Stiffness	-	-	-	<2	-	-
<b>Respiratory</b>						
Bronchitis	1 to 2	-	-	-	-	-
Cough	1 to 2	≤1	-	1 to 6	-	-
Dyspnea	1 to 2	<3	≤1	3 to 6	≤1	-
Epistaxis	1 to 2	-	-	1	-	-
Influenza/flu-like illness	1 to 2	-	-	-	-	1
Nasal congestion	1 to 2	≤1	✓	2 to 6	-	-
Nasopharyngitis	1 to 2	-	-	-	-	-
Pharyngitis	-	-	-	-	-	5
Pharyngolaryngeal pain	-	≤1	-	-	-	-
Shortness of breath	-	≤1	-	<2	1	-
Sinusitis	1 to 2	-	✓	1	-	3
Sore throat	-	-	✓	6	-	-
Upper respiratory tract infection	1 to 4	-	-	1	-	-
<b>Special Senses</b>						
Abnormal visual accommodation	-	-	✓	1	-	-
Blurred vision	-	-	✓	<2	-	-
Conjunctivitis	-	-	✓	-	-	-
Ear pain/disorder	-	-	✓	-	-	-
Taste perversion	-	-	-	1	-	-
Tinnitus	-	-	✓	<5	-	-
Visual disturbance	1 to 2	≤1	-	<5	-	-
<b>Other</b>						

Adverse Events	Felodipine	Isradipine	Nicardipine	Nifedipine	Nimodipine	Nisoldipine
Allergic reaction	-	-	✓	-	-	-
Angioedema	1 to 2	-	-	-	-	-
Cellulitis	-	-	-	-	-	1
Contusion	1 to 2	-	-	-	-	-
Facial edema	-	-	-	-	-	1
Fever	-	≤1	-	<2	-	1
Gingival hyperplasia	1 to 2	-	-	-	-	1
Glossitis	-	-	-	-	-	1
Hot flush	-	-	✓	-	-	-
Infection	-	-	✓	-	-	-
Rigors	-	-	-	1	-	-
Warm sensation	1 to 2	-	-	-	-	-
Weight gain	-	≤1	-	1	-	-

✓ Percent not specified  
- Event not reported

**Table 8. Boxed Warning for Amlodipine and Benazepril<sup>1</sup>**

WARNING
When pregnancy is detected, discontinue amlodipine and benazepril as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

**Table 9. Boxed Warning for Amlodipine and Olmesartan<sup>1</sup>**

WARNING
When pregnancy is detected, discontinue amlodipine and olmesartan as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

**Table 10. Boxed Warning for Amlodipine and Valsartan<sup>1</sup>**

WARNING
When pregnancy is detected, discontinue amlodipine and valsartan as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

**Table 11. Boxed Warning for Amlodipine and Valsartan and Hydrochlorothiazide<sup>1</sup>**

WARNING
When pregnancy is detected, discontinue amlodipine and valsartan and hydrochlorothiazide as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

**Table 12. Boxed Warning for Nimodipine<sup>1</sup>**

WARNING
Do not administer nimodipine intravenously or by other parenteral routes. Deaths and serious, life-threatening adverse reactions have occurred when the contents of nimodipine capsules have been injected parenterally.

## VII. Dosing and Administration

The usual dosing regimens for the dihydropyridines are listed in Table 14.

**Table 13. Usual Dosing Regimens for the Dihydropyridines<sup>1,2,6-17</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Amlodipine	<u>Angina pectoris (chronic stable and vasospastic):</u> Suspension, tablet: maintenance, 5 to 10 mg/day; maximum, 10 mg/day	<u>Hypertension in children six to 17 years of age:</u> Suspension, tablet: Initial, 2.5 mg/day; maintenance, 2.5 to 5	Suspension: 1 mg/mL  Tablet: 2.5 mg

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<p><u>Coronary artery disease:</u> Suspension, tablet: maintenance, 5 to 10 mg/day; maximum, 10 mg/day</p> <p><u>Hypertension:</u> Suspension, tablet: initial, 2.5 to 5 mg/day maintenance, 5 to 10 mg/day; maximum, 10 mg/day</p>	<p>mg/day; maximum, 5 mg/day</p> <p>Safety and efficacy in children &lt;6 years of age have not been established.</p>	<p>5 mg 10 mg</p>
Felodipine	<p><u>Hypertension:</u> Extended-release tablet: initial, 5 mg/day; maintenance, 2.5 to 10 mg/day</p>	<p>Safety and efficacy in children have not been established.</p>	<p>Extended-release tablet: 2.5 mg 5 mg 10 mg</p>
Isradipine	<p><u>Hypertension:</u> Capsule: initial, 2.5 mg twice daily; maximum, 20 mg/day</p>	<p>Safety and efficacy in children have not been established.</p>	<p>Capsule: 2.5 mg 5 mg</p>
Nicardipine	<p><u>Angina pectoris (chronic stable):</u> Capsule: initial, 20 mg three times daily; maintenance, 20 to 40 mg three times daily</p> <p><u>Hypertension:</u> Capsule: initial, 20 mg three times daily; maintenance, 20 to 40 mg three times daily Injection: titrate dose to achieve the desired blood pressure reduction; individualize dosage depending on the blood pressure to be obtained and the response of the patient</p>	<p>Safety and efficacy in children have not been established.</p>	<p>Capsule: 20 mg 30 mg</p> <p>Injection: 25 mg/10 mL</p>
Nifedipine	<p><u>Angina pectoris (chronic stable):</u> Capsule: initial, 10 mg three times daily; maintenance, 10 to 20 mg three times daily; maximum, 180 mg/day</p> <p>Extended-release tablet: initial, 30 or 60 mg/day; maintenance, 30 to 90 mg/day; maximum, 120 mg/day</p> <p><u>Angina pectoris (vasospastic):</u> Capsule: initial, 10 mg three times daily; maintenance, 20 to 30 mg three to four times daily; maximum, 180 mg/day</p> <p>Extended-release tablet: initial, 30 or 60 mg/day; maintenance, 30 to 90 mg/day; maximum, 120 mg/day</p> <p><u>Hypertension:</u></p>	<p>Safety and efficacy in children have not been established.</p>	<p>Capsule: 10 mg 20 mg</p> <p>Extended-release tablet: 30 mg 60 mg 90 mg</p>

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	Extended-release tablet: initial, 30 or 60 mg/day; maintenance, 30 to 90 mg/day; maximum, 120 mg/day		
Nimodipine	<u>Subarachnoid hemorrhage:</u> Capsule, solution: 60 mg every four hours for 21 consecutive days	Safety and efficacy in children have not been established.	Capsule: 30 mg  Solution: 60 mg/20 mL
Nisoldipine	<u>Hypertension:</u> Extended-release tablet: initial, 20 mg once daily; maintenance, 20 to 40 mg/day; maximum, 60 mg/day  Extended-release tablet (Sular <sup>®</sup> only): initial, 17 mg once daily; maintenance, 17 to 34 mg/day; maximum, 34 mg/day	Safety and efficacy in children have not been established.	Extended-release tablet: 8.5 mg 17 mg 20 mg 25.5 mg 30 mg 34 mg 40 mg
<b>Combination Products</b>			
Amlodipine and benazepril	<u>Hypertension:</u> Capsule: initial, 2.5-10 mg once daily; maintenance, individualize and adjust dosage according to clinical response, dose once daily	Safety and efficacy in children have not been established.	Capsule: 2.5-10 mg 5-10 mg 5-20 mg 5-40 mg 10-20 mg 10-40 mg
Amlodipine and olmesartan	<u>Hypertension:</u> Tablet: initial, 5-20 mg once daily; maximum, 10-40 mg once daily	Safety and efficacy in children have not been established.	Tablet: 5-20 mg 5-40 mg 10-20 mg 10-40 mg
Amlodipine and valsartan	<u>Hypertension:</u> Tablet: initial, 5-160 mg once daily; maximum, 10-320 mg once daily	Safety and efficacy in children have not been established.	Tablet: 5-160 mg 5-320 mg 10-160 mg 10-320 mg
Amlodipine and valsartan and HCTZ	<u>Hypertension:</u> Tablet: maximum, 10-320-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 5-160-12.5 mg 5-160-25 mg 10-160-12.5 mg 10-160-25 mg 10-320-25 mg

HCTZ=hydrochlorothiazide



## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the dihydropyridines are summarized in Table 14.

**Table 14. Comparative Clinical Trials with the Dihydropyridines**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Angina</b>				
Koenig et al. <sup>37</sup> (1997)  Amlodipine 5 to 10 mg QD for 4 weeks  vs  felodipine ER 5 to 10 mg QD for 4 weeks	DB, PRO, RCT, XO  Patients, age 30 to 80 years, who have a history of angina, a positive exercise-stress test or positive 24-hour ambulatory monitoring, and $\geq 6$ ischemic episodes in 24 hours	N=52  8 weeks	Primary: Number of ST-segment depressions in 24 hours of ambulatory monitoring  Secondary: Total and mean duration of each ST-segment depression episode, maximum ST depression, length of ischemic episode, adverse events	Primary: Significant reductions from baseline were seen in both groups for the number of ST-segment depressions, from 19.9 at baseline for both groups to 2.3 for amlodipine and 2.4 for felodipine ( $P < 0.001$ for both from baseline; $P = 0.83$ between treatments).  Secondary: Total and mean duration of each ST-segment depression episode, maximum ST depression and length of ischemic episode were significantly different from baseline for both treatment groups, but treatments were not significantly different ( $P < 0.001$ for all from baseline, $P = 0.53$ , $P = 0.40$ , $P = 0.68$ , $P = 0.35$ , respectively between treatments).  Adverse event rates similar between the treatments.
Savanitto et al. <sup>38</sup> (1996)  <u>Weeks 1 to 6:</u> Nifedipine 20 mg BID  vs  metoprolol ER 200 mg QD  <u>Weeks 7 to 10:</u>	DB, MC, RCT  Patients with typical anginal symptoms that had been stable for approximately 6 months, who showed a positive response to exercise stress testing with 23 min of exercise tolerance and were in sinus	N=280  6 weeks	Primary: Angina frequency, exercise tolerance, safety  Secondary: Not reported	Primary: At week six, both metoprolol (mean change, -1.95; 95 % CI, -1.25 to -2.64) and nifedipine (mean change, -1.57; 95 % CI, -0.69 to -2.45) significantly reduced the frequency of angina compared to baseline, but there was not a statistical difference between groups. At the end of 10 weeks, there was not a statistical difference observed between the groups.  At week six, both metoprolol and nifedipine significantly increased the mean exercise time to 1-mm ST-segment depression compared to baseline (both $P < 0.01$ ); but metoprolol was significantly more effective than nifedipine ( $P < 0.05$ ).  At week 10, the groups randomized to combination therapy had a further

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>nifedipine 20 mg BID plus placebo vs metoprolol ER 200 mg QD plus placebo vs metoprolol ER 200 mg QD and nifedipine 20 mg BID</p>	<p>rhythm and had an analyzable ST segment on ECG</p>			<p>increase in time to 1-mm ST-segment depression (P&lt;0.05 vs placebo).</p> <p>There were 14 cardiovascular events including one sudden death, three acute myocardial infarctions, eight cases of unstable angina, one of syncope and one of stroke and the incidence of these events did not differ among the treatment groups.</p> <p>Secondary: Not reported</p>
<b>Cardiovascular Outcomes Trials</b>				
<p>Pitt et al.<sup>39</sup> (2000) PREVENT  Amlodipine 5 to 10 mg QD  vs  placebo</p>	<p>DB, MC, PC, RCT  Men and women, age 30 to 80 years with angiographic evidence of CAD, DBP &lt;95 mm Hg, TC 325 mg/dL, FBG &lt;200 mg/dL</p>	<p>N=825  3 years</p>	<p>Primary: Change in mean minimal diameter with a quantitative coronary angiography</p> <p>Secondary: Progression of atherosclerosis in the carotid arteries assessed by B-mode ultrasonography for intimal-medial thicknesses, all-cause mortality, occurrence of major fatal/nonfatal vascular events or procedures, adverse events</p>	<p>Primary: Change, reduction, in the minimal diameter was similar between the amlodipine group and the placebo group (0.084 vs 0.0095 P=0.38).</p> <p>Secondary: Amlodipine treatment significantly decreased the progression of atherosclerosis as compared to placebo treatment, a 0.013 mm decrease for the amlodipine group vs a 0.033 mm increase with placebo (P=0.007).</p> <p>There was no difference in all-cause mortality between amlodipine and placebo.</p> <p>There was no difference in occurrence of fatal and nonfatal vascular events between the treatment groups (HR, 0.82; 95% CI, 0.47 to 1.42).</p> <p>Amlodipine treatment significantly reduced the occurrence of hospitalized CHF and unstable angina (HR, 0.65; 95% CI, 0.47 to 0.91) and coronary revascularizations (HR, 0.57; 95% CI, 0.41 to 0.81) and combined overall procedures (HR, 0.69; 95% CI, 0.52 to 0.92).</p> <p>There was no significant difference between groups in rates of adverse events: cancer rate (HR, 2.13; 95% CI, 0.90 to 5.21) and bleeding episode (HR, 1.42; 95% CI, 0.88 to 2.30).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Dahlöf et al.<sup>40</sup> (2005) ASCOT-BPLA</p> <p>Amlodipine 5 to 10 mg/day adding perindopril 4 to 8 mg/day as needed</p> <p>vs</p> <p>atenolol 50 to 100 mg/day adding bendroflumethiazide* 1.25 to 2.5 mg/day and potassium as needed</p> <p>If blood pressure was still not achieved, doxazosin 4 to 8 mg/day was added to the regimen.</p>	<p>MC, OL, RCT</p> <p>Patients 40 to 79 years of age with HTN and ≥3 other cardiovascular risk factors (left ventricular hypertrophy, other specified abnormalities on ECG, type 2 diabetes, PAD, history of stroke or TIA, male, age ≥55 years, microalbuminuria or proteinuria, smoking, TC:HDL-C ratio ≥6, or family history of CHD)</p>	<p>N=19,257</p> <p>5.5 years</p>	<p>Primary: Nonfatal MI (including silent MI) and fatal CHD</p> <p>Secondary: All-cause mortality, total stroke, primary end points minus silent MI, all coronary events, total cardiovascular events and procedures, cardiovascular mortality, nonfatal and fatal heart failure, effects on primary end point and on total cardiovascular events and procedures among prespecified subgroups</p> <p>Tertiary: Silent MI, unstable angina, chronic stable angina, PAD, life-threatening arrhythmias, development of diabetes, development of</p>	<p>Primary: No statistically significant difference in nonfatal MI and fatal CHD was reported between the amlodipine plus perindopril group compared to the atenolol plus bendroflumethiazide groups (HR, 0.90; 95% CI, 0.79 to 12; P=0.1052).</p> <p>Secondary: Significantly greater reductions in the following secondary end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: all- cause mortality (P=0.0247), total stroke (P=0.0003), primary end points minus silent MI (P=0.0458), all coronary events (P=0.0070), total cardiovascular events and procedures (P&lt;0.0001), and cardiovascular mortality (P=0.0010).</p> <p>There were no significant differences in nonfatal and fatal heart failure between the two treatment groups (P=0.1257).</p> <p>The study was terminated early due to higher mortality and worse outcomes on several secondary end points observed in the atenolol study group.</p> <p>Tertiary: Significantly greater reductions in the following end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: unstable angina (P=0.0115), PAD (P=0.0001), development of diabetes (P&lt;0.0001), and development of renal impairment (P=0.0187).</p> <p>There were no significant differences in the incidence of silent MI (P=0.3089), chronic stable angina (P=0.8323) or life-threatening arrhythmias (P=0.8009) between the two treatment groups.</p> <p>There was no significant difference in the percent of patients who stopped therapy because of an adverse event between the two treatment groups (overall 25%). There was, however, a significant difference in favor of amlodipine plus perindopril in the proportion of patients who stopped trial therapy because of a serious adverse events (2 vs 3%; P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Chapman et al.<sup>41</sup> (2007) ASCOT-BPLA</p> <p>Atenolol 50 to 100 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); bendroflumethiazide* plus potassium 1.25 to 2.5 mg plus doxazosin were added for additional blood pressure control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p> <p>vs</p> <p>amlodipine 5 to 10 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); perindopril 4 to 8 mg and doxazosin were added for</p>	<p>Sub analysis of ASCOT-BPLA evaluating effects of spironolactone on treatment-resistant HTN</p> <p>Patients 40 to 79 years of age with HTN and ≥3 cardiovascular risk factors, with SBP ≥160 mm Hg and/or DBP ≥100 mm Hg (not on antihypertensive therapy) or SBP ≥140 mm Hg and/or DBP ≥90 mm Hg (on antihypertensive therapy)</p>	<p>N=1,411</p> <p>1.3 years</p>	<p>renal impairment</p> <p>Primary: Change in DBP and SBP, adverse effects</p> <p>Secondary: Not reported</p>	<p>Primary: Spironolactone-treated patients lead to a significant 21.9 mm Hg reduction in SBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 20.8 to 23.0 mm Hg; P&lt;0.001).</p> <p>Spironolactone-treated patients lead to a significant 9.5 mm Hg reduction in DBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 9.0 to 10.1; P&lt;0.001).</p> <p>Spironolactone-treated patients exhibited small but significant decreases in sodium, LDL-C and TC as well as increases in potassium, glucose, creatinine and HDL-C (P&lt;0.05).</p> <p>The most common adverse effect reported in the trial was gynecomastia in men (P value not reported).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>additional control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p>				
<p>Nissen et al.<sup>42</sup> (2004) CAMELOT  Amlodipine 10 mg/day  vs  enalapril 20 mg/day  vs  placebo</p>	<p>DB, MC, PC, RCT  Patients 30 to 79 years of age requiring coronary angiography for evaluation for chest pain or PCI and a diastolic pressure &lt;100 mm Hg, with or without treatment</p>	<p>N=1,991  2 years</p>	<p>Primary: Composite of cardiovascular events (cardiovascular death, nonfatal MI, resuscitated cardiac arrest, coronary revascularization, hospitalization for angina pectoris, hospitalization for CHF, fatal or nonfatal stroke or TIA, and any new diagnosis of PVD), nominal change in percent atheroma volume (substudy)  Secondary: Incidence of adverse events; all-cause mortality, incidence of revascularization in vessels that had undergone</p>	<p>Primary: Cardiovascular events occurred in 23.1% of placebo-treated patients, 16.6% amlodipine-treated patients (HR, 0.69; 95% CI, 0.54 to 0.88; P=0.003) and 20.2% enalapril-treated patients (HR, 0.85; 95% CI, 0.67 to 17; P=0.16).  The primary end point comparison for enalapril vs amlodipine was not significant (HR, 0.81; 95% CI, 0.63 to 14; P=0.10).  Secondary: Coronary revascularization was reduced in the amlodipine group from 15.7 to 11.8% (HR, 0.73; 95% CI, 0.54 to 0.98; P=0.03). Hospitalization for angina was reduced in the amlodipine group from 12.8 to 7.7% (HR, 0.58; 95% CI, 0.41 to 0.82; P=0.002).  Individual components of the primary end point generally showed fewer events with enalapril treatment vs placebo, but none of the comparisons reached statistical significance.  For components of the primary end point, only the rate of hospitalization for angina showed a statistically significant difference between amlodipine and enalapril (HR, 0.59; 95% CI, 0.42 to 0.84; P=0.003). A trend toward fewer episodes of revascularization in patients undergoing intervention at baseline was observed for amlodipine vs enalapril (HR, 0.66; 95% CI, 0.40 to 16; P=0.09).  The mean change in percent atheroma volume was 0.5% for amlodipine (P=0.12 vs placebo), 0.8% for enalapril (P=0.32 vs placebo) and 1.3% for placebo. In patients with SBP greater than the mean, the amlodipine group</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			previous stent placement	<p>showed a significantly slower progression (0.2%) compared to placebo (2.3%; P=0.02). Compared to baseline, intravascular ultrasound showed progression in patients receiving placebo (P&lt;0.001), a trend toward progression with enalapril (P=0.08) and no progression in patients receiving amlodipine (P=0.31). For the amlodipine group, correlation between blood pressure reduction and progression was r=0.19 (P=0.07).</p> <p>Discontinuation from the study for treatment-emergent adverse events was low, averaging 0.4% and not statistically significant between the three treatment groups.</p> <p>The only statistically significant difference in secondary end points was that amlodipine demonstrated a significant reduction in revascularization after previous stent placement compared to placebo (4.1 vs 7.9%; HR, 0.49; 95% CI, 0.31 to 0.78; P=0.002). The rate of revascularization was lower than enalapril (6.2%) but not statistically significant (HR, 0.66; 95% CI, 0.40 to 1.6; P=0.09).</p>
<p>ALLHAT<sup>43</sup> (2002) ALLHAT Amlodipine 2.5 to 10 mg/day vs lisinopril 10 to 40 mg/day vs chlorthalidone 12.5 to 25 mg/day</p>	<p>DB, MC, RCT  Patients ≥55 years with HTN and ≥1 additional CHD risk factor</p>	<p>N=33,357  4.9 years (mean)</p>	<p>Primary: Combined fatal CHD or nonfatal MI  Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization, heart failure, and PAD)</p>	<p>Primary: There were no significant differences in the primary outcome between lisinopril (11.4%), amlodipine (11.3%), and chlorthalidone (11.5%).  Secondary: All-cause mortality did not differ between groups.  Five year SBPs were significantly higher in the lisinopril (2 mm Hg; P&lt;0.001) and amlodipine groups (0.8 mm Hg; P=0.03) compared to chlorthalidone, and five year DBPs were significantly lower with amlodipine (0.8 mm Hg; P&lt;0.001).  Amlodipine had a higher six year rate of heart failure compared to chlorthalidone (10.2 vs 7.7%; RR, 1.38; 95% CI, 1.25 to 1.52).  Lisinopril had a higher six year rate of combined cardiovascular disease (33.3 vs 30.9%; RR, 1.10; 95% CI, 1.05 to 1.16); stroke (6.3 vs 5.6%; RR, 1.15; 95% CI, 1.05 to 1.30) and heart failure (8.7 vs 7.7%; RR, 1.19; 95% CI, 1.07 to 1.31).</p>
Black et al. <sup>44</sup>	MC, RCT	N=17,515	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2008) ALLHAT</p> <p>Amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>lisinopril 10 to 40 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg QD</p>	<p>Men and women, age 55 years old and older, with HTN and metabolic syndrome</p>	<p>4.9 years (mean)</p>	<p>Fatal coronary heart disease and nonfatal MI</p> <p>Secondary: All cause mortality, fatal and nonfatal stroke, combined coronary heart disease, combined cardiovascular disease</p>	<p>For patients with metabolic syndrome, there was no significant difference in rates of coronary heart disease and nonfatal MI with amlodipine vs chlorthalidone (RR, 0.96; 95% CI, 0.79 to 1.16), or lisinopril vs chlorthalidone (RR, 1.15; 95% CI, 0.88 to 1.27).</p> <p>Secondary: For patients with metabolic syndrome, there were no significant differences found between amlodipine vs chlorthalidone in all secondary endpoints (P value not significant).</p> <p>For patients without metabolic syndrome, amlodipine treatment was associated with significantly more heart failure, but in patients with metabolic syndrome, there was no difference (P=0.03).</p> <p>Patients with metabolic syndrome who received lisinopril experienced more heart failure and cardiovascular disease than those who received chlorthalidone (RR, 1.31; 95% CI, 1.14 to 1.64 and RR, 1.19; 95% CI, 1.07 to 1.32).</p>
<p>Rahman et al.<sup>45</sup> (2012) ALLHAT</p> <p>Amlodipine 2.5 to 10 mg/day</p> <p>vs</p> <p>lisinopril 10 to 40 mg/day</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg/day</p>	<p>Long-term, post-trial, follow-up</p> <p>Patients in ALLHAT stratified based on eGFR</p>	<p>N=31,350</p> <p>4 to 8 years</p>	<p>Primary: Cardiovascular mortality</p> <p>Secondary: Total mortality, CHD, cardiovascular disease, stroke, heart failure, ESRD</p>	<p>Primary: After an average of 8.8 years of follow-up, total mortality was significantly higher in patients with moderate/severe eGFR reduction (eGFR &lt;60 mL/min/1.73 m<sup>2</sup>) compared to patients with normal/increased (eGFR ≥90 mL/min/1.73 m<sup>2</sup>) and mildly reduced eGFR (eGFR 60 to 89 mL/min/1.73 m<sup>2</sup>) (P&lt;0.001).</p> <p>In patients with moderate/severe eGFR reduction, there was no significant difference in cardiovascular mortality between chlorthalidone and amlodipine (P=0.64), or chlorthalidone and lisinopril (P=0.56).</p> <p>Secondary: No significant differences were observed for any of the secondary endpoints among eGFR reduction groups.</p>
<p>Muntner et al.<sup>46</sup> (2014) ALLHAT</p>	<p>Post-hoc analysis of ALLHAT</p>	<p>N=24,004</p> <p>6 to 28 months</p>	<p>Primary: Visit-to-visit variability (VVV)</p>	<p>Primary: Each measure of VVV of SBP was lower among participants randomized to chlorthalidone and amlodipine compared with those randomized to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Chlorthalidone 12.5 to 25 mg/day</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg/day</p> <p>vs</p> <p>lisinopril 10 to 40 mg/day</p>	<p>Patients in ALLHAT with 5, 6, or 7 visits in 6 to 28 months of follow-up</p>		<p>of blood pressure</p> <p>Secondary: Not reported</p>	<p>lisinopril. All four VVV of SBP metrics were lower among participants randomized to amlodipine vs chlorthalidone after full multivariable adjustment.</p> <p>After multivariable adjustment including mean SBP across visits and compared with participants randomized to chlorthalidone, participants randomized to amlodipine had a 0.36 (standard error [SE]: 0.07) lower standard deviation (SD) of SBP and participants randomized to lisinopril had a 0.77 (SE=0.08) higher SD of SBP. Results were consistent using other VVV of SBP metrics. These data suggest chlorthalidone and amlodipine are associated with lower VVV of SBP than lisinopril.</p> <p>Secondary: Not reported</p>
<p>Bangalore et al.<sup>47</sup> (2017) ALLHAT</p> <p>Amlodipine 2.5 to 10 mg/day</p> <p>vs</p> <p>lisinopril 10 to 40 mg/day</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg/day</p>	<p>Post-hoc analysis of ALLHAT</p> <p>Patients in ALLHAT with average blood pressure <math>\geq 140</math> mmHg systolic or <math>\geq 90</math> mm Hg diastolic on <math>\geq 3</math> antihypertensive medications, or blood pressure <math>&lt; 140/90</math> mmHg on <math>\geq 4</math> antihypertensive medications (i.e., identified as having apparent treatment-resistant hypertension) at 2-year follow up</p>	<p>N=14,684</p> <p>4.9 years (mean)</p>	<p>Primary: Combined fatal CHD or nonfatal MI</p> <p>Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization, heart failure, and PAD)</p>	<p>Primary: Of participants assigned to chlorthalidone, amlodipine, or lisinopril, 9.6%, 11.4%, and 19.7%, respectively, had treatment-resistant hypertension. During mean follow-up of 2.9 years, primary outcome incidence was similar for those assigned to chlorthalidone compared with amlodipine or lisinopril (amlodipine- vs chlorthalidone-adjusted HR, 0.86; 95% CI, 0.53 to 1.39; P=0.53; lisinopril- vs chlorthalidone-adjusted HR, 1.06; 95% CI, 0.70 to 1.60; P=0.78).</p> <p>Secondary: Secondary outcome risks were similar for most comparisons except coronary revascularization, which was higher with amlodipine than with chlorthalidone (HR, 1.86; 95% CI, 1.11 to 3.11; P=0.02). An as-treated analysis based on diuretic use produced similar results.</p>
<p>Ogihara et al.<sup>48</sup> (2008)</p>	<p>AC, MC, OL, RCT</p>	<p>N=4,703</p>	<p>Primary: First fatal or</p>	<p>Primary: A total of 134 patients experienced a cardiovascular event in each</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>CASE-J</p> <p>Amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>candesartan 4 to 12 mg QD</p>	<p>Patients with high risk HTN (SBP <math>\geq 140</math> mm Hg or DBP <math>\geq 90</math> mm Hg in patients <math>&lt; 70</math> years old or SBP <math>\geq 160</math> mm Hg or DBP <math>\geq 90</math> mm Hg in patients <math>\geq 70</math> years old), with either type 2 diabetes, history of stroke or ischemic attack, left ventricular hypertrophy, proteinuria or serum creatinine <math>\geq 1.3</math> mg/dL</p>	<p>Up to 4 years</p>	<p>nonfatal cardiovascular event</p> <p>Secondary: All-cause death, new-onset diabetes, discontinuation due to adverse events</p>	<p>treatment regimen (HR, 1.0; 95% CI, 0.78 to 1.27; <math>P=0.969</math>).</p> <p>Secondary: All-cause death rates did not differ between treatments, 73 deaths in the candesartan group and 86 in the amlodipine group.</p> <p>New-onset diabetes occurred in significantly fewer patients in the candesartan group than the amlodipine group (HR, 0.64; 95% CI, 0.43 to 0.97; <math>P=0.033</math>).</p> <p>A total of 125 (5.4%) patients in the candesartan group and 134 (5.8%) of patients in the amlodipine group discontinued due to adverse events.</p>
<p>Julius et al.<sup>49</sup> (2004)</p> <p>VALUE</p> <p>Amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>valsartan 80 to 160 mg QD</p>	<p>DB, PG, RCT</p> <p>Patients <math>\geq 50</math> years old with treated or untreated HTN and history of cardiovascular disease, stroke, or diabetes, previous medications were discontinued at trial onset</p>	<p>N=15,245</p> <p>4.2 years (mean)</p>	<p>Primary: Time to first cardiac event (cardiac morbidity and mortality)</p> <p>Secondary: Fatal and nonfatal MI, fatal and nonfatal heart failure and fatal and nonfatal stroke, all-cause mortality, new onset diabetes</p>	<p>Primary: There were no differences in the primary composite end point between the valsartan and amlodipine groups (10.6 vs 10.4%; <math>P=0.49</math>).</p> <p>Secondary: There was a higher incidence of myocardial infarction (4.8 vs 4.1%; <math>P=0.02</math>) in patients receiving valsartan than amlodipine.</p> <p>There was no difference in the incidence of heart failure (4.6 vs 5.3%; <math>P=0.12</math>), stroke (4.2 vs 3.7%; <math>P=0.08</math>), and all-cause mortality (11 vs 10.8%; <math>P=0.45</math>) between valsartan- and amlodipine-treated patients.</p> <p>New onset diabetes occurred less with valsartan (13.1%) vs amlodipine (16.4%; <math>P&lt;0.001</math>).</p> <p>Combined target blood pressure (<math>&lt; 140/90</math> mm Hg) was achieved in 58% and 62% of patients receiving valsartan and amlodipine, respectively.</p>
<p>Zanchetti et al.<sup>50</sup> (2006)</p>	<p>Subgroup analysis of VALUE</p>	<p>N=15,245</p>	<p>Primary: Time to first</p>	<p>Primary: The only significant result of the analyses by subgroup for time to first</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>VALUE</p> <p>Amlodipine 5 mg QD</p> <p>vs</p> <p>valsartan 80 mg QD</p>	<p>Patients with HTN</p>	<p>4.2 years</p>	<p>cardiac event, analyzed by subgroup</p> <p>Secondary: MI, heart failure and stroke</p>	<p>cardiac event was sex; women in the valsartan group experienced more cardiac events as compared to men in the valsartan group (HR for women, 1.21; 95% CI, 1.13 to 1.42; HR for men, 0.94; 95% CI, 0.82 to 1.17; P=0.016).</p> <p>The VALUE trial showed no difference in the primary outcome as well as in cardiac morbidity and mortality between amlodipine treatment and valsartan treatment. SBP and DBP were lower, as was incidence of MI, in the amlodipine treatment group as compared to the valsartan group.</p> <p>Secondary: Male patients treated with valsartan had a significantly lower incidence of heart failure than males treated with amlodipine (P&lt;0.001 for male vs female difference; for men, HF rates with valsartan were 4.1% vs amlodipine 5.8% [HR, 0.73; 95% CI, 0.60 to 0.88]; for women, rates were valsartan 5.3% vs amlodipine 4.6%, [HR, 1.18; 95% CI, 0.95 to 1.47]).</p> <p>Patients without a history of stroke had a greater reduction in stroke risk if treated with amlodipine (valsartan 3.4% vs amlodipine 2.6%; HR, 1.34; 95% CI, 1.19 to 1.65).</p>
<p>Jamerson et al.<sup>51</sup> (2008)</p> <p>ACCOMPLISH</p> <p>Amlodipine 5 mg QD and benazepril 20 mg QD</p> <p>vs</p> <p>benazepril 20 mg QD and HCTZ 12.5 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=11,506</p> <p>36 months (mean)</p>	<p>Primary: The composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for angina, resuscitation after sudden cardiac arrest, and coronary revascularization.</p> <p>Secondary: Death from</p>	<p>Primary: There were 552 primary-outcome events in the benazepril plus amlodipine group (9.6%) and 679 events in the benazepril plus HCTZ group (11.8%). The absolute risk reduction with benazepril plus amlodipine therapy was 2.2% and the relative risk reduction was 19.6% compared to benazepril plus HCTZ (HR, 0.80; 95% CI, 0.72 to 0.90; P&lt;0.001).</p> <p>Secondary: For the secondary end point of death from cardiovascular causes, nonfatal MI, and nonfatal stroke, there were 288 (5%) events in the benazepril plus amlodipine group compared to 364 (6.3%) events in the benazepril plus HCTZ group. The absolute risk reduction with benazepril plus amlodipine therapy was 1.3% and the RR reduction was 21.2% compared to benazepril plus HCTZ (HR, 0.79; 95% CI, 0.67 to 0.92; P=0.002).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			cardiovascular causes, nonfatal MI, and nonfatal stroke	
<p>Bakris et al.<sup>52</sup> (2010) ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to 40-10 mg/day, followed by forced titration after 1 month on benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p> <p>benazepril and HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after 1 month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)</p>	<p>Prespecified sub analysis of ACCOMPISH</p> <p>Men and women &gt;60 years of age with HTN and at high risk for cardiovascular events (history of coronary events, MI, revascularization, or stroke; impaired renal function; PAD, left ventricular hypertrophy; or diabetes)</p>	<p>N=11,482</p> <p>2.9 years (mean duration)</p>	<p>Primary: Time to first event of doubling of serum creatinine concentration or end stage renal disease (defined as eGFR &lt;15 mL/min/1.73 m<sup>2</sup> or need for chronic dialysis)</p> <p>Secondary: Progression of chronic kidney disease plus death, change in albuminuria, and change in eGFR</p>	<p>Primary: There were fewer chronic kidney disease events in the benazepril and amlodipine group (2.0% of patients) compared to the benazepril and HCTZ group (3.7%; HR, 0.52; 95% CI, 0.41 to 0.65; P&lt;0.0001).</p> <p>Secondary: The composite endpoint of progression of chronic kidney disease and all-cause mortality was lower in the benazepril and amlodipine group (6.0%) compared to the benazepril and HCTZ group (8.1%; HR, 0.73; 95% CI, 0.64 to 0.84; P&lt;0.0001). There was a slower decline in eGFR in the benazepril and amlodipine group compared to the benazepril and HCTZ group (-0.88 vs -4.22 mL/min/1.73 m<sup>2</sup>; P=0.01). Of the patients with baseline microalbuminuria, there was a reduction in the urinary albumin:creatinine in the benazepril and HCTZ group of -63.8% (median change) compared to a median change of -29.0% in the benazepril and amlodipine group (P&lt;0.0001).</p> <p>There was a higher percentage of patients reporting peripheral edema in the benazepril and amlodipine group compared to the benazepril and HCTZ group (P&lt;0.0001).</p>
<p>Weber et al.<sup>53</sup> (2010) ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to</p>	<p>Prespecified subanalysis of ACCOMPISH</p> <p>Men and women &gt;60 years of age</p>	<p>N=6,946</p> <p>Mean treatment duration 29.7 months for</p>	<p>Primary: Primary: Time to first event (composite of cardiovascular event and death</p>	<p>Primary: The primary endpoint occurred in 8.8% of diabetic patients in the benazepril and amlodipine group and 11.0% in the benazepril and HCTZ group (HR, 0.79; P=0.003; NNT, 46). In high risk diabetic patients, 13.6% of patients in the benazepril and amlodipine group and 17.3% in the benazepril and HCTZ group (HR, 0.77, P=0.007; NNT, 28).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>40-10 mg/day, followed by forced titration after 1 month on benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p> <p>benazepril and HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after one month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)</p>	<p>with HTN and at high risk for cardiovascular events (history of coronary events, MI, revascularization, or stroke; impaired renal function; peripheral arterial disease, left ventricular hypertrophy; or diabetes)</p> <p>(Subanalysis of patients with diabetes)</p>	<p>benazepril and amlodipine group and 29.5 months for benazepril and HCTZ group</p>	<p>from cardiovascular causes)</p> <p>Secondary: Composite of cardiovascular events (the primary endpoint excluding fatal events) and composite of death from cardiovascular disease, nonfatal stroke and nonfatal MI</p>	<p>Secondary: Due to early termination, the study had limited power to detect differences in the diabetic subgroups.</p> <p>Peripheral edema was higher in the benazepril and amlodipine group compared to the benazepril and HCTZ group.</p>
<p>Weber et al.<sup>54</sup> (2013) ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to 40-10 mg/day, followed by forced titration after 1 month on benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p>	<p>Subanalysis of ACCOMPLISH based on body size</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=11,482</p>	<p>Primary: Composite of cardiovascular death or nonfatal MI or stroke</p> <p>Secondary: Cardiovascular death, total MI, total stroke</p>	<p>Primary: In patients receiving benazepril and HCTZ, the primary endpoint (per 1,000 patient-years) was 30.7 in normal weight (BMI &lt;25), 21.9 in overweight (BMI ≥25 to &lt;30), and 18.2 in obese patients (BMI ≥30) (overall P=0.0034). In patients receiving benazepril and amlodipine, the primary endpoint did not differ between the three BMI groups (18.2, 16.9, and 16.5, respectively; P=0.9721). In obese patients, primary event rates were similar between the two treatments, but rates were significantly lower with benazepril and amlodipine in overweight patients (HR, 0.76; 95% CI, 0.59 to 0.94; P=0.0369) and normal weight patients (HR, 0.57; 95% CI, 0.39 to 0.84; P=0.0037).</p> <p>Secondary: Comparing obese and overweight patients, event rates were all numerically lower, but not significantly lower, in obese patients. Cardiovascular deaths were significantly lower in overweight patients</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>benazepril and HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after 1 month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)</p>				<p>compared to normal weight patients (HR, 57; 95% CI, 0.37 to 0.89; P=0.0125). Cardiovascular death (HR, 0.40; 95% CI, 0.25 to 0.63; P&lt;0.0001) and total stroke (HR, 0.60; 95% CI, 0.37 to 0.96; P=0.0335) were significantly lower in obese patients compared to normal weight patients.</p>
<p>Bakris et al.<sup>55</sup> (2013) ACCOMPLISH  HCTZ 12.5 to 25 mg QD and benazepril 20 to 40 mg QD (B+H)  vs  benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD (B+A)</p>	<p>Post hoc analysis  Patients included in the ACCOMPLISH trial (&gt;60 years of age with HTN and at high risk of cardiovascular events) stratified by presence of known CAD at baseline</p>	<p>N=11,506  36 months (mean)</p>	<p>Primary: The composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for angina, resuscitation after sudden cardiac arrest, and coronary revascularization.  Secondary: Death from cardiovascular causes, nonfatal MI, and nonfatal stroke</p>	<p>Primary: Among the patients with CAD, 13% in the B+A group and 16% in the B+H group reached the primary end point, representing an absolute risk reduction of 3% and a hazard reduction of 18%. The difference in event rates of the composite primary end point between the B+A and B+H groups was significant (HR, 0.80; 95% CI, 0.69 to 0.92; P=0.0016).  Among the patients without CAD, fewer patients in the B+A treatment arm (204 of 3,096) reached the primary end point compared with those in the B+H arm (251 of 3,095). The difference in event rates between the B+A and B+H groups was significant (HR, 0.81; 95% CI, 0.67 to 0.98; P=0.026).  A comparison of patients with and without CAD event rates for the primary end points demonstrated that the patients with CAD had a greater CV event rate than those without CAD (15 vs 7%; P&lt;0.0001).  Secondary: The composite secondary end point of CV mortality, MI, and stroke occurred in 5.74% in the B+A group and 8% in the B+H group, resulting in an absolute risk reduction of 1.95% and a hazard reduction of 25% (HR, 0.73; 95% CI, 0.59 to 0.9; P=0.033). The rate of all-cause mortality differed significantly between the treatment arms (HR, 0.77; 95% CI, 0.6 to 0.99; P=0.042). Among the patients without CAD, the rates of CV mortality, MI, and stroke did not differ between the two arms (HR, 0.86; 95% CI, 0.68 to 1.08). The secondary end point events were lower in the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				group of patients without CAD.
<p>Hansson et al.<sup>56</sup> (1999) STOP-Hypertension</p> <p>Felodipine 2.5 mg or isradipine 2.5 mg QD</p> <p>vs</p> <p>enalapril 10 mg or lisinopril 10 mg QD</p> <p>vs</p> <p>atenolol 50 mg or metoprolol 100 mg or pindolol 5 mg QD and/or HCTZ 25 mg with amiloride 2 to 5 mg QD</p>	<p>MC, OL, PRO, RCT</p> <p>Men and women, age 70 to 84 years with HTN (SBP <math>\geq</math>180 mm Hg or DBP <math>\geq</math>105 mm Hg or both)</p>	<p>N=6,614</p> <p>4 years</p>	<p>Primary: Fatal stroke, fatal MI, other fatal cardiovascular events</p> <p>Secondary: Blood pressure</p>	<p>Primary: The rate of prevention of cardiovascular deaths was similar in all groups (RR, 0.97 to 1.14; 95% CI, 0.86 to 1.26).</p> <p>Fatal cardiovascular events, including fatal stroke and fatal myocardial infarction MI, occurred in 19.8 per 1,000 patient-years in the <math>\beta</math>-blocker and/or HCTZ group, in the felodipine or isradipine group and in the enalapril or lisinopril group (RR, 0.99; 95% CI, 0.84 to 1.16).</p> <p>The RR of cardiovascular death in patients in the enalapril or lisinopril group as compared to the felodipine or isradipine group was 1.14 (95% CI, 0.86 to 1.26; P=0.67).</p> <p>Secondary: Decreases in blood pressure were similar among the groups.</p>
<p>Borhani et al.<sup>57</sup> (1996) MIDAS</p> <p>Isradipine 2.5 to 5 mg BID</p> <p>vs</p> <p>HCTZ 12.5 to 25 mg QD</p>	<p>DB, MC, positive-control, RCT</p> <p>Patients, average of 58.5 years old, with HTN</p>	<p>N=883</p> <p>3 years</p>	<p>Primary: Rate of progression of intimal-medial thickness in carotid arteries</p> <p>Secondary: Rate of cardiovascular events (MI, stroke, CHF, angina, sudden death), rate of non-major cardiovascular events and</p>	<p>Primary: There was no difference in the rate of progression of intimal-medial thickness between the treatment groups (P=0.68).</p> <p>Secondary: The rate of cardiovascular events was greater in the isradipine group than in the HCTZ group (5.65 vs 3.17%; P=0.07).</p> <p>The rate of non-major cardiovascular events was greater in the isradipine group than in the HCTZ group (9.05 vs 5.22%; P=0.02).</p> <p>There was a significant decrease in SBP in the HCTZ group as compared to isradipine (-19.5 vs -16.0 mm Hg; P=0.002).</p> <p>There was no difference in change in DBP (both groups, -13.0 mm Hg).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>National Intervention Cooperative Study<sup>58</sup> (1999) NICS-EH  Nicardipine SR 20 mg BID  vs  trichlor-methiazide* 2 mg QD</p>	<p>DB, RCT  Patients age 60 years old and older with a SBP between 160 to 220 mm Hg and a DBP &lt;115 mm Hg and no history of cardiovascular complications</p>	<p>N=414  5 years</p>	<p>procedures (TIAs, dysrhythmia, aortic valve replacement, femoral popliteal bypass graft), blood pressure  Primary: Cardiovascular complications  Secondary: Blood pressure, pulse, side effects, laboratory values</p>	<p>Primary: There was no difference in rate of cardiovascular complications during the study period (P=0.923).  There was no difference in the number of patients experiences left ventricular hypertrophy on ECG (P=0.975).  Secondary: Both groups experienced significant reductions in blood pressure from baseline (P=0.000).  There was no significant difference in pulse rate between the groups.  Side-effect rates did not differ between the groups (P=0.897).  More patients in the trichlormethiazide group than in the nicardipine group had abnormal lab results at the end of the study; differences were significant for serum sodium levels (decreased in the trichlormethiazide group) and uric acid levels (increased with trichlormethiazide).</p>
<p>Lichtlen et al.<sup>59</sup> (1990) INTACT  Nifedipine 80 mg QD  vs  placebo</p>	<p>DB, MC, PC, RCT  Patients, age 65 years and younger, demonstrating early CAD who were not candidates for invasive therapeutic procedures</p>	<p>N=348  3 years</p>	<p>Primary: Progression of coronary artery disease detected on angiogram (change in minimal diameter, percent stenosis, transition into occlusion, new stenosis)  Secondary:</p>	<p>Primary: In patients without study deviations, there were no significant differences in number of stenoses and occlusions per patient (nifedipine=3.7, placebo=3.88; P=0.437). The distribution among the arteries of the occlusions was not different between groups.  The progression of stenosis was significant from baseline but changes were not significantly different between the groups (P&lt;0.006 for all vs baseline; P&gt;0.585 for group comparisons).  Secondary: There was no difference between nifedipine treatment and placebo in</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Critical clinical events (cardiac death, nonfatal MI, unstable angina, need for procedure, heart failure, severe arrhythmias), progression of new lesions	number of critical events, 44 events in 24 patients receiving nifedipine vs 52 events in 35 patients in the placebo group (P=0.278).  The nifedipine group had significantly fewer new lesions as compared to the placebo group: 78 (0.58 lesions/patients) vs 118 (0.8 lesions/patient) (P=0.031).
Brown et al. <sup>60</sup> (2000) INSIGHT  Nifedipine 30 mg QD  vs  amiloride and HCTZ 2.5-25 mg QD (fixed-dose combination product)  Doses were doubled or atenolol 25 to 50 mg or enalapril 5 to 10 mg was added.	DB, MC, PRO, RCT  Patients, age 55 to 80 years old with HTN (blood pressure $\geq$ 150/95 mm Hg or SBP $\geq$ 160 mm Hg) and $\geq$ 1 cardiovascular risk factor	N=6,575  3 years	Primary: Composite death from any cardiovascular cause together with nonfatal stroke, MI, or heart failure  Secondary: Total mortality, death from a vascular cause, nonfatal vascular event	Primary: There was no difference in composite cardiovascular deaths between the groups. Events occurred in 200 (6.3%) patients in the nifedipine group and 182 (5.8%) of the amiloride and HCTZ group (18.2 vs 16.5 events per 1,000 patient-years; P=0.34).  Secondary: There was no difference in all-cause mortality (P=0.62), death from a vascular cause (P=0.67) and in nonfatal vascular events (P=0.50) between the treatment groups.
Estacio et al. <sup>61</sup> (1998) ABCD  Nisoldipine 10 to 60 mg/day  vs	DB, PRO, RCT  Patients between the ages of 40 and 74 years with NIDDM, baseline DBP $\geq$ 90 mm Hg and receiving no	N=470  67 months	Primary: Effect of intensive (target DBP of 75 mm Hg) or moderate (target DBP between 80 to 89 mm Hg) blood pressure control on	Primary: Analysis of the 470 patients in the trial who had HTN (DBP $\geq$ 90 mm Hg) showed similar control of blood pressure, blood glucose and lipid concentrations between the two study medications throughout the five years of follow-up.  Secondary: Nisoldipine was associated with a higher incidence of fatal and nonfatal



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
enalapril 5 to 40 mg/day	antihypertensive medications at the time of randomization		the incidence and progression of complications of diabetes; compare enalapril to nisoldipine as a first-line antihypertensive agent  Secondary: Incidence of MI	MI than enalapril (RR, 7.0; 95% CI, 2.3 to 21.4).
<b>Hypertension</b>				
Sheehy et al. <sup>62</sup> (2000)  Amlodipine 2.5 to 10 mg QD  vs  felodipine 2.5 to 10 mg QD	RETRO  Patients, age 65 years and older, with HTN	N=7,818  Duration not reported	Primary: Prescription renewal, drug switch rates, compliance rates, office visits  Secondary: Not reported	Primary: Patients prescribed amlodipine had a greater compliance rate, 67.9%, than those prescribed felodipine 66.2% (P<0.01).  Discontinuation rates were higher in the felodipine group by 27%.  Amlodipine treatment resulted in more continuous months of treatment (69.2), than felodipine treatment (57.8) (P<0.01).  Renewal rates were significantly larger in the amlodipine group (89.0%), than the felodipine group (85.6%) (P<0.01).  Switch rates were significantly larger, 5 times, in the felodipine group (10.2%) than the amlodipine group (1.9%) (P<0.01).  Visits to specialists occurred significantly more in patients treated with amlodipine than felodipine, (OR, 1.14; 95% CI, 1.8 to 1.20).  Secondary: Not reported
Van der Krogt et al. <sup>63</sup> (1996)	DB, MC, PG, RCT  Patients, age 18 to 75 years old, with	N=201  12 weeks	Primary: Number of responders (DBP ≤90 mm Hg after	Primary: Amlodipine treatment resulted in significantly more responders than felodipine treatment (P=0.046): 68% (69 of 101) of the amlodipine group were responders.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>felodipine ER 5 to 10 mg QD</p>	<p>mild to moderate HTN (DBP <math>\geq</math>95 mm Hg and <math>\leq</math>114 mm Hg)</p>		<p>12 weeks of monotherapy or decrease of <math>&gt;</math>10 mm Hg if baseline DBP <math>&gt;</math>100 mm Hg) who did not experience serious adverse events</p> <p>Secondary: Blood pressure, adverse events</p>	<p>53% (49 of 92) of the felodipine group were responders. 32% (32 of 101) of the amlodipine group were not responders. 47% (43 of 92) of the felodipine group were not responders.</p> <p>Secondary: The decreases in SBP and DBP from baseline were significant within each group, but were similar between the groups (amlodipine SBP and DBP 12 weeks vs baseline; <math>P&lt;</math>0.001, felodipine SBP and DBP 12 weeks vs baseline; <math>P&lt;</math>0.001, amlodipine 12 week change vs felodipine 12 week change; <math>P&gt;</math>0.05).</p> <p>Adverse events were experienced by 33% of the amlodipine group and 42% of the felodipine group.</p> <p>Significantly more patients in the felodipine group experienced serious adverse events (9 patients who experienced 17 serious events vs two patients who experienced three serious events; <math>P=</math>0.048).</p>
<p>Mounier-Vehier et al.<sup>64</sup> (2002)</p> <p>Amlodipine 5 mg QD</p> <p>vs</p> <p>nicardipine 60 mg/day, divided 2 to 3 times daily</p>	<p>DB, MC, PG RCT</p> <p>Men and women, age 60 years and older with isolated systolic HTN (SBP 160 to 208 mm Hg) and DBP <math>&lt;</math>90 mm Hg</p>	<p>N=133</p> <p>90 days</p>	<p>Primary: Mean difference in SBP from baseline to day 90</p> <p>Secondary: Mean difference in DBP, pulse pressure, heart rate, percent of patients with normal blood pressure (<math>&lt;</math>140/90 mm Hg), safety</p>	<p>Primary: The decrease in SBP from baseline was significant within each group, but were similar between the groups (amlodipine day 90 vs baseline; <math>P=</math>0.0001, nicardipine day 90 vs baseline; <math>P=</math>0.0001, amlodipine 90 day change vs nicardipine 90 day change; <math>P=</math>0.38).</p> <p>Secondary: The decrease in DBP from baseline was significant within each group but similar between the groups (amlodipine day 90 vs baseline; <math>P=</math>0.0001; nicardipine day 90 vs baseline; <math>P=</math>0.0003, amlodipine 90 day change vs nicardipine 90 day change; <math>P=</math>0.12).</p> <p>The decrease in pulse pressure from baseline was significant within each group but similar between the groups (amlodipine day 90 vs baseline, <math>P=</math>0.0001; nicardipine day 90 vs baseline, <math>P=</math>0.0001; amlodipine 90 day change vs nicardipine 90 day change, <math>P=</math>0.88). There was no difference between the groups in heart rate (<math>P=</math>0.60).</p> <p>At day 90, 25.9, and 23.4% of the amlodipine and nicardipine groups had achieved normal blood pressure (<math>P=</math>0.76).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				The numbers of people in each group reporting at least 1 adverse event were similar, 23 in the amlodipine group and 20 in the nifedipine group.
Kes et al. <sup>65</sup> (2003)  Amlodipine 5 to 10 mg QD  vs  nifedipine 30 to 60 mg QD	MC, OL, RCT  Patients with HTN	N=155  12 weeks	Primary: Change in DBP  Secondary: Not reported	Primary: There was no significant difference in DBP between the amlodipine group and nifedipine group at 12 weeks (P=0.436).  Secondary: Not reported
Ryuzaki et al. <sup>66</sup> (2007) i-TECHO  Amlodipine 2.5 to 10 mg QD  vs  nifedipine CR 20 to 80 mg QD	OL, RCT, XO  Patients treated for HTN (SBP >140 mm Hg or DBP >90 mm Hg)	N=55  12 weeks (6 weeks per treatment)	Primary: Average home blood pressure readings, pulse rates, clinic blood pressure and pulse readings  Secondary: Not reported	Primary: The morning home SBP and DBP readings were lower in the nifedipine group than the amlodipine group (SBP 131±8 vs 133±10 mm Hg; P<0.05, DBP 80±8 vs 81±8 mm Hg; P<0.05).  There were no significant differences in evening home blood pressure readings (P>0.05).  There was no significant difference in rates of achieving target blood pressure between the groups (P<0.05).  Morning home pulse rates were greater in the nifedipine group than the amlodipine group (70±9 vs 69±9 beats/min; P<0.05).  There were no significant differences between the groups in evening home pulse rates (P>0.05).  The clinic SBP and DBP readings were significantly lower in the nifedipine group than in the amlodipine group (P<0.05).  There were no significant differences between the groups in clinic pulse rates (P>0.05).  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
Saito et al. <sup>67</sup> (2007) ADVANCE-Combi  Amlodipine 2.5 to 5 mg QD  vs  nifedipine CR 20 to 40 mg QD  Valsartan 40 to 80 mg was added on if blood pressure goal not met.	DB, RCT  Patients with untreated essential HTN with sitting SBP $\geq$ 160 mm Hg or DBP $\geq$ 100 mm Hg; or previously treated with sitting SBP $\geq$ 150 mm Hg or DBP $\geq$ 95 mm Hg	N=514  16 weeks	Primary: Target blood pressure, achievement rate  Secondary: Safety	Primary: Target blood pressure achievement rates were higher for the nifedipine treatment group than the amlodipine group (P<0.001).  Patients in the amlodipine group were more likely to require additional treatment with valsartan or a dose increase of amlodipine (P<0.05).  The reduction in blood pressure from baseline was greater in the nifedipine group (-34.0/-20.1) than in the amlodipine group (-27.0/-15.9; P<0.05).  Secondary: Adverse event rates were not significantly different between the groups, 12.4% in the nifedipine group vs 7.6% of the amlodipine group (P=0.07).
Pepine et al. <sup>68</sup> (2003) CESNA-II  Amlodipine 5 to 10 mg QD  vs  nisoldipine ER 20 to 40 mg QD	DB, DD, PG, MC, RCT  Men and women with HTN (DBP 90-109 mm Hg) and CAD	N=not specified  6 weeks	Primary: Change from baseline in DBP at 6 weeks  Secondary: Exercise duration, antihypertensive responder rate (% of patients with DBP <90 mm Hg), exercise test responder rate (increase in time by 20% and 60 seconds)	Primary: At six weeks, the mean SBP and mean DBP for the two treatment groups were not significantly different from each other and mean reductions in blood pressure were similar: amlodipine SBP/DBP 138/83 mm Hg, a decrease of 13/11 mm Hg, vs nisoldipine 137/81 mm Hg, a decrease of 15/13 mm Hg (P values not significant).  Secondary: Both treatment groups experienced increases in exercise duration, increased by 21 seconds in the amlodipine group and 23 seconds in the nisoldipine group (P=0.268).  Antihypertensive and exercise responder rates were similar between the groups (antihypertensive rates: 78% for amlodipine and 87% for nisoldipine; P>0.05 for both).
Whitcomb et al. <sup>69</sup> (2000)  Amlodipine 2.5 to	DB, DD, MC, RCT  Men and women, age 21 to 75 years,	N=161  8 weeks	Primary: Between treatment comparison of change from	Primary: Treatment with amlodipine resulted in a significantly larger change from baseline in DBP (between-group difference 2.7 mm Hg; P=0.005). However, a pre-specified difference of greater than 5 mm Hg in least mean

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
10 mg QD  vs  nisoldipine ER 10 to 40 mg QD	with HTN		baseline in DBP  Secondary: Change from baseline in SBP, heart rate, percent of patients who responded	squares, here 1.1 to 4.3 mm Hg, showed that the treatments were similar in reduction of DBP.  Secondary: Amlodipine treatment resulted in a significantly larger change from baseline in SBP than nisoldipine treatment (P value not reported, least mean square difference >5 mm Hg).  At week eight, more patients in the amlodipine group were responders, 79%, as compared to the nisoldipine group, 60% (P=0.004).
White et al. <sup>70</sup> (2003) CESNA-III  Amlodipine 5 to 10 mg QD  vs  nisoldipine ER 20 to 60 mg QD	DB, MC, PRO, RCT  African American patients with HTN (blood pressure of 92 mm Hg to 114 mm Hg and SBP <200 mm Hg)	N=192  12 weeks	Primary: ABPM change from baseline in DBP in mean 24 hour period  Secondary: ABPM change in SBP, awake and asleep blood pressure, changes in clinic blood pressure and pulse	Primary: The decrease from baseline in DBP was similar between the groups: -16.0±2.3 mm Hg for nisoldipine and -15.0±2.3 mm Hg for amlodipine (P=0.500).  Secondary: The decrease from baseline in SBP was similar between the groups: -23.0±2.7 mm Hg for nisoldipine and -19.9±2.7 mm Hg for amlodipine (P=0.067).  The changes from baseline in awake and asleep SBP and DBP were not significantly different between the groups except for awake SBP, for which the nisoldipine group had a larger reduction, -19.2 vs -15.9 mm Hg (P=0.045).  The changes from baseline in clinic blood pressure and pulse were similar between the groups (P>0.05 for SBP and DBP; P=0.362).
Lenz et al. <sup>71</sup> (2001)  Amlodipine 5 to 10 mg QD  vs  nisoldipine 10 to 20 mg QD	OL, XO  Patients, 35 to 70 years old, with HTN, (SBP 140 to 179 mm Hg and DBP 90 to 109 mm Hg), stable on amlodipine for ≥3 months prior to	N=21  10 weeks	Primary: 24-hr ABPM  Secondary: Not reported	Primary: No significant difference in ABPM was found after patients switched from amlodipine to nisoldipine for the following: systolic nighttime, daytime and 24-hr blood pressure, diastolic nighttime and daytime blood pressure (P>0.05 for all).  24-hr DBP was significantly lower with amlodipine treatment than with nisoldipine treatment (75±10 vs 77±8.5 mm Hg; P=0.017).  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	switch to nisoldipine			Not reported
Drummond et al. <sup>72</sup> (2007)  Amlodipine 5 mg QD  vs  amlodipine 10 mg QD  vs  aliskiren and amlodipine 150-5 mg QD (fixed-dose combination product)  Patients not responding to amlodipine 5 mg QD at the end of 4 week single-blind run-in period received combination therapy, continuation of amlodipine 5 mg QD or titration to amlodipine 10 mg QD.	AC, DB, MC, PG, RCT  Patients 18 years of age and older with mild to moderate HTN	N=545  6 weeks	Primary: Change in DBP at 6 weeks  Secondary: SBP, comparison of SBP and DBP reductions between combination therapy group and amlodipine 10 mg group, proportion of patients responding to treatment, and proportion of patients achieving blood pressure control	Primary: DBP reduction was significantly greater in the combination therapy group compared to those in the amlodipine 5 mg group (P<0.0001).  Secondary: SBP reduction was significantly greater in the combination therapy group compared to those in the amlodipine 5 mg group (P<0.0001).  No significant differences were observed in DBP or SBP reduction between the combination therapy group and the amlodipine 10 mg group (P=0.6167 and P=0.2666 respectively).  The proportion of patients responding to treatment was significantly higher in the combination therapy group compared to the amlodipine 5 mg group (P<0.0001). No significant difference was observed between the combination therapy group and the amlodipine 10 mg group (P value not reported).  The proportion of patients achieving blood pressure control was significantly higher in the combination therapy group compared to the amlodipine 5 mg group (P<0.0001). No significant difference was observed between the combination therapy group and the amlodipine 10 mg group (P=0.5229).
Benetos et al. <sup>73</sup> (2000)	DB, MC, PG, RCT	N=164	Primary: Changes in blood	Primary: Both bisoprolol and HCTZ and amlodipine significantly reduced SBP

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Amlodipine 5 mg QD</p> <p>vs</p> <p>bisoprolol and HCTZ 2.5-6.25 mg QD (fixed-dose combination product)</p>	<p>Patients over 60 years with supine SBP 160 to 210 mm Hg and DBP &lt;90 mm Hg</p>	<p>12 weeks</p>	<p>pressure, heart rate, adverse events, QOL scores</p> <p>Secondary: Not reported</p>	<p>(-20.0±13.7 and -19.6±14.2 mm Hg, respectively; P&lt;0.001) and DBP (-4.5±7.4 and -2.4±8.4 mm Hg, respectively from baseline to week 12, but there was not a significant difference between the agents (SBP; P=0.85 and DBP; P=0.09).</p> <p>Bisoprolol and HCTZ significantly reduced heart rate from baseline, but amlodipine did not (-7.6±8.4 [P&lt;0.001] and -0.2±11.4 bpm, respectively).</p> <p>Bisoprolol and HCTZ significantly reduced heart rate when compared to amlodipine (P=0.0001).</p> <p>Overall adverse events were not significantly different between the amlodipine and the bisoprolol and HCTZ group (39 and 40%, respectively). Adverse events reported included headache, leg edema, fatigue and bradycardia but severity of events was not reported.</p> <p>Overall QOL scores were not significantly different between the amlodipine and the bisoprolol and HCTZ group.</p> <p>Secondary: Not reported</p>
<p>Prisant et al.<sup>74</sup> (1995)</p> <p>Amlodipine 2.5, 5, or 10 mg</p> <p>vs</p> <p>bisoprolol and HCTZ 2.5-6.25, 5-6.25, or 10-6.25 mg/day (fixed-dose combination product)</p> <p>vs</p>	<p>DB, MC, PG, RCT</p> <p>Patients ≥21 years with mild to moderate essential HTN, (average sitting DBP 95 to 114 mm Hg) each treatment was once daily and titrated to effect</p>	<p>N=218</p> <p>17 weeks</p>	<p>Primary: Mean change from baseline in SBP and DBP, lab measurements, adverse events, QOL questionnaire</p> <p>Secondary: Not reported</p>	<p>Primary: Mean decreases in SBP and DBP from baseline were 13.4/10.7 mm Hg for bisoprolol and HCTZ patients, 12.8/10.2 mm Hg for amlodipine patients, and 7.3/6.6 mm Hg for enalapril patients. The hypotensive effects were significant for all three groups (P&lt;0.001).</p> <p>SBP and DBP mean changes from baseline for the bisoprolol and HCTZ group and the amlodipine group were greater than the change from baseline for the enalapril group (P&lt;0.01).</p> <p>Response rates (DBP ≤90 mm Hg or ≥10 mm Hg decrease from baseline) were 71% for the bisoprolol and HCTZ group, 69% for the amlodipine group, and 45% for the enalapril group. The response rates for the bisoprolol and HCTZ and the amlodipine groups differed significantly from the enalapril group (P&lt;0.01).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
enalapril 5, 10, or 20 mg				<p>Twenty nine percent of bisoprolol patients had adverse experiences compared to 42% of amlodipine patients (P=0.12). Nearly 47% of enalapril patients had adverse experience compared to bisoprolol (P=0.04). Adverse events reported included headache, fatigue, peripheral edema, and dizziness.</p> <p>Drug related adverse events were 16% for the bisoprolol and HCTZ patients, 21% for the amlodipine patients, and 23% for the enalapril patients. There was no significant difference between the groups.</p> <p>Enalapril demonstrated a mean decrease from baseline of 7.9 mg/dL for TC (P=0.02 vs amlodipine) and 6.6 mg/dL for LDL-C (P=0.04 vs amlodipine) which were not significantly different from the increase from the bisoprolol and HCTZ group of 1.7 mg/dL (P=0.07 vs enalapril) for TC and +0.6 mg/dL in LDL-C. However, the increase in TGs was highest for bisoprolol and HCTZ-treated patients compared to amlodipine- and enalapril-treated patients (P=0.08, for bisoprolol and HCTZ vs enalapril).</p> <p>There was not a significant difference from baseline or between treatment groups in QOL scores: 0.9 for the bisoprolol and HCTZ group, 0.5 for the amlodipine group, and 2.3 for the enalapril group.</p>
<p>Mazza et al.<sup>75</sup> (2002)</p> <p>Amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>nebivolol 2.5 to 5 mg QD</p>	<p>DB, MC, PG, RCT</p> <p>Patients between 65 to 89 years of age with mild to moderate essential HTN and DBP ranging from 95 to 114 mm Hg</p>	<p>N=168</p> <p>16 weeks</p>	<p>Primary: Change in sitting blood pressure, response rates</p> <p>Secondary: Standing blood pressure changes, standing and sitting heart rate changes</p>	<p>Primary: There was not a significant difference observed between the amlodipine and nebivolol treatments groups in changes in sitting DBP (blood pressure values and P values not reported). At weeks four and eight, a slightly lower sitting SBP was observed in per-protocol patients in the amlodipine groups vs those in the nebivolol group (blood pressure values not reported, P&lt;0.005).</p> <p>Response rates were not significantly difference between the amlodipine group and the nebivolol group (86 vs 88%, respectively). The percentage of patients who reached normalization (blood pressure &lt;140/90 mm Hg) was no significant between the amlodipine and the nebivolol groups (47 vs 50%).</p> <p>Secondary: There were significant differences in standing blood pressure observed</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>between the groups.</p> <p>Heart rate was significantly lower in the nebivolol group compared to the amlodipine group at all treatment visits (P&lt;0.001).</p> <p>Patients in the amlodipine group experienced a significantly greater rate of headache (seven vs five patients) and ankle edema (12 vs zero patients) compared to the patients in the nebivolol group (P&lt;0.05 for both).</p>
<p>Hollenberg et al.<sup>76</sup> (2003)</p> <p>Amlodipine 2.5 mg/day</p> <p>vs</p> <p>eplerenone 50 mg/day</p> <p>Both medications were titrated to a maximum of 200 (eplerenone) or 10 (amlodipine) mg/day to achieve a SBP&lt;140 mm Hg.</p>	<p>RCT</p> <p>Patients ≥50 years of age, with untreated SBP between 140 to 190 mm Hg</p>	<p>N=269</p> <p>24 weeks</p>	<p>Primary: Change in SBP and DBP, discontinuation rate, symptom distress index, SF-36 Health Survey</p> <p>Secondary: Not reported</p>	<p>Primary: Both treatments exhibited similar reductions in SBP and DBP from baseline (P=0.01).</p> <p>The dropout rate was 50% greater in amlodipine-treated patients compared to eplerenone-treated patients (P value not reported).</p> <p>Symptom distress (technique used to assess the influence of drug treatment on QOL) index was assessed and results favored eplerenone therapy (P=0.03).</p> <p>SF-36 Health Survey showed no significant difference between the two treatments (P value not reported).</p> <p>Both treatments experienced similar incidences of adverse effects (P value not reported). Eplerenone-treated patients did not experience breast pain/tenderness, breast enlargement, changes in menstruation, gynecomastia or loss of libido.</p> <p>Secondary: Not reported</p>
<p>White et al.<sup>77</sup> (2003)</p> <p>Amlodipine 2.5 mg/day</p> <p>vs</p>	<p>AC, DB, MC, RCT</p> <p>Patients ≥50 years of age with systolic HTN (seated clinic SBP 150 to 165 mm Hg with a pulse pressure ≥70 mm</p>	<p>N=269</p> <p>24 weeks</p>	<p>Primary: Mean change from baseline in SBP, DBP, 24 hour ambulatory BP, pulse pressure, and heart rate at week 24; urine albumin/</p>	<p>Primary: Mean reduction in SBP from baseline was comparable in eplerenone- and amlodipine-treated patients (P=0.83).</p> <p>Eplerenone-treated patients exhibited significant reductions in DBP from baseline at 24 weeks of therapy compared to amlodipine-treated patients (P=0.014).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>eplerenone 50 mg/day</p> <p>Both medications were titrated to a maximum of 200 (eplerenone) or 10 (amlodipine) mg/day to achieve a SBP&lt;140 mm Hg.</p>	<p>Hg or 165 to 200 mm Hg with a DBP ≤95 mm Hg)</p>		<p>creatinine ratio; adverse events</p> <p>Secondary: Not reported</p>	<p>The two treatments exhibited comparable decreases in 24 hour ambulatory BP, pulse pressure and heart rate after 24 weeks of therapy (P&gt;0.05).</p> <p>Eplerenone-treated patients exhibited a significant reduction from baseline in the urine albumin/creatinine ratio compared to amlodipine-treated patients (P=0.002).</p> <p>Treatment-emergent adverse events were reported in 64 and 70% of eplerenone- and amlodipine-treated patients. The only adverse event that was significant between the two treatments was the incidence of edema (3.7 vs 25.5%; P&lt;0.05). There were no reports of gynecomastia, breast tenderness or menstrual irregularities with either treatment.</p> <p>Secondary: Not reported</p>
<p>Jordan et al.<sup>78</sup> (2007)</p> <p>Amlodipine 5 to 10 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>aliskiren 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>irbesartan 150 to 300 mg QD, added to existing HCTZ</p>	<p>DB, DD, MC, PG, RCT</p> <p>Obese men and women (BMI ≥30 kg/m<sup>2</sup>) ≥18 years with essential HTN (mean sitting DBP 95 to 109 mm Hg and SBP &lt;180 mm Hg) who had not responded to 4 weeks of treatment with HCTZ 25 mg</p>	<p>N=489</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting DBP with aliskiren 300 mg plus HCTZ vs HCTZ alone at 8 weeks</p> <p>Secondary: Comparisons of mean sitting DBP and SBP with aliskiren plus HCTZ vs the other treatment groups, percentage of responders (mean sitting DBP &lt;90 mm Hg or ≥10 mm Hg reduction from baseline), proportion of</p>	<p>Primary: Aliskiren 300 mg added to HCTZ 25 mg significantly reduced mean sitting DBP compared to HCTZ alone at week eight (mean difference, -4.0; P&lt;0.0001).</p> <p>Secondary: Aliskiren 300 mg added to HCTZ caused numerically larger reductions in mean sitting DBP and SBP compared to amlodipine 10 mg plus HCTZ and irbesartan 300 mg plus HCTZ at week eight, but there were no statistically significant differences between treatment groups (P&gt;0.05).</p> <p>Responder rates were significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0193) and week 12 (P=0.004) but comparable to responder rates observed with amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>The proportion of patients achieving blood pressure control was significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0005) and week 12 (P=0.0001) but not statistically different than amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>Plasma renin activity significantly increased (P&lt;0.05) during four weeks</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
therapy (single entity products)  vs  HCTZ 25 mg QD (existing therapy)			patients achieving blood pressure control (mean sitting blood pressure <140/90 mm Hg), plasma renin activity, safety and tolerability	of HCTZ monotherapy. Combination with aliskiren neutralized this increase and led to an overall significant reduction in plasma renin activity compared to pretreatment baseline (P<0.05) whereas amlodipine and irbesartan led to further significant increases (P<0.05).  All of the study treatments were generally well tolerated. Amlodipine plus HCTZ (45.2%) was associated with a higher incidence of adverse events than the other treatment groups (36.1 to 39.3%), largely due to a higher rate of peripheral edema (11.1 vs 0.8 to 1.6%).
Messerli et al. <sup>79</sup> (2002)  Amlodipine and benazepril 5-10 mg to 5-20 mg QD (fixed-dose combination product)	OL  Patients ≥18 years with mild-to-moderate HTN taking amlodipine 5 to 10 mg with inadequate blood pressure (DBP ≥90 mm Hg, Group 1) or intolerance with amlodipine (DBP ≤90 mm Hg with edema, Group 2)	N=7,912  4 weeks	Primary: Change in mean sitting DBP (group 1), and percentage of patients whose edema improved (group 2)  Secondary: Group 1-change in mean sitting SBP	Primary: In Group 1, mean reduction in DBP at week four was 11.5 mm Hg (95% CI, -11.8 to -11.3 mm Hg; P<0.001). Mean DBP declined from 96.5 (baseline) to 84.9 mm Hg (at 4 weeks).  In Group 2, 85% of patients saw improvement in edema with 42% of patients experiencing complete resolution after receiving combination therapy (95% CI, 83 to 87).  Secondary: In Group 1, mean reduction in SBP at week four was 15.6 mm Hg (95% CI, -16.0 to -15.2 mm Hg; P<0.001).
Chrysant et al. <sup>80</sup> (2012)  Study 1: Benazepril 40 mg/day (Group 1)  vs  amlodipine and benazepril 5-40 mg/day, up titrated to 10-40 mg/day after 4 weeks.	Post-hoc analysis of 2 trials  Patients with HTN	N=1,013  14 weeks	Primary: Change in baseline mean sitting DBP and mean sitting SBP, rate of blood pressure control (<140/90 mm Hg), rate of blood pressure control (mean sitting DBP <90 mm Hg or ≥10 mm Hg decrease from baseline)	Primary: Pooled results demonstrate that combination therapy resulted in significantly greater lowering of mean sitting DBP and mean seated SBP compared to benazepril or amlodipine (P<0.001). Amlodipine and benazepril 10-20 mg/day resulted in significantly greater blood pressure reductions in White patients (mean sitting DBP: 12.99 mm Hg; mean sitting SBP: 13.72 mm Hg) compared to Black patients (8.80 and 8.72 mm Hg) (P<0.004). Amlodipine and benazepril 10-40 mg/day resulted in similar reductions in blood pressure in both White and Black patients.  The proportion of patients who achieved blood pressure control with amlodipine and benazepril 10-40 mg/day was similar between White and Black patients (60.7%), whereas with amlodipine and benazepril 10-20 mg/day the rate of control was higher with White patients (61.2 vs 39.4%);

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(fixed-dose combination product) (Group 2)</p> <p>Study 2: Amlodipine and benazepril 10-20 mg/day, uptitrated to 10-40 mg/day after 2 weeks (Group 3)</p> <p>vs</p> <p>amlodipine and benazepril 10-20 mg/day (fixed-dose combination product) (Group 4)</p> <p>vs</p> <p>amlodipine 10 mg/day (Group 5)</p>			<p>Secondary: Safety</p>	<p>P&lt;0.023).</p> <p>There was no difference in the proportion of patients who responded to treatment between Black and White patients with amlodipine and benazepril 10-40 mg/day (74.8 vs 77%; P&lt;0.639). The proportion of patients who responded to amlodipine and benazepril 10-20 mg/day was significantly lower in Black patients (50.7 vs 73.5%; P&lt;0.007).</p> <p>Secondary: There were no serious clinical or metabolic side effects reported, with the exception of pedal edema which occurred more frequently with amlodipine monotherapy.</p>
<p>Messerli et al.<sup>81</sup> (2000)</p> <p><u>Study 1:</u> Amlodipine and benazepril 5-10 mg to 5-20 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>nifedipine 30 to 60</p>	<p>2 DB, MC, RCT</p> <p>Patients 18 to 80 years of age with uncomplicated essential HTN</p>	<p>N=1,079</p> <p>8 weeks</p>	<p>Primary: Change in DBP from baseline</p> <p>Secondary: Change from baseline in SBP and heart rate</p>	<p>Primary: Study 1 Significant reductions in DBP were observed with benazepril and amlodipine 10-5 and 20-5 mg (-9.4 and -9.7 mm Hg, respectively) compared to nifedipine 30 mg (-7.0 mm Hg; P&lt;0.05), but not nifedipine 60 mg (-8.5; P&gt;0.05).</p> <p>Study 2 Benazepril and amlodipine 10-5 (-8.9 mm Hg) and 20-5 mg (-9.1 mm Hg) produced significantly greater reductions in DBP than amlodipine 5 mg (-6.8 mm Hg; P&lt;0.05), but not amlodipine 10 mg (-8.7 mm Hg; P&gt;0.05).</p> <p>Secondary: Study 1</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg/day</p> <p><u>Study 2:</u> Amlodipine and benazepril 5-10 mg to 5-20 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>				<p>Significant reductions in SBP were observed with benazepril and amlodipine 20-5 mg (-11.6 mm Hg) compared to nifedipine 30 mg (-7.9 mm Hg; P&lt;0.05).</p> <p>Significantly less edema was reported with combination therapies (3.1 to 3.8%; P≤0.001) compared to nifedipine 60 mg (15.5%; P=0.008) but not nifedipine 30 mg (5.4%).</p> <p>Study 2 Significant reductions in SBP were observed with benazepril and amlodipine 20-5 mg (-9.1 mm Hg) compared to amlodipine 5 mg (-5.3 mm Hg; P&lt;0.05). There were no significant difference in SBP between amlodipine 10 mg and the combination therapies.</p> <p>Significantly less edema (P&lt;0.001) was reported with amlodipine 5 mg (4.9%) and combination therapies (1.5 to 2.2%) compared to amlodipine 10 mg (23.6%).</p>
<p>Jamerson et al.<sup>82</sup> (2004)</p> <p>Amlodipine and benazepril 5-20 and 10-20 mg/day (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 to 10 mg/day</p>	<p>DB, MC, PG, RCT</p> <p>Men and women 18 to 80 years of age with stage 2 HTN</p>	<p>N=364</p> <p>12 weeks</p>	<p>Primary: Percentage of patients with SBP reduction ≥25 mm Hg (if baseline &lt;180 mm Hg) or ≥32 mm Hg (if baseline ≥180 mm Hg)</p> <p>Secondary: Percentage of patients with DBP reduction ≥15 mm Hg (if baseline &lt;110 mm Hg) or ≥20 mm Hg (if baseline ≥110 mm Hg), percentage of patients meeting</p>	<p>Primary: Significantly more patients on combination therapy (74.2%) met the primary end point than patients on amlodipine monotherapy (53.9%; P&lt;0.0001). The time by which 50% of patients attained the primary end point was four weeks shorter among patients randomized to combination therapy compared to those randomized to monotherapy (P&lt;0.0001).</p> <p>Secondary: Significantly more patients on combination therapy met the DBP end point than patients on amlodipine monotherapy (67.0 vs 48.3%; P=0.0003).</p> <p>Patients on combination therapy had significantly greater mean SBP reductions (-25.5 vs -20.5 mm Hg; P=0.0003) and DBP reductions (-14.3 vs -10.4 mm Hg; P=0.0001) than patients on amlodipine monotherapy.</p> <p>Significantly more patients on combination therapy met the BP goal of &lt;140/90 mm Hg than patients on amlodipine monotherapy (61.0 vs 43.3%; P=0.0007).</p> <p>Significantly more patients on combination therapy met the BP goal of</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			goal of 140/90 and $\leq$ 130/85 mm Hg, mean reduction in SBP and DBP and incidence of edema	<p>&lt;130/85 mm Hg than patients on amlodipine monotherapy (35.7 vs 19.1%; P=0.0004).</p> <p>The incidence of peripheral edema was significantly higher in the amlodipine monotherapy group (23.3 vs 12.6%; P=0.0102).</p> <p>There was no significant difference in the incidence of other adverse events.</p>
<p>Neutel et al.<sup>83</sup> (2005) SELECT</p> <p>Amlodipine and benazepril 5-20 mg/day (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 mg/day</p> <p>vs</p> <p>benazepril 20 mg/day</p>	<p>DB, RCT</p> <p>Patients with stage 2 systolic HTN</p>	<p>N=443</p> <p>8 weeks</p>	<p>Primary: Reduction in SBP, proportion of patients achieving blood pressure control</p> <p>Secondary: Not reported</p>	<p>Primary: Significantly greater SBP reductions were achieved with combination therapy compared to amlodipine or benazepril monotherapy (P&lt;0.0001).</p> <p>Significantly more patients on combination therapy met blood pressure goals than on monotherapy (P&lt;0.0001).</p> <p>No significant difference was noted in the incidence of adverse events. Adverse events were low in all three treatment arms, with less peripheral edema in the combination group than in the amlodipine-treated group.</p> <p>Secondary: Not reported</p>
<p>Kuschnir et al.<sup>84</sup> (1996)</p> <p>Amlodipine-benazepril 5/20 mg QD (fixed-dose combination)</p> <p>vs</p>	<p>DB, MC, PC, PG, RCT</p> <p>Men and women 21 to 80 years of age with uncomplicated primary HTN</p>	<p>N=308</p> <p>8 weeks</p>	<p>Primary: Reduction in mean sitting DBP, SBP and percentage of patients with DBP &lt;90 mm Hg or <math>\geq</math>10 mm Hg reduction</p>	<p>Primary: All treatment groups significantly reduced mean sitting DBP compared to placebo (P&lt;0.001).</p> <p>Combination amlodipine/benazepril had significantly greater reductions in DBP (-13.2 mm Hg; P&lt;0.001) compared to amlodipine (-8.8 mm Hg) and benazepril (-6.7 mm Hg) monotherapy.</p> <p>Combination amlodipine and benazepril had significantly greater reductions in SBP (-24.7 mm Hg; P&lt;0.001) compared to amlodipine (-</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
amlodipine 5 mg QD vs benazepril 20 mg QD vs placebo				16.2 mm Hg) and benazepril (-12.4 mm Hg).  Significantly more patients on combination amlodipine and benazepril reached DBP <90 mm Hg or ≥10 mm Hg reduction (87.0%; P≤0.005) compared to amlodipine (67.5%) and benazepril (53.3%).  Adverse events considered to be drug related occurred in 15.6% of patients receiving amlodipine and benazepril, 24.7% of patients receiving amlodipine, 6.5% of patients on benazepril and 11.7% of patients on placebo.
Chrysant et al. <sup>85</sup> (2007)  Amlodipine and benazepril 10-40 mg QD for 6 weeks (fixed-dose combination product) vs amlodipine and benazepril 10-40 mg QD for 2 weeks, followed by 20-40 mg QD for 4 weeks (fixed-dose combination product) vs amlodipine 10 mg QD for 6 weeks	DB, MC, RCT  Men and women ≥18 years of age with mean sitting DBP ≥95 mm Hg not adequately controlled with amlodipine 10 mg/day monotherapy	N=812  6 weeks	Primary: Reduction in mean sitting DBP and SBP, reductions in ambulatory blood pressure, successful response (mean sitting DBP <90 mm Hg or decrease of ≥10 mm Hg from baseline), safety  Secondary: Not reported	Primary: Treatment with benazepril 40 mg and amlodipine 10 and benazepril 20 mg and amlodipine 10 mg resulted in a decrease of mean sitting SBP and DBP by 13.3/12.7 and 12.1/11.6 mm Hg, respectively, compared to monotherapy (6.6/8.5 mm Hg; P<0.0001).  Benazepril 40 mg and amlodipine 10 mg and benazepril 40 mg and amlodipine 20 mg decreased ambulatory SBP and DBP by 9.9/6.7 and 7.4/5.2 mm Hg, respectively, compared to monotherapy (P<0.0001).  Both combination therapy groups resulted in more responders than monotherapy (74 and 65 vs 54%; P<0.0001 and P<0.0085, respectively). Combination therapy had significantly greater reductions in sitting SBP (-17 mm Hg; P<0.0001) compared to amlodipine monotherapy (-5 mm Hg).  The incidence of pedal edema was lower but not significantly different in the combination therapy groups compared to monotherapy (4.5, 5.5 vs 9.2%, respectively; P value not significant). No significant metabolic side effects were noted among the combination therapy groups.  Secondary: Not reported
Chrysant et al. <sup>86</sup>	DB, RCT	N=329	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2004)</p> <p>Amlodipine and benazepril 5-40 mg QD for 4 weeks, followed by 10-40 mg QD for 4 weeks (fixed-dose combination product)</p> <p>vs</p> <p>benazepril 40 mg QD for 8 weeks</p>	<p>Men and women (mean age 53 years) with mean sitting DBP <math>\geq</math>95 mm Hg not adequately controlled with benazepril 40 mg/day monotherapy</p>	<p>8 weeks</p>	<p>Reduction in mean sitting DBP and SBP, reduction in standing DBP and SBP, and change in heart rate, safety</p> <p>Secondary: Not reported</p>	<p>Combination therapy had significantly greater reductions in sitting SBP (-17 mm Hg; <math>P &lt; 0.0001</math>) compared monotherapy (-5 mm Hg).</p> <p>Combination therapy had significantly greater reductions in sitting DBP (-14 mm Hg; <math>P &lt; 0.0001</math>) compared to monotherapy (-7 mm Hg).</p> <p>Combination therapy had significantly greater reductions in standing SBP (-17 mm Hg; <math>P &lt; 0.0001</math>) compared to monotherapy (-6 mm Hg).</p> <p>Combination therapy had significantly greater reductions in standing DBP (-14 mm Hg; <math>P &lt; 0.0001</math>) compared to monotherapy (-7 mm Hg).</p> <p>No significant differences in heart rate were observed (<math>P &gt; 0.05</math>).</p> <p>No significant differences in adverse events were reported (<math>P &gt; 0.05</math>).</p> <p>Secondary: Not reported</p>
<p>Fogari et al.<sup>87</sup> (1997)</p> <p>Amlodipine and benazepril 2.5-10 to 5-10 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>benazepril 10 mg QD</p>	<p>DB, MC, PC, RCT</p> <p>Men and women 24 to 73 years of age (mean 55 years) with HTN inadequately controlled with ACE inhibitor monotherapy</p>	<p>N=448</p> <p>8 weeks</p>	<p>Primary: Reduction in mean sitting DBP</p> <p>Secondary: Reduction in sitting SBP, standing DBP and SBP, and percentage of patients with DBP <math>&lt;</math>90 mm Hg (deemed excellent response) or a <math>\geq</math>10 mm Hg reduction (deemed good response)</p>	<p>Primary: Significantly greater reductions in sitting DBP were observed with benazepril 10 mg and amlodipine 2.5 mg (-5.3 mm Hg, 97.5% CI, -8.3 to -2.4; <math>P = 0.0001</math>) and benazepril 10 mg and amlodipine 5 mg (-4.5 mm Hg, 97.5% CI, -7.4 to -1.6; <math>P = 0.0006</math>) compared to benazepril monotherapy.</p> <p>Secondary: Significantly greater reductions in sitting SBP were seen with benazepril 10 mg and amlodipine 2.5 mg (-7.9 mm Hg, 97.5% CI, -12.3 to -3.5; <math>P = 0.0001</math>) and benazepril 10 mg and amlodipine 5 mg (-7.9 mm Hg, 97.5% CI, -12.2 to -3.6; <math>P = 0.0000</math>) compared to benazepril monotherapy.</p> <p>Significantly greater reductions in standing DBP and SBP were also reported with the combination therapy compared to benazepril monotherapy (<math>P \leq 0.001</math>).</p> <p>Significantly more patients had excellent or good response with benazepril 10 mg and amlodipine 2.5 mg (69.2%; <math>P = 0.0004</math>) and 10-5 mg (65.8%; <math>P = 0.02</math>) compared to benazepril monotherapy (40.5%).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Minami et al.<sup>88</sup> (2007)</p> <p>Losartan 50 mg/day and HCTZ 12.5 mg/day</p> <p>vs</p> <p>candesartan 8 mg QD or amlodipine 5 mg QD</p>	<p>OL</p> <p>Japanese outpatients with essential HTN treated for <math>\geq 2</math> months with either candesartan or amlodipine and 24-hour ambulatory blood pressure <math>\geq 135/80</math> mm Hg</p>	<p>N=15</p> <p>12 months</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Tolerability was good in the three treatment groups and no significant abnormal laboratory data was detected.</p> <p>Primary: In patients who had previously received candesartan, 24-hr blood pressure decreased significantly from 137/89 mm Hg to 126/81 mm Hg after three months (<math>P&lt;0.05/P&lt;0.001</math>) and to 123/81 mm Hg after 12 months (<math>P&lt;0.01/P&lt;0.001</math>) of treatment with losartan and HCTZ.</p> <p>In patients who had previously received amlodipine, 24-hr blood pressure decreased significantly from 137/81 to 125/75 mm Hg after three months (<math>P&lt;0.05/P&lt;0.05</math>) and to 124/77 mm Hg after 12 months (<math>P&lt;0.05/P</math> value not significant) of treatment with losartan and HCTZ.</p> <p>There were significant decreases in SBP during the daytime, nighttime and early morning after 12 months in both groups.</p> <p>No adverse changes in the indices of glucose or lipid metabolism were observed in either group.</p> <p>Secondary: Not reported</p>
<p>Hilleman et al.<sup>89</sup> (1999)</p> <p>Amlodipine-benazepril (fixed-dose combination)</p> <p>vs</p> <p>monotherapy (atenolol, HCTZ, captopril, enalapril, lisinopril, amlodipine,</p>	<p>MA</p> <p>Patients with mild-to-moderate essential hypertension</p>	<p>82 trials</p> <p><math>\geq 4</math> weeks</p>	<p>Primary: Absolute change in supine DBP from baseline</p> <p>Secondary: Percent of patients who achieved blood pressure control, safety</p>	<p>Primary: The mean absolute decrease in supine DBP ranged from 9.7 to 13.3 mm Hg with verapamil showing the greatest effect and captopril the least. When studies were weighted by sample size, amlodipine and benazepril, atenolol, lisinopril, and verapamil showed the greatest blood pressure effect.</p> <p>Secondary: The average percentage of patients defined as controlled after treatment varied from 53.5 to 79.0%, with amlodipine and benazepril (74.3%) and lisinopril (79.0%) showing the highest percentage control (<math>P=0.096</math>).</p> <p>The incidence of adverse events ranged from 12.1 to 41.8%, with lisinopril and verapamil showing the lowest incidences (12.1% and 14.1%, respectively) and nifedipine the highest incidence. Lisinopril demonstrated</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>diltiazem, nifedipine, verapamil)</p>				<p>significantly less overall side effects compared to nifedipine (P=0.030).  Nifedipine demonstrated a higher withdrawal rate due to side effects compared to atenolol, HCTZ, enalapril, amlodipine, and diltiazem (P=0.002). Although amlodipine and benazepril had the lowest rate of withdrawals due to adverse events, lack of significant change was due to the low number of cohorts available for analysis.</p>
<p>Jamerson et al.<sup>90</sup> (2007) ACCOMPLISH  Amlodipine 5 mg QD plus benazepril 20 mg QD  vs  benazepril 20 mg QD plus HCTZ 12.5 mg QD</p>	<p>DB, MC, RCT  Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=10,704  Analysis performed at 6 months (complete trial duration 5 years)</p>	<p>Primary: Changes in mean SBP from baseline to 6 months, blood pressure control rates (SBP/DBP &lt;140/90 mm Hg or &lt;130/89 mm Hg for patients with diabetes and chronic kidney disease)  Secondary: Not reported</p>	<p>Primary: At baseline, 97% of subjects were treated with antihypertensive medications at entry, but only 37% of participants had blood pressure control.  Mean blood pressure fell from 145/80 to 132/74 mm Hg after six months of treatment with either combination regimen (P&lt;0.001).  The six month blood pressure control rate was 73% in the overall trial (78% in the United States), 43% in diabetics, and 40% in patients with renal disease. Of the patients uncontrolled, 61% were not on maximal medications.  Secondary: Not reported</p>
<p>Malacco et al.<sup>91</sup> (2002)  Amlodipine and benazepril 5-10 mg QD (fixed-dose combination product)  vs  captopril and HCTZ 50-25 mg QD (fixed-dose</p>	<p>DB, MC, RCT  Patients with mild to moderate arterial HTN (sitting DBP &gt;95 mm Hg and/or SBP &gt;160 mm Hg) inadequately controlled by monotherapy with an ACE inhibitor, calcium-channel blocking agent or diuretic</p>	<p>N=397  12 weeks</p>	<p>Primary: Reduction in sitting DBP and SBP  Secondary: Percentage of patients responding to therapy (DBP&lt;90 mm Hg, reduction in DBP ≥10 mm Hg or SBP ≥20 mm Hg, or SBP &lt;150 mm</p>	<p>Primary: Significantly lower sitting DBP (-2.7 mm Hg; P&lt;0.001) and SBP (-3.7 mm Hg; P&lt;0.001) were achieved with amlodipine and benazepril compared to captopril and HCTZ.  Secondary: Significantly more amlodipine and benazepril patients responded to therapy (94.8%) compared to captopril and HCTZ (86.0%; P=0.004).  No differences in adverse events were reported between the two treatment groups.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
combination product)			Hg)	
<p>Kereiakes et al.<sup>92</sup> (2007)</p> <p>Benazepril 10 mg/day for 2 weeks, then 20 mg/day for 2 weeks, then benazepril 20 mg/day plus amlodipine 5 mg/day for 4 weeks, then benazepril 20 mg/day plus amlodipine 10 mg/day for 4 weeks</p> <p>vs</p> <p>olmesartan 20 mg/day for 2 weeks, then 40 mg/day for 2 weeks then olmesartan and HCTZ 40-12.5 mg/day for 4 weeks increased to 40-25 mg for 4 weeks</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients with stage 2 HTN</p>	<p>N=190</p> <p>12 weeks</p>	<p>Primary: Change in mean seated SBP at the end of week 12</p> <p>Secondary: DBP at the end of week 12, percent of patients attaining blood pressure goals of &lt;140/90, &lt;130/85, and &lt;130/80 mm Hg</p>	<p>Primary: Patients treated with olmesartan and HCTZ experienced significantly greater reductions in mean seated SBP at week 12 than patients treated with benazepril plus amlodipine (least square mean change, -32.5 vs -26.5 mm Hg; P=0.024; least square mean treatment difference, -6.0 mm Hg; 95% CI, -11.1 to -0.8).</p> <p>Secondary: The least square mean change for reduction in DBP approached statistical significance with olmesartan and HCTZ compared to benazepril plus amlodipine at week 12 (P=0.056).</p> <p>The percentage of patients achieving goal rates at the end of the study for olmesartan and HCTZ and benazepril plus amlodipine were 66.3 and 44.7% (P=0.006) for &lt;140/90 mm Hg, 44.9 vs 21.2% (P=0.001) for &lt;130/85 mm Hg, and 32.6 and 14.1% (P=0.006) for &lt;130/80 mm Hg.</p> <p>Both treatments were well tolerated.</p>
<p>Tatti et al.<sup>93</sup> (1998) FACET</p> <p>Amlodipine 10 mg QD</p> <p>vs</p>	<p>OL, PRO, RCT</p> <p>Men and women, diagnosed with HTN (SBP &gt;140 mm Hg or DBP &gt;90 mm Hg) and non-insulin dependent</p>	<p>N=380</p> <p>Up to 3.5 years</p>	<p>Primary: Blood pressure</p> <p>Secondary: Fasting serum glucose, serum creatinine, plasma insulin, HbA<sub>1c</sub>, TC,</p>	<p>Primary: Both treatment groups significantly lowered SBP and DBP from baseline (P&lt;0.05).</p> <p>SBP was lower in the amlodipine group by 4 mm Hg than in the fosinopril group (P&lt;0.01). There was no difference in DBP, both groups decreased by 8 mm Hg.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>fosinopril 20 mg QD</p> <p>If blood pressure was not controlled on monotherapy, the other study drug was added.</p>	<p>diabetes</p>		<p>HDL-C, TG, fibrinogen, microalbuminuria</p>	<p>Amlodipine was added by 30.7% of the fosinopril group and fosinopril was added by 26.2% of the amlodipine group (P&gt;0.1).</p> <p>Secondary: No difference between the groups was found for serum creatinine, HbA<sub>1c</sub>, and triglycerides at the endpoint (P&gt;0.05).</p> <p>Fasting serum glucose, serum insulin and microalbuminuria were significantly lower at endpoint for both groups but not significantly different from each other (P&gt;0.05).</p> <p>Total cholesterol increased in both groups, and high-density lipoprotein cholesterol increased significantly in the fosinopril group (P&lt;0.05).</p> <p>No difference in fibrinogen levels was observed between the groups at the end of the trial (P&gt;0.05).</p>
<p>Miranda et al.<sup>94</sup>(2008)</p> <p>Amlodipine 2.5 to 10 mg and ramipril 2.5 to 10 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Adults 40 to 79 years of age with stage 1 or 2 essential HTN</p>	<p>N=222</p> <p>18 weeks</p>	<p>Primary: Change in SBP and DBP</p> <p>Secondary: Safety and tolerability</p>	<p>Primary: The mean changes in ambulatory BP were greater with amlodipine and ramipril compared to amlodipine monotherapy (SBP, -20.21 vs -15.31 mm Hg and DBP, -11.61 vs -8.42 mm Hg, respectively; both, P=0.002]. There was no significant difference among the treatment groups in office BP (SBP, -26.60 vs -22.97 mm Hg and DBP, -16.48 vs -14.48 mm Hg; both, P value not significant).</p> <p>Secondary: Twenty-nine patients (22.1%) treated with combination therapy and 41 patients (30.6%) treated with monotherapy experienced ≥1 adverse event considered possibly related to study drug. The combination-therapy group had lower prevalence of edema (7.6 vs 18.7%; P=0.011) and a similar prevalence of dry cough (3.8 vs 0.8%; P value not significant).</p>
<p>Fogari et al.<sup>95</sup> (2007)</p> <p>CANDIA</p> <p>Amlodipine 10 mg QD</p>	<p>DB, MC, RCT</p> <p>Patients, 20 to 80 years old, with mild to moderate uncomplicated HTN not controlled on</p>	<p>N=203</p> <p>8 weeks</p>	<p>Primary: Decrease in DBP</p> <p>Secondary: Sitting SBP, reduction of the orthostatic blood</p>	<p>Primary: There was no significant difference in the mean decrease in DBP between treatment groups; the difference in final DBP was -0.02 mm Hg (95% CI, -1.48 to 1.52f; P=0.979).</p> <p>Secondary: There was no significant difference between the groups at week eight for</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs candesartan 16 mg and HCTZ 12.5 mg QD	monotherapy with an antihypertensive (SBP <180 mm Hg and DBP 90 to 110 mm Hg)		pressure at least two minutes after standing, change in heart rate, percentage of patients normalized (DBP <90 mm Hg and SBP <140 mm Hg), percentage of responders (reduction in DBP ≥5 mm Hg)	the following: sitting SBP (P=0.835), heart rate (P<0.500), orthostatic SBP (P=0.883), orthostatic DBP (P=0.264), percentage of patients normalized (P=10), percentage of responders (P=0.900).  The number of patients reporting an adverse event was greater in the amlodipine group (P=0.001).  The number of patients reporting an adverse drug-related event was greater in the amlodipine group (P<0.001).  Changes in blood chemistry and other secondary measurements were not significantly different between the treatment groups.
Ribeiro et al. <sup>96</sup> (2007) LAMHYST  Amlodipine 5 to 10 mg QD  vs  losartan 50 to 100 mg QD	DB, DD, RCT  Males and females, age 18 to 79 years old, with diagnosis of mild (>95 mm Hg but <115 mm Hg) to moderate essential HTN and not taking an antihypertensive medication (within last 4 weeks)	N=194  12 weeks	Primary: Difference between treatment groups in mean change in ABPM for last 9 hours of treatment and during drug holiday  Secondary: Not reported	Primary: After 12 weeks, mean reductions in SBP were significantly larger in the amlodipine group than the losartan group (-18.1 vs -10.1 mm Hg; P<0.001). Mean reductions in DBP were significantly larger in the amlodipine group than the losartan group (-18.1 vs -10.1 mm Hg; P<0.05).  Mean increases in SBP were similar between the groups during the two day drug holiday (P>0.05).  After the two day drug holiday, SBP was lower than baseline in both groups (P<0.001), with the amlodipine group SBP remaining significantly lower (P<0.01).  Mean increases in DBP were similar between the groups during the two day drug holiday (P>0.05). After the two day drug holiday, DBP was lower than baseline in both groups (P=0.0001), with the amlodipine group DBP remaining significantly lower (P<0.05).  Secondary: Not reported
Oparil et al. <sup>97</sup> (1996)  Amlodipine 5 to 10	DB, DD, MC, RCT  Patients with HTN	N=900  12 weeks	Primary: Efficacy, tolerability, effects on QOL	Primary: DBP reductions after 4, 8, and 12 weeks of therapy were clinically comparable (losartan group: 7.3, 10.4, and 11.1 mm Hg, respectively; amlodipine group: 7.9, 11.2, and 11.8 mm Hg, respectively; P value not

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD</p> <p>vs</p> <p>losartan 50 to 100 mg QD</p> <p>If goal DBP (<math>\leq 90</math> mm Hg) was not attained, drug doses could be doubled and/or HCTZ mg was added.</p>			<p>Secondary: Not reported</p>	<p>significant).</p> <p>Similar reductions in SBP were seen for both treatment groups (P value not significant).</p> <p>The percentage of patients reaching goal DBP (<math>\leq 90</math> mm Hg) or DBP <math>\geq 90</math> mm Hg with a <math>\geq 10</math> mm Hg decrease from baseline) was comparable for the two groups, with 68% of patients in the losartan group and 71% of patients in the amlodipine group reaching goal.</p> <p>Significantly more patients in the amlodipine group had drug-related adverse experiences (27 vs 13%; <math>P=0.029</math>). Edema was more common in patients receiving the amlodipine regimen than in those receiving the losartan regimen (11 vs 1%; <math>P=0.004</math>).</p> <p>Overall QOL was not different in the two treatment groups.</p> <p>Secondary: Not reported</p>
<p>Chrysant et al.<sup>98</sup> (2008) COACH</p> <p>Amlodipine 5 to 10 mg QD and olmesartan 10 to 40 mg</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>olmesartan 10 to 40 mg QD</p>	<p>DB, MC, PC, RCT</p> <p>Patients, age 18 years and older, with seated DBP of 95 to 120 mm Hg</p>	<p>N=1,940</p> <p>8 weeks</p>	<p>Primary: Change from baseline in seated DBP at week 8</p> <p>Secondary: Change from baseline in seated SBP at week 8; mean change from baseline in seated DBP and SBP at weeks 2, 4, 6 and 8 without last observation carried forward; proportion of patients achieving</p>	<p>Primary: All active treatments and placebo resulted in significant decreases in seated DBP at week eight (<math>P&lt;0.001</math>). Reductions in seated DBP with monotherapy treatment ranged from -8.3 to -12.7 mm Hg; reductions with combination therapy ranged from -13.8 to -19.0 mm Hg. All combinations reduced seated DBP significantly greater than either component as monotherapy at the same dosage (<math>P&lt;0.001</math>).</p> <p>Secondary: All active treatments and placebo resulted in significant decreases in seated SBP at week eight (<math>P&lt;0.001</math> for treatment, <math>P=0.024</math> for placebo). All combinations reduced seated SBP significantly greater either component as monotherapy at the same dosage (<math>P&lt;0.001</math>).</p> <p>The proportion of patients achieving goal blood pressures were: 20.0 to 36.3% of patients receiving olmesartan monotherapy, 21.1 to 32.5% of patients receiving amlodipine monotherapy, 35.0 to 53.2% of patients receiving combination therapy, and 8.8% of patients receiving placebo.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			blood pressure goal (<140/90 mm Hg or <130/80 mm Hg); safety	<p>Combination therapy resulted in significantly greater achievement of goal blood pressure than monotherapy (P&lt;0.005).</p> <p>No difference in overall rates of adverse events across the different treatment groups was seen. Nearly 27% of patients experienced a drug-related adverse event.</p> <p>Changes in laboratory values were not considered clinically significant nor followed a consistent pattern with treatment: none of the changes were considered clinically significant. Platelet counts increased significantly from baseline (statistically) for patients receiving amlodipine, however the increase was &lt;10% and not deemed clinically relevant.</p>
<p>Chrysant et al.<sup>99</sup> (2009) COACH</p> <p>Amlodipine 5 to 10 mg QD and olmesartan 10 to 40 mg</p> <p>HCTZ 12.5 to 25 mg could be added if blood pressure was not controlled (&lt;140/90 mm Hg or &lt;130/80 mm Hg in patients with diabetes).</p>	<p>OL, ES</p> <p>Patients ≥ 18 years of age with essential HTN (seated DBP ≥95 and &lt;120 mm Hg)</p>	<p>N=1,684</p> <p>44 weeks OL therapy (52 weeks total study duration including 8 week DB phase)</p>	<p>Primary: Reduction in mean sitting SBP DBP, change in mean sitting SBP and DBP, percentage of patients achieving blood pressure goal (&lt;140/90 mm Hg or &lt;130/80 mm Hg for patients with diabetes)</p>	<p>Primary: Mean sitting DBP decreased from 101.5 mm Hg at baseline to 81.9 mm Hg and mean sitting SBP decreased from 163.6 mm Hg at baseline to 131.2 mm Hg at week 52.</p> <p>Approximately 31% of patients remained on amlodipine 5 mg and olmesartan 40 mg. Increasing the dose of amlodipine to 10 mg in combination with olmesartan 40 mg produced further decreases in mean sitting DBP of 4.8 mm Hg and mean sitting SBP of 7.3 mm Hg. Addition of HCTZ 12.5 mg to amlodipine 10 mg and olmesartan 40 mg decreased mean sitting DBP by 4.5 mm Hg and mean sitting SBP by 7.7 mm Hg. Doubling the HCTZ dose from 12.5 to 25 mg decreased mean sitting DBP and mean sitting SBP by an additional 6.0 mm Hg and 9.9 mm Hg, respectively. Patients who received the triple therapy had the greatest mean sitting SBP reduction (36.1 mm Hg).</p> <p>Approximately 67% of patients achieved blood pressure goal by week 52. The blood pressure goal achievement was 80% for amlodipine and olmesartan 5/40 mg, 70.6% for amlodipine and olmesartan 10/40 mg, 66.6% for amlodipine and olmesartan and HCTZ 10/40/12.5 mg, and 46.3% for amlodipine and olmesartan and HCTZ 10/40/25 mg.</p> <p>The addition of HCTZ 25 mg enabled more patients to achieve blood pressure targets of &lt;140/90 mm Hg (77.7%), &lt;130/85 mm Hg (47.5%),</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>and &lt;130/80 mm Hg (36.4%) compared to the other treatment regimens.</p> <p>No major safety issues emerged with long-term therapy. The frequency of edema ranged from 8.9% in patients treated with amlodipine 5 mg and olmesartan 40 mg to 14.5% in patients treated with amlodipine 10 mg and olmesartan 40 mg plus HCTZ 25 mg. Other treatment-emergent adverse events experienced by <math>\geq 3\%</math> of patients included upper respiratory tract infection (6.5%), nasopharyngitis (5.2%), extremity pain (4.1%), sinusitis (3.6%), arthralgia (3.3%), and back pain (3.1%). headache (2.0%), hypotension (1.8%), and fatigue (1.6%). The incidence of cough was 0.4%.</p>
<p>Oparil et al.<sup>100</sup> (2009) COACH</p> <p>Amlodipine 5 to 10 mg QD and olmesartan 10 to 40 mg</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>olmesartan 10 to 40 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, factorial, MC, PC, RCT</p> <p>Patients <math>\geq 18</math> years of age with seated DBP 95 to 120 mm Hg, with a subgroup analysis based on HTN (stage 1: SBP 140 to 159 mm Hg or DBP 90 to 99 mm Hg; stage 2: SBP <math>\geq 160</math> mm Hg or DBP <math>\geq 100</math> mm Hg) and no prior antihypertensive medication</p>	<p>N=1,940</p> <p>8 weeks</p>	<p>Primary: Mean change in DBP and SBP at week 8 for each subgroup</p> <p>Secondary: Proportion of patients achieving blood pressure goal (&lt;140/90 mm Hg or &lt;130/80 mm Hg)</p>	<p>Primary: Reductions in mean DBP as a result of combination treatment were similar between subgroups. Patients with stage 1 HTN achieved reductions of 14.8 to 15.8 mm Hg and patients with stage 2 HTN achieved reductions of 13.6 to 19.8 mm Hg. Reductions in mean SBP as a result of combination treatment resulted in greater reductions in patients with stage 2 HTN (25.1 to 32.7 mm Hg) compared to stage 1 HTN (17.7 to 23.7 mm Hg) (P value not reported).</p> <p>Reductions in mean DBP and SBP were similar between those with no prior antihypertensive treatment and those with prior hypertensive treatment.</p> <p>Secondary: The proportion of patients with stage 1 HTN who received combination treatment and achieved blood pressure goal was 65.6 to 80.0%, compared to 40.5 to 66.7% of those who received monotherapy (P&lt;0.0001 across treatments).</p> <p>The proportion of patients with stage 2 HTN who received combination treatment and achieved BP goal was 40.5 to 49.2%, compared to 13.1 to 29.2% of those who received monotherapy (P&lt;0.0001).</p> <p>Results of patients with baseline SBP <math>\geq 180</math> mm Hg were similar to other subgroups.</p>
<p>Braun et al.<sup>101</sup></p>	<p>OL, PRO</p>	<p>N=257</p>	<p>Primary:</p>	<p>Primary:</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(abstract) (2009)</p> <p>Amlodipine 10 mg plus olmesartan 20 mg QD</p> <p>If patients were uncontrolled after 4 weeks, they were changed to amlodipine and valsartan 10-160 mg QD.</p>	<p>Patients with DBP 100 to 109 mm Hg</p>	<p>8 weeks</p>	<p>Reduction in SBP and DBP</p> <p>Secondary: Adverse events</p>	<p>Following treatment with amlodipine and olmesartan, SBP/DBP decreased by 19.2±12.4/14.4±7.4 mm Hg.</p> <p>The number of patients who progressed to treatment with amlodipine and valsartan was 175. Additional reductions in SBP of 7.9 mm Hg and DBP of 3.9 mm Hg were seen (P&lt;0.0001 for both).</p> <p>Secondary: Both treatments were well tolerated and reported adverse events were consistent with drug profiles.</p>
<p>Elliott et al.<sup>102</sup> (2015)</p> <p>Amlodipine 10 mg QD</p> <p>vs</p> <p>perindopril 16 mg QD</p> <p>vs</p> <p>perindopril + amlodipine 14-10 mg QD</p>	<p>PRO, RCT</p> <p>Hypertensive patients 18 to 75 years of age</p>	<p>N=820</p> <p>42 days</p>	<p>Primary: Change in mean seated office trough DBP from baseline to day 42</p> <p>Secondary: Mean seated office SBP, responder rates (those achieving target BP of (&lt;140/90 for non-diabetics, &lt;130/80 for diabetics) , safety</p>	<p>Primary: Least square mean BP changes over the 42 days of treatment: -12.7/-9.1 mmHg for perindopril, -18.8/-12.9 mmHg for amlodipine, and -22.8/-15.4 mmHg for the combination of perindopril + amlodipine. Changes in both office systolic and diastolic BPs were significantly greater with combination therapy, at 42 days of therapy (-10.1/-6.3 mmHg vs perindopril, both P&lt;0.0001, and -3.9/2.5 mmHg vs amlodipine, both P&lt;0.002). Analogous analyses restricted to the per-protocol population provided similar results.</p> <p>Secondary: Regardless of the duration of therapy, or the intent-to-treat or “per protocol” population, a significantly greater proportion of subjects achieved “target BP” in the perindopril + amlodipine group, compared with perindopril or amlodipine alone: in the pre-specified analysis, the proportions were 52.4% versus 37.1% versus 25.9%, respectively (P&lt;0.0001).</p> <p>The combination showed a lower incidence of pedal edema and adverse events compared with amlodipine. No deaths or significant differences across groups in early discontinuation, serum potassium, or rates of total or serious adverse events or glomerular filtration, were observed.</p>
<p>Manolis et al.<sup>103</sup></p>	<p>OBS, PRO</p>	<p>N=2,231</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2015)</p> <p>Fixed-dose combination perindopril-amlodipine (available in dosages of 5/5, 5/10, 10/5, or 10/10 mg)</p>	<p>Ambulatory men or women <math>\geq 18</math> years of age with diagnosed essential hypertension that was treated with daily fixed-dose combination perindopril arginine-amlodipine</p>	<p>6 months</p>	<p>BP reduction over six months</p> <p>Secondary: BP control after six months</p>	<p>SBP and DBP of patients who received perindopril-amlodipine decreased significantly versus baseline after three months and six months in the per protocol set (<math>P &lt; 0.001</math>). Mean systolic BP decreased from <math>157.0 \pm 15.4</math> mmHg to <math>129.0 \pm 7.9</math> mmHg after six months, and diastolic BP from <math>91.5 \pm 10.1</math> to <math>78.8 \pm 6.7</math> mmHg (both <math>P &lt; 0.001</math>).</p> <p>Secondary: BP control (<math>&lt; 140/90</math> mmHg) was achieved in 84.8% of the per protocol set.</p>
<p>Littlejohn et al.<sup>104</sup> (2009)</p> <p>Amlodipine 2.5 to 10 mg QD and telmisartan 20 to 80 mg QD</p> <p>vs</p> <p>telmisartan 20 to 80 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients <math>\geq 18</math> years of age with Stage 1 or 2 HTN (DBP <math>\geq 95</math> and <math>\leq 119</math> mm Hg)</p>	<p>N=2,607</p> <p>8 weeks</p>	<p>Primary: Change in the in-clinic seated diastolic BP</p> <p>Secondary: Change in the in-clinic seated SBP, DBP and SBP response (DBP <math>&lt; 90</math> mm Hg, decrease in DBP <math>\geq 10</math> mm Hg, SBP <math>&lt; 140</math> mm Hg, decrease in SBP <math>\geq 15</math> mm Hg), and BP control (DBP <math>&lt; 90</math> mm Hg and SBP <math>&lt; 140</math> mm Hg)</p>	<p>Primary: Both telmisartan (irrespective of amlodipine dosage; <math>P &lt; 0.0001</math>) and amlodipine (irrespective of telmisartan dosage; <math>P &lt; 0.0001</math>) significantly lowered the in-clinic DBP.</p> <p>The greatest reduction in blood pressure was with telmisartan 80 mg plus amlodipine 10 mg (SBP/DBP <math>-26.4/-20.1</math> mm Hg; <math>P &lt; 0.05</math> vs both monotherapies).</p> <p>DBP and SBP response was achieved by 91.2 and 90.4% of patients in the telmisartan 80 mg plus amlodipine 10 mg group, respectively.</p> <p>More than 50% of patients treated with combination therapy achieved blood pressure control, with the highest percentages (76.5% [overall control] and 85.3% [DBP control]) being achieved by patients treated with telmisartan 80 mg plus amlodipine 10 mg.</p> <p>A total of 37.3% of patients reported at least one adverse event. The most commonly reported adverse events were headache (5.4%) and peripheral edema (4.4%). Headache was more frequent in the placebo group (10.9%) compared to the telmisartan monotherapy (5.9%), amlodipine monotherapy (6.0%), and combination therapy (4.7%). The incidence of peripheral edema was highest in the amlodipine 10 mg group (17.8%); however, this rate was lower when amlodipine was used in combination with telmisartan: 11.4% (telmisartan 20 mg and amlodipine 10 mg), 6.2% (telmisartan 40 mg and amlodipine 10 mg), and 11.3% (telmisartan 80 mg</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Littlejohn et al.<sup>105</sup> (2009)</p> <p>Telmisartan and amlodipine 40-5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 40-10 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 80-5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 80-10 mg QD (fixed-dose combination product)</p> <p>vs</p>	<p>DB, DD, MC, PC, PG, RCT</p> <p>Patients ≥18 years of age with stage 1 or 2 HTN (DBP ≥95 and ≤119 mm Hg), with a subgroup analysis including patients with DBP ≥100 mm Hg at baseline</p>	<p>N=1,078</p> <p>8 weeks</p>	<p>Primary: Change in DBP from baseline to study end point</p> <p>Secondary: Change from baseline to study end in SBP; percent of patients achieving a DBP response (DBP &lt;90 mm Hg) and SBP response (SBP &lt;140 mm Hg or reduction from baseline ≥15 mm Hg); percent of patients achieving BP control (SBP/DBP &lt;140/&lt;90 mm Hg) and DBP control (&lt;90 mm Hg) and safety</p>	<p>and amlodipine 10 mg).</p> <p>Primary: Significant reductions in DBP were seen from baseline to study end for both dual therapy and monotherapy (P values not reported).</p> <p>Amlodipine 5 and 10 mg with telmisartan 40 and 80 mg significantly reduced DBP compared to respective monotherapies (P values not reported).</p> <p>Secondary: Amlodipine 5 and 10 mg with telmisartan 40 and 80 mg significantly reduced SBP compared to respective monotherapies (P values not reported).</p> <p>Combination therapy resulted in a greater DBP and SBP response than monotherapy (P values not reported).</p> <p>The highest rate of BP control was achieved with amlodipine 10 mg with telmisartan 80 mg.</p> <p>Rates of adverse events were similar between dual therapy and monotherapy. Incidences of adverse events were 4.40% with telmisartan monotherapy, 11.00% with amlodipine monotherapy and 11.75% with combination therapy. The most commonly reported events were headache and peripheral edema. Patients receiving amlodipine 10 mg had the highest incidence of peripheral edema; however rates were lower when amlodipine was used in combination with telmisartan.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
respective monotherapies, dosing frequency not specified				
Sharma et al. <sup>106</sup> (2007)  Telmisartan and amlodipine 40-5 mg QD (fixed-dose combination)  vs  amlodipine 5 mg QD	DB, MC, RCT  Patients 18 to 65 years of age with established stage 2 uncomplicated essential HTN	N=210  12 weeks	Primary: SBP/DBP reductions and responder rates (SBP/DBP <130/<80 mm Hg)  Secondary: Not reported	Primary: There was a significant reduction from baseline in mean SBP in both groups (telmisartan and amlodipine, from 176.3 to 128.0 mm Hg; amlodipine, from 171.8 to 143.4 mm Hg; both, P<0.05 vs baseline). There was a significant reduction in SBP from baseline in the telmisartan and amlodipine and amlodipine groups (-27.4 and -16.6%, respectively; P<0.05 within group and between groups).  There was a significant reduction from baseline in mean DBP in both treatment groups (telmisartan and amlodipine, from 100.9 to 93.8 mm Hg; amlodipine, from 99.7 to 94.3 mm Hg; both, P<0.05). There was a 20.2% reduction in mean DBP in the telmisartan and amlodipine group, which was significantly greater compared to the reduction of 12.7% observed in the amlodipine group (P<0.05 between groups and within both groups).  A total of 87.3% of patients receiving telmisartan and amlodipine reached the target SBP/DBP goal, compared to 69.3% of patients receiving amlodipine (P<0.05).  A total of 16.0% of patients in the telmisartan and amlodipine group experienced adverse events compared to 15.4% of patients in the amlodipine group (P value not significant). The most common adverse events in the telmisartan and amlodipine group were peripheral edema (8.5%), headache (5.7%), dizziness and cough (3.8%), and diarrhea (1.9%).  Secondary: Not reported
Neutel et al. <sup>107</sup> (2012) TEAMSTA  Telmisartan and	DB, MC, PG, RCT  Patients ≥18 years of age with severe HTN	N=858  8 weeks	Primary: Change in baseline blood pressure, blood pressure goal and response rates	Primary: Reductions in seated trough cuff blood pressure (-47.5/-18.7 mm Hg) were significantly greater with combination therapy compared to telmisartan (P<0.001) or amlodipine (P=0.002). Significant reductions with combination therapy were observed at one, two, four, and six weeks.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
amlodipine 80-10 mg QD (fixed-dose combination product)  vs  telmisartan 80 mg QD  vs  amlodipine 10 mg QD			Secondary: Safety	Blood pressure goal and response rates were consistently higher with combination therapy (50.4 and 91.4 to 99.7%) compared to monotherapy with either agent (24.1 and 69.3 to 91.5% and 35.6 and 83.9 to 98.5%).  Secondary: Combination therapy was well tolerated and fewer adverse events were reported with combination therapy compared to amlodipine (12.6 vs 16.4%). Peripheral edema was reported more frequently with amlodipine compared to combination therapy (13.2 vs 9.3%).
Maciejewski et al. <sup>108</sup> (2006)  Amlodipine 5 to 10 mg QD  vs  valsartan 80 to 160 mg QD  If blood pressure exceeded 140/90 while on highest treatment dose, HCTZ 12.5mg/day was added to the regimen.	DB, PRO, RCT, XO  African-Americans, older than 35 years, with baseline blood pressure >140/90 mm Hg and not on antihypertensive treatment	N=20  8 to 10 weeks for each arm with 2 week washout period before crossover	Primary: Comparison of 24-hr ABPM recordings  Secondary: Magnitude of change from baseline in SBP and DBP with each treatment, percent of patients who achieved goal <140/<90 with each treatment based on clinic blood pressure measurements	Primary: There was no difference between the groups based on 24-hr ABPM: SBP amlodipine 130±8 vs valsartan 127±17 (P=0.350) and DBP amlodipine 82±5 vs valsartan 84±16 (P=0.430).  Secondary: There was no difference between groups in magnitude of change from baseline in blood pressure (amlodipine -25±8/-18±7 vs valsartan -25±9/-16±7; P=0.61), and in percent of patients achieving goal blood pressure, 70% in the valsartan group and 75% in the amlodipine group (P=0.62).
Ichihara et al. <sup>109</sup> (2006)  Amlodipine 2.5 to 10 mg QD	RCT  Patients with untreated HTN (clinic SBP >140	N=100  12 months	Primary: ABPM and clinic blood pressure  Secondary:	Primary: Both treatments resulted in significant decreases in blood pressure, both ambulatory and clinic, over 12 months from baseline; blood pressure decreases were similar between treatment groups (between treatments: clinic SBP P=0.34; clinic DBP P=0.85; 24 hour ABPM P=0.14).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>valsartan 40 to 160 mg QD</p>	<p>mm Hg and/or DBP &gt;90 mm Hg; or ABPM SBP &gt;135 mm Hg and/or DBP &gt;98 mm Hg)</p>		<p>Pulse wave velocity, carotid intima-media thickness, urinary albumin excretion</p>	<p>Blood pressure variability decreased significantly in the amlodipine group compared to the valsartan group, where there was no change in blood pressure variability (P&lt;0.01).</p> <p>Secondary: The decrease in pulse wave velocity was significant from baseline for both groups, but not significantly different from each other (P&lt;0.05 from baseline).</p> <p>Intima-media thickness was not changed significantly from baseline for either treatment (P&gt;0.05 for both from baseline).</p> <p>Urinary albumin excretion in the valsartan group decreased significantly both from baseline and compared to amlodipine treatment (P&lt;0.05 from baseline, P value for comparison not reported).</p>
<p>Karpov et al.<sup>110</sup> (2012)</p> <p>Amlodipine and valsartan 5-80, 5-160, 10-160 mg QD (fixed-dose combination product)</p>	<p>OL, OS, PRO</p> <p>Patients with HTN</p>	<p>N=8,336</p> <p>3 months</p>	<p>Primary: Baseline reductions in blood pressure, blood pressure control (&lt;140/90 mm Hg)</p> <p>Secondary: Safety</p>	<p>Primary: Reductions in blood pressure were dose related. Overall, mean reductions in blood pressure ranged from 165.0/99.3 mm Hg at baseline to 128.7/80.4 mm Hg at 12 weeks (-36.3/-18.9 mm Hg; P&lt;0.0001).</p> <p>A total of 77.7% of patients achieved blood pressure control.</p> <p>Secondary: A total of 5.3% of patients reported adverse events. The incidence of edema declined from 10.4% at baseline to 8.5% at trial end.</p>
<p>Philipp et al.<sup>111</sup> (2007)</p> <p><u>Study 1</u></p> <p>Amlodipine 2.5 to 5 mg and valsartan 40 to 320 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 5</p>	<p>DB, MC, PC, RCT</p> <p>Males and females, ages 18 years and older with HTN (mean sitting DBP ≥95 mm Hg and &lt;110 mm Hg)</p>	<p>N=1,911</p> <p>8 weeks</p>	<p>Primary: Mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, response rate (proportion of patients with mean sitting DBP &lt;90 mm Hg or a ≥10</p>	<p>Primary: All treatments significantly decreased mean sitting DBP from baseline (P&lt;0.05).</p> <p>Combination treatment resulted in significantly greater blood pressure reduction than either monotherapy (P&lt;0.05 for all combinations compared to respective doses of monotherapy except amlodipine 2.5 mg and valsartan 40 mg QD).</p> <p>Secondary: All treatments significantly decreased mean sitting SBP from baseline</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD</p> <p>vs</p> <p>valsartan 40 to 320 mg QD</p> <p>vs</p> <p>placebo</p>			<p>mm Hg reduction from baseline), control rate (proportion of patients with mean sitting DBP &lt;90 mm Hg), adverse events (combined with study 2)</p>	<p>(P&lt;0.05).</p> <p>Combination treatment resulted in significantly greater blood pressure reduction than either monotherapy (P&lt;0.05 for all combinations compared to respective doses of monotherapy).</p> <p>Response rates were significantly different from placebo for all treatment groups (P&lt;0.05).</p> <p>Response rates for combination products were significantly different than each monotherapy for the following combinations: amlodipine 5 mg plus valsartan 80 mg, amlodipine 5 mg plus valsartan 40 mg and amlodipine 2.5 mg plus valsartan 80 mg (P&lt;0.05 for each combination compared to both monotherapy).</p> <p>Response rates for all combinations produced significantly improved compared to either one of the monotherapies except amlodipine 2.5 mg plus valsartan 40 mg (P&lt;0.05 for each combination compared to one of the respective monotherapy).</p> <p>Control rates with therapy were significantly better than placebo, with the highest control rate achieved with amlodipine 5 mg plus valsartan 320 mg (P&lt;0.05 compared to placebo, P value not reported for others).</p> <p>Adverse event rates were not significantly different among combination treatment, amlodipine treatment, and placebo.</p> <p>Adverse event rates were significantly different between amlodipine plus valsartan and valsartan monotherapy (P&lt;0.05).</p> <p>The most commonly reported adverse events for combination treatment were: peripheral edema, headache, nasopharyngitis, upper respiratory tract infection and dizziness. Peripheral edema occurred significantly less frequently in the combination treatment group than the amlodipine monotherapy group (5.4 vs 8.7%; P=0.014) and significantly more frequently than in the valsartan monotherapy group (5.4 vs 2.1%; P&lt;0.001). Peripheral edema occurrence in the valsartan group was similar</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Philipp et al.<sup>112</sup> (2007)</p> <p><u>Study 2</u> Amlodipine 10 mg and valsartan 160 or 320 mg QD</p> <p>vs</p> <p>amlodipine 10 mg QD</p> <p>vs</p> <p>valsartan 160 to 320 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Male and females, ages 18 years and older with hypertension (mean sitting DBP <math>\geq</math>95 mm Hg and <math>&lt;</math>110 mm Hg)</p>	<p>N=1,250</p> <p>8 weeks</p>	<p>Primary: Mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, response rate (proportion of patients with mean sitting DBP <math>&lt;</math>90 mm Hg or a <math>\geq</math>10 mm Hg reduction from baseline), control rate (proportion of patients with mean sitting DBP <math>&lt;</math>90 mm Hg), adverse events (combined with study 1)</p>	<p>to the rate in the placebo group.</p> <p>Primary: Mean sitting DBP was significantly reduced for both combination as compared to the individual components and to placebo (P<math>&lt;</math>0.05).</p> <p>Secondary: Response rates and control rates for combination treatments were significantly greater than valsartan monotherapy therapy and placebo therapy, but not different from amlodipine monotherapy (P<math>&lt;</math>0.05).</p> <p>Adverse event rates were not significantly different between combination treatment, amlodipine treatment and placebo.</p> <p>Adverse event rates were significantly different between amlodipine plus valsartan and valsartan monotherapy (P<math>&lt;</math>0.05).</p>
<p>Philipp et al.<sup>113</sup> (abstract) (2011)</p> <p>Amlodipine and valsartan 10-160 or 10-320 mg/day (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 10 mg/day</p>	<p>Post-hoc analysis</p> <p>Patients with HTN</p>	<p>N=834</p> <p>Not reported</p>	<p>Primary: Rate of blood pressure control (<math>&lt;</math>140/90 mm Hg), change in baseline blood pressure</p> <p>Secondary: Safety</p>	<p>Primary: Two weeks after starting therapy, blood pressure control rates were greater with amlodipine and valsartan 10-320 mg/day (49%) vs monotherapies (32 to 38%) and placebo (16%). Consistent results were observed in patients with stage 1 and 2 HTN. Among patients receiving combination therapy, statistically significant differences were observed at endpoint vs comparators. At all baseline blood pressure levels, the probability of achieving a blood pressure <math>&lt;</math>140/90 or <math>&lt;</math>130/80 mm Hg was greater with combination therapy compared to monotherapies and placebo.</p> <p>Secondary: Overall adverse events incidence was similar with combination therapy vs monotherapies and placebo.</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs valsartan 160 or 320 mg/day vs placebo				
Schunkert et al. <sup>114</sup> (2009) Amlodipine and valsartan 10-160 mg QD (fixed-dose combination product) vs amlodipine 10 mg QD	RCT, MC, DB, AC Patients ≥18 years of age with mild to moderate essential HTN (mean sitting DBP ≥90 mm Hg and <110 mm Hg) who were inadequately controlled on amlodipine 10 mg	N=944 8 weeks	Primary: Change from baseline in mean sitting DBP Secondary: Change from baseline in mean sitting SBP, responder rate (mean sitting DBP <90 mm Hg or ≥10 mm Hg reduction from baseline) and DBP control rate (mean sitting DBP <90 mm Hg)	Primary: At week eighth, a significantly greater reduction from baseline in msDBP was observed with amlodipine and valsartan (11.4 mm Hg) compared to amlodipine monotherapy (9.3 mm Hg; P<0.0001). Secondary: At week eight, a significantly greater reduction from baseline in msSBP was observed with amlodipine and valsartan (12.9 mm Hg) compared to amlodipine monotherapy (10.0 mm Hg; P<0.0001). The mean reductions in mean sitting SBP/mean sitting DBP were 24.4/17.2 and 21.6/15.0 mm Hg for the amlodipine and valsartan and amlodipine monotherapy, respectively The responder rate was significantly greater with amlodipine and valsartan (79.0%) than with amlodipine monotherapy (70.1%; P=0.0011). The percentage of patients with controlled DBP was significantly higher with amlodipine and valsartan (77.8%) compared to amlodipine monotherapy (66.5%; P<0.0001). The incidence of peripheral edema was higher with amlodipine monotherapy (9.4%) compared to amlodipine and valsartan (7.6%).
Ke et al. <sup>115</sup> (2010) Amlodipine and valsartan 5-80 mg	AC, DB, MC, RCT Hypertensive patients 18 to 86 years of age with	N=698 8 weeks	Primary: Change in mean sitting DBP Secondary:	Primary: At week eight, the reduction in mean sitting DBP was greater with amlodipine and valsartan (11.4/9.7 mm Hg) compared to amlodipine (7.4/7.1 mm Hg; P<0.0001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD (fixed-dose combination)</p> <p>vs</p> <p>amlodipine 5 mg QD</p>	<p>mean sitting DBP <math>\geq 95</math> and <math>&lt; 110</math> mm Hg who were inadequately controlled on amlodipine 5 mg for 4 weeks</p>		<p>Change in mean sitting SBP, diastolic response rate (mean sitting DBP <math>&lt; 90</math> mm Hg or <math>\geq 10</math> mm Hg decrease from baseline), diastolic control rate (mean sitting DBP <math>&lt; 90</math> mmHg) and overall BP control rate (mean sitting SBP/DBP <math>&lt; 140/90</math> mmHg)</p>	<p>Secondary:</p> <p>At week eight, the diastolic control and response rates were significantly greater in the amlodipine and valsartan compared to amlodipine monotherapy (diastolic control, 75.5 vs. 64.5%; <math>P=0.0002</math> and response rates, 79.3 vs. 66.8% [<math>P&lt;0.0001</math>], respectively).</p> <p>The proportion of patients achieving overall blood pressure control was greater in the amlodipine and valsartan group compared to amlodipine monotherapy (69.2 vs. 57.6%, <math>P=0.0013</math>). More than 50% of patients not adequately controlled on amlodipine monotherapy achieved blood pressure control after two weeks of therapy with amlodipine and valsartan.</p> <p>In a subgroup of patients, there was a significant reduction in 24-hr mean blood pressure from baseline with amlodipine and valsartan (-7.3/-6.3 mm Hg; <math>P&lt;0.0001</math>). There was no significant difference with amlodipine from baseline (-0.2/+0.3 mm Hg; <math>P&gt;0.05</math>).</p>
<p>Destro et al.<sup>116</sup> (2008) Ex-EFFeCTS</p> <p>Amlodipine and valsartan 5-160 mg QD for 2 weeks, followed by 10-160 mg QD for 6 weeks (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 mg QD for 2 weeks, followed by 10 mg QD for 6 weeks</p> <p>HCTZ 12.5 mg</p>	<p>DB, MC, RCT</p> <p>Patients <math>\geq 18</math> years of age with stage 2 HTN (mean sitting SBP <math>\geq 160</math> mm Hg)</p>	<p>N=646</p> <p>8 weeks</p>	<p>Primary:</p> <p>Mean changes in mean sitting SBP at week 4</p> <p>Secondary:</p> <p>Change from baseline in mean sitting DBP at week 4; change in mean sitting blood pressure at weeks 2, 4, and 8; overall blood pressure control rate at week 8 (mean sitting SBP/DBP <math>&lt; 140/90</math> mm Hg)</p>	<p>Primary:</p> <p>At week four, reductions in mean sitting SBP were significantly greater in patients receiving amlodipine and valsartan (30.1 mm Hg) than in those receiving amlodipine (23.5 mm Hg; <math>P&lt;0.0001</math>).</p> <p>At week four, mean sitting SBP reductions in patients with baseline mean sitting SBP <math>\geq 180</math> mm Hg were greater for amlodipine and valsartan (40.1 mm Hg) than for those receiving amlodipine (-31.7 mm Hg; <math>P=0.0018</math>).</p> <p>Secondary:</p> <p>At week four, reductions in mean sitting DBP were significantly greater in patients receiving amlodipine and valsartan (12.5 mm Hg) than in those receiving amlodipine (8.6 mm Hg; <math>P&lt;0.0001</math>) and all other time points (data not provided).</p> <p>At week four, 45.3% of patients were controlled on amlodipine and valsartan compared to 23.8% on amlodipine monotherapy. At week eight, corresponding control rates were 53.0 and 31.1%, respectively (<math>P&lt;0.0001</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
could be added at week 4 if mean sitting SBP was $\geq 130$ mm Hg.				
<p>Flack et al.<sup>117</sup> (2009) EX-STAND</p> <p>Amlodipine and valsartan 5-160 mg QD for 2 weeks, followed by 10-160 mg QD for 10 weeks</p> <p>vs</p> <p>amlodipine 5 mg QD for 2 weeks, then 10 mg QD for 10 weeks</p> <p>If SBP was <math>\geq 130</math> mm Hg at week 4, amlodipine and valsartan could be titrated to 10-320 mg dose. At week 8, HCTZ 12.5 mg was optionally added to both amlodipine and valsartan and amlodipine if SBP <math>\geq 130</math> mm Hg.</p>	<p>AC, DB, MC, RCT</p> <p>African American patients <math>\geq 18</math> years of age with stage 2 HTN (mean sitting SBP <math>\geq 160</math> and <math>&lt; 200</math> mm Hg)</p>	<p>N=572</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting SBP from baseline to week 8</p> <p>Secondary: Change in mean sitting SBP from baseline to week 8; change from baseline in mean sitting SBP and DBP after 2, 4, 8 and 12 weeks of treatment; blood pressure control (<math>&lt; 140/90</math> mmHg) after 12 weeks of therapy</p>	<p>Primary: At week eight, treatment with amlodipine and valsartan significantly decreased mean sitting SBP (33.3 mm Hg) compared to amlodipine monotherapy (26.6 mm Hg; <math>P &lt; 0.0001</math>).</p> <p>Secondary: Amlodipine and valsartan produced significantly greater reductions in mean sitting DBP from baseline compared to amlodipine monotherapy throughout the study: week two (9.7 vs 6.9 mm Hg; <math>P = 0.0001</math>), week four (13.2 vs 10.7 mm Hg; <math>P = 0.0008</math>), week eight (14.0 vs 11.2 mm Hg; <math>P = 0.0002</math>), and week 12 (16.1 vs 12.8 mm Hg; <math>P &lt; 0.0001</math>).</p> <p>At week eight, 49.8% of patients in the amlodipine and valsartan group and 30.2% in the amlodipine monotherapy group had their blood pressure controlled to <math>&lt; 140/90</math> mm Hg (OR, 2.4; <math>P &lt; 0.0001</math>). At week 12, 57.2% of patients in the amlodipine and valsartan group and 35.9% in the amlodipine monotherapy group attained blood pressure <math>&lt; 140/90</math> mm Hg (OR, 2.5; <math>P &lt; 0.0001</math>).</p>
<p>Schrader et al.<sup>118</sup> (2009)</p> <p>Amlodipine and</p>	<p>DB, MC, RCT</p> <p>Hypertensive patients who were</p>	<p>N=1,183</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting systolic SBP</p>	<p>Primary: At week eight, there was a greater reduction in mean sitting SBP with amlodipine and valsartan (-8.01 mm Hg) than with amlodipine (-5.95 mm Hg; <math>P &lt; 0.001</math> for non-inferiority and <math>P = 0.002</math> for superiority).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>valsartan 5-160 mg QD for 12 weeks (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 10 mg QD for 8 weeks, followed by amlodipine and valsartan 5-160 mg QD for 4 weeks (fixed-dose combination product)</p>	<p>≥55 years of age with mean sitting SBP ≥130 and ≤160 mm Hg who were inadequately controlled on amlodipine 5 mg for 4 weeks</p>		<p>Secondary: Change in mean sitting SBP and DBP, SBP control rate (mean sitting SBP &lt;130 mm Hg), overall blood pressure control rate (blood pressure &lt;140/90 mm Hg for nondiabetic patients and &lt;130/80 mm Hg for diabetic patients), and SBP response (mean sitting SBP &lt;130 mm Hg or ≥20 mm Hg reduction from baseline)</p>	<p>Secondary: Non-inferiority was also observed at week four (-8.29 vs -6.29; P&lt;0.001) and week eight (-8.23 vs -6.13; P&lt;0.001) in mean sitting SBP, at week 4 (-5.02 vs -4.23; P&lt;0.001) and week eight (-4.70 vs -4.06; P&lt;0.001) in mean sitting DBP, and at week 12 after the switch from amlodipine to amlodipine and valsartan (-9.13 vs -8.16; P&lt;0.001 for mean sitting SBP and -5.52 vs -4.90; P&lt;0.001 for mean sitting DBP).</p> <p>Systolic control with amlodipine and valsartan was greater than with amlodipine at week four (34.98 vs 24.83%; P&lt;0.001) and week eight (34.28 vs 26.21%; P=0.019), and similar after the switch from amlodipine 10 mg to amlodipine and valsartan at week 12 (38.04 vs 31.81%; P=0.162).</p> <p>SBP response rates were higher with amlodipine and valsartan than with amlodipine at week four (37.20 vs 26.72%, P&lt;0.001) and week eight [36.57 vs 27.77%; P=0.009], and similar after the switch from amlodipine to amlodipine and valsartan at week 12 (40.36 vs 35.76%; P=0.347).</p> <p>The incidence of peripheral edema was significantly lower with amlodipine and valsartan than with amlodipine (6.6 vs 31.1%, P&lt;0.001). Peripheral edema resolved in 56% patients who switched from amlodipine and valsartan without the loss of effect on blood pressure reduction.</p>
<p>Sinkiewicz et al.<sup>119</sup> (2009)</p> <p>Amlodipine and valsartan 10-160 mg or 5-160 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>valsartan 160 mg</p>	<p>AC, DB, MC, RCT</p> <p>Patients ≥18 years of age with essential HTN (mean sitting DBP ≥90 mm Hg and &lt;110 mm Hg) who were inadequately controlled on valsartan 160 mg</p>	<p>N=947</p> <p>8 weeks</p>	<p>Primary: Change from baseline in mean DBP</p> <p>Secondary: Change from baseline in mean sitting SBP, responder rate (mean DBP &lt;90 mm Hg or ≥10 mm Hg reduction from</p>	<p>Primary: At week eight, a significantly greater reduction in mean DBP was observed with both amlodipine and valsartan combinations (10-160 mg: -11.5 mm Hg, 5-160 mg: -9.6 mm Hg; P&lt;0.0001 for both) compared to valsartan monotherapy (-6.7 mm Hg).</p> <p>Secondary: At week eight, a significantly greater reduction in mean SBP was observed in both amlodipine and valsartan combinations (10-160 mg: -14.3 mm Hg, 5-160 mg: -12.2 mm Hg; P&lt;0.0001 for both) compared to valsartan monotherapy (-8.3 mm Hg).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
QD			baseline), and DBP control rate (mean DBP < 90 mm Hg)	<p>Overall mean SBP/DBP reductions of 22.5/15.5 and 21.3/13.7 mm Hg were observed in the amlodipine and valsartan 10-160 and 5-160 mg treatment groups, respectively compared to 16.7/11.4 mm Hg in the valsartan 160 mg group. The amlodipine and valsartan 10-160 mg combination showed a significantly greater reduction in mean SBP/DBP compared to amlodipine and valsartan 5-160 mg (P&lt;0.001).</p> <p>Responder rates were higher in both amlodipine and valsartan groups (10-160 mg: 81% [P&lt;0.0001]; 5-160 mg: 68% [P=0.0018], respectively) compared to valsartan monotherapy (57%).</p> <p>Peripheral edema was the most frequent adverse event, which was reported in 9.1% of patients receiving amlodipine and valsartan (10-160 mg), 0.9% of patients receiving amlodipine and valsartan (5-160 mg), and 1.3% of patients receiving valsartan monotherapy.</p>
<p>Fogari et al.<sup>120</sup> (2009)</p> <p>Amlodipine and valsartan 5 to 10-160 mg/day (fixed-dose combination)</p> <p>vs</p> <p>irbesartan and HCTZ 300-12.5 to 25 mg/day (fixed-dose combination product)</p>	<p>Blind end endpoint, OL, PG, PRO, RCT</p> <p>Patients 75 to 89 years of age with moderate essential HTN (SBP ≥160, DBP &gt;95 to &lt;110 mm Hg)</p>	<p>N=94</p> <p>24 weeks</p>	<p>Primary: Proportion of patients achieving DBP &lt;90 mm Hg</p> <p>Secondary: Changes in ambulatory blood pressure, lying and standing changes in blood pressure, safety</p>	<p>Primary: The proportion of patients receiving valsartan and amlodipine and irbesartan and HCTZ who achieved blood pressure &lt;140/&lt;90 mm Hg was 82.9 and 85.1% (P value not significant between groups).</p> <p>Secondary: Both treatment combinations resulted in a significant decrease in ambulatory blood pressure without any differences between treatment groups (P&lt;0.001 from baseline, P&gt;0.05 between groups).</p> <p>Results were similar between groups for lying SBP/DBP but patients receiving irbesartan and HCTZ experienced greater changes in ambulatory blood pressure than those receiving valsartan and amlodipine (17.2/9.0 vs 10.1/1.9 mm Hg; P&lt;0.05 for SBP and P&lt;0.01 for DBP).</p> <p>Changes from baseline in serum potassium (decrease) and uric acid (increase) were significant for those receiving irbesartan and HCTZ, but not valsartan and amlodipine (P&lt;0.05 for irbesartan and HCTZ).</p>
<p>Poldermans et al.<sup>121</sup> (2007)</p> <p>Amlodipine 5 to 10</p>	<p>AC, DB, MC, PG, RCT</p> <p>Males and females,</p>	<p>N=130</p> <p>6 weeks</p>	<p>Primary: Safety/adverse events, vital signs, hematology,</p>	<p>Primary: Both treatments were well tolerated, 26 (40.6%) of patients receiving amlodipine and valsartan and 21 (31.8%) of patients receiving lisinopril and HCTZ reported an adverse events and most were not considered drug</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD and valsartan 160 mg QD</p> <p>vs</p> <p>lisinopril 10 to 20 mg and HCTZ 12.5 mg QD</p>	<p>ages 18 years and older with HTN (mean DBP <math>\geq</math>110 mm Hg and <math>&lt;</math>120 mm Hg)</p>		<p>biochemistry variables</p> <p>Secondary: Efficacy (mean DBP, response rate, proportion of patients with mean DBP <math>&lt;</math>90 mm Hg or a <math>\geq</math>10 mm Hg reduction from baseline)</p>	<p>related.</p> <p>Peripheral edema was reported more often in the amlodipine and valsartan group than the lisinopril and HCTZ group (7.7 vs 1.5%) and cough was reported less often in the amlodipine and valsartan group than the receiving lisinopril and hydrochlorothiazide group (1.6 vs 3.0%).</p> <p>No difference was found between the treatments in changes in laboratory values or biochemistry variables.</p> <p>Secondary: Both treatments led to a reduction in mean SBP and DBP (<math>P &lt; 0.0001</math> for both from baseline) but were not significantly different from each other. Mean blood pressure for each group at study end: amlodipine and valsartan 135.0/83.6 mm Hg and lisinopril and HCTZ 138.7/85.2 mm Hg.</p> <p>The response rate was similar among the groups (100 vs 95.5%; <math>P</math> value not significant).</p>
<p>Calhoun et al.<sup>122</sup> (2009)</p> <p>Amlodipine and valsartan and HCTZ 10-320-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>valsartan and HCTZ 320-25 mg QD (fixed-dose combination product)</p> <p>vs</p>	<p>DB, MC, RCT</p> <p>Patients 18 to 85 years of age with moderate to severe essential HTN</p>	<p>N=2,271</p> <p>8 weeks</p>	<p>Primary: Difference in mean sitting diastolic blood pressure and mean sitting systolic blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: At each assessment after week three, a significantly greater proportion of patients receiving triple therapy achieved overall blood pressure control (<math>&lt;</math>140/90 mm Hg) compared to those receiving any of the dual therapies (<math>P &lt; 0.0001</math> for all).</p> <p>At end point, 70.8% of patients in the triple therapy group achieved control, compared to 48.3% for valsartan and HCTZ, 54.1% for amlodipine and valsartan, and 44.8% for amlodipine and HCTZ (<math>P &lt; 0.0001</math> for all).</p> <p>Triple therapy improved blood pressure control significantly better than any of the dual therapies.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amlodipine and valsartan 10-320 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and HCTZ 10-25 mg QD (fixed-dose combination product)</p>				
<p>Calhoun et al.<sup>123</sup> (2009)</p> <p>Amlodipine and valsartan and HCTZ 10-320-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>valsartan and HCTZ 320-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and valsartan 10-320 mg QD (fixed-dose</p>	<p>Secondary analysis</p> <p>Patients 18 to 85 years of age with moderate to severe HTN (mean SBP/DBP <math>\geq 145/\geq 100</math> mm Hg)</p>	<p>N=2,271</p> <p>8 weeks</p>	<p>Primary: Proportion and mean SBP of patients with mean SBP reductions <math>\geq 60</math>, <math>\geq 50</math>, <math>\geq 40</math>, <math>\geq 30</math> and <math>\geq 20</math> mm Hg at week three and at the end of the study</p> <p>Secondary: Changes from baseline in mean SBP based upon baseline severity, SBP control rates, safety</p>	<p>Primary: The proportion of patients with mean SBP reductions <math>\geq 20</math> mm Hg was greater with triple therapy than dual therapy at week three (74.5 vs 58.8 to 65.5%) and at study endpoint (87.6 vs 75.8 to 81.5%).</p> <p>More patients who received triple therapy, as compared to dual therapy, achieved mean SBP reductions of <math>\geq 30</math>, <math>\geq 40</math>, <math>\geq 50</math> and <math>\geq 60</math> mm Hg at week three and at study endpoint (P value not reported).</p> <p>In patients with severe SBP (<math>\geq 180</math> mm Hg), triple therapy resulted in significantly greater reductions than those for each dual therapy at week three (P&lt;0.01), except for amlodipine/valsartan (P=0.11).</p> <p>Secondary: Patients with higher baseline mean SBP had greater reductions in mean SBP than those with lower baseline mean SBP. Changes in mean SBP were significantly greater for triple therapy than dual therapy for all baseline SBP (P&lt;0.05), except for valsartan and HCTZ and amlodipine and HCTZ in patients with baseline mean SBP 150 to &lt;160 mm Hg (P value not reported).</p> <p>Significantly more patients (91.8%) receiving triple therapy achieved SBP control (<math>\geq 20</math> mm Hg reduction or mean SBP &lt;140 mm Hg) compared to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
combination product)  vs  amlodipine and HCTZ 10-25 mg QD (fixed-dose combination product)				those receiving amlodipine and HCTZ (80.1%), valsartan and HCTZ (80.8%) or valsartan and amlodipine (85.7%) (P<0.01 for all).  The overall incidence of adverse events was comparable across treatments, regardless of baseline blood pressure severity.
Pareek et al. <sup>124</sup> (2010)  Amlodipine 2.5 to 5 mg and atenolol 25 to 50 mg QD  vs  atenolol 25 to 50 mg QD	AC, MC, OL, RCT  Adults with either untreated or pretreated essential HTN	N=190  12 weeks	Primary: Change in SBP and DBP  Secondary: Not reported	Primary: At the end of four weeks, the mean change in SBP (-30.0±10.4 vs -25.08±9.05; P=0.008) and DBP (-18.10±7.45 vs -14.78±7.48; P=0.021) was significantly greater in the low-dose combination therapy as compared to the low-dose monotherapy.  At the end of 12 weeks, the mean SBP (127.82±8.90 vs 138.0±14.4; P=0.001) and mean DBP (81.73±8.78 vs 87.35±5.50; P=0.011) were significantly lower in the high-dose combination group as compared to the high-dose monotherapy group.  Secondary: Not reported
Gustin et al. <sup>125</sup> (1996)  Felodipine 5 to 10 mg QD  vs  nifedipine 30 to 60 mg QD	XO  Patients with HTN, stable on nifedipine for ≥3 months were switched to felodipine	N=127  2 months	Primary: Blood pressure  Secondary: Side effects and use of supplemental antihypertensive agents	Primary: There was no difference in SBP before and after switching agents. However, there was a difference in DBP, which was slightly lower (-2±2 mm Hg) with felodipine than with nifedipine treatment (P<0.05).  Secondary: Reported adverse events by patients and providers did not differ between the agents, with the most commonly reported side effect for both groups being leg swelling/edema.  There was no difference in use of supplemental antihypertensive agents and heart rate between treatments (P>0.05 for both).
Karotsis et al. <sup>126</sup> (2006)	RCT	N=211	Primary: Blood pressure	Primary: There was a significant decline in both office and home SBP and DBP



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Felodipine 5 mg QD vs lisinopril 10 mg QD vs chlorthalidone 12.5 mg QD vs valsartan 80 mg QD</p> <p>All patients also received diltiazem 240 mg QD.</p>	<p>Patients 25 to 79 years of age with uncontrolled HTN (average office blood pressure &gt;140/90 mm Hg for all or &gt;153/85 mm Hg for diabetics or patients &lt;65 years of age, confirmed on 2 office visits <math>\geq</math>1 week apart) after <math>\geq</math>4 weeks of OL monotherapy with diltiazem at 240 mg QD</p>	<p>8 weeks</p>	<p>Secondary: Not reported</p>	<p>during the trial with all treatments. The antihypertensive effect was more pronounced and reached significance when home blood pressure monitoring was used in comparison to office blood pressure without the white-coat effect (P&lt;0.001 for all blood pressure changes). With or without the white-coat effect, blood pressure still declined and the differences were significant (P&lt;0.0001 for all blood pressure changes).</p> <p>Secondary: Not reported</p>
<p>Manyemba et al.<sup>127</sup> (1997)</p> <p>reserpine 0.25 mg QD plus HCTZ 25 mg QD vs nifedipine SR 20 mg BID plus HCTZ 25 mg QD plus</p>	<p>OL, RCT, XO</p> <p>African American patients aged 21 to 65 years with HTN (blood pressure &gt;140/95 mm Hg) after 4 weeks of daily HCTZ therapy</p>	<p>N=32</p> <p>10 weeks</p>	<p>Primary: The change in blood pressure from baseline to the end of each 4-week treatment period</p> <p>Secondary: Not reported</p>	<p>Primary: Reserpine reduced SBP by 15.9 mm Hg (95% CI, 8.4 to 23.4) and DBP by 11.1 mm Hg (95% CI, 7.5 to 14.6).</p> <p>Nifedipine SR reduced SBP by 18.9 mm Hg (95% CI, 12.1 to 25.7) and DBP by 9.6 mm Hg (95% CI, 7.2 to 12.0).</p> <p>There was no significant difference between the two groups.</p> <p>Secondary: Not reported</p>
<p>Lindholm et al.<sup>128</sup> (2005)</p> <p>Other antihypertensive therapies (amiloride,</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a <math>\beta</math>-blocker as first-</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary: The RR of stroke was 16% higher with <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non <math>\beta</math>-blockers (RR, 1.26%; 95% CI, 15 to 38; P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)</p> <p>or</p> <p>placebo</p> <p>vs</p> <p>β-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol, or propranolol)</p>	<p>line treatment (in ≥50% of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>			<p>The relative risk of MI was 2% higher for β-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).</p> <p>The RR of all-cause mortality was 3% higher for β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).</p> <p>Secondary: Not reported</p>
<p>Van Bortel et al.<sup>129</sup> (2008)</p> <p>ACE inhibitor, ARB, β-blocker, calcium channel blocker, or placebo</p> <p>vs</p> <p>nebivolol</p>	<p>MA</p> <p>12 RCTs involving &gt;25 patients with essential HTN where nebivolol 5 mg QD was compared to placebo or other active drugs for &gt;1 month</p>	<p>N=2,653</p> <p>Duration varied</p>	<p>Primary: Antihypertensive effect and tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: Overall, higher response rates were observed with nebivolol than all other antihypertensive agents combined (OR, 1.41; 95% CI, 1.15 to 1.73; P=0.001) and compared to the ACE inhibitors (OR, 1.92; 1.30 to 2.85; P=0.001), but response rates to nebivolol were similar to β-blockers (OR, 1.29; 95% CI, 0.81 to 2.04; P=0.283), calcium channel blockers (OR, 1.19; 95% CI, 0.83 to 1.70; P=0.350) and losartan (OR, 1.35; 95% CI, 0.84 to 2.15; P=0.212).</p> <p>Overall, a higher percentage of patients obtained normalized BP with nebivolol compared to the other antihypertensive agents combined (OR, 1.35; 95% CI, 1.07 to 1.72; P=0.012). A higher percentage of patient receiving nebivolol obtained normalized BP compared to losartan (OR, 1.98; 95% CI, 1.24 to 3.15; P=0.004) and calcium channel blockers (OR, 1.96; 95% CI, 1.05 to 1.96; P=0.024), but not when compared to other β-blockers (OR, 1.29; 95% CI, 0.81 to 1.65; P=0.473).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Overall, the percentage of adverse events was significantly lower with nebivolol compared to the other antihypertensive agents combined (OR, 0.59; 95% CI, 0.48 to 0.72; P&lt;0.001) and similar to placebo (OR, 1.16; 95% CI, 0.76 to 1.67; P=0.482). In comparing nebivolol to the individual treatments, nebivolol had a lower percentage of adverse events compared to losartan (OR, 0.52; 95% CI, 0.30 to 0.89; P=0.016), the other <math>\beta</math>-blockers (OR, 0.56; 95% CI, 0.36 to 0.85; P=0.007) and calcium channel blockers (OR, 0.49; 95% CI 0.33 to 0.72; P&lt;0.001), but was similar to ACE inhibitors (OR, 0.75; 95% CI 0.52 to 1.08).</p> <p>Secondary: Not reported</p>
<p>Wiysonge et al.<sup>130</sup> (2007)</p> <p>Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)</p> <p>vs</p> <p><math>\beta</math>-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)</p>	<p>MA</p> <p>13 RCTs evaluating patients <math>\geq 18</math> years of age with HTN</p>	<p>N=91,561</p> <p>Duration varied</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions</p>	<p>Primary: There was not a significant difference observed in all-cause mortality between <math>\beta</math>-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with <math>\beta</math>-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).</p> <p>Secondary: There was a significant decrease in stroke observed with <math>\beta</math>-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with <math>\beta</math>-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between <math>\beta</math>-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with <math>\beta</math>-blocker therapy</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of <math>\beta</math>-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with <math>\beta</math>-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>
<p>Baguet et al.<sup>131</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)</p> <p>Drugs were used as monotherapy, either at a fixed daily</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the <math>\beta</math>-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).</p> <p>The weighted average reduction of SBP and DBP for each drug class were as follows: Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively. <math>\beta</math>-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively. Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively. ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively. Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary: Not reported</p>
<p>Laurent et al.<sup>132</sup> (2018)</p> <p>Perindopril 3.5 mg/amlodipine 2.5 mg</p> <p>vs</p> <p>RAS-inhibitor monotherapy (perindopril 5 mg, irbesartan 150 mg, or valsartan 80 mg)</p>	<p>MA</p> <p>Patients &gt;18 years of age with essential HTN (SBP <math>\geq</math>140 mm Hg and/or DBP <math>\geq</math>90 mm Hg) with an SBP assessment at baseline and at month one</p>	<p>N=5,496</p> <p>two to nine months</p>	<p>Primary: Change in SBP and DBP from baseline after one month of treatment</p> <p>Secondary: Emergent adverse events</p>	<p>Primary: <u>Perindopril/amlodipine versus perindopril</u> Perindopril/amlodipine reduced SBP by 20.3 mm Hg and perindopril reduced SBP by 16.9 mm Hg (P=0.009).</p> <p>Perindopril/amlodipine reduced DBP by 11.9 mm Hg and perindopril reduced DBP by 10.2 mm Hg (P=0.018).</p> <p><u>Perindopril/amlodipine versus irbesartan</u> Perindopril/amlodipine reduced SBP by 14.1 mm Hg and irbesartan reduced SBP by 12.7 mm Hg (P=0.003).</p> <p>Perindopril/amlodipine reduced DBP by 5.8 mm Hg and irbesartan reduced DBP by 5.2 mm Hg (P=0.008).</p> <p><u>Perindopril/amlodipine versus valsartan</u> Perindopril/amlodipine reduced SBP by 18 mm Hg and valsartan reduced SBP by 14.6 Hg (P&lt;0.001).</p> <p>Perindopril/amlodipine reduced DBP by 13 mm Hg and valsartan reduced DBP by 11.2 mm Hg (P&lt;0.001).</p> <p>There was a significant difference observed in the estimated treatment difference for SBP between perindopril/amlodipine and RAS-inhibitor</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>monotherapy (estimated treatment difference, -2.36; 95% CI, -2.36 to -0.89; P=0.002).</p> <p>There was a significant difference observed in the estimated treatment difference for DBP between perindopril/amlodipine and RAS-inhibitor monotherapy (estimated treatment difference, -1.25; 95% CI, -2.12 to -0.38; P=0.005).</p> <p>Secondary: The proportion of patients who experienced emergent adverse events was similar for both perindopril/amlodipine (28.4%) and RAS-inhibitor monotherapy (28.2%) groups (P=0.929).</p>
<b>Renal Effects</b>				
<p>Esnault et al.<sup>133</sup>(2008)</p> <p>Amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>enalapril 5 to 20 mg/day</p>	<p>MC, DB, PC, RCT</p> <p>Nondiabetic, adult patients with estimated creatinine clearance of 20 to 60 ml/min</p>	<p>N=263</p> <p>3 years</p>	<p>Primary: Change in GFR measured yearly by blood clearance</p> <p>Secondary: Composite of renal events and tolerability</p>	<p>Primary: No statistically significant difference was found between amlodipine and enalapril in GFR decline (-4.92 and -3.98 mL/min., respectively, at last observation).</p> <p>Secondary: No statistically significant difference was found between amlodipine and enalapril in the composite secondary end point after a median follow-up of 2.9 years, including in the subgroup of patients with proteinuria &gt;1 g/d at baseline.</p>
<p>Agodoa et al.<sup>134</sup> (2001)</p> <p>AASK</p> <p>Amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>ramipril 2.5 to 10 mg QD</p>	<p>DB, MC, RCT</p> <p>African American patients, age 18 to 70 years old, with hypertensive renal disease (GFR 20 to 65 mL/min)</p>	<p>N=1,094</p> <p>4 years</p>	<p>Primary: Rate of change in GFR (GFR slope)</p> <p>Secondary: Composite of: confirmed reduction GFR by 50% or by 25 mL/min for baseline, ESRD</p>	<p>Primary: The average decline in GFR was slower, by 36% in the ramipril group as compared to the amlodipine group (P=0.002). However, during the first three months, GFR increased more in the amlodipine group than the ramipril group (P&lt;0.001). The mean total slope did not differ between the groups (P=0.38).</p> <p>Secondary: The risk reduction for the composite secondary outcome was significantly greater for the ramipril group than the amlodipine group (P=0.005). The rate of ESRD was significantly lower in the ramipril group (P=0.01).</p>
<p>Wright et al.<sup>135</sup> (2002)</p> <p>AASK</p>	<p>DB, MC, RCT</p> <p>Patients were self-</p>	<p>N=1,094</p> <p>3 to 6.4 years</p>	<p>Primary: Rate of change in GFR (grouped by</p>	<p>Primary: No significant difference in primary outcome was reported between the usual blood pressure group compared to the lower blood pressure group</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Amlodipine 5 to 10 mg/day vs metoprolol 50 to 200 mg/day vs ramipril 2.5 to 10 mg/day</p>	<p>identified African Americans aged 18 to 70 years with HTN and a GFR between 20 and 65 mL/min/ 1.73 m<sup>2</sup> and no other identified cause of renal insufficiency</p>		<p>usual blood pressure [MAP goal 102 to 107 mm Hg] vs lower blood pressure [<math>\leq</math>92 mm Hg])  Secondary: Clinical composite outcome (reduction in GFR by 50% or more, ESRD, or death)</p>	<p>(P=0.24).  None of the drug group comparisons showed consistently significant differences in the GFR slope.  Secondary: The lower blood pressure goal did not significantly reduce the rate of the clinical composite outcome (risk reduction for lower blood pressure group, 2%; 95% CI, -22 to 21; P=0.85).  Ramipril resulted in significant risk reductions in the clinical composite outcomes compared to amlodipine (38%; 95% CI, 14 to 56; P=0.004) and metoprolol (22%; 95% CI, 1 to 38; P=0.04).  There was no significant difference in the clinical composite outcome between the amlodipine and metoprolol groups.</p>
<p>Lewis et al.<sup>136</sup> (2001) IDNT  Amlodipine 10 mg/day vs irbesartan 300 mg/day vs placebo</p>	<p>DB, MC, PC, PRO, RCT  Patients 30 to 70 years old, with type 2 diabetes mellitus, HTN, and nephropathy</p>	<p>N=1,715  2.6 years</p>	<p>Primary: Composite of risk of doubling serum creatinine, ESRD, or death from any cause  Secondary: Composite of death from cardiovascular causes, nonfatal MI, heart failure requiring hospitalization, permanent neurologic deficit caused by a cerebrovascular event, or lower limb amputation</p>	<p>Primary: Compared to placebo, irbesartan 300 mg/day resulted in a 20% lower relative risk of the composite primary outcome (P=0.02). Irbesartan treatment was associated with a 33% lower risk of doubling serum creatinine (P=0.003) and 23% trend towards lower risk of ESRD (P=0.07) compared to placebo. There was no significant difference in risk of death from any cause for irbesartan compared to placebo (P=0.57).  Compared to amlodipine, irbesartan treatment resulted in a 23% lower risk of composite primary outcome (P=0.006). Irbesartan treatment was associated with a 37% lower risk of doubling serum creatinine vs amlodipine (P&lt;0.001) and 23% trend towards lower risk of ESRD vs amlodipine (P=0.07). There was no significant difference in risk of death from any cause (P=0.80).  Secondary: There were no significant differences in the secondary cardiovascular composite end point (P=0.40 and P=0.79 for irbesartan vs placebo and amlodipine, respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Viberti et al.<sup>137</sup> (2002) MARVAL</p> <p>Amlodipine 5 mg QD</p> <p>vs</p> <p>valsartan 80 mg QD</p> <p>A target blood pressure of 135/85 mm Hg was aimed for by dose-doubling followed by the addition of bendrofluzide* and doxazosin whenever needed.</p>	<p>AC, DB, RCT</p> <p>Patients 35 to 75 years old with type 2 diabetes mellitus and microalbuminuria, with or without HTN</p>	<p>N=332</p> <p>24 weeks</p>	<p>Primary: Change in UAER; proportion of patients who returned to normal albuminuria</p> <p>Secondary: Proportion of patients returning to normoalbuminuria</p>	<p>Primary: Valsartan resulted in a UAER reduction of 44% at 24 weeks compared to baseline vs an 8% reduction with amlodipine (P&lt;0.001). Valsartan lowered UAER similarly in both the hypertensive and normotensive groups.</p> <p>Over the study period, blood pressure reductions were similar between the two treatments and at no time point was there a between-group significant difference in blood pressure values in either the hypertensive or the normotensive subgroup.</p> <p>Secondary: The proportion of patients returning to normal albuminuria was greater with valsartan (29.9%) vs amlodipine (14.5%; P=0.001).</p>
<p>Bakris et al.<sup>138</sup> (2008) GUARD</p> <p>Amlodipine and benazepril (fixed-dose combination product)</p> <p>vs</p> <p>benazepril and HCTZ (fixed-dose combination product)</p>	<p>DB, RCT</p> <p>Hypertensive, albuminuric type 2 diabetic patients, mean age 58 years were randomized to receive either initial fixed-dose combination product</p>	<p>N=322</p> <p>52 weeks</p>	<p>Primary: Change in urinary albumin to creatinine ratio after 1 year of initial treatment with either fixed-dose combination, blood pressure reductions</p> <p>Secondary: Proportion who progressed to overt diabetic nephropathy, safety</p>	<p>Primary: Both combinations significantly reduced the urinary albumin to creatinine ratio compared to baseline (P&lt;0.0001). The median percent change was -72.1% for benazepril and HCTZ and -40.5% for amlodipine and benazepril (P&lt;0.0001).</p> <p>Both regimens significantly reduced SBP and DBP compared to baseline (P&lt;0.0001). The mean reduction in both SBP and DBP was greater in the amlodipine-based arm than in the HCTZ-based arm; however, significance in favor of the amlodipine regimen was observed only for DBP (SBP -20.5 vs -18.8; P=0.19; DPB -13.1 vs -9.97; P=0.02).</p> <p>A greater proportion of patients who had microalbuminuria at baseline and treated with benazepril and HCTZ compared to amlodipine and benazepril attained normalization of the urinary albumin to creatinine ratio, defined as &lt;30 mg/g (69.2 vs 47.8%; P=0.0004).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Secondary: The percentage of patients progressing to overt proteinuria was similar for both groups.</p> <p>Overall, both study drugs were well tolerated. Adverse reactions possibly related to the study medications occurred in 11.4 and 3.6% of patients receiving amlodipine and benazepril and benazepril and HCTZ, respectively. They included peripheral edema (7.8 vs 2.4%, respectively), fatigue (1.2% in each group), pitting edema (1.2 vs 0.0%), face edema (0.6 vs 0.0%) and thirst (0.6 vs 0.0%). More patients receiving the HCTZ-based regimen (10.8%) discontinued study drug than with the amlodipine-based regimen due to side effects (5.4%).</p>
<p>Casas et al.<sup>139</sup> (2005)</p> <p>ACE inhibitor or ARBs compared to placebo</p> <p>vs</p> <p>ACE inhibitor or ARBs compared to other antihypertensive drugs (<math>\beta</math>-adrenergic blocking agents, <math>\alpha</math>-adrenergic blocking agents, calcium-channel blocking agents, or combinations)</p> <p>Specific agents and doses were not specified.</p>	<p>MA (127 trials)</p> <p>Studies in adults that examined the effect of any drug treatment with a blood pressure lowering action on progression of renal disease</p>	<p>N=not reported</p> <p>4.2 years (mean)</p>	<p>Primary: Doubling of serum creatinine, and ESRD</p> <p>Secondary: Serum creatinine, urine albumin excretion and GFR</p>	<p>Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>Secondary: Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).</p> <p>Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).</p> <p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Miscellaneous</b>				
Rosendorff et al. <sup>140</sup> (2009)  Amlodipine 5 to 10 mg QD  vs  olmesartan 20 to 40 mg QD	AC, DB, RCT  Adults with HTN and left ventricular hypertrophy	N=102  52 weeks	Primary: Change in left ventricular mass from baseline to 52 weeks  Secondary: Change in left ventricular mass after 26 weeks of treatment	Primary: Mean±SD left ventricular masses of 252.9±73.06 g in the olmesartan group and 236.9±59.94 g in the amlodipine group at baseline were decreased to 248.2±69.31 and 223.9±53.18 g, respectively, after 52 weeks of therapy. Neither of these changes was significantly different from baseline, and the difference between the two treatment groups was not significant.  Secondary: At 26 weeks, adjusted percent changes in left ventricular mass were 8.0% with olmesartan and 6.0% with amlodipine. Changes occurring at the 26-week assessment were not significantly different from baseline or from each other.
Luscher et al. <sup>141</sup> (2009) ENCORE II  Nifedipine 30 to 60 mg QD  vs  placebo	DB, MC, PC, RCT  Adults undergoing coronary angiography with or without PCI	N=226  18 to 24 months	Primary: The effect of nifedipine compared to placebo on acetylcholine-induced coronary vascular response at the highest dose of acetylcholine at baseline and follow-up  Secondary: Effect of nifedipine on the percent change in plaque volume as assessed by intravascular ultrasound	Primary: The change in mean luminal diameter averaged 13.9±16.5% with nifedipine and 7.7±18% with placebo. The difference between groups was 6.3% (95% CI, 1.6 to 10.9; P=0.0088).  Secondary: Neither the difference in absolute nor relative changes in mean plaque volume as measure by intravascular ultrasound between treatments was significant (P=0.84 and 0.66, respectively).
Schmid-Elsaesser et al. <sup>142</sup> (2006)	RCT  Patients with	N=104  7 days	Primary: Incidence of clinical vasospasm	Primary: There was no significant difference between the groups in number of patients experiencing clinical vasospasm or transcranial

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Nimodipine continuous infusion of 1 mg/hr for 6 hours, followed by 2.0 mg/hr</p> <p>Vs</p> <p>magnesium sulfate bolus infusion 10 mg/kg, followed by continuous infusion of 30 mg/kg QD</p>	<p>aneurismal subarachnoid hemorrhage</p>		<p>and transcranial Doppler angiographic vasospasm, and infarction attributable to vasospasm</p> <p>Secondary: Incidence of angiographic vasospasm</p>	<p>doppler/angiographic vasospasm: 14 patients (27%) in the nimodipine group vs eight patients (15%) in the magnesium group (P=0.193); 17 (33%) in the nimodipine group vs 20 (38%) in the magnesium group (P=0.792).</p> <p>No difference between the groups was found in incidence of cerebral infarction, 11 (22%) in the nimodipine group vs 10 (19%) in the magnesium group.</p> <p>Secondary: There were no significant differences in incidence of angiographic vasospasm, neuronal markers or Glasgow outcome scores (all values: P&gt;0.05).</p>
<p>Liu et al.<sup>143</sup> (abstract) (2011)</p> <p>Nimodipine vs placebo</p>	<p>MA (8 trials)</p> <p>Patients receiving prophylactic nimodipine for aneurismal subarachnoid hemorrhage</p>	<p>N=1,514</p> <p>Not reported</p>	<p>Primary: Not reported</p> <p>Secondary: Not reported</p>	<p>Primary: Not reported</p> <p>Secondary: Not reported</p> <p>Compared to placebo, fully recovered (all cases) patients increased 64% with nimodipine (OR, 1.64; 95% CI, 1.26 to 2.13; P=0.002; NNT, -1.048), fully recovered or moderately disabled (all cases) patients increased 79% (OR, 1.79; 95% CI, 1.28 to 2.51; P=0.0007; NNT, -5.889), patient death (in cerebral vasospasm cases) decreased 74% (OR, 0.26; 95% CI, 0.09 to 0.71; P=0.008; NNT, 2.298), the incidence of symptomatic cerebral vasospasm decreased 46% (OR, 0.54; 95% CI, 0.42 to 0.69; P&lt;0.00001; NNT, 1.952), the incidence of delayed neurological function deficits (all cases) decreased 38% (OR, 0.62; 95% CI 0.50 to 0.78; P&lt;0.0001; NNT, 1.078), the occurrence of cerebral infarction (on CT scan) decreased 58% (OR, 0.58; 95% CI, 0.42 to 0.81; P=0.001; NNT, 3.314), the occurrence of cerebral infarction (in cerebral vasospasm cases) decreased 65% (OR, 0.35; 95% CI, 0.17 to 0.69; P=0.003; NNT, 3.688), and the occurrence of cerebral infarction (all cases) decreased 48% (OR, 0.52; 95% CI, 0.41 to 0.66; P&lt;0.00001; NNT, 1.196). The difference in recurrent hemorrhage and adverse reactions between the nimodipine and placebo was not statistically significant (recurrent hemorrhage: OR, 0.75; 95% CI, 0.50 to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				1.11; P=0.15; adverse reaction: OR, 1.13; 95% CI, 0.71 to 1.81; P = 0.59).  Secondary: Not reported

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, CR=controlled-release, ER=extended-release, QD=once daily, SR=sustained-release

Study design abbreviations: AC=active comparator, DB=double blind, DD=double dummy, ES=extended-release, MA=meta analysis, MC=multicenter, OL=open label, OS=observational, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, RETRO=retrospective, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitor, ABPM=ambulatory blood pressure monitoring, ARB=angiotensin II receptor blocker, BMI=body mass index, CAD=coronary artery disease, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, CT=computed tomography, DBP=diastolic blood pressure, ECG=electrocardiogram, ESRD=end stage renal disease, FBG=fasting blood glucose, GFR=glomerular filtration rate, HbA<sub>1c</sub>=glycosylated hemoglobin, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HTN=hypertension, HR=hazard ratio, LDL-C=low-density lipoprotein cholesterol, MI=myocardial infarction, MMSE=Mini Mental State Examination, NIDDM=non-insulin dependent diabetes mellitus, NNT=number needed to treat, OR=odds ratio, PAD=peripheral artery disease, PCI=percutaneous coronary intervention, PVD=peripheral vascular disease, QOL=quality of life, RAS=renin-angiotensin system, RR=relative risk, SBP=systolic blood pressure, SD=standard deviation, TC=total cholesterol, TG=triglyceride, TIA=transient ischemic attack, UAER=urinary albumin excretion rate

## **Additional Evidence**

### Dose Simplification

Taylor et al. evaluated adherence rates in patients receiving a fixed-dose combination of amlodipine and benazepril compared to patients receiving an ACE inhibitor and a long-acting dihydropyridine calcium-channel blocking agent as separate formulations. There was no significant difference in adherence in younger subjects (18 to 39 year olds); however, overall adherence was higher in patients receiving amlodipine/benazepril fixed-dose combination product compared to those receiving separate formulations (80.8 vs 73.8%;  $P < 0.001$ ).<sup>144</sup> Dickson et al. also evaluated adherence rates with the fixed-dose combination of amlodipine/benazepril compared to the administration of an ACE inhibitor and dihydropyridine calcium-channel blocking agent as separate formulations in an elderly Medicaid population. Over a 12 month period, adherence rates were higher in patients receiving the fixed-dose combination product compared to those receiving separate formulations (63.4 vs 49.0%;  $P < 0.0001$ ).<sup>145</sup> Gerbino et al. assessed adherence rates in patients receiving the fixed-dose combination of amlodipine/benazepril or an ACE inhibitor and dihydropyridine calcium-channel blocking agent administered as separate formulations. Adherence rates were 69.2% for patients who received the antihypertensive agents as separate formulations compared to 87.9% for patients receiving the fixed-dose combination product ( $P < 0.0001$ ).<sup>146</sup>

### Stable Therapy

Lenz et al. compared the 24-hour blood pressure control in patients stabilized on amlodipine who were then converted to nisoldipine. After three months, blood pressure control was similar between treatments, except for average 24-hour diastolic blood pressure, where nisoldipine treatment resulted in slightly greater readings (by 2 mm Hg).<sup>71</sup> Gustin et al. reviewed medical records of hypertensive patients who were switched from long-acting nifedipine to felodipine. This resulted in slightly lower diastolic blood pressure measurements (78 vs 80 mm Hg;  $P < 0.05$ ). Adverse events and supplemental medication use were similar between the agents.<sup>125</sup> Sapienza et al. measured the impact of converting long-term care patients previously on high dose calcium-channel blocking agents or dual therapy with an ACE inhibitor and calcium-channel blocking agents to the fixed-dose combination of amlodipine/benazepril. There was no significant change in blood pressure following the conversion; however, there was a significant reduction in the number of patients reporting  $\geq 1$  drug-related adverse event (22 vs 4;  $P < 0.05$ ).<sup>147</sup>

### Impact on Physician Visits

Sheehy et al. conducted a comparative review of patients receiving amlodipine or felodipine. The investigators found an increased number of specialist visits in the amlodipine group (odds ratio, 1.14; 95% confidence interval, 1.8 to 1.20); however, this same group of patients receiving amlodipine had significantly better compliance and refill rates and fewer medication switches.<sup>62</sup>

## **IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 15. Relative Cost of the Dihydropyridines**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Amlodipine	suspension, tablet	Katerzia <sup>®</sup> , Norvasc <sup>®</sup> *	\$\$\$\$\$	\$
Felodipine	extended-release tablet	N/A	N/A	\$\$
Isradipine	capsule	N/A	N/A	\$\$\$\$\$
Nicardipine	capsule*, injection*	N/A	N/A	\$\$\$\$\$
Nifedipine	capsule, extended-release tablet	Adalat CC <sup>®</sup> *, Procardia XL <sup>®</sup> *	\$\$\$\$\$	\$
Nimodipine	capsule*, solution	Nymalize <sup>®</sup>	\$\$\$\$\$	\$\$\$\$\$
Nisoldipine	extended-release tablet*	Sular ER <sup>®</sup> *	\$\$\$\$\$	\$\$\$\$\$
<b>Combination Products</b>				
Amlodipine and benazepril	capsule	Lotrel <sup>®</sup> *	\$\$\$\$\$	\$
Amlodipine and olmesartan	tablet	Azor <sup>®</sup> *	\$\$\$\$\$	\$
Amlodipine and valsartan	tablet	Exforge <sup>®</sup> *	\$\$\$\$\$	\$\$
Amlodipine, valsartan, and HCTZ	tablet	Exforge HCT <sup>®</sup> *	\$\$\$\$\$	\$

\*Generic is available in at least one dosage form or strength.

HCTZ=hydrochlorothiazide, N/A=not available

## X. Conclusions

All of the dihydropyridines, with the exception of nimodipine, are approved for the treatment of hypertension. Amlodipine, nicardipine, and nifedipine are also indicated for the treatment of angina. Additionally, amlodipine reduces the risk of hospitalization due to angina and reduces the risk of coronary revascularization procedures in patients with recently documented coronary artery disease.<sup>1,2,6-17</sup> Amlodipine is available in combination with benazepril, olmesartan, valsartan, or valsartan-hydrochlorothiazide. It should be noted that the amlodipine and telmisartan fixed-dose combination product and the amlodipine, olmesartan, and hydrochlorothiazide fixed-dose combination product are included in the angiotensin II receptor antagonists class review (AHFS Class 243208). All of the products with the exception of clevidipine is available in a generic formulation.

There are several national and international guidelines that provide recommendations regarding the use of calcium-channel blocking agents.<sup>18-36</sup> For the treatment of chronic angina,  $\beta$ -blockers are recommended as initial therapy; however, long-acting calcium-channel blocking agents may be used if  $\beta$ -blockers are contraindicated or if additional therapy is required.<sup>18-24</sup> Calcium-channel blocking agents are recommended as initial therapy in patients with variant/vasospastic angina.<sup>19,22</sup> For the treatment of heart failure, ACE inhibitors, ARBs, aldosterone antagonists, and isosorbide dinitrate/hydralazine are recommended as initial therapy. In general, calcium-channel blocking agents are not recommended in the management of heart failure.<sup>26-27</sup> There are several published guidelines on the treatment of hypertension. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>28-34</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High

Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>29</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>28-34,36</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>28-34,36</sup>

Numerous clinical trials have shown that the dihydropyridines can effectively lower systolic and diastolic blood pressure when administered alone or in combination with other agents. In trials comparing combination therapy to monotherapy, the more aggressive treatment regimens lowered blood pressure to a greater extent than the less-intensive treatment regimens. Some comparative trials have demonstrated slight differences in blood pressure effects among the various dihydropyridines; however, the clinical significance of these differences remains to be established.<sup>62-131</sup> Most patients will require more than one antihypertensive agent to achieve blood pressure goals.<sup>28-34</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>30,31,144-146</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations. The dihydropyridines have been shown to favorably affect cardiovascular morbidity and mortality, and several studies have demonstrated comparable efficacy with  $\beta$ -blockers, diuretics, ACE inhibitors, ARBs.<sup>39-61</sup>

There is insufficient evidence to support that one brand dihydropyridine is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand dihydropyridines within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand dihydropyridine is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. Kannam JP, Gersh BJ. Calcium channel blockers in the management of chronic coronary syndrome. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): 2021 [cited 2022 Jan].
4. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
5. Kaplan NM, Victor RG, Flynn JT. Kaplan's clinical hypertension. 11<sup>th</sup> ed. Philadelphia (PA): Lippincott, Williams, and Wilkins; 2015.
6. Norvasc® [package insert]. New York (NY): Pfizer Inc; 2019 Jan.
7. Procardia® [package insert]. New York (NY): Pfizer Inc; 2016 Jul.
8. Procardia XL® [package insert]. New York (NY): Pfizer Inc; 2021 May.
9. Adalat® CC [package insert]. Pine Brook (NJ): Almatica, Inc.; 2018 Nov.
10. Sular® [package insert]. Florham Park (NJ): Shionogi Pharma, Inc.; 2015 Mar.
11. Nymalize® [package insert]. Atlanta (GA): Arbor Pharmaceuticals, Inc.; 2019 Dec.
12. Azor® [package insert]. Basking Ridge (NJ): Daiichi-Sankyo, Inc; 2020 Dec.
13. Lotrel® [package insert]. East Hanover (NJ): Novartis Pharmaceuticals Corporation; 2021 Apr.
14. Prestalia® [package insert]. Durham (NC): Adhera Therapeutics, Inc.; 2019 Aug.
15. Exforge® [package insert]. East Hanover (NJ): Novartis Pharmaceuticals Corp; 2021 Apr.
16. Exforge HCT® [package insert]. East Hanover (NJ): Novartis Pharmaceuticals Corp; 2021 Feb.
17. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
18. Fraker T, Fihn S, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of chronic stable angina: a report of the ACC/AHA task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *Circulation*. 2007 Dec 4;116(23):2762-72.
19. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425.
20. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline from the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
21. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
22. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>.
23. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.
24. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
25. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].



26. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
27. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
28. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
29. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
30. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
31. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
32. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
33. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
34. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
35. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
36. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
37. Koenig W, Hoher M. Felodipine and amlodipine in stable angina pectoris: results of a randomized double-blind crossover trial. *J Cardiovasc Pharmacol*. 1997;29(4):520-4.
38. Savonitto S, Ardissio D, Egstrup K, Rasmussen K, Bae EA, Omland T, et al. Combination therapy with metoprolol and nifedipine versus monotherapy in patients with stable angina pectoris. Results of the International Multicenter Angina Exercise (IMAGE) Study. *J Am Coll Cardiol*. 1996 Feb;27(2):311-6.
39. Pitt B, Byington RP, Furberg CD, et al. Effect of amlodipine on the progression of atherosclerosis and the occurrence of clinical events. PREVENT Investigators. *Circulation*. 2000;102(13):1503-10.
40. Dahlöf B, Sever PS, Poulter NR, et al. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomized controlled trial. *Lancet*. 2005 Sep 10-16;366(9489):895-906.
41. Chapman N, Dobson J, Wilson S, Dahlöf B, Sever PS, Wedel H, Poulter NR; Anglo-Scandinavian Cardiac Outcomes Trial Investigators. Effect of spironolactone on blood pressure in subjects with resistant hypertension. *Hypertension*. 2007 Apr;49(4):839-45.
42. Nissen SE, Tuzcu EM, Libby P, et al; CAMELOT Investigators. Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: a randomized controlled trial. *JAMA*. 2004;292(18):2217-25.
43. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium-channel blocker vs diuretic: The

- Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA*. 2002 Dec 18;288(23):2981-97.
44. Black HR, Davis B, Barzilay J, et al; Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Metabolic and clinical outcomes in nondiabetic individuals with the metabolic syndrome assigned to chlorthalidone, amlodipine, or lisinopril as initial treatment for hypertension: a report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Diabetes Care*. 2008 Feb;31(2):353-60.
  45. Rahman M, Ford CE, Cutler JA, Davis BR, Piller LB, Whelton PK, et al. Long-term renal and cardiovascular outcomes in antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT) participants by baseline estimated GFR. *Clin J Am Soc Nephrol*. 2012;7:989-1002.
  46. Muntner P, Levitan EB, Lynch AI, et al. Effect of chlorthalidone, amlodipine, and lisinopril on visit-to-visit variability of blood pressure: results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. *J Clin Hypertens (Greenwich)*. 2014 May;16(5):323-330.
  47. Bangalore S, Davis BR, Cushman WC, Pressel SL, Muntner PM, Calhoun DA, et al. Treatment-Resistant Hypertension and Outcomes Based on Randomized Treatment Group in ALLHAT. *Am J Med*. 2017 Apr;130(4):439-448.e9.
  48. Ogihara T, Nakao K, Fukui T, et al; Candesartan Antihypertensive Survival Evaluation in Japan Trial Group. Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. *Hypertension*. 2008 Feb;51(2):393-8.
  49. Julius S, Kjeldsen SE, Weber M, Brunner HR, Ekman S, Hansson L, et al; VALUE trial group. Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomised trial. *Lancet*. 2004 Jun 19;363(9426):2022-31.
  50. Zanchetti A, Julius S, Kjeldsen S, et al. Outcomes in subgroups of hypertensive patients treated with regimens based on valsartan and amlodipine: An analysis of findings from the VALUE trial. *J Hypertens*. 2006 Nov;24(11):2163-8.
  51. Jamerson K, Weber MA, Bakris GL, et al. Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high risk patients. *N Engl J Med* 2008;359:2417-28.
  52. Bakris GL, Sarafidis PA, Weir MR, Dahlöf B, Pitt B, Jamerson K, et al.; ACCOMPLISH Trial investigators. Renal outcomes with different fixed-dose combination therapies in patients with hypertension at high risk for cardiovascular events (ACCOMPLISH): a prespecified secondary analysis of a randomized controlled trial. *Lancet*. 2010 Apr 3;375(9721):1173-81.
  53. Weber MA, Bakris GL, Jamerson K, Weir M, Kjeldsen SE, Devereux RB, et al.; ACCOMPLISH Investigators. Cardiovascular events during differing hypertension therapies in patients with diabetes. *J Am Coll Cardiol*. 2010 Jun 29;56(1):77-85.
  54. Weber MA, Jamerson K, Bakris G, Weir MR, Zappe D, Zhang Y, et al. Effects of body size and hypertension treatments on cardiovascular event rates: subanalysis of the ACCOMPLISH randomised controlled trial. *Lancet*. 2013;381:537-45.
  55. Bakris G, Briassoulis A, Dahlöf B, et al. Comparison of benazepril plus amlodipine or hydrochlorothiazide in high-risk patients with hypertension and coronary artery disease. *Am J Cardiol*. 2013 Jul 15;112(2):255-259.
  56. Hansson L, Lindholm LH, Ekblom T, et al. Randomized trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet*. 1999 Nov 20;354(9192):1751-6.
  57. Borhani NO, Mercuri M, Borhani PA, et al. Final outcome results of the Multicenter Isradipine Diuretic Atherosclerosis Study (MIDAS): a randomized controlled trial. *JAMA*. 1996; 276:785-791.
  58. Randomized double-blind comparison of a calcium antagonist and a diuretic in elderly hypertensives. National Intervention Cooperative Study in Elderly Hypertensives Study Group. *Hypertension*. 1999 Nov;34(5):1129-33.
  59. Lichtlen PR, Hugenholz PG, Rafflenbeul W, et al. Retardation of coronary artery disease in humans by the calcium-channel blocker nifedipine: results of the INTACT study (International Nifedipine Trial on Antiatherosclerotic Therapy Cardiovasc Ther. 1990 Aug;4 Suppl 5:1047-68.
  60. Brown MJ, Palmer CR, Castaigne A, et al. Morbidity and mortality in patients randomized to double-blind treatment with a long-acting calcium-channel blocker or diuretic in the international nifedipine GTS study: Intervention as a Goal in Hypertension Treatment (INSIGHT). *Lancet*. 2000; 356:366-72.
  61. Estacio RO, Schrier RW. Antihypertensive therapy in type 2 diabetes: implications of the appropriate blood pressure control in diabetes (ABCD) trial. *Am J Cardiol*. 1998;82(9B):9R-14R.
  62. Sheehy O, LeLorier J. Patterns of amlodipine and felodipine use in an elderly Quebec population. *Can J Cardiol*. 2000;16(9):1109-17.

63. Van Der Krogt J, Brand R, Dawson E. Amlodipine versus extended-release felodipine in general practice: A randomized, parallel-group study in patients with mild-to-moderate hypertension. *Curr Therap Res.* 1996; 57(3):145-58.
64. Mounier-Vehier C, Jaboureck O, Emeriau JP, Bernaud C, Clerson P, Carre A. Randomized, comparative, double-blind study of amlodipine vs nifedipine as a treatment of isolated systolic hypertension in the elderly. *Fundam Clin Pharmacol.* 2002;16(6):537-44.
65. Kes S, Caglar N, Canberk A, et al. Treatment of mild-to-moderate hypertension with calcium-channel blockers: a multicentre comparison of once-daily nifedipine GITS with once-daily amlodipine. *Curr Med Res Opin.* 2003; 19(3):226-37.
66. Ryuzaki M, Nakamoto H, Nishida E et al. Crossover study of amlodipine versus nifedipine CR with home blood pressure monitoring via cellular phone: internet-mediated open-label crossover trial of calcium-channel blockers for hypertension (I-TECHO trial). *J Hypertens.* 2007 Nov;25(11):2352-8.
67. Saito I, Saruta T. Controlled release nifedipine and valsartan combination therapy in patients with essential hypertension: The Adalat CR and valsartan cost-effectiveness combination (ADVANCE-Combi) study. *J Hypertens.* 2007 Nov;25(11):2352-8.
68. Pepine CJ, Cooper-DeHoff RM, et al. Comparative Efficacy and Safety of Nisoldipine and Amlodipine (CESNA-II) Study Investigators. Comparison of effects of nisoldipine-extended-release and amlodipine in patients with systemic hypertension and chronic stable angina pectoris. *Am J Cardiol.* 2003; 91(3):274-9.
69. Whitcomb C, Enzmann G, Pershadsingh HA, et al. A comparison of nisoldipine ER and amlodipine for the treatment of mild to moderate hypertension. *Int J Clin Pract.* 2000;54(8):509-13.
70. White WB, Saunders E, Noveck RJ, Ferdinand K. Comparative efficacy and safety of nisoldipine extended-release (ER) and amlodipine (CESNA-III study) in African American patients with hypertension. *Am J Hypertens.* 2003;16(9 Pt 1):739-45.
71. Lenz TL, Wurdeman RL, Hilleman DE. Comparison of 24-hour blood pressure profiles in patients with hypertension who were switched from amlodipine to nisoldipine. *Pharmacotherapy.* 2001;21(8):898-903.
72. Drummond W, Munger M, Essop M, Maboudian M, Khan M, Keefe D. Antihypertensive efficacy of the oral direct renin inhibitor aliskiren as add-on therapy in patients not responding to amlodipine monotherapy. *J Clin Hypertens.* 2007;9:742-50.
73. Benetos A, Consoli S, Safavian A, Dubanchet A, Safar M. Efficacy, safety, and effects on quality of life of bisoprolol/hydrochlorothiazide versus amlodipine in elderly patients with systolic hypertension. *Am Heart J.* 2000 Oct;140(4):E11.
74. Prisant LM, Weir MR, Papademetriou V, et al. Low-dose combination therapy: an alternative first-line approach to hypertension treatment. *Am Heart J.* 1995 Aug;130(2):359-66.
75. Mazza A, Gil-Extremera B, Maldonato A, Toutouzas T, Pessina AC. Nebivolol vs amlodipine as first-line treatment of essential arterial hypertension in the elderly. *Blood Press.* 2002;11(3):182-8.
76. Hollenberg HK, Williams GH, Anderson H et al. Symptoms and the distress they cause: comparison of an aldosterone antagonist and a calcium-channel blocking agent in patients with systolic hypertension. *Arch Intern Med.* 2003;163(13):1543-8.
77. White WB, Duprez D, St Hillaire R et al. Effects of the selective aldosterone blocker eplerenone versus the calcium antagonist amlodipine in systolic hypertension. *Hypertension* 2003;41(5):1021-6.
78. Jordan J, Engeli S, Boye S, et al. Direct renin inhibition with aliskiren in obese patients with arterial hypertension. *Hypertension.* 2007 May;49:1047-55.
79. Messerli FH, Weir MR, Neutel JM. Combination therapy of amlodipine and benazepril versus monotherapy of amlodipine in a practice-based setting. *Am J Hypertens.* 2002 Jun;15(6):550-6.
80. Chrysant SG. Blood pressure effects of high-dose amlodipine-benazepril combination in black and white hypertensive patients not controlled on monotherapy. *Drugs RD.* 2012;12(2):57-64.
81. Messerli FH, Oparil S, Feng Z. Comparison of efficacy and side effects of combination therapy of angiotensin-converting enzyme inhibitor (benazepril) with calcium antagonist (either nifedipine or amlodipine) versus high-dose calcium antagonist monotherapy for systemic hypertension. *Am J Cardiol.* 2000 Dec 1;86:1182-7.
82. Jamerson KA, Nwose O, Jean-Louis L, et al. Initial angiotensin-converting enzyme inhibitor/calcium-channel blocker combination therapy achieves superior blood pressure control compared with calcium-channel blocker monotherapy in patients with stage 2 hypertension. *Am J Hypertens.* 2004 Jun;17(6):495-501.
83. Neutel JM, Smith DH, Weber MA, et al. Efficacy of combination therapy for systolic blood pressure in patients with severe systolic hypertension: the Systolic Evaluation of Lotrel Efficacy and Comparative Therapies (SELECT) study. *J Clin Hypertens.* 2005 Nov;7(11):641-6.
84. Kuschnir E, Acuna E, Sevilla D, et al. Treatment of patients with essential hypertension: amlodipine 5 mg/benazepril 20 mg compared with amlodipine 5 mg, benazepril 20 mg, and placebo. *Clin Ther.* 1996;18(6):1213-24.

85. Chrysant SG, Sugimoto DH, Lefkowitz M, et al. The effects of high-dose amlodipine/benazepril combination therapies on blood pressure reduction in patients not adequately controlled with amlodipine monotherapy. *Blood Press Suppl.* 2007 Mar;1:10-7.
86. Chrysant SG, Bakris GL. Amlodipine/benazepril combination therapy for hypertensive patients nonresponsive to benazepril monotherapy. *Am J Hypertens.* 2004 Jul;17(7):590-6.
87. Fogari R, Corea L, Cardoni O, et al. Combined therapy with benazepril and amlodipine in the treatment of hypertension inadequately controlled by an ACE inhibitor alone. *J Cardiovasc Pharmacol.* 1997(Oct);30(4):497-503.
88. Minami J, Abe C, Akashiba A, Takahashi T, Kameda T, Ishimitsu T, Matsuoka H. Long-term efficacy of combination therapy with losartan and low-dose hydrochlorothiazide in patients with uncontrolled hypertension. *Int Heart J.* 2007 Mar;48(2):177-86.
89. Hilleman DE, Ryschon KL, Mohiuddin SM, Wurdeman RL. Fixed-dose combination vs monotherapy in hypertension: a meta-analysis evaluation. *J Hum Hypertens.* 1999;13:477-83.
90. Jamerson K, Bakris GL, Dahlof B, et al; for the ACCOMPLISH Investigators. Exceptional early blood pressure control rates: the ACCOMPLISH trial. *Blood Press.* 2007;16(2):80-6.
91. Malacco E, Piazza S, Carretta R, et al; Italian Blood Pressure Study Group. Comparison of benazepril-amlodipine and captopril-thiazide combinations in the management of mild-to-moderate hypertension. *Int J Clin Pharmacol Ther.* 2002 Jun;40(6):263-9.
92. Kereiakes DJ, Neutel JM, Punzi HA, et al. Efficacy and safety of olmesartan medoxomil and hydrochlorothiazide compared with benazepril and amlodipine besylate. *Am J Cardiovasc Drugs.* 2007;7(5):36-72.
93. Tatti P, Pahor M, Byington RP, et al. Outcome results of the Fosinopril versus Amlodipine Cardiovascular Events Trial (FACET) in patients with hypertension and non-insulin dependent diabetes mellitus. *Diabetes Care.* 1998; 21:597-603.
94. Miranda RD, Mion D, Rocha JC, et al. An 18-week, prospective, randomized, double-blind, multicenter study of amlodipine/ramipril combination versus amlodipine monotherapy in the treatment of hypertension: The assessment of combination therapy of amlodipine/ramipril (ATAR) study. *Clin Ther* 2008;30:1618-28.
95. Fogari R, Mugellini A, Derosa G; CANDIA (CANdesartan and DLuretic vs Amlodipine in hypertensive patients) Study Group. Efficacy and tolerability of candesartan cilexetil/hydrochlorothiazide and amlodipine in patients with poorly controlled mild-to-moderate essential hypertension. *J Renin Angiotensin Aldosterone Syst.* 2007 Sep;8(3):139-44.
96. Ribeiro AB, Mion D Jr, Marin MJ, et al; Latin American Hypertension Study (LAMHYST) Group. Antihypertensive efficacy of amlodipine and losartan after two 'missed' doses in patients with mild to moderate essential hypertension. *J Int Med Res.* 2007 Nov-Dec;35(6):762-72.
97. Oparil S, Barr E, Elkins M, Liss C, Vrecenak A, Edelman J. Efficacy, tolerability, and effects on quality of life of losartan, alone or with hydrochlorothiazide, versus amlodipine, alone or with hydrochlorothiazide, in patients with essential hypertension. *Clin Ther.* 1996 Jul-Aug;18(4):608-25.
98. Chrysant SG, Melino M, Karki S, Lee J, Heyrman R. The combination of olmesartan medoxomil and amlodipine besylate in controlling high blood pressure: COACH, a randomized, double-blind, placebo-controlled, 8-week factorial efficacy and safety study. *Clin Ther.* 2008 Apr;30(4):587-604.
99. Chrysant SG, Oparil S, Melino M, et al. Efficacy and safety of long-term treatment with the combination of amlodipine besylate and olmesartan medoxomil in patients with hypertension. *J Clin Hypertens (Greenwich)* 2009;11:475-82.
100. Oparil S, Lee J, Karki S, Melino M. Subgroup analyses of an efficacy and safety study of concomitant administration of amlodipine besylate and olmesartan medoxomil: evaluation by baseline hypertension stage and prior antihypertensive medication use. *J Cardiovasc Pharmacol.* 2009;54(5):427-36.
101. Braun N, Ulmer HJ, Handrock R, Klebs S. Efficacy and safety of the single pill combination of amlodipine 10 mg plus valsartan 160 mg in hypertensive patients not controlled by amlodipine 10 mg plus olmesartan 20 mg in free combination (abstract). *Current Medical Research & Opinion.* 2009;25(2):421-30.
102. Elliott WJ, Whitmore J, Feldstein JD, Bakris GL. Efficacy and safety of perindopril arginine + amlodipine in hypertension. *J Am Soc Hypertens.* 2015 Apr;9(4):266-74.
103. Manolis A, Grammatikou V, Kallistratos M, Zarifis J, Tsioufis K. Blood pressure reduction and control with fixed-dose combination perindopril/amlodipine: A Pan-Hellenic prospective observational study. *J Renin Angiotensin Aldosterone Syst.* 2015 Dec;16(4):930-5.
104. Littlejohn TW 3rd, Majul CR, Olvera R, et al. Results of treatment with telmisartan-amlodipine in hypertensive patients. *J Clin Hypertens (Greenwich)* 2009;11:207-13.
105. Littlejohn T, Majul C, Olver R, Seeber M, Kobe M, Guthrie R et al. Telmisartan plus amlodipine in patients with moderate or severe hypertension: results from a subgroup analysis of a randomized, placebo-controlled, parallel-group, 4x4 factorial study. *Postgrad Med.* 2009;121(2):5-14.

106. Sharma A, Bagchi A, Kinagi SB, et al. Results of a comparative, phase III, 12-week, multicenter, prospective, randomized, double-blind assessment of the efficacy and tolerability of a fixed-dose combination of telmisartan and amlodipine versus amlodipine monotherapy in Indian adults with stage II hypertension. *Clin Ther* 2007;29:2667-76.
107. Neutel JM, Mancia G, Black HR, Dahlof B, Defeo H, Ley L, et al. Single-pill combination of telmisartan/amlodipine in patients with severe hypertension: results from the TEAMSTA severe HTN study. *J Clin Hypertens (Greenwich)*. 2012 Apr;14(4):206-215.
108. Maciejewski S, Mohiuddin SM, Packard KA, et al. Randomized, double-blind, crossover comparison of amlodipine and valsartan in African-Americans with hypertension using 24-hour ambulatory blood pressure monitoring. *Pharmacotherapy*. 2006 Jul;26(7):889-95.
109. Ichihara A, Kaneshiro Y, Takemitsu T, Sakoda M. Effects of amlodipine and valsartan on vascular damage and ambulatory blood pressure in untreated hypertensive patients. *J Hum Hypertens*. 2006 Oct;20(10):787-94.
110. Karpov Y, Dongre N, Vigdorichik A, Sastravaha K. Amlodipine/valsartan single-pill combination: a prospective, observational evaluation of the real-life safety and effectiveness in the routine treatment of hypertension. *Adv Ther*. 2012;29(2):134-147.
111. Philipp T, Smith TR, Glazer R, et al. Two multicenter, 8-week, randomized, double-blind, placebo-controlled, parallel-group studies evaluating the efficacy and tolerability of amlodipine and valsartan in combination and as monotherapy in adult patients with mild to moderate essential hypertension. *Clin Ther*. 2007 Apr;29(4):563-80.
112. Philipp T, Smith TR, Glazer R, et al. Two multicenter, 8-week, randomized, double-blind, placebo-controlled, parallel-group studies evaluating the efficacy and tolerability of amlodipine and valsartan in combination and as monotherapy in adult patients with mild to moderate essential hypertension. *Clin Ther*. 2007 Apr;29(4):563-80.
113. Philipp T, Glazer RD, Wernsing M, Yen J. Initial combination therapy with amlodipine/valsartan compared with monotherapy in the treatment of hypertension (abstract). *J Amer Soc Hyperten*. 2011;5(5):417-424.
114. Schunkert H, Glazer RD, Wernsing M, et al. Efficacy and tolerability of amlodipine/valsartan combination therapy in hypertensive patients not adequately controlled on amlodipine monotherapy. *Curr Med Res Opin* 2009;25:2655-62.
115. Ke Y, Zhu D, Hong H, et al. Efficacy and safety of a single-pill combination of amlodipine/valsartan in Asian hypertensive patients inadequately controlled with amlodipine monotherapy. *Curr Med Res Opin* 2010;26:1705-13.
116. Destro M, Luckow A, Samson M, et al. Efficacy and safety of amlodipine/valsartan compared with amlodipine monotherapy in patients with stage 2 hypertension: a randomized, double-blind, multicenter study: the EX-EFFeCTS study. *J Am Society Hypertension* 2008;2:294-302.
117. Flack JM, Calhoun DA, Satlin L, et al. Efficacy and safety of initial combination therapy with amlodipine/valsartan compared with amlodipine monotherapy in black patients with stage 2 hypertension: the EX-STAND study. *J Hum Hypertens* 2009;23:479-89.
118. Schrader J, Salvetti A, Calvo C, et al. The combination of amlodipine/valsartan 5/160 mg produces less peripheral oedema than amlodipine 10 mg in hypertensive patients not adequately controlled with amlodipine 5 mg. *Int J Clin Pract* 2009;63:217-25.
119. Sinkiewicz W, Glazer RD, Kavoliuniene A, et al. Efficacy and tolerability of amlodipine/valsartan combination therapy in hypertensive patients not adequately controlled on valsartan monotherapy. *Curr Med Res Opin* 2009;25:315-24.
120. Fogari R, Zoppi A, Mugellini A, Corradi L, Lazzari P, Preti P et al. Efficacy and safety of two treatment combinations of hypertension in very elderly patients. *Archives of Gerontology and Geriatrics*. 2009;48:401-5.
121. Poldermans D, Glazer R, Karagiannis S, et al. Tolerability and blood pressure-lowering efficacy of the combination of amlodipine plus valsartan compared with lisinopril plus hydrochlorothiazide in adult patients with stage 2 hypertension. *Clin Ther*. 2007 Feb;29(2):279-89.
122. Calhoun DA, Lacourciere Y, Chiang YT, Glazer RD. Triple antihypertensive therapy with amlodipine, valsartan, and hydrochlorothiazide: A randomized controlled trial. *Hypertension* 2009;54:32-9.
123. Calhoun D, Crikelair N, Yen J, Glazer R. Amlodipine/valsartan/hydrochlorothiazide triple combination therapy in moderate/severe hypertension: secondary analyses evaluating efficacy and safety. *Adv Ther*. 2009;26(11):1012-23.
124. Pareek A, Salkar H, Mulay P, Desai S, Chandurkar N, Redkar N. A randomized, comparative, multicenter, evaluation of atenolol/amlodipine combination with atenolol alone in essential hypertension patients. *Am J Ther* 2010;17:46-52.
125. Gustin G, White WB, Taylor S, Daragjati C. Clinical outcome of a mandatory formulary switch for dihydropyridine calcium-channel blocker therapy at a Veteran's Administration Medical Center. *Am J Hypertens*. 1996;9(4 Pt 1):312-6.

126. Karotsis AK, Symeonidis A, Mastorantonakis SE, Stergiou GS. Additional antihypertensive effect of drugs in hypertensive subjects uncontrolled on diltiazem monotherapy: a randomized controlled trial using office and home blood pressure monitoring. *Clin Exp Hypertens*. 2006;28(7):655-62.
127. Manyemba J. A randomized crossover comparison of reserpine and sustained-release nifedipine in hypertension. *Cent Afr J Med*. 1997 Dec;43(12):344-9.
128. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*. 2005 Oct 29-Nov 4;366(9496):1545-53.
129. Van Bortel LM, Fici F, Mascagni F. Efficacy and tolerability of nebivolol compared with other antihypertensive drugs: a meta-analysis. *Am J Cardiovasc Drugs*. 2008;8(1):35-44.
130. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
131. Baguet JP, Legallier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig*. 2007;27(11):735-53.
132. Laurent S, Mancia G, Poulter N. Perindopril 3.5 mg/amlodipine 2.5 mg versus renin-angiotensin system inhibitor monotherapy as first-line treatment in hypertension: a combined analysis. *J Hypertens*. 2018 March 31; 36(9):1915-20.
133. Esnault VL, Brown EA, Apetrei E, et al. The effects of amlodipine and enalapril on renal function in adults with hypertension and nondiabetic nephropathies: A 3-year, randomized, multicenter, double-blind, placebo-controlled study. *Clin Ther* 2008;30:482-98.
134. Agodoa LY, Appel L, Bakris GL, et al; African American Study of Kidney Disease and Hypertension (AASK) Study Group. Effect of ramipril vs amlodipine on renal outcomes in hypertensive nephrosclerosis: a randomized controlled trial. *JAMA*. 2001 Jun 6;285(21):2719-28.
135. Wright JT Jr, Bakris G, Green T, et al. Effect of blood pressure lowering and antihypertensive kidney disease: results from the AASK trial. *JAMA*. 2002; 288:2421-31.
136. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, et al; Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med*. 2001 Sep 20;345(12):851-60.
137. Viberti G, Wheeldon NM; MicroAlbuminuria Reduction With VALsartan (MARVAL) Study Investigators. Microalbuminuria reduction with valsartan in patients with type 2 diabetes mellitus: a blood pressure-independent effect. *Circulation*. 2002 Aug 6;106(6):672-8.
138. Bakris GL, Toto RD, McCullough PA, et al; on behalf of GUARD (Gauging Albuminuria Reduction With Lotrel in Diabetic Patients With Hypertension) Study Investigators. Effects of different ACE inhibitor combinations on albuminuria: results of the GUARD study. *Kidney Int*. 2008 Jun;73(11):1303-9.
139. Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet*. 2005 Dec 10;366:2026-33.
140. Rosendorff C, Dubiel R, Xu J, Chavanu KJ. Comparison of olmesartan medoxomil versus amlodipine besylate on regression of ventricular and vascular hypertrophy. *Am J Cardiol* 2009;104:359-65.
141. Luscher TF, Pieper M, Tendera M, et al. A randomized, placebo-controlled study on the effect of nifedipine on coronary endothelial function and plaque formation in patients with coronary artery disease: the ENCORE II study. *Euro Heart J* 2009;30:1590-7.
142. Schmid-Elsaesser R, Kunz M, Zausinger S, et al. Intravenous magnesium versus nimodipine in the treatment of patients with aneurysmal subarachnoid hemorrhage: a randomized study. *Neurosurgery*. 2006 Jun;58(6):1054-65.
143. Liu GJ, Zhang LLP, Wang ZJ, Xu LL, He GH, Zeng YJ, et al. Meta-analysis of the effectiveness and safety of prophylactic use of nimodipine in patients with an aneurysmal subarachnoid haemorrhage. *CNS Neurol Disord Drug Targets*. 2011;10(7):834-44.
144. Taylor AA, Shoheiber O. Adherence to antihypertensive therapy with fixed-dose amlodipine besylate/benazepril HCl versus comparable component-based therapy. *Congest Heart Fail*. 2003 Nov-Dec;9(6):324-32.
145. Dickson M, Plauschinat CA. Compliance with antihypertensive therapy in the elderly: a comparison of fixed-dose combination amlodipine/benazepril versus component-based free-combination therapy. *Am J Cardiovasc Drugs* 2008;1:45-50.
146. Gerbino PP, Shoheiber O. Adherence patterns among patients treated with fixed-dose combination versus separate antihypertensive agents. *Am J Health Syst Pharm* 2007;64:1279-83.
147. Sapienza S, Sacco P, Floyd K, et al. Results of a pilot pharmacotherapy quality improvement program using fixed-dose, combination amlodipine/benazepril antihypertensive therapy in a long-term care setting. *Clin Ther* 2003;25:1872-87.



**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Calcium-Channel Blocking Agents, Miscellaneous  
AHFS Class 242892  
May 4, 2022**

**I. Overview**

The movement of calcium ions is essential for the function of all types of muscle, including cardiac and vascular smooth muscle. When this flow is reduced, the result is a weakening of muscle contraction and relaxation of muscle tissue.<sup>1-3</sup> Relaxation of coronary vascular smooth muscle increases the flow of oxygenated blood into the myocardium, while relaxation of arteriolar smooth muscle decreases peripheral vascular resistance. Both coronary and systemic vasodilation serve to reduce cardiac workload. The calcium-channel blocking agents include dihydropyridines and miscellaneous agents (nondihydropyridines). Although they have different binding sites on the L-type calcium channel, both block the transmembrane influx of calcium ions into cardiac and vascular smooth muscle. The nondihydropyridines also block the T-type calcium channel in the atrioventricular node.<sup>1-5</sup>

The miscellaneous calcium-channel blocking agents include diltiazem and verapamil, which are approved for the treatment of angina, arrhythmias, and hypertension.<sup>1,2,6-14</sup> Diltiazem is a potent coronary vasodilator, but is only a mild arterial vasodilator. Although it decreases atrioventricular (AV) node conduction, diltiazem does not have negative inotropic properties.<sup>1,2,6-14</sup> Verapamil dilates coronary and peripheral arteries. It also slows conduction through the AV node, and has negative inotropic and chronotropic effects.<sup>1,2,6-14</sup> Both diltiazem and verapamil are available in a variety of modified-release delivery systems that alter their pharmacokinetic properties, including onset and duration of action.<sup>1,2</sup>

The miscellaneous calcium-channel blocking agents that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. Diltiazem and verapamil are available in generic formulations. This class was last reviewed in February 2020.

**Table 1. Calcium-Channel Blocking Agents, Miscellaneous Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Diltiazem	extended-release capsule, extended-release tablet, injection, tablet	Cardizem <sup>®</sup> *, Cardizem CD <sup>®</sup> *, Cardizem LA <sup>®</sup> *, Matzim LA <sup>®</sup> *, Tiazac <sup>®</sup> *	diltiazem
Verapamil	extended-release capsule, extended-release tablet, injection, tablet	Calan SR <sup>®</sup> *, Verelan <sup>®</sup> *, Verelan PM <sup>®</sup> *	verapamil

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the miscellaneous calcium-channel blocking agents are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Calcium-Channel Blocking Agents, Miscellaneous**

Clinical Guideline	Recommendations
American College of Cardiology/ American Heart Association: <b>2007 Chronic Angina Focused Update of the 2002 Guidelines for the</b>	<ul style="list-style-type: none"> <li>Aspirin should be started at 75 to 162 mg/day and continued indefinitely in all patients, unless contraindicated.</li> <li>Use of warfarin in conjunction with aspirin and/or clopidogrel is associated with an increased risk of bleeding and should be monitored closely.</li> <li>Patients with hypertension and established coronary artery disease (CAD) should be treated with blood pressure medication(s) as tolerated, including angiotensin converting enzyme (ACE) inhibitors and/or <math>\beta</math>-adrenergic antagonists (<math>\beta</math>-blockers) with the addition of other medications as needed to achieve blood pressure goals of</li> </ul>



Clinical Guideline	Recommendations
<b>Management of Patients With Chronic Stable Angina (2007)<sup>15</sup></b>	<p>&lt;140/90 or &lt;130/80 mm Hg for patients with chronic kidney disease or diabetes.</p> <ul style="list-style-type: none"> <li>• Long-acting calcium channel blocking agents or long-acting nitrates may be used if <math>\beta</math>-blockers are contraindicated. Immediate-release and short-acting dihydropyridine calcium channel blockers can increase adverse cardiac events and should not be used.</li> <li>• Long-acting calcium channel blockers or long-acting nitrates may be used with <math>\beta</math>-blockers if initial treatment is not successful.</li> <li>• ACE inhibitors should be used indefinitely in patients with a left ventricular ejection fraction (LVEF) <math>\leq 40\%</math> and in those with hypertension, diabetes or chronic kidney disease, unless contraindicated.</li> <li>• ACE inhibitors should also be used indefinitely in patients at lower risk (mildly reduced or normal LVEF in whom cardiovascular risk factors remain well controlled and revascularization has been performed), unless contraindicated.</li> <li>• Angiotensin II receptor blockers (ARBs) are recommended in patients with hypertension, those who have an indication for an ACE inhibitor and are intolerant to them, who have heart failure, or who have had a myocardial infarction (MI) and have a LVEF <math>\leq 40\%</math>.</li> <li>• ARBs may be considered in combination with an ACE inhibitor for heart failure due to left ventricular systolic dysfunction.</li> <li>• Aldosterone blockade is recommended in patients post-MI without significant renal dysfunction or hyperkalemia who are already receiving therapeutic doses of an ACE inhibitor and a <math>\beta</math>-blocker, have a LVEF <math>\leq 40\%</math> and have either diabetes or heart failure.</li> <li>• It is beneficial to start and continue <math>\beta</math>-blocker therapy indefinitely in all patients who have had a MI, acute coronary syndrome or left ventricular dysfunction with or without heart failure symptoms, unless contraindicated.</li> <li>• Annual influenza vaccination is recommended in patients with cardiovascular disease.</li> </ul>
<b>European Society of Cardiology: Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)<sup>16</sup></b>	<p><b>Pharmacological management of stable coronary artery disease (CAD) patients</b></p> <ul style="list-style-type: none"> <li>• The two aims of the pharmacological management of stable CAD patients are to obtain relief of symptoms and to prevent CV events.</li> <li>• Optimal medical treatment indicates at least one drug for angina/ischaemia relief plus drugs for event prevention.</li> <li>• It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>• It is indicated to review the patient's response soon after starting therapy.</li> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin, consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</li> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief:             <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a beta-blocker and/or a non-DHP-calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</li> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</p> <ul style="list-style-type: none"> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischaemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <ul style="list-style-type: none"> <li>● Event prevention: <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><u>Treatment in patients with microvascular angina</u></p> <ul style="list-style-type: none"> <li>● It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>● <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>● Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>● ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>● Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul> <p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>● Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy (DAPT).</li> <li>● Aspirin is recommended for elective stenting.</li> <li>● Clopidogrel is recommended for elective stenting.</li> <li>● Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>● GP IIb/IIIa antagonists should be considered for bailout situation only.</li> <li>● Platelet function testing or genetic testing may be considered in specific or high-risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>● Prasugrel or ticagrelor may be considered in specific high-risk situations of elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</li> <li>● Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>● Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>● Prasugrel or ticagrelor is not recommended in low-risk elective stenting.</li> <li>● After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low</li> <li>● Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months</li> </ul> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>● It is recommended that all revascularized patients receive a secondary prevention</li> </ul>

Clinical Guideline	Recommendations
	<p>and be scheduled for follow-up visit.</p> <ul style="list-style-type: none"> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after second generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/          American College of Cardiology Foundation/          American Heart Association/          American Association for Thoracic Surgery/          Preventive Cardiovascular Nurses Association/          Society of Thoracic Surgeons:  <b>Management of Stable Ischemic Heart Disease (2012)</b><sup>17</sup></p>	<p><u>Medical therapy to prevent MI and death in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</li> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</li> <li>• Patients should receive an annual influenza vaccine.</li> </ul> <p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> <li>• Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/          American Heart Association:</p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTE-ACS) with arterial oxygen saturation <math>&lt; 90\%</math>, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications</li> </ul>

Clinical Guideline	Recommendations
<p><b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes (2014)<sup>18</sup></b></p>	<ul style="list-style-type: none"> <li>○ Nitrates <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use.</li> </ul> </li> <li>○ Beta-adrenergic blockers <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval <math>&gt;0.24</math> second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</li> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> </li> <li>○ Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval <math>&gt;0.24</math> seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</li> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> </li> <li>○ Other anti-ischemic interventions <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>mortality, need for myocardial revascularization, and stroke.</p> <ul style="list-style-type: none"> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> <ul style="list-style-type: none"> <li>• Inhibitors of renin-angiotensin-aldosterone system <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF &lt;0.40 and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF &lt;0.40 who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine &gt;2.5 mg/dL in men or &gt;2.0 mg/dL in women) or hyperkalemia (K &gt;5.0 mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF &lt;0.40, diabetes mellitus, or heart failure.</li> </ul> </li> <li>• Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include: <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatide or tirofiban.</li> </ul> </li> </ul> </li> </ul> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet agents <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatide, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Anticoagulant therapy               <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>• Timing of CABG in relation to use of antiplatelet agents               <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatid or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> </li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>• Medications at discharge               <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</li> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post-NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> <li>● Late hospital and post-hospital oral antiplatelet therapy               <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS               <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation</b> (2020)<sup>19</sup></p>	<p><b>Pharmacological treatment of ischemia</b></p> <ul style="list-style-type: none"> <li>● Sublingual or intravenous nitrates and early initiation of β-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>● Continuation of chronic β-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>● Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>● In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and β-blockers avoided.</li> </ul> <p><b>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</b></p> <ul style="list-style-type: none"> <li>● Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> <li>● A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds.           <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>● P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>● Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>● It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>● It is not recommended to administer prasugrel in patients whom coronary anatomy</li> </ul>

Clinical Guideline	Recommendations
	<p>is not known.</p> <ul style="list-style-type: none"> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul> <p><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new</li> </ul>



Clinical Guideline	Recommendations
	<p>P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</p> <ul style="list-style-type: none"> <li>• If at low bleeding risk (HAS-BLED ≤2), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED ≥3), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED ≥3 and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>
<p>American College of Cardiology/ American Heart Association: <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>20</sup></p>	<p><u>Routine medical therapies: calcium channel blockers</u></p> <ul style="list-style-type: none"> <li>• Evidence demonstrates that beneficial effect on infarct size or the rate of reinfarction when calcium channel blocker therapy was initiated during either the acute or convalescent phase of ST-segment elevation myocardial infarction (STEMI). However, calcium channel blockers may be useful to relieve ischemia, lower blood pressure, or control the ventricular response rate to atrial fibrillation in patients who are intolerant to β-blockers.</li> <li>• Use of immediate-release nifedipine is contraindicated in patients with STEMI due to hypotension and reflex sympathetic activation with tachycardia.</li> </ul> <p><u>Routine medical therapies: β-blockers</u></p> <ul style="list-style-type: none"> <li>• Oral β-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral β-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li>• β-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>• Patients with initial contraindications to the use of β-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>• It is reasonable to administer intravenous β-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or</li> </ul>

Clinical Guideline	Recommendations
	<p>have ongoing ischemia.</p> <p><u>Routine medical therapies: Renin-Angiotensin-Aldosterone System Inhibitors</u></p> <ul style="list-style-type: none"> <li>• An angiotensin-converting enzyme (ACE) inhibitor should be administered within the first 24 hours to all patients with STEMI with anterior location, HF, or ejection fraction (EF) <math>\leq 40\%</math>, unless contraindicated.</li> <li>• An angiotensin receptor blocker (ARB) should be given to patients with STEMI who have indications for but are intolerant of ACE inhibitors.</li> <li>• An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq 40\%</math> and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>• High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>• It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>21</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of ST-elevation myocardial infarction (STEMI)</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>• Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>• A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>• In patients with an indication for oral anticoagulation, oral anticoagulants are indicated in addition to antiplatelet therapy.</li> <li>• In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the estimated risk of recurrent coronary events and bleeding).</li> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age <math>\geq 50</math> years, and at least one of the following risk factors: age <math>\geq 65</math> years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR <math>&lt; 60</math> mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</li> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with <math>\beta</math>-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with <math>\beta</math>-blockers is indicated in patients with heart failure or left ventricular dysfunction, LVEF <math>\leq 40\%</math> unless contraindicated.</li> <li>• Intravenous <math>\beta</math>-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Intravenous <math>\beta</math>-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of <math>&lt;1.8</math> mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C <math>&gt;1.8</math> mmol/L (<math>&gt;70</math> mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction <math>\leq 40\%</math> and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
<p>American College of Cardiology/ American Heart Association: <b>Guideline on the Primary Prevention of Cardiovascular Disease (2019)</b><sup>22</sup></p>	<p><u>Top 10 messages for the primary prevention of cardiovascular disease</u></p> <ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</li> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</li> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with</li> </ul>

Clinical Guideline	Recommendations
	<p>diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</p> <ul style="list-style-type: none"> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be &lt;130/80 mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul> <p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a decision is made for statin therapy, a moderate-intensity statin should be recommended.</li> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) patients, LDL-C levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</li> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND             <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> </li> <li>• In patients at borderline risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>• In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include:               <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>• In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>• In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>• In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>• In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than 130/80 mm Hg.</li> <li>• In adults with an estimated 10-year ASCVD risk &lt;10% and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering medication are recommended.</li> <li>• In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul> <p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>• All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>• To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>• In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>• In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> <li>• To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>• All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>• Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>• Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</li> <li>• Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/Americ</p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>• Hypertension should be controlled in accordance with guideline-directed</li> </ul>

Clinical Guideline	Recommendations
<p>an College of Cardiology/Heart Failure Society of America: <b>2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)</b><sup>23</sup></p>	<p>medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>• In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>• Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>• In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>• In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><u>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</u></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36</li> </ul>

Clinical Guideline	Recommendations
	<p>hours of the last dose of an ACE inhibitor. (LoE: B)</p> <ul style="list-style-type: none"> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq 35\%</math>) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq 70</math> beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on</li> </ul>

Clinical Guideline	Recommendations
	<p>guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</p> <p><b>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</b></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><b>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</b></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><b>Treatment of Stage D (advanced/refractory) HF</b></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>24</sup></p>	<p><b>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</b></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to</li> </ul>



Clinical Guideline	Recommendations
	<p>further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g., with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>American Heart Association/ American College of Cardiology/ Heart Rhythm Society: <b>2019</b> <b>AHA/ACC/HRS Focused update of the 2014 AHA/ACC/ARS Guideline for the Management of</b></p>	<p><u>Recommendations for risk-based anticoagulant therapy:</u> Class I</p> <ul style="list-style-type: none"> <li>• In patients with atrial fibrillation (AF), anticoagulant therapy should be individualized based on shared decision-making after discussion of the absolute and relative risks of stroke, bleeding and the patient’s values and preferences (Level of Evidence: C).</li> <li>• NOACs (dabigatran, rivaroxaban, apixaban, and edoxaban) are recommended over warfarin in NOAC-eligible patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) (Level of Evidence: A).</li> <li>• Selection of anticoagulant therapy should be based on the risk of thromboembolism irrespective of whether the AF patter is paroxysmal, persistent, or permanent (Level of Evidence: B).</li> </ul>

Clinical Guideline	Recommendations
<p><b>Patients with Atrial Fibrillation (2019)<sup>25</sup></b></p>	<ul style="list-style-type: none"> <li>• In patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve), the CHA<sub>2</sub>DS<sub>2</sub>-VASc score is recommended for assessment of stroke risk (Level of Evidence: B). For patients with AF who have mechanical heart valves, warfarin is recommended. (Level of Evidence: B).</li> <li>• Selection of anticoagulant therapy should be based on the risk of thromboembolism, irrespective of whether the AF pattern is paroxysmal, persistent, or permanent.</li> <li>• For patients with AF and an elevated CHA<sub>2</sub>DS<sub>2</sub>-VASc score <math>\geq 2</math> in men or <math>\geq 3</math> in women, oral anticoagulants are recommended. Options include warfarin (Level of Evidence: A), dabigatran, rivaroxaban, apixaban, or edoxaban. (Level of Evidence: B).</li> <li>• Among patients treated with warfarin, the INR should be determined at least weekly during initiation of anticoagulant therapy and at least monthly when anticoagulation (INR in range) is stable (Level of Evidence: A)</li> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) who are unable to maintain a therapeutic INR level with warfarin, use of a non-vitamin K oral anticoagulant (NOAC) is recommended (Level of Evidence: C-EO).</li> <li>• Re-evaluation of the need for and choice of anticoagulant therapy at periodic intervals is recommended to reassess stroke and bleeding risks (Level of Evidence: C).</li> <li>• Renal function should be evaluated prior to initiation of a NOAC and should be re-evaluated when clinically indicated and at least annually (Level of Evidence: B-NR).</li> <li>• For patients with atrial flutter, anticoagulant therapy is recommended according to the same risk profile used for AF (Level of Evidence: C).</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 0 in men or 1 in women, it is reasonable to omit antithrombotic therapy (Level of Evidence: B).</li> <li>• For patients with AF with a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of <math>\geq 2</math> in men or <math>\geq 3</math> in women and who have end-stage chronic kidney disease (creatinine clearance <math>&lt; 15</math> mL/min) or who are on dialysis, it is reasonable to prescribe warfarin (INR 2.0 to 3.0) or apixaban for oral anticoagulation (Level of Evidence: B).</li> </ul> <p><b>Class IIb</b></p> <ul style="list-style-type: none"> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 1 in men and 2 in women, prescribing an oral anticoagulant to reduce thromboembolic stroke risk may be considered. (Level of Evidence: C-LD).</li> <li>• For patients with AF (except with moderate-to-severe mitral stenosis or a mechanical heart valve) and moderate-to-severe chronic kidney disease with elevated CHA<sub>2</sub>DS<sub>2</sub>-VASc, treatment with reduced doses of direct thrombin or factor Xa inhibitors may be considered (e.g., dabigatran, rivaroxaban, apixaban, or edoxaban. (Level of Evidence: C).</li> </ul> <p><b>Class III: No Benefit</b></p> <ul style="list-style-type: none"> <li>• In patients with AF and end-stage CKD or on dialysis, the direct thrombin inhibitor dabigatran or the factor Xa inhibitors rivaroxaban or edoxaban are not recommended because of the lack of evidence from clinical trials that benefit exceeds risk (Level of Evidence: C-EO).</li> </ul> <p><b>Class III: Harm</b></p> <ul style="list-style-type: none"> <li>• The direct thrombin inhibitor, dabigatran, should not be used in patients with AF and a mechanical heart valve (Level of Evidence: B).</li> </ul> <p><u>Recommendations for Interruption and Bridging Anticoagulation</u></p> <p><b>Class I</b></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Bridging therapy with UFH or LMWH is recommended for patients with AF and a mechanical heart valve undergoing procedures that require interruption of warfarin. Decisions regarding bridging therapy should balance the risks of stroke and bleeding (Level of Evidence: C).</li> <li>• For patients with AF without mechanical heart valves who require interruption of warfarin or newer anticoagulants for procedures, decisions about bridging therapy (LMWH or UFH) should balance the risks of stroke and bleeding and the duration of time a patient will not be anticoagulated (Level of Evidence: B-R).</li> <li>• Idarucizumab is recommended for the reversal of dabigatran in the event of life-threatening bleeding or an urgent procedure (Level of Evidence: B-NR)</li> </ul> <p>Class IIa</p> <ul style="list-style-type: none"> <li>• Andexanet alfa can be useful for the reversal of rivaroxaban and apixaban in the event of life-threatening or uncontrolled bleeding (Level of Evidence: B-NR).</li> </ul> <p><u>Recommendations for Thromboembolism Prevention:</u></p> <p>Class I</p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of 48-hour duration or longer, or when the duration of AF is unknown, anticoagulation with warfarin (INR 2.0 to 3.0), a factor Xa inhibitor, or direct thrombin inhibitor is recommended for at least three weeks prior to and four weeks after cardioversion, regardless of the CHA<sub>2</sub>DS<sub>2</sub>-VASc score and the method used to restore sinus rhythm (Level of Evidence: B).</li> <li>• For patients with AF or atrial flutter of more than 48 hours duration that requires immediate cardioversion for hemodynamic instability, anticoagulation should be initiated as soon as possible and continued for at least four weeks after cardioversion unless contraindicated (Level of Evidence: C).</li> <li>• After cardioversion for AF of any duration, the decision about long-term anticoagulation therapy should be based on the thromboembolic risk profile and bleeding risk profile (Level of Evidence: C-EO).</li> </ul> <p>Class IIa</p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of less than 48-hour duration and a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of <math>\geq 2</math> in men or <math>\geq 3</math> in women, administration of, heparin, a factor Xa or direct thrombin inhibitor, is recommended as soon as possible before cardioversion, followed by long-term anticoagulation therapy (Level of Evidence: B-NR).</li> <li>• After cardioversion for AF of any duration, the decision regarding long-term anticoagulation therapy should be based on the thromboembolic risk profile (Level of Evidence: C-EO).</li> <li>• For patients with AF or atrial flutter of 48-hour duration or longer or of unknown duration who have not been anticoagulated for the preceding three weeks, it is reasonable to perform a TEE prior to cardioversion and proceed with cardioversion if no LA thrombus is identified, including in the LAA, provided that anticoagulation is achieved before TEE and maintained after cardioversion for at least four weeks (Level of Evidence: B).</li> <li>• For patients with AF or atrial flutter of 48-hour duration or longer, or when the duration of AF is unknown, anticoagulation with dabigatran, rivaroxaban, or apixaban is reasonable for at least three weeks prior to and four weeks after cardioversion (Level of Evidence: C).</li> </ul> <p>Class IIb</p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of less than 48 hours' duration with a CHA<sub>2</sub>DS<sub>2</sub>-VASc score of 0 in men or 1 in women, administration of heparin, a factor Xa inhibitor, or a direct thrombin inhibitor, versus no anticoagulant therapy, may be considered before cardioversion, without the need for post cardioversion oral anticoagulation (Level of Evidence: B-NR).</li> </ul> <p><u>Recommendations for pharmacological cardioversion</u></p>

Clinical Guideline	Recommendations
	<p><b>Class I</b></p> <ul style="list-style-type: none"> <li>• For patients with AF or atrial flutter of 48 hours' duration or longer, or when the duration of AF is unknown, anticoagulation with warfarin (INR 2.0 to 3.0), a factor Xa inhibitor, or direct thrombin inhibitor is recommended for at least 3 weeks before and at least 4 weeks after cardioversion, regardless of the CHA<sub>2</sub>DS<sub>2</sub>-VASc score or the method (electrical or pharmacological) used to restore sinus rhythm (Level of Evidence: B-R).</li> <li>• For patients with AF or atrial flutter of more than 48 hours' duration or unknown duration that requires immediate cardioversion for hemodynamic instability, anticoagulation should be initiated as soon as possible and continued for at least 4 weeks after cardioversion unless contraindicated (Level of Evidence: C).</li> <li>• After cardioversion for AF of any duration, the decision about long-term anticoagulation therapy should be based on the thromboembolic risk profile and bleeding risk profile (Level of Evidence: C-EO).</li> </ul> <p><u>Recommendations for AF complicating ACS</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>• For patients with ACS and AF at increased risk of systemic thromboembolism (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater), anticoagulation is recommended unless the bleeding risk exceeds the expected benefit (Level of Evidence: B-R).</li> <li>• Urgent direct-current cardioversion of new-onset AF in the setting of ACS is recommended for patients with hemodynamic compromise, ongoing ischemia, or inadequate rate control (Level of Evidence: C).</li> <li>• Intravenous β blockers are recommended to slow a rapid ventricular response to AF in patients with ACS who do not display HF, hemodynamic instability, or bronchospasm (Level of Evidence: C).</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>• If triple therapy (oral anticoagulant, aspirin, and P2Y<sub>12</sub> inhibitor) is prescribed for patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone percutaneous coronary intervention (PCI) with stenting for ACS, it is reasonable to choose clopidogrel in preference to prasugrel (Level of Evidence: B-NR).</li> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with a P2Y<sub>12</sub> inhibitor (clopidogrel or ticagrelor) and dose-adjusted vitamin K antagonist is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with P2Y<sub>12</sub> inhibitors (clopidogrel) and low-dose rivaroxaban 15 mg daily is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> <li>• In patients with AF at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) who have undergone PCI with stenting for ACS, double therapy with a P2Y<sub>12</sub> inhibitor (clopidogrel) and dabigatran 150 mg twice daily is reasonable to reduce the risk of bleeding as compared with triple therapy (Level of Evidence: B-R).</li> </ul> <p><b>Class IIb</b></p> <ul style="list-style-type: none"> <li>• If triple therapy (oral anticoagulant, aspirin, and P2Y<sub>12</sub> inhibitor) is prescribed for patients with AF who are at increased risk of stroke (based on CHA<sub>2</sub>DS<sub>2</sub>-VASc risk score of 2 or greater) and who have undergone PCI with stenting (drug eluting or bare metal) for ACS, a transition to double therapy (oral anticoagulant and P2Y<sub>12</sub> inhibitor) at four to six weeks may be considered (Level of Evidence: B-R).</li> <li>• Administration of amiodarone or digoxin may be considered to slow a rapid</li> </ul>

Clinical Guideline	Recommendations
	<p>ventricular response in patients with ACS and AF associated with severe LV dysfunction and HF or hemodynamic instability (Level of Evidence: C).</p> <ul style="list-style-type: none"> <li>Administration of amiodarone or digoxin may be considered to slow a rapid ventricular response in patients with ACS and AF associated with severe LV dysfunction and HF or hemodynamic instability (Level of Evidence: C).</li> </ul> <p><u>Recommendations for Device Detection of AF and Atrial Flutter</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>In patients with cardiac implantable electronic devices (pacemakers or implanted cardioverter-defibrillators), the presence of recorded atrial high-rate episodes (AHREs) should prompt further evaluation to document clinically relevant AF to guide treatment decisions (Level of Evidence: B-NR).</li> </ul> <p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>In patients with cryptogenic stroke (i.e., stroke of unknown cause) in whom external ambulatory monitoring is inconclusive, implantation of a cardiac monitor (loop recorder) is reasonable to optimize detection of silent AF (Level of Evidence: B-R).</li> </ul> <p><u>Recommendations for Weight loss</u></p> <p><b>Class I</b></p> <ul style="list-style-type: none"> <li>For overweight and obese patients with AF, weight loss, combined with risk factor modification, is recommended (Level of Evidence: B-R).</li> </ul>
<p>National Institute for Health and Care Excellence: <b>Atrial Fibrillation: Diagnosis and Management</b> (2021)<sup>26</sup></p>	<p><u>Rate control</u></p> <ul style="list-style-type: none"> <li>Offer rate control as the first-line treatment strategy for atrial fibrillation except in people: <ul style="list-style-type: none"> <li>whose atrial fibrillation has a reversible cause</li> <li>who have heart failure thought to be primarily caused by atrial fibrillation</li> <li>with new-onset atrial fibrillation</li> <li>with atrial flutter whose condition is considered suitable for an ablation strategy to restore sinus rhythm</li> <li>for whom a rhythm-control strategy would be more suitable based on clinical judgement.</li> </ul> </li> <li>Offer either a standard <math>\beta</math>-blocker (that is, a <math>\beta</math>-blocker other than sotalol) or a rate-limiting calcium channel blocker (CCB) (diltiazem or verapamil) as initial monotherapy to people with atrial fibrillation (AF) unless the person has the features described above. Base the choice of drug on the person's symptoms, heart rate, comorbidities and preferences..</li> <li>Consider digoxin monotherapy for initial rate control for people with non-paroxysmal atrial fibrillation if the person does no or very little physical exercise or if other rate-limiting drug options are ruled out because of comorbidities or the person's preferences..</li> <li>If monotherapy does not control symptoms, and if continuing symptoms are thought to be due to poor ventricular rate control, consider combination therapy with any two of the following: a <math>\beta</math>-blocker, diltiazem, and digoxin.</li> <li>Do not offer amiodarone for long-term rate control.</li> </ul> <p><u>Rhythm control</u></p> <ul style="list-style-type: none"> <li>Consider pharmacological and/or electrical rhythm control for people with AF whose symptoms continue after heart rate has been controlled or for whom a rate-control strategy has not been successful.</li> </ul> <p><u>Drug treatment for long-term rhythm control</u></p> <ul style="list-style-type: none"> <li>Assess the need for drug treatment for long-term rhythm control, taking into account the person's preferences, associated comorbidities, risks of treatment, and likelihood of recurrence of AF.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Do not offer class 1c antiarrhythmic drugs such as flecainide or propafenone to people with known ischaemic or structural heart disease.</li> <li>• If drug treatment for long-term rhythm control is needed, consider a standard <math>\beta</math>-blocker as first-line treatment unless there are contraindications.</li> <li>• If <math>\beta</math>-blockers are contraindicated or unsuccessful, assess the suitability of alternative drugs for rhythm control, taking comorbidities into account.</li> <li>• Dronedarone is recommended as an option for the maintenance of sinus rhythm after successful cardioversion in people with paroxysmal or persistent atrial fibrillation:             <ul style="list-style-type: none"> <li>○ Whose AF is not controlled by first-line therapy (usually including <math>\beta</math>-blockers), that is, as a second-line treatment option and after alternative options have been considered AND</li> <li>○ Who have at least one of the following cardiovascular risk factors:                 <ul style="list-style-type: none"> <li>▪ Hypertension requiring drugs of at least two different classes.</li> <li>▪ Diabetes mellitus.</li> <li>▪ Previous TIA, stroke, or systemic embolism.</li> <li>▪ Left atrial diameter of 50 mm or greater, OR</li> <li>▪ Age <math>\geq 70</math> years, AND</li> </ul> </li> <li>○ Who do not have left ventricular systolic dysfunction, AND</li> <li>○ Who do not have a history of, or current, heart failure.</li> </ul> </li> <li>• People who do not meet the criteria above who are currently receiving dronedarone should have the option to continue treatment until they and their clinicians consider it appropriate to stop.</li> <li>• Consider amiodarone for people with left ventricular impairment or heart failure.</li> <li>• Do not offer class 1c antiarrhythmic drugs such as flecainide or propafenone to people with known ischemic or structural heart disease.</li> <li>• Where people have infrequent paroxysms and few symptoms, or where symptoms are induced by known precipitants (such as alcohol, caffeine), a 'no drug treatment' strategy or a 'pill-in-the-pocket' strategy should be considered and discussed with the person.</li> </ul> <p><u>Preventing postoperative atrial fibrillation</u></p> <ul style="list-style-type: none"> <li>• In people having cardiothoracic surgery:             <ul style="list-style-type: none"> <li>○ reduce the risk of postoperative atrial fibrillation by offering one of the following: amiodarone; a standard <math>\beta</math>-blocker (that is, a <math>\beta</math>-blocker other than sotalol); a rate-limiting calcium-channel blocker (diltiazem or verapamil)</li> <li>○ do not offer digoxin.</li> </ul> </li> <li>• In people having cardiothoracic surgery who are already on <math>\beta</math>-blocker therapy, continue this treatment unless contraindications develop (such as postoperative bradycardia or hypotension).</li> <li>• Do not start statins in people having cardiothoracic surgery solely to prevent postoperative atrial fibrillation.</li> <li>• In people having cardiothoracic surgery who are already on statins, continue this treatment. For further advice on statins for the prevention of cardiovascular disease, see NICE's guideline on cardiovascular disease: risk assessment and reduction.</li> </ul>
<p>American Association for Thoracic Surgery: <b>2014 AATS Guidelines for the Prevention and Management of Peri-Operative Atrial Fibrillation</b></p>	<p><u>Recommended prevention strategies for all postoperative atrial fibrillation (POAF) patients</u></p> <ul style="list-style-type: none"> <li>• Patients taking <math>\beta</math>-blockers prior to thoracic surgery should continue them in the postoperative period to avoid <math>\beta</math>-blockade withdrawal.</li> <li>• Intravenous magnesium supplementation may be considered to prevent postoperative AF when serum magnesium level is low or it is suspected that total body magnesium is depleted.</li> <li>• Digoxin should not be used for prophylaxis against AF.</li> </ul>

Clinical Guideline	Recommendations
<p><b>and Flutter (POAF) for Thoracic Surgical Procedures (2014)<sup>27</sup></b></p>	<p><u>Recommended prevention strategies for intermediate to high-risk POAF patients</u></p> <ul style="list-style-type: none"> <li>• It is reasonable to administer diltiazem to those patients with preserved cardiac function who are not taking <math>\beta</math>-blockers preoperatively in order to prevent POAF.</li> <li>• It is reasonable to consider the postoperative administration of amiodarone to reduce the incidence of POAF for intermediate and high risk patients undergoing pulmonary resection.</li> <li>• Postoperative administration of intravenous amiodarone may be considered to prevent POAF in patients undergoing esophagectomy.</li> <li>• Atorvastatin may be considered to prevent POAF for statin naïve patients scheduled for intermediate and high risk thoracic surgical procedures.</li> </ul> <p><u>Rate control recommendations for patients with new onset POAF</u></p> <ul style="list-style-type: none"> <li>• Intravenous administration of <math>\beta</math>-blockers (e.g., esmolol or metoprolol) or nondihydropyridine calcium channel blockers (diltiazem or verapamil) is recommended to achieve rate control (heart rate <math>\leq 110</math> bpm) for patients who develop POAF with rapid ventricular response.</li> <li>• Caution should be used with patients with hypotension, left ventricular (LV) dysfunction, or heart failure.</li> <li>• Combination use of atrioventricular (AV) nodal blocking agents, such as <math>\beta</math>-blockers (e.g., esmolol or metoprolol), nondihydropyridine calcium channel antagonists (e.g., diltiazem or verapamil), or digoxin, can be useful to control heart rates when a single agent fails to control rates of POAF. The choice should be individualized and doses modified to avoid bradycardia.</li> <li>• For patients with hypotension, heart failure or LV dysfunction, or when other measures are unsuccessful or contraindicated, intravenous amiodarone can be useful for control of heart rate. Amiodarone could result in conversion to sinus rhythm, and if it is initiated after 48 hours of AF, both a transesophageal echocardiography (TEE) when possible, to rule out left atrial/LA appendage (LA/LAA) thrombus, and full anticoagulation should be considered.</li> <li>• For patients with heart failure, LV dysfunction or hypotension, intravenous digoxin may be considered for rate control of POAF.</li> <li>• For patients with ventricular preexcitation (i.e., Wolff-Parkinson-White syndrome) and POAF, use of AV nodal blocking agents, such as <math>\beta</math>-blockers (e.g., esmolol or metoprolol), intravenous amiodarone, nondihydropyridine calcium channel antagonists (e.g., diltiazem or verapamil), or digoxin, should be avoided.</li> </ul> <p><u>Recommendations for the use of antiarrhythmic drugs for pharmacologic cardioversion of POAF</u></p> <ul style="list-style-type: none"> <li>• Restoration of sinus rhythm with pharmacologic cardioversion is reasonable in patients with symptomatic, hemodynamically stable POAF. Intravenous amiodarone can be useful for pharmacologic cardioversion of POAF.</li> <li>• It is reasonable to administer antiarrhythmic medications in an attempt to maintain sinus rhythm for patients with recurrent or refractory POAF.</li> <li>• Amiodarone, sotalol, flecainide, propafenone, or dofetilide can be useful to maintain sinus rhythm in patients with POAF, depending on underlying heart disease, renal status and other comorbidities.</li> <li>• Flecainide or propafenone may be considered for pharmacologic cardioversion of POAF and maintenance of sinus rhythm if the patient has had no prior history of myocardial infarction, coronary artery disease, impaired LV function, significant LV hypertrophy, or valvular heart disease that is considered moderate or greater. These agents may need to be combined with an AV nodal blocking agent.</li> <li>• Intravenous ibutilide or procainamide may be considered for pharmacologic conversion of POAF for patients with structural heart disease and new onset POAF, but no hypotension or manifestations of congestive heart failure. Serum electrolytes and QTc interval must be within a normal range and patients must be closely</li> </ul>



Clinical Guideline	Recommendations
	<p>monitored during and for at least six hours after the infusion if either ibutilide or procainamide.</p> <ul style="list-style-type: none"> <li>• Intravenous ibutilide or procainamide may be considered for patients with POAF and an accessory pathway.</li> <li>• Flecainide and propafenone should not be used to treat POAF in patients with a history of a prior myocardial infarction, coronary artery disease, and/or severe structural heart disease, including severe left ventricular hypertrophy, or significantly reduced left ventricular ejection fraction.</li> <li>• Dronedarone should not be used for treatment of POAF in patients with heart failure.</li> </ul> <p><u>Recommendations for prevention of thromboembolism for patients with stable atrial fibrillation/flutter undergoing direct current cardioversion</u></p> <ul style="list-style-type: none"> <li>• For stable patients with POAF of 48-hours duration or longer, anticoagulation (with warfarin for INR 2.0 to 3.0, a novel oral anti-coagulant [NOAC] or LMWH) is recommended for at least three weeks prior to and four weeks after cardioversion, regardless of the method (electrical or pharmacological) used to restore sinus rhythm.</li> <li>• During the first 48 hours after the onset of POAF, the need for anticoagulation before and after direct current (DC) cardioversion may be based on the patient's risk of thromboembolism (CHA2DS2-VASc score) balanced by the risk of postoperative bleeding.</li> <li>• For POAF lasting longer than 48 hours, as an alternative to three weeks of therapeutic anticoagulation prior to cardioversion of POAF, it is reasonable to perform TEE in search of thrombus in the LA or LA appendage, preferably with full anticoagulation at the time of TEE in anticipation of DC cardioversion after the TEE.</li> <li>• For POAF lasting longer than 48 hours in patients who are not candidates for TEE (e.g., post-esophageal surgery), an initial rate control strategy combined with therapeutic anticoagulation using warfarin (aiming for INR 2.0 to 3.0), a direct thrombin inhibitor (e.g. dabigatran), factor Xa inhibitor (e.g. rivaroxaban, apixaban), or LMWH is recommended for at least three weeks prior to and four weeks after cardioversion.</li> <li>• Anticoagulation recommendations for cardioversion of atrial flutter are similar to those for atrial fibrillation.</li> <li>• For patients with an identified thrombus, cardioversion should not be performed until a longer period of anticoagulation is achieved (usually at least three weeks) and in accordance with established AF guidelines.</li> </ul> <p><u>Management of anticoagulation for new onset POAF</u></p> <ul style="list-style-type: none"> <li>• For the prevention of strokes for patients who develop POAF lasting longer than 48 hours, it is recommended to administer antithrombotic medications similarly to non-surgical patients. Anticoagulation within the first 48-hours of POAF should be considered based on the CHA2DS2-VASc risk score of the patient for stroke weighed against the risk of postoperative bleeding.</li> <li>• New oral anticoagulants (dabigatran, rivaroxaban, apixaban) are reasonable as an alternative to warfarin for patients who do not have a prosthetic heart valve, hemodynamically significant valve disease, and/or severe renal impairment or risk of GI bleeding.</li> <li>• It is reasonable to continue anticoagulation therapy for four weeks after the return of sinus rhythm because of the possibility of slowly resolving impairment of atrial contraction with an associated ongoing risk for thrombus formation and for delayed embolic events.</li> <li>• New oral anticoagulants should be avoided for patients at risk for serious bleeding (including GI bleeding) as they cannot be readily reversed. However, their use may</li> </ul>

Clinical Guideline	Recommendations
	<p>be recommended in situations where achievement of a therapeutic INR with warfarin has proved to be difficult.</p>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>28</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic blood pressure <math>&lt; 140</math> mm Hg.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 140</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>29</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist</li> </ul>

Clinical Guideline	Recommendations
	<p>circumference should be used. Alternatively, a waist-to-height ratio <math>&lt;0.5</math> is recommended for all populations.</p> <ul style="list-style-type: none"> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><u>Pharmacological Treatment</u></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment             <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment             <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq 160/100</math>): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy             <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq 80</math> years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq 80</math> years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;45</math> mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;4.5</math> mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indication for their use, e.g., heart failure, angina, post-MI, atrial</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
<p>Hypertension            Canada:  <b>2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)</b><sup>30</sup></p>	<p>fibrillation, or younger women planning pregnancy or pregnant.</p> <p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq 100</math> mmHg or average systolic blood pressure (SBP) measurements of <math>\geq 160</math> mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq 90</math> mmHg or for average SBP readings <math>\geq 140</math> mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC).               <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are:                   <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or</li> </ul>

Clinical Guideline	Recommendations
	<p>there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</p> <ul style="list-style-type: none"> <li>Possible reasons for poor response to therapy should be considered.</li> <li><math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>Short-acting nifedipine should not be used.</li> <li>When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>In patients with systolic dysfunction (ejection fraction <math>&lt; 40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal</li> </ul>

Clinical Guideline	Recommendations
	<p>function. Additional therapies may also include dihydropyridine CCBs.</p> <ul style="list-style-type: none"> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq</math>30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in</li> </ul>

Clinical Guideline	Recommendations
	<p>cases of complex lesions less amenable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</p> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric</li> </ul>



Clinical Guideline	Recommendations
	<p>kidney disease).</p> <ul style="list-style-type: none"> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/          European Society of Cardiology:  <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>31</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection</li> </ul>

Clinical Guideline	Recommendations
	<p>fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</p> <ul style="list-style-type: none"> <li>• In patients with HF<sub>r</sub>EF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HF<sub>p</sub>EF, BP treatment threshold and target values should be the same as for HF<sub>r</sub>EF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>32</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on “Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence”.</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African–Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person’s medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks:  <b>Management of High Blood Pressure in Blacks</b></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or</li> </ul>

Clinical Guideline	Recommendations
(2010) <sup>33</sup>	<p>an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</p> <ul style="list-style-type: none"> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)<sup>34</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>• The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney</li> </ul>

Clinical Guideline	Recommendations
	<p><b>transplant recipients.</b></p> <p><b>BP management in children with CKD</b></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>35</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>In adults with increased risk of HF, the optimal BP in those with hypertension should be <math>&lt; 130</math> mmHg.</li> <li>Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP <math>&lt; 130/80</math> mmHg.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFpEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq</math>65 years of age) with an average SBP of <math>\geq</math>130 mmHg.</li> <li>• For older adults (<math>\geq</math>65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP ≥180 mmHg or DBP ≥110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>36</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (≥140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure ≥180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk ≥15%), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure ≥140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated</li> </ul>

Clinical Guideline	Recommendations
	<p>urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</p> <ul style="list-style-type: none"> <li>Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>

\*Agent not available in the United States.

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the miscellaneous calcium-channel blocking agents are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Calcium-Channel Blocking Agents, Miscellaneous<sup>1,2,6-14</sup>**

Indication	Diltiazem	Verapamil
<b>Angina Pectoris</b>		
Angina due to coronary artery spasm	✓ (tablet, ER capsule [Cardizem CD <sup>®</sup> ])	
Chronic stable angina	✓	✓ (tablet)
Unstable angina		✓ (tablet)
Vasospastic angina		✓ (tablet)
<b>Arrhythmias</b>		
Control of ventricular rate at rest and during stress in patients with chronic atrial flutter and/or atrial fibrillation in association with digitalis		✓ (tablet)
Prophylaxis of repetitive paroxysmal supraventricular tachycardia		✓ (tablet)
Rapid conversion to sinus rhythm of paroxysmal supraventricular tachycardias	✓ (injection)	✓ (injection)
Temporary control of rapid ventricular rate in atrial flutter or atrial fibrillation	✓ (injection)	✓ (injection)
<b>Hypertension</b>		
Hypertension	✓ * (ER)	✓

\*May be used alone or in combination with other antihypertensive agents.  
ER=extended-release

### IV. Pharmacokinetics

The pharmacokinetic parameters of the miscellaneous calcium-channel blocking agents are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Calcium-Channel Blocking Agents, Miscellaneous<sup>2</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)

Diltiazem*	35 to 40	77 to 93	Liver, extensive (% not reported)	Renal (35) Feces (60 to 65)	3 to 10
Verapamil*	20 to 35	88 to 94	Liver (65 to 80)	Renal (70) Feces (9 to 16)	4 to 12

\*Immediate-release

## V. Drug Interactions

Major drug interactions with the miscellaneous calcium-channel blocking agents are listed in Table 5.

**Table 5. Major Drug Interactions with the Calcium-Channel Blocking Agents, Miscellaneous<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Macrolides	Increased serum levels of macrolide antibiotics may result if administered with calcium-channel blocking agents, miscellaneous, due to the inhibitory effect on CYP3A4. Coadministration should be avoided.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Narcotic Analgesics	Calcium-channel blocking agents, miscellaneous may increase plasma concentrations of narcotic analgesics, increasing the potential for enhanced pharmacologic effects and toxicity. Inhibition of CYP3A4 isoenzyme by Calcium-channel blocking agents, miscellaneous may decrease the metabolic elimination of narcotic analgesics.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Vasopressin Receptor Antagonists	Plasma concentrations and pharmacologic effects of vasopressin receptor antagonists may be increased by diltiazem. Inhibition of CYP3A isoenzymes by diltiazem may decrease the metabolic elimination of vasopressin receptor antagonists.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Amiodarone	Concurrent use of amiodarone and calcium channel blockers may result in bradycardia, atrioventricular block and/or sinus arrest.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Colchicine	Plasma concentrations of colchicine may be increased. Colchicine toxicity may occur. Inhibition of CYP3A4 and/or efflux transporter P-glycoprotein diltiazem may increase the absorption and decrease the metabolic elimination of colchicine.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Carbamazepine	Increased serum levels of carbamazepine may result if administered with diltiazem, increasing the risk of greater effect and toxicity, due to inhibition of carbamazepine metabolism by diltiazem.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Cyclosporine	Increased serum levels of cyclosporine may result if administered with diltiazem, due to inhibition of cyclosporine metabolism by diltiazem.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Everolimus	Pharmacologic effects and plasma concentrations of everolimus may be increased by diltiazem. Inhibition of CYP3A4 and P-glycoprotein by diltiazem may decrease the metabolic elimination of everolimus.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Ibrutinib	Diltiazem inhibits CYP3A4 metabolism of ibrutinib, thereby increasing plasma concentrations, pharmacologic effects, and risk of toxicity (e.g., hemorrhage, renal toxicity).
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Lomitapide	Diltiazem inhibits CYP3A4 metabolism of lomitapide, thereby increasing plasma concentrations, pharmacologic effects, and risk of adverse reactions, including hepatotoxicity.

Generic Name(s)	Interaction	Mechanism
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	Ranolazine	Increased serum levels of ranolazine may result if administered with diltiazem, due to diltiazem's inhibitory effect on CYP3A4. Coadministration should be avoided due to the increased risk of QTc prolongation, torsades de pointes arrhythmias and death.
Calcium-channel blocking agents, miscellaneous (diltiazem, verapamil)	HMG CoA reductase inhibitors	Plasma concentrations and pharmacologic effects of statins may be increased by co-administration. The risk of myopathy and rhabdomyolysis may be increased. Inhibition of CYP3A4 isoenzymes by miscellaneous calcium-channel blockers may decrease the metabolic elimination of statins.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Benzodiazepines	Increased serum levels of benzodiazepines may result if administered with diltiazem, increasing the risk of central nervous system depression, due to decreased metabolism of benzodiazepines.
Calcium-channel blocking agents, miscellaneous (diltiazem)	$\beta$ -Blockers	Increased serum levels of $\beta$ -blockers may result if administered with diltiazem, increasing the risk of symptomatic bradycardia, due to decreased metabolism of $\beta$ -blockers and additive pharmacologic effects.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Cilostazol	Pharmacologic effects of cilostazol may be increased by diltiazem. Elevated plasma concentrations with toxicity may occur. Inhibition of CYP3A4 isoenzymes by diltiazem may decrease the metabolic elimination of cilostazol.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Cisapride	Concurrent use may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Calcium-channel blocking agents, miscellaneous (diltiazem)	Corticosteroids	Diltiazem may increase the pharmacologic effects of corticosteroids. Inhibition of CYP3A4 isoenzymes by diltiazem may decrease the metabolic elimination of corticosteroids.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Digoxin	Increased serum levels of digoxin may result, increasing the risk of digoxin toxicity, if administered with diltiazem, due to decreased renal clearance of digoxin.
Calcium-channel blocking agents, miscellaneous (diltiazem)	HIV Protease Inhibitors	Plasma concentrations and pharmacologic effects of diltiazem may be increased by HIV protease inhibitors. An additive effect on the PR interval has also been demonstrated. Plasma concentrations and pharmacologic effects of diltiazem may be increased by HIV protease inhibitors.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Macrolide immuno-suppressives	Plasma trough concentrations of macrolide immunosuppressives may be increased by diltiazem. Neurologic toxicity may occur. Diltiazem may increase the plasma trough concentrations of macrolide immunosuppressives. Neurologic toxicity may occur.
Calcium-channel blocking agents, miscellaneous (diltiazem)	Theophyllines	The pharmacologic and toxic effects of theophyllines may be increased due to the inhibition of metabolism of theophylline by diltiazem.
Calcium-channel blocking agents, miscellaneous (verapamil)	$\beta$ -Blockers	Effects of $\beta$ -blockers and diltiazem may be increased, close monitoring of cardiac function is recommended. Diltiazem may inhibit the metabolism of some $\beta$ -blockers (atenolol, metoprolol and propranolol), leading to increased effects of these $\beta$ -blockers.
Calcium-channel	Digoxin	Verapamil may alter the pharmacokinetics and increase

Generic Name(s)	Interaction	Mechanism
blocking agents, miscellaneous (verapamil)		serum concentrations of digoxin. Verapamil may decrease nonrenal and total digoxin clearance.
Calcium-channel blocking agents, miscellaneous (verapamil)	Dofetilide	Increase serum levels and effects of dofetilide may occur if coadministered with verapamil, increasing the risk of arrhythmia.
Calcium-channel blocking agents, miscellaneous (verapamil)	Quinidine	Pharmacologic effects of quinidine may be increased. This combination may produce marked hypotension. Verapamil inhibits the hepatic metabolism of quinidine.
Calcium-channel blocking agents, miscellaneous (verapamil)	Aldosterone Blockers	Verapamil may increase plasma concentrations and pharmacologic or toxic effects of aldosterone blockers. Inhibition of CYP3A4 isoenzymes by verapamil may decrease the metabolic elimination of aldosterone blockers.
Calcium-channel blocking agents, miscellaneous (verapamil)	Clonidine	Sinus bradycardia, atrioventricular block and severe hypotension may occur with coadministration of clonidine and verapamil.
Calcium-channel blocking agents, miscellaneous (verapamil)	Dronedarone	Plasma concentrations and pharmacologic effects of dronedarone may be increased by verapamil. Dronedarone may also increase the plasma concentrations and pharmacologic effects of verapamil. Additionally, verapamil may enhance the electrophysiologic effects of dronedarone.
Calcium-channel blocking agents, miscellaneous (verapamil)	Flecainide	Increased risk of cardiotoxic effects may occur when flecainide and verapamil are coadministered. Cardiogenic shock or asystole may develop. Pharmacologic effects may be additive or synergistic.
Calcium-channel blocking agents, miscellaneous (verapamil)	Nondepolarizing muscle relaxants	Increased serum levels of nondepolarizing muscle relaxants may result, increasing the risk of respiratory depression, if coadministered with verapamil, due to calcium's role on muscle contraction.
Calcium-channel blocking agents, miscellaneous (verapamil)	Quinazolines	The combination of verapamil and quinazolines may produce an acute hypotensive effect which is greater than when either drug is taken alone. Verapamil may decrease the first-pass hepatic metabolism and increase the bioavailability of quinazolines.
Calcium-channel blocking agents, miscellaneous (verapamil)	Rifampin	Decreased serum levels of verapamil may result if coadministered with rifampin, due to increased metabolism of verapamil.

CYP=cytochrome P450 isoenzymes, HIV=human immunodeficiency virus, HMG CoA=3-hydroxy-3-methyl-glutaryl-CoA

## VI. Adverse Drug Events

The most common adverse drug events reported with the miscellaneous calcium-channel blocking agents are listed in Table 6.

**Table 6. Adverse Drug Events (%) Reported with the Calcium-Channel Blocking Agents, Miscellaneous<sup>1,2,6-14</sup>**

Adverse Events	Diltiazem	Verapamil
<b>Cardiovascular</b>		
Angina	-	<1

<b>Adverse Events</b>	<b>Diltiazem</b>	<b>Verapamil</b>
Arrhythmia	<2	-
Atrial fibrillation	-	✓
Atrioventricular dissociation	-	<1
Atrioventricular block	2 to 8	1 to 2
Bradycardia	2 to 6	1
Bundle branch block	<2	-
Chest pain	-	<1
Claudication	-	<1
Congestive heart failure	<2	2
Edema	2 to 15	-
Extrasystoles	2	-
Flushing	1 to 2	1
Hypotension	<4	3
Myocardial infarction	-	<1
Palpitations	1 to 2	<1
Peripheral edema	2 to 8	2 to 4
Postural hypotension	-	<1
Syncope	<2	<1
Tachycardia	<2	-
Vasodilation	2 to 3	-
Ventricular fibrillation	-	✓
<b>Central Nervous System</b>		
Cerebrovascular accident	-	<1
Confusion	-	<1
Depression	<2	-
Dizziness	3 to 10	1 to 5
Fatigue	-	2 to 5
Headache	5 to 12	1 to 12
Insomnia	-	<1
Lethargy	-	3
Nervousness	2	-
Paresthesia	-	1
Psychotic symptoms	-	<1
Sleep disturbance	-	1
Somnolence	-	<1
Tremor	<2	<1
Vertigo	-	<1
<b>Dermatologic</b>		
Alopecia	-	<1
Ecchymosis	-	<1
Erythema multiforme	-	<1
Hair color change	-	✓
Hyperhidrosis	-	<1
Hyperkeratosis	-	<1
Petechiae	<2	-
Photosensitivity	<2	-
Rash	1 to 4	1 to 2
Stevens-Johnson syndrome	<2	-
Toxic epidermal necrolysis	<2	-
<b>Endocrine and Metabolic</b>		
Gout	1 to 2	-
Gynecomastia	-	<1
Hyperprolactinemia/galactorrhea	-	<1
<b>Gastrointestinal</b>		

<b>Adverse Events</b>	<b>Diltiazem</b>	<b>Verapamil</b>
Abdominal discomfort	-	<1
Constipation	<4	7 to 12
Diarrhea	1 to 2	2
Dry mouth	-	<1
Dysgeusia	<2	-
Dyspepsia	1 to 6	3
Gingival hyperplasia	<2	<19
Nausea	-	1 to 3
Vomiting	2	-
<b>Genitourinary</b>		
Acute renal failure	-	✓
Albuminuria	-	-
Crystalluria	-	-
Impotence	-	<1
Nocturia	-	-
Polyuria	-	<1
Sexual dysfunction	-	-
Spotty menstruation	-	<1
<b>Hematological</b>		
Hemolytic anemia	<2	-
Purpura	-	<1
Thrombocytopenia	<2	-
<b>Laboratory Test Abnormalities</b>		
Alkaline phosphatase increase	<2	-
ALT increased	<2	-
AST increased	<2	-
Liver enzyme elevations	-	1
<b>Musculoskeletal</b>		
Arthralgia	-	<1
Extrapyramidal symptoms	<2	-
Muscle cramps	-	<1
Myalgia	2	1
Pain	6	2
Paresthesia	-	1
Weakness	1 to 4	-
<b>Respiratory</b>		
Bronchitis	1 to 4	-
Cough	≤3	✓
Dyspnea	1 to 6	1
Pharyngitis	2 to 6	-
Rhinitis	<10	-
Sinus congestion	1 to 2	-
<b>Other</b>		
Abnormal visual accommodation	-	<1
Allergic reaction	<2	-
Amblyopia	<2	-
Amnesia	<2	-
Blurred vision	-	<1
Flu-like syndrome	-	4
Parkinsonian syndrome	-	✓
Tinnitus	-	<1

✓ Percent not specified  
- Event not reported



## VII. Dosing and Administration

The usual dosing regimens for the miscellaneous calcium-channel blocking agents are listed in Table 7.

**Table 7. Usual Dosing Regimens for the Calcium-Channel Blocking Agents, Miscellaneous<sup>1,2,6-14</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Diltiazem	<p><u>Angina pectoris (chronic stable):</u>            Extended-release capsule: initial, 120 mg/day; maintenance, 180 to 480 mg/day; maximum, 480 mg/day</p> <p>Extended-release tablet: initial, 180 mg once daily; maximum, 360 mg/day</p> <p>Tablet: initial, 30 mg four times daily; maintenance, 180 to 360 mg/day</p> <p><u>Angina pectoris (due to coronary artery spasm):</u>            Extended-release capsule (Cardizem CD<sup>®</sup>): initial, 120 or 180 mg once daily; maintenance, adjust dosage to each patient's needs</p> <p>Tablet: initial, 30 mg four times daily; maintenance, 180 to 360 mg/day</p> <p><u>Arrhythmias:</u>            Injection: weight based dosing administered intravenously</p> <p><u>Hypertension:</u>            Extended-release capsule: initial, 180 to 240 mg once daily; maintenance, 180 to 480 mg/day; maximum, 540 mg/day</p> <p>Extended-release tablet: initial, 180 to 240 mg once daily; maintenance, 120 to 540 mg/day; maximum, 540 mg/day</p>	Safety and efficacy in children have not been established.	<p>Extended-release capsule:            60 mg            90 mg            120 mg            180 mg            240 mg            300 mg            360 mg            420 mg</p> <p>Extended-release tablet:            120 mg            180 mg            240 mg            300 mg            360 mg            420 mg</p> <p>Injection:            5 mg/mL            100 mg</p> <p>Tablet:            30 mg            60 mg            90 mg            120 mg</p>

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Verapamil	<p><u>Angina pectoris (chronic stable, unstable, and vasospastic):</u>            Tablet: maintenance, 80 to 120 mg three times a day</p> <p><u>Arrhythmias:</u>            Injection: weight based dosing administered by slow intravenous injection</p> <p>Tablet: maintenance, 240 to 480 mg/day, divided (three to four times daily)</p> <p><u>Hypertension:</u>            Tablet: initial, 80 mg three times daily; maintenance, 360 to 480 mg/day divided (three to four times daily); maximum, 480 mg/day</p> <p>Extended-release tablet: maintenance, 180 to 480 mg/day</p>	<p>Safety and efficacy of oral verapamil in children have not been established.</p> <p><u>Arrhythmias in children 0 to 15 years of age:</u>            Injection: weight based dosing administered by slow intravenous injection</p>	<p>Extended-release capsule            100 mg            120 mg            180 mg            200 mg            240 mg            300 mg            360 mg</p> <p>Extended-release tablet:            120 mg            180 mg            240 mg</p> <p>Injection:            2.5 mg/mL</p> <p>Tablet:            40 mg            80 mg            120 mg</p>

**VIII. Effectiveness**

Clinical studies evaluating the safety and efficacy of the miscellaneous calcium-channel blocking agents are summarized in Table 8.

**Table 8. Comparative Clinical Trials with the Calcium-Channel Blocking Agents, Miscellaneous**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Angina</b>				
<p>De Rosa et al.<sup>37</sup> (1998)</p> <p>Diltiazem SR 300 mg QD</p> <p>vs</p> <p>verapamil SR 240 mg QD</p>	<p>DB, XO</p> <p>Men and women 48 to 72 years of age, with stable exertional angina, a positive test for myocardial ischemia and documented coronary artery disease</p>	<p>N=20</p> <p>12 weeks</p>	<p>Primary: Exercise tolerance test: time to onset of angina, time to 1-mm ST-segment depression and total exercise duration</p> <p>Secondary: Heart rate, angina frequency, nitroglycerin use and adverse events</p>	<p>Primary: Time to onset of angina increased significantly in both groups compared to the placebo group (verapamil vs placebo; P&lt;0.05 and diltiazem vs placebo; P&lt;0.005).</p> <p>Time to 1-mm ST-segment depression increased significantly in both groups compared to the placebo group (verapamil vs placebo; P&lt;0.05 and diltiazem vs placebo; P&lt;0.005).</p> <p>Total exercise duration increased significantly in both groups compared to the placebo group (verapamil vs placebo; P&lt;0.05 and diltiazem vs placebo; P&lt;0.005).</p> <p>For each primary endpoint, there was no significant difference between the treatment groups.</p> <p>Secondary: Heart rates were similar between the treatment groups, except resting heart rate was significantly lower in the diltiazem group as compared to the verapamil group (68.5 vs 75.9; P&lt;0.05).</p> <p>Angina frequency and nitroglycerin use decreased significantly in the diltiazem group compared to the placebo group (P&lt;0.05) and to the verapamil group (P&lt;0.05).</p> <p>Edema and flushing were most frequently reported. Similar rates of adverse events were reported for both treatments.</p>
<p>Chugh et al.<sup>38</sup> (2001)</p> <p>Diltiazem 240 mg</p>	<p>DB, DD, PG, RCT</p> <p>Patients with stable angina, blood</p>	<p>N=67</p> <p>4 weeks</p>	<p>Primary: Treadmill exercise test: time to onset of angina, time to</p>	<p>Primary: Both treatment groups, and all doses, had significant increases in time to onset of angina from baseline (P&lt;0.001 for all). There was no significant difference between the treatment groups (P=0.838) and between dose</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD for 2 weeks then 360 mg QD for 2 weeks</p> <p>vs</p> <p>amlodipine 5 mg QD for 2 weeks then 10 mg QD for 2 weeks</p>	<p>pressure in the range of 100/60 to 170/110 mm Hg and a positive ischemic response on a treadmill test, history of angiography</p>		<p>1-mm ST-segment depression</p> <p>Secondary: Heart rate, blood pressure, number of angina episodes and use of nitrates</p>	<p>levels (P=0.144) in time to onset of angina.</p> <p>Both treatment groups, and all doses, had significant increases in time to 1-mm ST-segment depression from baseline, except the low-dose amlodipine group (P&lt;0.004, except P=0.063). There was no significant difference between the treatment groups and between dose levels (P=0.114) in time to 1-mm ST-segment depression (P=0.691).</p> <p>Secondary: There was no significant difference between the groups in heart rate at rest or maximal exercise.</p> <p>There was no significant difference between the groups in blood pressure at rest or maximal exercise, except SBP at rest was higher in the diltiazem group (137 to 143 vs 129 to 135 mm Hg; P=0.029).</p> <p>Both treatments reduced the number of angina episodes and the use of nitrates, but these results were not statistically different between the groups (P value not reported).</p>
<p>van Kesteren et al.<sup>39</sup> (1998)</p> <p>Diltiazem CR 90 to 120 mg BID</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>DB, MC</p> <p>Men and women 41 to 77 years of age with a history of stable angina pectoris, a positive exercise tolerance test, and positive thallium scan or positive coronary angiogram</p>	<p>N=132</p> <p>8 weeks</p>	<p>Primary: Exercise tolerance test: time to 1-mm ST-segment depression, time to onset of chest pain, time to end of exercise (exercise duration)</p> <p>Secondary: Safety</p>	<p>Primary: Diltiazem and amlodipine treatment resulted in significant increases in time to 1-mm ST-segment depression as compared to baseline (P&lt;0.0001). Treatments were not significantly different from each other (P&gt;0.05).</p> <p>Diltiazem and amlodipine treatment resulted in significant increases in time to onset of chest pain at four and eight weeks, (10 and 13% for amlodipine; P&lt;0.0001; 5 and 7% for diltiazem; P=0.009). Treatments were not significantly different from each other (P&gt;0.05).</p> <p>Amlodipine treatment resulted in a significant increase in total exercise duration as compared to baseline (P=0.0002), however the change from baseline for diltiazem was not significantly increased (P=0.43). There was no significant difference between the treatment groups at endpoint.</p> <p>Secondary: Ten patients (15.2%) in the amlodipine group and 17 patients (25.8%) in the diltiazem group reported an adverse event; two patients from the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				amlodipine group and six patients from the diltiazem group subsequently withdrew from the study.
<p>Frishman et al.<sup>40</sup> (1999)</p> <p>Diltiazem 240 to 480 mg at bedtime</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD plus atenolol 50 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients 30 to 80 years of age with chronic stable angina pectoris, evidence of exercise-induced ST-segment depression <math>\geq 1</math> mm and other evidence of cardiac disease</p>	<p>N=551</p> <p>4 week</p>	<p>Primary: Exercise tolerance test (symptom-limited exercise duration, time <math>\geq 1</math>-mm ST-segment depression and time to moderate angina)</p> <p>Secondary: 48-hour Holter-determined number of ischemic episodes, mean and total duration of ischemia, maximal depth of ST depression, heart rate at onset of ischemia</p>	<p>Primary: Treatment with verapamil, amlodipine, and amlodipine plus atenolol resulted in significantly better results than patients treated with placebo in: symptom-limited exercise duration, time <math>\geq 1</math>-mm ST-segment depression and time to moderate angina (<math>P \leq 0.01</math> for all vs placebo).</p> <p>Secondary: Treatment with verapamil, amlodipine, and amlodipine plus atenolol resulted in significantly fewer ischemic episodes in 48-hour Holter monitoring (<math>P = 0.003</math> for verapamil vs placebo).</p> <p>Treatment with amlodipine monotherapy resulted in a significant increase in duration of ischemic episode (<math>P \leq 0.05</math> vs verapamil vs amlodipine plus atenolol and vs placebo).</p> <p>Treatment with verapamil and amlodipine plus atenolol resulted in a decrease in duration of ischemic episodes as compared to treatment with amlodipine and placebo (<math>P \leq 0.05</math> for each).</p> <p>Heart rate at the onset of ischemic episode was significantly lower in the verapamil group and in the amlodipine plus atenolol group (<math>P \leq 0.05</math> vs amlodipine) and higher in the amlodipine group (<math>P \leq 0.05</math> vs verapamil, vs amlodipine plus atenolol and vs placebo).</p>
<p>Hauf-Zachariou et al.<sup>41</sup> (1997)</p> <p>Verapamil 120 mg TID</p> <p>vs</p> <p>carvedilol 25 mg BID</p>	<p>DB, MC, PG, RCT</p> <p>Patients 18 to 75 years with a confirmed diagnosis of CAD, exertional chest pain relieved by rest or glyceryl trinitrate for <math>\geq 2</math> months and 2 exercise tests with signs and symptoms</p>	<p>N=313</p> <p>12 weeks</p>	<p>Primary: Total exercise time, time to onset of angina, and time to 1 mm ST-segment depression, blood pressure, heart rate, rate pressure product</p> <p>Secondary:</p>	<p>Primary: There was not a significant difference in total exercise time observed between the carvedilol (increased from 378 s to 436 s) and verapamil (increased from 386 s to 438 s) groups (RR, 1.14; 90% CI, 0.85±1.52).</p> <p>There was not a significant difference observed between the carvedilol and verapamil groups in time to onset of angina (increase from 296 s to 325 s vs 285 s to 326 s) and in time to 1 mm ST-segment depression (increase from 267 s to 298 s vs 286 s to 302 s).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced SBP (from 175 to 166 mm Hg) compared to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	of ischemia		Not reported	<p>verapamil (from 173 to 173 mm Hg)).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced heart rate (from 123 to 112 mm Hg) compared to verapamil (from 124 to 120 mm Hg)).</p> <p>At peak exercise and at maximum comparable workload, carvedilol significantly reduced rate pressure product (from 21564 to 18802 mm Hg) compared to verapamil (from 21488 to 20992 mm Hg)).</p> <p>Secondary: Not reported</p>
<b>Cardiovascular Outcomes</b>				
<p>Boden et al.<sup>42</sup> (2002) INTERCEPT</p> <p>Diltiazem 300 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PG, PRO, RCT</p> <p>Patients 75 years of age and younger, with acute MI, without CHF and who received a thrombolytic agent</p>	<p>N=874</p> <p>Up to 6 months</p>	<p>Primary: Composite first-event rate of: cardiac death, nonfatal reinfarction or refractory ischemia</p> <p>Secondary: Composite of first occurrence of cardiac death, nonfatal reinfarction, recurrent ischemia, composite of cardiac death, nonfatal reinfarction, need for myocardial revascularization, safety</p>	<p>Primary: There was no significant difference between diltiazem treatment and placebo treatment in composite event rate (131 primary outcome events occurred in the placebo group and 97 occurred in the diltiazem group; P=0.07).</p> <p>Secondary: Rates of all composite nonfatal cardiac events (nonfatal reinfarction combined with refractory ischemia or all recurrent ischemia or need for revascularization) significantly favored the diltiazem group over the placebo group (P=0.05, P=0.05, P=0.03 respectively).</p> <p>Rates of cardiac death, nonfatal reinfarction, refractory ischemia and all recurrent ischemia were similar between the diltiazem group and the placebo group, however the need for revascularization favored the diltiazem group (P=0.67, P=0.47, P=0.07, P=0.07, P=0.03).</p> <p>There was no increase in rates of CHF, bleeding, cancer or cerebrovascular accidents in the diltiazem group.</p>
<p>Gibson et al.<sup>43</sup> (2000)</p>	<p>RETRO combined subgroup analysis</p>	<p>N=817</p>	<p>Primary: All cause mortality</p>	<p>Primary: Patients receiving treatment (either agent) had a 42% lower mortality rate</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Diltiazem 60 mg QID or verapamil 120 mg TID  vs  placebo	of 2 RCT  Patients suffering acute non-Q-wave MI	12 to 18 months	Secondary: Combined cardiac events	than those receiving placebo (P=0.010).  Secondary: Patients receiving treatment (either agent) had a 31% lower event rate (death or recurrent MI) than those receiving placebo (P<0.006).
Hansson et al. <sup>44</sup> (2000) NORDIL  Diltiazem 180 to 360 mg QD  vs  conventional therapy (diuretic, β-blocker or both)	BE, MC, OL, PRO, RCT  Patients 50 to 74 years of age with DBP ≥100 mm Hg and previously untreated	N=10,881  4.5 years	Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death  Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI	Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/β-blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).  Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/β-blocker patients (P=0.04).  Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/β-blocker patients (P=0.17).  Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57 and congestive heart failure (P=0.42).
Pepine et al. <sup>45</sup> (2003) INVEST  Verapamil SR 240 mg/day (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist	MC, OL, RCT  Patients with essential HTN	N=22,576  24 months	Primary: First occurrence of death (all cause), nonfatal MI or stroke  Secondary: Cardiovascular death, angina, cardiovascular hospitalization, angina, blood pressure control (SBP/DBP	Primary: At 24 months, in the calcium antagonist strategy subgroup, 81.5% of patients were taking verapamil SR, 62.9% trandolapril, and 43.7% HCTZ. In the non-calcium antagonist strategy, 77.5% of patients were taking atenolol, 60.3% HCTZ, and 52.4% trandolapril.  After a follow-up of 61,835 patient-years (mean, 2.7 years per patient), 2,269 patients had a primary outcome event with no statistically significant difference between treatment strategies (9.93% in calcium antagonist strategy vs 10.17% in non-calcium antagonist strategy; RR, 0.98; 95% CI, 0.90 to 1.16; P=0.57).  Secondary: There was no significant difference in the rate of cardiovascular death

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>strategy)</p> <p>vs</p> <p>atenolol 50 mg/day (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p> <p>Trandolapril was recommended for all patients with heart failure, diabetes, or renal insufficiency.</p>			<p>&lt;140/90 mm Hg or &lt;130/85 mm Hg if diabetic or renal impairment), safety</p>	<p>(P=0.94) or cardiovascular hospitalization (P=0.59) between the two treatment groups.</p> <p>At 24 months, angina episodes decreased in both groups, but the mean frequency was lower in the calcium antagonist strategy group (0.77 episodes/week) compared to the non-calcium antagonist strategy group (0.88 episodes/week; P=0.02).</p> <p>Two-year blood pressure control was similar between groups. The blood pressure goals were achieved by 65.0% (systolic) and 88.5% (diastolic) of calcium antagonist strategy patients and 64.0% (systolic) and 88.1% (diastolic) of non-calcium antagonist strategy patients. A total of 71.7% of calcium antagonist strategy patients and 70.7% of non-calcium antagonist strategy patients achieved an SBP &lt;140 mm Hg and DBP &lt;90 mm Hg.</p> <p>Both regimens were generally well tolerated. Patients in the calcium antagonist strategy group reported constipation and cough more frequently than patients in the non-calcium antagonist strategy group, while non-calcium antagonist strategy patients experienced more dyspnea, lightheadedness, symptomatic bradycardia and wheezing (all were statistically significant with P≤0.05).</p>
<p>Mancia et al.<sup>46</sup> (2007) INVEST</p> <p>Verapamil SR 120 to 480 mg QD</p> <p>vs</p> <p>atenolol 25 to 200 mg QD</p>	<p>MC, open blinded endpoint, PRO, RCT</p> <p>Patients with HTN, requiring drug therapy (BP&gt;140/90 or &gt;130/80 mm Hg if diabetic or with renal impairment), and CAD</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Occurrence of death, nonfatal MI and nonfatal stroke</p> <p>Secondary: Blood pressure control rates</p>	<p>Primary: Rates (death, nonfatal MI and nonfatal stroke) were similar for both treatment groups (P value not reported).</p> <p>Secondary: Rates of death, MI and stroke declined as the number of office visits for which blood pressure was controlled increased (P&lt;0.001).</p>
<p>Pepine et al.<sup>47</sup> (2006) INVEST</p>	<p>Post hoc analysis of INVEST</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Risk for adverse outcome associated</p>	<p>Primary: Previous heart failure (adjusted HR, 1.96), as well as diabetes (HR, 1.77), increased age (HR, 1.63), United States residency (HR, 1.61), renal</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Verapamil SR (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p> <p>vs</p> <p>atenolol (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p>	<p>Patients with essential HTN</p>		<p>with baseline factors, follow-up blood pressure and drug treatments</p> <p>Secondary: Not reported</p>	<p>impairment (HR, 1.50), stroke/TIA (HR, 1.43), smoking (HR, 1.41), MI (HR, 1.34), PVD (HR, 1.27), and revascularization (HR, 1.15) predicted increased risk.</p> <p>Follow-up SBP &lt;140 mm Hg (HR, 0.82) or DBP &lt;90 mm Hg (HR, 0.70) and trandolapril with verapamil SR (HR, 0.78 and 0.79) were associated with reduced risk.</p> <p>Secondary: Not reported</p>
<p>Bangalore et al.<sup>48</sup> (2008) INVEST</p> <p>Verapamil SR 120 to 480 mg QD</p> <p>vs</p> <p>atenolol 25 to 200 mg QD</p> <p>Trandolapril</p>	<p>INVEST substudy</p> <p>Patients 50 years of age and older with hypertension requiring drug therapy (blood pressure &gt;140/90 or &gt;130/80 mm Hg if diabetic or with renal impairment), and documented coronary artery</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: First occurrence of death, nonfatal MI, nonfatal stroke</p> <p>Secondary: Death, total MI, total stroke</p>	<p>Primary: No significant difference was observed between groups in the primary endpoint (P=0.30).</p> <p>Among patients with the primary outcome, no significant difference was observed between groups in the risk of death (P=0.94).</p> <p>There was no significant difference between groups in the risk of nonfatal MI (P=0.41).</p> <p>There was a trend toward a 29% reduction in the risk of nonfatal stroke in the verapamil group compared to the atenolol group (P=0.06).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
and/or HCTZ were added to control blood pressure.	disease			<p>Secondary: The risks of fatal and nonfatal MI were similar between groups.</p> <p>No significant differences were observed between groups in fatal and nonfatal stroke (P=0.18).</p>
<p>Brunner et al.<sup>49</sup> (2007) INVEST</p> <p>Verapamil SR 240 mg and trandolapril 1 to 4 mg</p>	<p>Post hoc analysis of INVEST</p> <p>Patients with essential HTN</p>	<p>N=1,832</p> <p>24 months</p>	<p>Primary: Factors influencing blood pressure response to trandolapril add-on therapy</p> <p>Secondary: Not reported</p>	<p>Primary: Trandolapril decreased mean unadjusted SBP and DBP by -9.1 and -4.1 mm Hg, respectively. The percentage of patients with blood pressure under control (&lt;140/90 mm Hg) increased from 6.7 to 41.3% (P&lt;0.0001).</p> <p>Adjusted blood pressure response was significantly associated with age and baseline SBP and DBP (P&lt;0.0001). Whereas the decrease in SBP was more pronounced in younger patients, the opposite was observed for DBP decrease.</p> <p>DBP response was significantly associated with race. Specifically, the adjusted DBP decrease was significantly smaller in Hispanics and African Americans than whites (P=0.0032 and P=0.0069, respectively). However, Hispanics achieved a decrease in SBP and an increase in blood pressure control similar to the other ethnic groups.</p> <p>Secondary: Not reported</p>
<p>Black et al.<sup>50</sup> (2003) CONVINCE</p> <p>Verapamil ER 180 mg QD</p> <p>vs</p> <p>atenolol 50 mg QD</p> <p>vs</p> <p>HCTZ 12.5 mg</p>	<p>AC, DB, MC, RCT</p> <p>Patients 55 years of age and older with HTN and ≥1 risk factor for cardiovascular disease</p>	<p>N=16,476</p> <p>3 years</p>	<p>Primary: Composite first occurrence of acute MI, stroke or cardiovascular disease-related death</p> <p>Secondary: Cardiovascular endpoints expanded, all-cause mortality, cancer,</p>	<p>Primary: There was no significant difference between the verapamil treatment group and the atenolol or HCTZ treatment groups in the composite primary endpoint (HR, 1.02; 95% CI, 0.88 to 1.18; P=0.77).</p> <p>Secondary: There was no significant difference between the verapamil treatment group and the atenolol or HCTZ treatment group in rates of cardiovascular-related hospitalization (P=0.31), death (all-cause mortality) (P=0.32) and cancer rates (P=0.46).</p> <p>Patients treated with verapamil experienced a significantly higher rate of death or bleeding unrelated to stroke (HR, 1.54; 95% CI, 1.15 to 2.04; P=0.003).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
QD			hospitalization for bleeding, incidence of primary endpoints between 6AM and noon, adverse events	<p>Primary endpoints did not differ significantly based on time of day (P=0.43).</p> <p>Patients treated with verapamil were more likely to withdraw for adverse events or symptoms than those treated with atenolol or HCTZ (P=0.02).</p>
<p>Lindholm et al.<sup>51</sup> (2005)</p> <p>Other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)</p> <p>or</p> <p>placebo</p> <p>vs</p> <p>β-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol, or propranolol)</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a β-blocker as first-line treatment (in ≥50% of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>The RR of stroke was 16% higher with β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non β-blockers (RR, 1.26%; 95% CI, 1.15 to 1.38; P&lt;0.0001).</p> <p>The relative risk of MI was 2% higher for β-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).</p> <p>The RR of all-cause mortality was 3% higher for β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Wysong et al.<sup>52</sup> (2007)</p> <p>Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)</p> <p>vs</p> <p>β-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)</p>	<p>MA</p> <p>13 RCTs evaluating patients ≥18 years of age with HTN</p>	<p>N=91,561</p> <p>Duration varied</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions</p>	<p>Primary: There was not a significant difference observed in all-cause mortality between β-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with β-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).</p> <p>Secondary: There was a significant decrease in stroke observed with β-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between β-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with β-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of β-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with β-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Hypertension</b>				
<p>Wright et al.<sup>53</sup> (2004)</p> <p>Diltiazem graded-release 360 to 540 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>AC, DB, MC, PG, RCT</p> <p>Male and female African Americans patients 18 to 80 years of age with hypertension (DBP 85 to 109 mm Hg and SBP &lt;180 mm Hg)</p>	<p>N=268</p> <p>12 weeks</p>	<p>Primary: Change from baseline in DBP during first 4 hours of awakening as recorded by ambulatory blood pressure monitoring</p> <p>Secondary: Changes from baseline in BP, heart rate, rate-pressure product, safety</p>	<p>Primary: Reductions in DBP during the first four hours after awakening, and from 6AM to noon, were significantly greater in the diltiazem group than in the amlodipine group (-13.12 vs -9.65 mm Hg; P=0.0049 and -11.97 vs -8.75 mm Hg; P=0.0019).</p> <p>Secondary: Reductions in SBP during the first four hours after awakening and between 6AM and noon, were similar between the groups (P&lt;0.0768 and P&lt;0.9470).</p> <p>Mean 24-hour SBP reductions were significantly greater in the amlodipine group than in the diltiazem group (-14.08 vs -10.64; P=0.0022).</p> <p>Reductions in heart rate were significantly greater in the diltiazem group than in the amlodipine group (24 hour mean: -4.88 vs 1.77; P&lt;0.0001).</p> <p>Reductions in rate-pressure product were significantly greater in the diltiazem group than in the amlodipine group (24 hour mean: -1,493 vs -881; P&lt;0.0008).</p> <p>In the diltiazem and amlodipine groups respectively, 1.5 and 2.2% discontinued early due to adverse events.</p>
<p>White et al.<sup>54</sup> (2004)</p> <p>Diltiazem ER 240 to 540 mg at bedtime</p> <p>vs</p> <p>ramipril 5 to 20 mg at bedtime</p>	<p>DB, MC, PG, RCT</p> <p>Men and women, with hypertension: DBP 90 to 110 mm Hg</p>	<p>N=261</p> <p>10 weeks</p>	<p>Primary: Change in early morning DBP from baseline</p> <p>Secondary: Change in SBP from baseline, heart rate, heart rate × systolic blood pressure product, 24-hr ambulatory</p>	<p>Primary: Changes in early morning DBP were significantly larger in the diltiazem group than in the ramipril group (-15 vs -8 mm Hg; P&lt;0.001).</p> <p>Secondary: Changes in early morning SBP were significantly larger in the diltiazem group than in the ramipril group (-18 vs -13 mm Hg; P=0.002).</p> <p>Decreases in heart rate and heart-rate systolic BP product were significantly larger in the diltiazem group than in the ramipril group (-8.9 vs -2.7 beats/min; P&lt;0.0001 and -2518 vs -1393; P&lt;0.0001).</p> <p>Reductions in DBP and heart rate and increases in the rate-pressure</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			monitoring, safety	<p>product measured by 24-hr ambulatory monitoring and clinic monitoring were significantly greater for diltiazem than for ramipril (P&lt;0.0001 for all).</p> <p>50% of diltiazem patients and 40% of ramipril patients reported experiencing any adverse event; edema and cough respectively were most frequently reported for each treatment. Withdrawal rates from the study were low and similar between the groups.</p>
<p>Rosei et al.<sup>55</sup> (1997) VHAS</p> <p>Verapamil SR 240 mg QD</p> <p>vs</p> <p>chlorthalidone 25 mg QD</p>	<p>DB (1st 6 months), MC, PG, RCT</p> <p>Patients 40 to 65 years of age, with HTN (SBP ≥160 mm Hg and DBP ≥95 mm Hg)</p>	<p>N=1,414</p> <p>2 years</p>	<p>Primary: Blood pressure</p> <p>Secondary: Cardiovascular events, adverse events</p>	<p>Primary: Both treatments significantly reduced SBP and DBP compared to baseline, however reductions did not significantly differ between treatments (verapamil reduction, 27.6/17.0 mm Hg vs chlorthalidone reduction, 28.6/16.6 mm Hg; P&lt;0.01 for each vs baseline).</p> <p>Goal DBP was achieved in 69.3% of patients receiving verapamil and 66.9% of patients receiving chlorthalidone (P value not reported).</p> <p>Secondary: Serum TC levels and heart rate decreased significantly in the verapamil group as compared to baseline and the chlorthalidone group (TC; P&lt;0.01 for both, heart rate; P&lt;0.05).</p> <p>The number of nonfatal cardiovascular events was similar between the groups, 37 in the verapamil group and 39 in the chlorthalidone group (P value not reported).</p> <p>The number of cardiovascular deaths was similar between the groups, five in the verapamil group and four in the chlorthalidone group (P value not reported).</p> <p>Hypokalemia and hyperuricemia occurred significantly more frequently in the chlorthalidone group than in the verapamil group (P&lt;0.01 for both).</p> <p>Two hundred and thirty six patients reported 403 adverse events in the chlorthalidone group and 230 patients reported 387 adverse events in the verapamil group. Asthenia was the most commonly reported adverse event in the chlorthalidone group and constipation was the most commonly</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Ruggenenti et al.<sup>56</sup> (2004) BENEDICT</p> <p>Trandolapril 2 mg/day</p> <p>vs</p> <p>verapamil SR 240 mg/day</p> <p>vs</p> <p>trandolapril and verapamil SR 2-180 mg/day (fixed-dose combination)</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, RCT</p> <p>Patients ≥40 years with type 2 diabetes (not exceeding 25 years) and HTN (SBP ≥130 mm Hg and/or DBP ≥85 mm Hg ) but with normoalbuminuria (urinary albumin excretion rate of &lt;20 mcg/minute)</p>	<p>N=1,204</p> <p>3.6 years (median)</p>	<p>Primary: Development of persistent microalbuminuria comparing combination therapy to placebo, acceleration factor</p> <p>Secondary: Primary end point comparing trandolapril and verapamil monotherapy to placebo, blood pressure, adverse events</p>	<p>reported adverse event in the verapamil group.</p> <p>Primary: The primary outcome was reached in 5.7% of patients receiving combination therapy vs 10.0% for patients receiving placebo. The estimated acceleration factor (which quantifies the effect of one treatment relative to another in accelerating or slowing disease progression) adjusted for predefined baseline characteristics was 0.39 for the comparison between verapamil plus trandolapril and placebo (P=0.01).</p> <p>Secondary: The primary outcome was reached in 6.0% of patients receiving trandolapril, 11.9% receiving verapamil, and 10.0% receiving placebo. The estimated acceleration factor was 0.47 for trandolapril vs placebo (P=0.01) and 0.83 for verapamil vs placebo (P=0.54).</p> <p>Trandolapril plus verapamil and trandolapril alone delayed the onset of microalbuminuria by factors of 2.6 and 2.1, respectively.</p> <p>Throughout the study the average trough SBP/DBP was 139/80 mm Hg for patients receiving trandolapril plus verapamil, 139/81 mm Hg for trandolapril, 141/82 mm Hg for verapamil and 142/83 mm Hg for placebo. The comparison was significant (P≤0.002) between trandolapril plus verapamil or trandolapril alone vs placebo, but not for verapamil vs placebo.</p> <p>Serious adverse events were similar in all treatment groups.</p>
<p>Messerli et al.<sup>57</sup> (2006)</p> <p>Verapamil SR 240 mg QD</p> <p>vs</p> <p>trandolapril 4 mg QD</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients, 21 years old and older with DBP of 95 to 114 mm Hg</p>	<p>N=581</p> <p>6 weeks</p>	<p>Primary: Blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: All 3 treatment groups had significant blood pressure reductions from baseline (P&lt;0.01 for all).</p> <p>Patients receiving the combination of trandolapril and verapamil had significantly greater reductions in blood pressure as compared to patients receiving trandolapril or verapamil alone (P&lt;0.01 for both comparisons).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  verapamil SR 240 mg and trandolapril 4 mg QD (separate entities)				
Karlberg et al. <sup>58</sup> (2000)  Trandolapril 2 mg/day  vs  verapamil 240 mg/day  vs  trandolapril and verapamil 2-180 mg/day (fixed-dose combination)	DB, MC, PRO, RCT, XO  Patients with uncomplicated primary HTN (sitting DBP between 95 and 115 mm Hg) between the ages of 20 to 80 years	N=226  2 months	Primary: Change in blood pressure and rate pressure product  Secondary: Predictive value of plasma concentrations of active renin regarding the blood pressure response to the different treatment regimens, safety	Primary: The mean fall in blood pressure was significantly greater with the combination (20/15 mm Hg; P<0.00054), as compared to trandolapril (14/11 mm Hg) or verapamil (13/11) mm Hg. The difference between verapamil and trandolapril was not significant.  Rate pressure product decreased significantly more on the combination (P<0.001) than on trandolapril or verapamil alone.  Secondary: There was a significant positive correlation between blood pressure fall and plasma concentrations of active renin (e.g., the higher the initial active renin, the better the blood pressure response to trandolapril [P<0.045 for SBP and P<0.004 for DBP]). No relationships were found for either verapamil or the combination.  All treatments were well tolerated and safe.
Van Bortel et al. <sup>59</sup> (2008)  ACE inhibitor, ARB, β-blocker, calcium channel blocker, or placebo  vs  nebivolol	MA  12 RCTs involving >25 patients with essential HTN where nebivolol 5 mg QD was compared to placebo or other active drugs for >1 month	N=2,653  Duration varied	Primary: Antihypertensive effect and tolerability  Secondary: Not reported	Primary: Overall, higher response rates were observed with nebivolol than all other antihypertensive agents combined (OR, 1.41; 95% CI, 1.15 to 1.73; P=0.001) and compared to the ACE inhibitors (OR, 1.92; 1.30 to 2.85; P=0.001), but response rates to nebivolol were similar to β-blockers (OR, 1.29; 95% CI, 0.81 to 2.04; P=0.283), calcium channel blockers (OR, 1.19; 95% CI, 0.83 to 1.70; P=0.350) and losartan (OR, 1.35; 95% CI, 0.84 to 2.15; P=0.212).  Overall, a higher percentage of patients obtained normalized blood pressure with nebivolol compared to the other antihypertensive agents combined (OR, 1.35; 95% CI, 1.07 to 1.72; P=0.012). A higher percentage



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>of patient receiving nebivolol obtained normalized blood pressure compared to losartan (OR, 1.98; 95% CI, 1.24 to 3.15; P=0.004) and calcium channel blockers (OR, 1.96; 95% CI, 1.05 to 1.96; P=0.024), but not when compared to other <math>\beta</math>-blockers (OR, 1.29; 95% CI, 0.81 to 1.65; P=0.473).</p> <p>Overall, the percentage of adverse events was significantly lower with nebivolol compared to the other antihypertensive agents combined (OR, 0.59; 95% CI, 0.48 to 0.72; P&lt;0.001) and similar to placebo (OR, 1.16; 95% CI, 0.76 to 1.67; P=0.482). In comparing nebivolol to the individual treatments, nebivolol had a lower percentage of adverse events compared to losartan (OR, 0.52; 95% CI, 0.30 to 0.89; P=0.016), the other <math>\beta</math>-blockers (OR, 0.56; 95% CI, 0.36 to 0.85; P=0.007) and calcium channel blockers (OR, 0.49; 95% CI 0.33 to 0.72; P&lt;0.001), but was similar to ACE inhibitors (OR, 0.75; 95% CI 0.52 to 1.08).</p> <p>Secondary: Not reported</p>
<p>Hilleman et al.<sup>60</sup> (1999)</p> <p>Amlodipine-benazepril (fixed-dose combination)</p> <p>vs</p> <p>monotherapy (atenolol, HCTZ, captopril, enalapril, lisinopril, amlodipine, diltiazem, nifedipine, verapamil)</p>	<p>MA</p> <p>Patients with mild to moderate essential HTN</p>	<p>82 trials</p> <p><math>\geq 4</math> weeks</p>	<p>Primary: Absolute change in supine DBP from baseline</p> <p>Secondary: Percent of patients who achieved blood pressure control, safety</p>	<p>Primary: The mean absolute decrease in supine DBP ranged from 9.7 to 13.3 mm Hg with verapamil showing the greatest effect and captopril the least. When studies were weighted by sample size, amlodipine and benazepril, atenolol, lisinopril, and verapamil showed the greatest blood pressure effect.</p> <p>Secondary: The average percentage of patients defined as controlled after treatment varied from 53.5 to 79.0%, with amlodipine and benazepril (74.3%) and lisinopril (79.0%) showing the highest percentage control (P=0.096).</p> <p>The incidence of adverse events ranged from 12.1 to 41.8%, with lisinopril and verapamil showing the lowest incidences (12.1% and 14.1%, respectively) and nifedipine the highest incidence. Lisinopril demonstrated significantly less overall side effects compared to nifedipine (P=0.030).</p> <p>Nifedipine demonstrated a higher withdrawal rate due to side effects compared to atenolol, HCTZ, enalapril, amlodipine, and diltiazem</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				(P=0.002). Although amlodipine and benazepril had the lowest rate of withdrawals due to adverse events, lack of significant change was due to the low number of cohorts available for analysis.
<p>Casas et al.<sup>61</sup> (2005)</p> <p>ACE inhibitor or ARBs compared to placebo</p> <p>vs</p> <p>ACE inhibitor or ARBs compared to other antihypertensive drugs (<math>\beta</math>-adrenergic blocking agents, <math>\alpha</math>-adrenergic blocking agents, calcium-channel blocking agents, or combinations)</p> <p>Specific agents and doses were not specified.</p>	<p>MA (127 trials)</p> <p>Studies in adults that examined the effect of any drug treatment with a blood pressure lowering action on progression of renal disease</p>	<p>N=not reported</p> <p>4.2 years (mean)</p>	<p>Primary: Doubling of serum creatinine, and ESRD</p> <p>Secondary: Serum creatinine, urine albumin excretion and GFR</p>	<p>Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>Secondary: Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).</p> <p>Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).</p> <p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>
<b>Miscellaneous</b>				
<p>Siu et al.<sup>62</sup> (2009)</p> <p>Diltiazem IV 0.25 mg/kg to 10 mg/kg</p> <p>vs</p>	<p>OL, RCT</p> <p>Patients who presented to the emergency room with symptomatic acute atrial fibrillation for &lt;48</p>	<p>N=150</p> <p>3 years</p>	<p>Primary: Sustained ventricular rate control (&lt;bpm) within 24 hours</p> <p>Secondary: Time to ventricular</p>	<p>Primary: The time to ventricular control for the 45 patients assigned to diltiazem was achieved 90% of the time compared to digoxin (74%) and amiodarone (74%) (P&lt;0.0001).</p> <p>Secondary: The median time to ventricular control was significantly shorter in the diltiazem group (3 hours, 1-21 hours) compared to the digoxin (6 hours, 3</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
digoxin IV 0.5 mg to 0.25 mg  vs  amiodarone IV 300 mg to 10 mg/kg	hours and rapid ventricular rate >120 bpm necessitating hospitalization		control, atrial fibrillation symptom improvement, hospital stay, and adverse events	to 15 hours, P<0.001) and amiodarone groups (7 hours, 1 to 18 hours, P=0.003).  The diltiazem group had the largest reduction in atrial fibrillation frequency score and severity score (P<0.0001).  Length of hospital stay was significantly shorter in the diltiazem group (3.9±1.6 days) compared to digoxin (4.7±2.1 days, P=0.023) and amiodarone groups (4.7±2.2 days, P=0.038).

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, CR=controlled-release, ER=extended-release, IV=intravenous, QD=once daily, SR=sustained-release, TID=three times daily

Study design abbreviations: AC=active comparator, BE=blinded endpoint, DB=double blind, DD=double dummy, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, RETRO=retrospective, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor blocker, CAD=coronary artery disease, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, ESRD=end stage renal disease, GFR=glomerular filtration rate, HCTZ=hydrochlorothiazide, HTN=hypertension, HR=hazard ratio, MI=myocardial infarction, OR=odds ratio, PVD=peripheral vascular disease, RR=relative risk, SBP=systolic blood pressure, TC=total cholesterol, TIA=transient ischemic attack

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 9. Relative Cost of the Calcium-Channel Blocking Agents, Miscellaneous**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Diltiazem	extended-release capsule, extended-release tablet, injection, tablet	Cardizem <sup>®*</sup> , Cardizem CD <sup>®*</sup> , Cardizem LA <sup>®</sup> , Matzim LA <sup>®</sup> , Tiazac ER <sup>®*</sup>	\$\$\$	\$\$
Verapamil	extended-release capsule, extended-release tablet, injection, tablet	Calan SR <sup>®*</sup> , Verelan <sup>®*</sup> , Verelan PM <sup>®*</sup>	\$\$\$\$ to \$\$\$\$\$	\$\$\$

\*Generic is available in at least one dosage form or strength.

**X. Conclusions**

The miscellaneous calcium-channel blocking agents are approved for the treatment of angina, arrhythmias and hypertension.<sup>1,2,6-14</sup> Diltiazem and verapamil are available in a variety of modified-release delivery systems that alter their pharmacokinetic properties, including onset and duration of action.<sup>1,2</sup> Both drugs are available in a generic formulation.

There are several national and international guidelines that provide recommendations regarding the use of calcium-channel blocking agents.<sup>15-36</sup> For the treatment of chronic angina,  $\beta$ -blockers are recommended as initial therapy; however, long-acting calcium-channel blocking agents may be used if  $\beta$ -blockers are contraindicated or if additional therapy is required.<sup>15-20</sup> Calcium-channel blocking agents are recommended as initial therapy in patients

with variant/vasospastic angina.<sup>16,19</sup> Verapamil may be considered for secondary prevention of cardiovascular disease in patients with no heart failure in whom  $\beta$ -blockers are contraindicated.<sup>21</sup> Treatment options for atrial fibrillation include ventricular rate control or drug therapy to maintain sinus rhythm. The AFFIRM, RACE, and HOT CAFE trials demonstrated similar outcomes with rate control compared to rhythm control strategies.  $\beta$ -blockers and nondihydropyridine calcium-channel blocking agents are recommended for patients with persistent or permanent atrial fibrillation, either alone or in combination with digoxin.<sup>25-27</sup> For the treatment of heart failure, ACE inhibitors, ARBs, aldosterone antagonists, and isosorbide dinitrate/hydralazine are recommended as initial therapy. In general, calcium-channel blocking agents are not recommended for the routine treatment of heart failure.<sup>22-23</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>28-33</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>28</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>28-35</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>28-35</sup>

Clinical trials demonstrate that diltiazem and verapamil can effectively treat angina and improve blood pressure.<sup>37-41,53-61</sup> Both agents have been shown to reduce mortality and cardiovascular event rates compared to placebo.<sup>43</sup> Evidence suggests that there is no overall difference between diltiazem and verapamil compared to other antihypertensive agents ( $\beta$ -blockers, atenolol, diuretics) in reducing cardiovascular events and mortality in patients with hypertension.<sup>44-50</sup>

There is insufficient evidence to support that one brand miscellaneous calcium-channel blocking agent is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand miscellaneous calcium-channel blocking agents within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand miscellaneous calcium-channel blocking agent is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. *Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.*
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. Kannam JP, Gersh BJ. Calcium channel blockers in the management of chronic coronary syndrome. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Jan]. Available from: <http://www.utdol.com/utd/index.do>.
4. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
5. Kaplan NM, Victor RG, Flynn JT. Kaplan's clinical hypertension. 11<sup>th</sup> ed. Philadelphia (PA): Lippincott, Williams, and Wilkins; 2015.
6. Cardizem® [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Jun.
7. Cardizem CD® [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Apr.
8. Cardizem LA® [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Apr.
9. Tiazac® [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Oct.
10. Calan® [package insert]. New York (NY): Pfizer Inc; 2017 Sep.
11. Calan® SR [package insert]. New York (NY): Pfizer Inc; 2022 Jan.
12. Verelan® [package insert]. Philadelphia (PA): Lannett Company, Inc.; 2019 Oct.
13. Verelan PM® [package insert]. Philadelphia (PA): Lannett Company, Inc.; 2019 Oct.
14. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
15. Fraker T, Fihn S, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of chronic stable angina: a report of the American College of Cardiology/American Heart Association task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *Circulation*. 2007 Dec 4;116(23):2762-72.
16. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425
17. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline From the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
18. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
19. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>
20. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.
21. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
22. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].

23. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
24. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
25. January CT, Wann L, Calkins H, et al. 2019 Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2019;140:e125-e151. doi:10.1161/CIR. 000000000000665.
26. National Institute for Health and Care Excellence. Atrial fibrillation: diagnosis and management. NG196. 2021. <https://www.nice.org.uk/guidance/ng196>.
27. Frendl G, Sodickson AC, Chung MK, et al. 2014 AATS Guidelines for the Prevention and Management of Peri-Operative Atrial Fibrillation and Flutter (POAF) for Thoracic Surgical Procedures. *The Journal of thoracic and cardiovascular surgery*. 2014;148(3):e153-e193. doi:10.1016/j.jtcvs.2014.06.036.
28. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
29. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
30. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi:10.1016/j.cjca.2020.02.086.
31. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
32. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
33. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
34. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
35. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
36. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
37. De Rosa ML, Giordano A, Melfi M, Della Guardia D, Ciaburri F, Rengo F. Antianginal efficacy over 24 hours and exercise hemodynamic effects of once daily sustained-release 300 mg diltiazem and 240 mg verapamil in stable angina pectoris. *Int J Cardiol*. 1998;63(1):27-35.
38. Chugh SK, Digpal K, Hutchinson T, McDonald CJ, Miller AJ, Lahiri A. A randomized, double-blind comparison of the efficacy and tolerability of once-daily modified-release diltiazem capsules with once-daily amlodipine tablets in patients with stable angina. *J Cardiovasc Pharmacol*. 2001 38(3):356-64.
39. van Kesteren HA, Withagen AJ. A comparative study of once-daily amlodipine versus twice-daily diltiazem controlled-release (CR) in the treatment of stable angina pectoris. *Amlodipine Study Group. Cardiovasc Drugs Ther*. 1998;12 Suppl 3:233-7.
40. Frishman WH, Glasser S, Stone P, Deedwania PC, Johnson M, Fakouhi TD. Comparison of controlled-onset, extended-release verapamil with amlodipine and amlodipine plus atenolol on exercise performance

- and ambulatory ischemia in patients with chronic stable angina pectoris. *Am J Cardiol.* 1999 Feb 15;83(4):507-14.
41. Hauf-Zachariou U, Blackwood RA, Gunawardena KA, O'Donnell JG, Garnham S, Pfarr E. Carvedilol versus verapamil in chronic stable angina: a multicentre trial. *Eur J Clin Pharmacol.* 1997;52(2):95-100.
  42. Boden WE, van Gilst WH, Scheldewaert RG, et al. Diltiazem in acute myocardial infarction treated with thrombolytic agents: a randomized placebo-controlled trial. Incomplete Infarction Trial of European Research Collaborators Evaluating Prognosis post-Thrombolysis (INTERCEPT) *Lancet.* 2002; 355(9217):1751-6.
  43. Gibson RS, Hansen JF, Messerli F, Schechtman KB, Boden WE. Long-term effects of diltiazem and verapamil on mortality and cardiac events in non-Q-wave acute myocardial infarction without pulmonary congestion: post hoc subset analysis of the multicenter diltiazem post infarction trial and the second Danish verapamil infarction trial studies. *Am J Cardiol.* 2000; 86(3):275-9.
  44. Hansson L, Hedner T, Lund-Johansen P, et al. Randomized trial of effects of calcium antagonists compared with diuretics and beta-blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet.* 2000;356(9227):359-65.
  45. Pepine CJ, Handberg EM, Cooper-DeHoff RM, et al. A calcium antagonist vs a non-calcium antagonist hypertension treatment strategy for patients with coronary artery disease: the international verapamil-trandolapril study (INVEST): a randomized controlled trial. *JAMA.* 2003 Dec 3;290(21):2805-16.
  46. Mancia G, Messerli F, Bakris G et al. Blood Pressure Control and Improved Cardiovascular Outcomes in the International Verapamil SR-Trandolapril Study. *Hypertension* 2007;50:299-305.
  47. Pepine CJ, Kowey PR, Kupfer S, et al. Predictors of adverse outcome among patients with hypertension and coronary artery disease. *J Am Coll Cardiol.* 2006;47(3):547-51.
  48. Bangalore S, Messerli F, Cohen J, Bacher P, Sleight P, Mancia G, et al. Verapamil-sustained release-based treatment strategy at reducing cardiovascular events in patients with prior myocardial infarction: an International Verapamil SR-Trandolapril (INVEST) substudy. *Am Heart J.* 2008;156:241-7.
  49. Brunner M, Cooper-DeHoff RM, Gong Y, et al; for the INVEST Investigators. Factors influencing blood pressure response to trandolapril add-on therapy in patients taking verapamil SR (from the International Verapamil SR/Trandolapril [INVEST] Study). *Am J Cardiol.* 2007;99:1549-54.
  50. Black HR, Elliott WJ, Grandits G, et al. Principal results of the Controlled Onset Verapamil Investigation of Cardiovascular End Points (CONVINCE) trial. *JAMA.* 2003;289(16):2073-82.
  51. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet.* 2005 Oct 29-Nov 4;366(9496):1545-53.
  52. Wysong CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev.* 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
  53. Wright JT Jr, Sica DA, Gana TJ, Bohannon K, Pascual LG, Albert KS. Antihypertensive efficacy of night-time graded-release diltiazem vs morning amlodipine in African Americans. *Am J Hypertens.* 2004;17(9):734-42.
  54. White WB, Lacourciere Y, Gana T, Pascual MG, Smith DH, Albert KS. Effects of graded-release diltiazem versus ramipril, dosed at bedtime, on early morning blood pressure, heart rate, and the rate-pressure product. *Am Heart J.* 2004;148(4):628-34.
  55. Rosei EA, Dal Palu C, Leonetti G, Magnani B, Pessina A, Zanchetti A. Clinical results of the Verapamil in Hypertension and Atherosclerosis Study. VHAS Investigators. *J Hypertens.* 1997;15(11):1337-44.
  56. Ruggenti P, Fassi A, Ilieva AP, et al; Bergamo Nephrologic Diabetes Complications Trial (BENEDICT) Investigators. *N Engl J Med.* 2004 Nov 4;351(19):1941-51.
  57. Messerli F, Frishman Wh, Elliott W et al. Antihypertensive properties of a high-dose combination of trandolapril and verapamil-SR. *Blood Pressure* 2006;16:6-9.
  58. Karlberg BE, Andrup M, Oden A; on behalf of the Swedish Tarka Trialists. Efficacy and safety of a new long-acting drug combination, trandolapril/verapamil as compared to monotherapy in primary hypertension. *Blood Pressure* 2000;9:140-5.
  59. Van Bortel LM, Fici F, Mascagni F. Efficacy and tolerability of nebivolol compared with other antihypertensive drugs: a meta-analysis. *Am J Cardiovasc Drugs.* 2008;8(1):35-44.
  60. Hilleman DE, Ryschon KL, Mohiuddin SM, Wurdeman RL. Fixed-dose combination vs monotherapy in hypertension: a meta-analysis evaluation. *J Hum Hypertens.* 1999;13:477-83.
  61. Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet.* 2005 Dec 10;366:2026-33.



62. Siu CW, Lau CP, Lee WL, Lam KF, Tse HF. Intravenous diltiazem is superior to intravenous amiodarone or digoxin for achieving ventricular rate control in patients with acute uncomplicated atrial fibrillation. *Crit Care Med* 2009;37:2174-79.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Angiotensin-Converting Enzyme Inhibitors  
AHFS Class 243204  
May 4, 2022**

**I. Overview**

The renin-angiotensin-aldosterone system (RAAS) is the most important component in the homeostatic regulation of blood pressure. Excessive activity of the RAAS may lead to hypertension, as well as fluid and electrolyte disorders. Renin catalyzes the conversion of angiotensinogen to angiotensin I. Angiotensin I is then cleaved to angiotensin II by angiotensin-converting enzyme (ACE). Angiotensin II may also be generated through other pathways (angiotensin I convertase). Angiotensin II can increase blood pressure by direct vasoconstriction, as well as through actions on the brain and autonomic nervous system. In addition, angiotensin II stimulates aldosterone synthesis from the adrenal cortex, leading to sodium and water reabsorption. Angiotensin II exerts other detrimental effects, which include ventricular hypertrophy and remodeling and myocyte apoptosis.<sup>1-2</sup>

The ACE inhibitors are approved for the treatment of diabetic nephropathy, heart failure, hypertension, and post-myocardial infarction.<sup>3-19</sup> They block the conversion of angiotensin I to angiotensin II, and also inhibit the breakdown of bradykinin, which is a potent vasodilator. However, this increase in bradykinin also leads to an increase in adverse effects, including cough.<sup>3-19</sup> The ACE inhibitors are available as single entity products, as well as in combination with hydrochlorothiazide or verapamil. Hydrochlorothiazide inhibits the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride. Verapamil dilates coronary and peripheral arteries. It also slows conduction through the AV node, and has negative inotropic and chronotropic effects.<sup>18,19</sup>

The angiotensin-converting enzyme inhibitors that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the products are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Angiotensin-Converting Enzyme Inhibitors Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Benazepril	tablet	Lotensin®*	benazepril
Captopril	tablet	N/A	captopril
Enalapril	solution, tablet	Epaned®*, Vasotec®*	enalapril
Enalaprilat	injection^	N/A	enalaprilat dihydrate
Fosinopril	tablet	N/A	fosinopril
Lisinopril	solution, tablet	Prinivil®*, Qbrelis®, Zestril®*	lisinopril
Moexipril	tablet	N/A	moexipril
Perindopril	tablet	N/A	perindopril
Quinapril	tablet	Accupril®*	quinapril
Ramipril	capsule	Altace®*	ramipril
Trandolapril	tablet	N/A	trandolapril
<b>Combination Products</b>			
Benazepril and hydrochlorothiazide	tablet	Lotensin HCT®*	benazepril and hydrochlorothiazide
Captopril and hydrochlorothiazide	tablet	N/A	captopril and hydrochlorothiazide
Enalapril and hydrochlorothiazide	tablet	Vaseretic®*	enalapril and hydrochlorothiazide
Fosinopril and hydrochlorothiazide	tablet	N/A	fosinopril and hydrochlorothiazide

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Lisinopril and hydrochlorothiazide	tablet	Prinzide <sup>®*</sup> , Zestoretic <sup>®*</sup>	lisinopril and hydrochlorothiazide
Quinapril and hydrochlorothiazide	tablet	Accuretic <sup>®*</sup>	quinapril and hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.

^Product is primarily administered in an institution.

PDL=Preferred Drug List

N/A=Not available

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the angiotensin-converting enzyme inhibitors are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Angiotensin-Converting Enzyme Inhibitors**

Clinical Guideline	Recommendations
American College of Cardiology/American Heart Association: <b>2007 Chronic Angina Focused Update of the 2002 Guidelines for the Management of Patients With Chronic Stable Angina (2007)</b> <sup>20</sup>	<ul style="list-style-type: none"> <li>Aspirin should be started at 75 to 162 mg/day and continued indefinitely in all patients, unless contraindicated.</li> <li>Use of warfarin in conjunction with aspirin and/or clopidogrel is associated with an increased risk of bleeding and should be monitored closely.</li> <li>Patients with hypertension and established coronary artery disease (CAD) should be treated with blood pressure medication(s) as tolerated, including angiotensin-converting enzyme inhibitors (ACE inhibitors) and/or <math>\beta</math>-adrenergic antagonists (<math>\beta</math>-blockers) with the addition of other medications as needed to achieve blood pressure goals of &lt;140/90 or &lt;130/80 mm Hg for patients with chronic kidney disease or diabetes.</li> <li>Long-acting calcium-channel blocking agents or long-acting nitrates may be used if <math>\beta</math>-blockers are contraindicated. Immediate-release and short-acting dihydropyridine calcium channel blockers can increase adverse cardiac events and should not be used.</li> <li>Long-acting calcium channel blockers or long-acting nitrates may be used with <math>\beta</math>-blockers if initial treatment is not successful.</li> <li>ACE inhibitors should be used indefinitely in patients with a left ventricular ejection fraction (LVEF) of <math>\leq 40\%</math> and in those with hypertension, diabetes or chronic kidney disease, unless contraindicated.</li> <li>ACE inhibitors should also be used indefinitely in patients at lower risk (mildly reduced or normal LVEF in whom cardiovascular risk factors remain well controlled and revascularization has been performed), unless contraindicated.</li> <li>Angiotensin II receptor blockers (ARBs) are recommended in patients with hypertension, those who have an indication for an ACE inhibitor and are intolerant to them, who have heart failure, or who have had a myocardial infarction (MI) and have a LVEF of <math>\leq 40\%</math>.</li> <li>ARBs may be considered in combination with an ACE inhibitor for heart failure due to left ventricular systolic dysfunction.</li> <li>Aldosterone blockade is recommended in patients post-MI without significant renal dysfunction or hyperkalemia who are already receiving therapeutic doses of an ACE inhibitor and a <math>\beta</math>-blocker, have a LVEF <math>\leq 40\%</math> and have either diabetes or heart failure.</li> <li>It is beneficial to start and continue <math>\beta</math>-blocker therapy indefinitely in all patients who have had a MI, acute coronary syndrome or left ventricular dysfunction with or without heart failure symptoms, unless contraindicated.</li> <li>Annual influenza vaccination is recommended in patients with cardiovascular disease.</li> </ul>
European Society of Cardiology:	<p><b>Pharmacological management of stable coronary artery disease (CAD) patients</b></p> <ul style="list-style-type: none"> <li>The two aims of the pharmacological management of stable CAD patients are to</li> </ul>

Clinical Guideline	Recommendations
<p><b>Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)<sup>21</sup></b></p>	<p>obtain relief of symptoms and to prevent CV events.</p> <ul style="list-style-type: none"> <li>• Optimal medical treatment indicates at least one drug for angina/ischemia relief plus drugs for event prevention.</li> <li>• It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>• It is indicated to review the patient's response soon after starting therapy.</li> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin, consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</li> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief: <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a beta-blocker and/or a non-DHP-calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</li> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</li> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and beta-blockers avoided.</li> </ul> </li> <li>• Event prevention: <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><u>Treatment in patients with microvascular angina</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>• <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>• Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>• ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>• Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul> <p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>• Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy</li> </ul>

Clinical Guideline	Recommendations
	<p>(DAPT).</p> <ul style="list-style-type: none"> <li>• Aspirin is recommended for elective stenting.</li> <li>• Clopidogrel is recommended for elective stenting.</li> <li>• Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>• GP IIb/IIIa antagonists should be considered for bailout situation only.</li> <li>• Platelet function testing or genetic testing may be considered in specific or high-risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>• Prasugrel or ticagrelor may be considered in specific high-risk situations of elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</li> <li>• Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>• Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>• Prasugrel or ticagrelor is not recommended in low-risk elective stenting.</li> <li>• After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low.</li> <li>• Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months.</li> </ul> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all revascularized patients receive a secondary prevention and be scheduled for follow-up visit.</li> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after second generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk.</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/ American College of Cardiology Foundation/ American Heart Association/ American Association for Thoracic Surgery/ Preventive Cardiovascular Nurses Association/ Society</p>	<p><u>Medical therapy to prevent MI and death in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients</li> </ul>

Clinical Guideline	Recommendations
<p>of Thoracic Surgeons: <b>Management of Stable Ischemic Heart Disease (2012)</b><sup>22</sup></p>	<p>with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</p> <ul style="list-style-type: none"> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</li> <li>• Patients should receive an annual influenza vaccine.</li> </ul> <p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> <li>• Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes (2014)</b><sup>23</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation <math>&lt; 90\%</math>, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications             <ul style="list-style-type: none"> <li>○ Nitrates                 <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy                 <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>○ Beta-adrenergic blockers                 <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval <math>&gt; 0.24</math> second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</li> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</p> <ul style="list-style-type: none"> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> <ul style="list-style-type: none"> <li>○ Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval <math>&gt;0.24</math> seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</li> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> </li> <li>○ Other anti-ischemic interventions <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> </li> </ul> <ul style="list-style-type: none"> <li>• Inhibitors of renin-angiotensin-aldosterone system <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF <math>&lt;0.40</math> and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF <math>&lt;0.40</math> who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine <math>&gt;2.5</math> mg/dL in men or <math>&gt;2.0</math> mg/dL in women) or hyperkalemia (<math>K &gt;5.0</math> mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF <math>&lt;0.40</math>, diabetes mellitus, or heart failure.</li> </ul> </li> <li>• Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include:</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatid or tirofiban.</li> </ul> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet agents <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatid, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> <li>• Anticoagulant therapy <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>• Timing of CABG in relation to use of antiplatelet agents <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</p> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>• Medications at discharge <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</li> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post-NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> </li> <li>• Late hospital and post-hospital oral antiplatelet therapy <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>• Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment</b></p>	<p><b>Pharmacological treatment of ischemia</b></p> <ul style="list-style-type: none"> <li>• Sublingual or intravenous nitrates and early initiation of <math>\beta</math>-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>• Continuation of chronic <math>\beta</math>-blocker therapy is recommended unless the patient is in overt heart failure.</li> <li>• Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> </ul>

Clinical Guideline	Recommendations
<p><b>Elevation (2020)<sup>24</sup></b></p>	<ul style="list-style-type: none"> <li>• In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> <li>• A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds.             <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul> <p><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li>• If at low bleeding risk (HAS-BLED <math>\leq 2</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED <math>\geq 3</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED <math>\geq 3</math> and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>
<p>American College of Cardiology/American Heart Association: <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>25</sup></p>	<p><u>Routine medical therapies: <math>\beta</math>-blockers</u></p> <ul style="list-style-type: none"> <li>• Oral <math>\beta</math>-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral <math>\beta</math>-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li>• <math>\beta</math>-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>• Patients with initial contraindications to the use of <math>\beta</math>-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>• It is reasonable to administer intravenous <math>\beta</math>-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia.</li> </ul> <p><u>Routine medical therapies: renin-angiotensin-aldosterone system inhibitors</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor should be administered within the first 24 hours to all patients with ST-segment elevation myocardial infarction with anterior location, heart failure, or ejection fraction <math>\leq 40\%</math>, unless contraindicated.</li> <li>• An ARB should be given to patients who have indications for but are intolerant of ACE inhibitors.</li> <li>• ACE inhibitors are reasonable for all patients with no contraindications to their use.</li> <li>• An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq 40\%</math> and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>• High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>• It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>26</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of ST-elevation myocardial infarction (STEMI)</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>• Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>• A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>• In patients with an indication for oral anticoagulation, oral anticoagulants are indicated in addition to antiplatelet therapy.</li> <li>• In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the</li> </ul>

Clinical Guideline	Recommendations
	<p>estimated risk of recurrent coronary events and bleeding).</p> <ul style="list-style-type: none"> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age <math>\geq 50</math> years, and at least one of the following risk factors: age <math>\geq 65</math> years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR <math>&lt; 60</math> mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</li> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with <math>\beta</math>-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with <math>\beta</math>-blockers is indicated in patients with heart failure or left ventricular dysfunction, LVEF <math>\leq 40\%</math> unless contraindicated.</li> <li>• Intravenous <math>\beta</math>-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> <li>• Intravenous <math>\beta</math>-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of <math>&lt; 1.8</math> mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C <math>&gt; 1.8</math> mmol/L (<math>&gt; 70</math> mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction <math>\leq 40\%</math> and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
<p>American College of Cardiology/ American Heart Association: <b>Guideline on the Primary Prevention of Cardiovascular Disease</b></p>	<p><u>Top 10 messages for the primary prevention of cardiovascular disease</u></p> <ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular</li> </ul>

Clinical Guideline	Recommendations
(2019) <sup>27</sup>	<p>disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</p> <ul style="list-style-type: none"> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</li> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</li> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be <math>&lt; 130/80</math> mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul> <p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a decision is made for statin therapy, a moderate-intensity statin should be recommended.</li> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) patients, LDL-C</li> </ul>

Clinical Guideline	Recommendations
	<p>levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</p> <ul style="list-style-type: none"> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> </li> <li>• In patients at borderline risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul> <p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>• In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include: <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>• In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>• In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>• In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>• In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than 130/80 mm Hg.</li> <li>• In adults with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering medication are recommended.</li> <li>• In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>• All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>• To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>• In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>• In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> <li>• To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>• All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>• Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>• Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</li> <li>• Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: <b>2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)</b><sup>28</sup></p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>• Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>• In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>• In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>• Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>• In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>• In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with LVEF <math>\leq</math>40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF <math>\leq</math>50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF <math>\leq</math>50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;</math>30 mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;</math>5.0 mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;</math>5.5 mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide</li> </ul>

Clinical Guideline	Recommendations
	<p>dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</p> <ul style="list-style-type: none"> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>29</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III–IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and</li> </ul>

Clinical Guideline	Recommendations
	<p>electrolytes during use of intravenous diuretics.</p> <ul style="list-style-type: none"> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g., with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>30</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>

Clinical Guideline	Recommendations
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>31</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP ≥160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (≥80 years)</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</p> <ul style="list-style-type: none"> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (≥80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or β-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR &lt;45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> &gt;4.5 mmol/L.</li> <li>○ Consider β-blockers at any treatment step when there is a specific indication for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>32</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of ≥100 mmHg or average systolic blood pressure (SBP) measurements of ≥160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings ≥90 mmHg or for average SBP readings ≥140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>• For high-risk patients, aged 50 years or older, with SBP levels ≥130 mmHg, intensive management to target a SBP &lt;120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A β-blocker (in patients &lt;60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or β-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a β-blocker. The combination of an</li> </ul>

Clinical Guideline	Recommendations
	<p>ACE inhibitor and an ARB is not recommended.</p> <ul style="list-style-type: none"> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination</li> </ul>



Clinical Guideline	Recommendations
	<p>or radiography.</p> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction &lt;40%), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary</li> </ul>

Clinical Guideline	Recommendations
	<p>protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</p> <ul style="list-style-type: none"> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyl dopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyl dopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>33</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed</li> </ul>

Clinical Guideline	Recommendations
	<p>by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</p> <ul style="list-style-type: none"> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing</li> </ul>

Clinical Guideline	Recommendations
	<p>proteinuria, is not recommended.</p> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq</math>220 mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq</math>140/90 mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq</math>130 mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>34</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq</math>160 mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with</li> </ul>

Clinical Guideline	Recommendations
	<p>diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</p> <ul style="list-style-type: none"> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>35</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease</b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease</u></p>



Clinical Guideline	Recommendations
<p><b>(2021)</b><sup>36</sup></p>	<p><u>(CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults</b> (2017)<sup>37</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>• Adults with SIHD and hypertension (BP <math>\geq</math>130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq</math>300 mg/d, or <math>\geq</math>300 mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and</li> </ul>

Clinical Guideline	Recommendations
	<p>close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</p> <ul style="list-style-type: none"> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq</math>220/120 mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq</math>140/90 mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with planned elective major surgery and SBP <math>\geq</math>180 mmHg or DBP <math>\geq</math>110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>38</sup></p>	<p><b>Hypertension/blood pressure control</b></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq</math>140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq</math>180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq</math>15%), a blood pressure target of <math>&lt;</math>130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;</math>15%), treat to blood pressure target of <math>&lt;</math>140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;</math>120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq</math>300 mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or</li> </ul>

Clinical Guideline	Recommendations
	<p>diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</p> <ul style="list-style-type: none"> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>

\*Agent is not available in the United States.

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the angiotensin-converting enzyme inhibitors are noted in Tables 3 and 4. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Angiotensin-Converting Enzyme Inhibitors<sup>3-18</sup>**

Indication(s)	Single Entity Agents									
	Benaze- pril	Capto- pril	Enala- pril	Fosino- pril	Lisino- pril	Moexi- pril	Perindo- pril	Quina- pril	Rami- pril	Trandola- pril
<b>Cardiovascular Risk Reduction</b>										
In patients 55 years or older at high risk of developing a major cardiovascular event because of a history of coronary artery disease, stroke, peripheral vascular disease, or diabetes that is accompanied by at least one other cardiovascular risk factor to reduce the risk of myocardial infarction, stroke, or death from cardiovascular causes									✓	
Stable coronary artery disease to reduce the risk of cardiovascular mortality or nonfatal myocardial infarction							✓			
<b>Diabetic Nephropathy</b>										
Treatment of diabetic nephropathy in patients with type 1 insulin-dependent diabetes and retinopathy		✓								
<b>Heart Failure</b>										
Congestive heart failure		✓ *	✓ *							
Heart failure				✓ †	✓ ‡			✓ †		
<b>Hypertension</b>										
Hypertension	✓ §	✓	✓	✓ §	✓	✓ §	✓	✓ §	✓ §	✓
<b>Left Ventricular Dysfunction</b>										
Decrease the rate of the development of overt heart failure and decrease the incidence of hospitalization for heart failure in clinically stable asymptomatic patients with left ventricular dysfunction (ejection fraction ≤35%)			✓							
<b>Myocardial Infarction</b>										
Hemodynamically stable patients within 24 hours of acute myocardial infarction to improve survival					✓					
Improve survival following myocardial infarction in clinically stable patients with left ventricular dysfunction manifested as an ejection fraction ≤40%		✓								

Indication(s)	Single Entity Agents									
	Benazepril	Captopril	Enalapril	Fosinopril	Lisinopril	Moexipril	Perindopril	Quinapril	Ramipril	Trandolapril
and to reduce the incidence of overt heart failure and subsequent hospitalizations for congestive heart failure in these patients										
Stable patients who have demonstrated clinical signs of congestive heart failure within the first few days after sustaining acute myocardial infarction									✓	
Stable patients who have evidence of left ventricular systolic dysfunction or who are symptomatic from congestive heart failure within the first few days after sustaining acute myocardial infarction										✓

\*Usually in combination with diuretics and digitalis.

† As adjunctive therapy when added to conventional therapy including diuretics with or without digitalis.

‡ As adjunctive therapy in patients who are not responding adequately to diuretics and digitalis.

§ May be used alone or in combination with thiazide diuretics.

|| May be used alone or in combination with other antihypertensive agents.

**Table 4. FDA-Approved Indications for the Angiotensin-Converting Enzyme Inhibitors<sup>3-18</sup>**

Indication(s)	Combination Products					
	Benazepril and HCTZ	Captopril and HCTZ	Enalapril and HCTZ	Fosinopril and HCTZ	Lisinopril and HCTZ	Quinapril and HCTZ
<b>Hypertension</b>						
Hypertension	✓ *	✓	✓ *	✓ *	✓ *	✓ *

\*This fixed combination product is not indicated for the initial therapy of hypertension.

HCTZ=hydrochlorothiazide



#### IV. Pharmacokinetics

The pharmacokinetic parameters of the angiotensin-converting enzyme inhibitors are listed in Table 5.

**Table 5. Pharmacokinetic Parameters of the Angiotensin-Converting Enzyme Inhibitors<sup>19</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Benazepril	37	96.7	Liver, extensive (% not reported)	Renal (33) Bile (12)	22*
Captopril	70 to 75	25 to 30	Liver (50)	Renal (95)	1.9†
Enalapril	60	50 to 60	Liver (70)	Renal (61) Feces (33)	11*
Fosinopril	30 to 36	89 to 100	Liver, extensive (% not reported)	Renal (44) Feces (46)	12*
Lisinopril	25	Minimal (% not reported)	Liver (7)	Renal (29) Feces (69)	12†
Moexipril	13 to 22	50 to 70	Liver, extensive (% not reported)	Renal (13) Feces (50)	2 to 10*
Perindopril	20 to 30	60	Liver (88 to 96)	Renal (75) Feces (25)	3 to 10*
Quinapril	50	97	Liver, extensive (% not reported)	Renal (50 to 60) Feces (33)	2 to 25*
Ramipril	60	73	Liver, extensive (% not reported)	Renal (40 to 60) Feces (40)	13 to 17*
Trandolapril	10	80	Liver, extensive (% not reported)	Feces (66) Renal (33)	16 to 24*
<b>Combination Products</b>					
Benazepril and HCTZ	37/70	96.7/40 to 70	Liver, extensive (% not reported)/ not reported	Feces (11 to 12/ Renal (70)	22*/10
Captopril and HCTZ	70 to 75/70	25 to 30/ not reported	Liver (50%)/ not reported	Renal (>95)/ Renal (% not reported)	<3/2.5
Enalapril and HCTZ	60/70	Not reported/40	Not reported/ Liver, minimal (% not reported)	Renal (61)/ Renal (60)	11/5.6 to 14.8
Fosinopril and HCTZ	36/50 to 80	95/67.9	Liver (% not reported)/ Not reported	Not reported/ Renal (61)	Not reported/ 5 to 15
Lisinopril and HCTZ	25/not reported	Not reported/ Not reported	Not reported/ Not reported	Not reported/ Not reported	Not reported/ Not reported
Quinapril and HCTZ	60/50 to 80	97/67.9	Liver (% not reported)/ Not metabolized	Renal (96)/ Renal (61)	2 to 25*/ 4 to 15

\*Metabolites

†Parent compound

HCTZ=hydrochlorothiazide

#### V. Drug Interactions

Major drug interactions with the angiotensin-converting enzyme inhibitors are listed in Table 6.

**Table 6. Major Drug Interactions with the Angiotensin-Converting Enzyme Inhibitors<sup>19</sup>**

<b>Generic Name(s)</b>	<b>Interaction</b>	<b>Mechanism</b>
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Potassium-sparing diuretics	Combining ACE inhibitors and potassium-sparing diuretics may result in elevated serum potassium concentrations in certain high-risk patients.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Sacubitril	Concurrent use of sacubitril and ACE Inhibitors may result in Increased risk of angioedema.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Aliskiren	The risk of hyperkalemia may be increased when ACE inhibitors are combined with aliskiren.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Angiotensin II receptor antagonists	The risk of hyperkalemia may be increased when ACE inhibitors are combined with angiotensin II receptor antagonists.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Indomethacin	Indomethacin inhibits prostaglandin synthesis. The hypotensive effect of ACE inhibitors may be reduced.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	NSAIDs and salicylates	NSAIDs and salicylates inhibit prostaglandin synthesis. The hypotensive and vasodilator effects of the ACE inhibitor may be reduced.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Potassium preparations	Hyperkalemia, possibly with cardiac arrhythmias or cardiac arrest, may occur with the combination of ACE inhibitors and potassium preparations.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Everolimus, sirolimus	Concurrent use of ACE inhibitors and MTOR inhibitors may result in increased risk of angioedema.
ACE inhibitors (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril,trandolapril)	Trimethoprim	Hyperkalemia, possibly with cardiac arrhythmias or cardiac arrest, may occur with the combination of ACE inhibitors and trimethoprim.
ACE inhibitors	Lithium	Through an unknown mechanism, ACE inhibitors

Generic Name(s)	Interaction	Mechanism
(benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, quinapril, ramipril, trandolapril)		may increase lithium levels, which results in neurotoxicity.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics increase potassium excretion. Hypokalemia may occur, increasing the risk of torsades de pointes.
Thiazide diuretics (HCTZ)	Lithium	Thiazide diuretics decrease the renal clearance of lithium which leads to increased serum lithium levels. Lithium toxicity has occurred.
Thiazide diuretics (HCTZ)	Diazoxide	Hyperglycemia may occur with symptoms similar to diabetes. The mechanism is unknown.
Thiazide diuretics (HCTZ)	Digitalis glycosides	Diuretic-induced electrolyte disturbances may predispose the patient to digitalis-induced cardiac arrhythmias.

ACE inhibitor=angiotensin converting enzyme inhibitor, HCTZ=hydrochlorothiazide, HMG CoA=3-hydroxy-3-methyl-glutaryl-CoA, NSAID=nonsteroidal anti-inflammatory drug

## VI. Adverse Drug Events

The most common adverse drug events reported with the angiotensin-converting enzyme inhibitors are listed in Tables 7 and 8. The boxed warning for the angiotensin-converting enzyme inhibitors is listed in Table 9.

**Table 7. Adverse Drug Events (%) Reported with the Angiotensin-Converting Enzyme Inhibitors-Single Entity Agents<sup>3-19</sup>**

Adverse Events	Benazepril	Captopril	Enalapril	Fosinopril	Lisinopril	Moexipril	Perindopril	Quinapril	Ramipril	Trandolapril
<b>Cardiovascular</b>										
Angina	<1	<1	2	<1	-	<1	-	<1	<1 to 3	-
Bradycardia	-	-	<1	<1	<1	-	-	-	<1	<5
Cardiac arrest	-	✓	<1	✓	<1	-	✓	-	✓	-
Cerebrovascular accident	-	✓	<1	<1	<1	<1	<1	<1	<1	-
Chest pain	-	1	2	<2	3	>1	2	2	<1	<1
Hypotension	<1	✓	1 to 7	1 to 4	1 to 10	<1	<1	3	<1	<1
Myocardial infarction	-	<1	<1	<1	<1	<1	<1	<1	<1	-
Orthostatic hypotension	<1	✓	1 to 2	<2	<1	<1	<1	<1	2	-
Palpitations	<1	1	<1	<1	<1	<1	<1	<1	<1	<1
Peripheral edema	<1	-	-	-	<1	>1	-	-	-	-
Rhythm disturbances	-	✓	<1	<1	-	<1	-	<1	-	-
Tachycardia	-	1	<1	<1	<1	-	-	<1	<1	-
<b>Central Nervous System</b>										
Anxiety	<1	-	-	-	-	<1	<1	-	<1	<1
Ataxia	-	✓	<1	-	<1	-	-	-	-	-
Depression	-	✓	<1	<1	-	-	2	<1	<1	-
Dizziness	4	-	1 to 8	2 to 12	5 to 12	4	8	4 to 8	2 to 4	1 to 23
Fatigue	2	-	1 to 3	≥1	3	2	-	3	2	-
Headache	6	-	2 to 5	≥1	4 to 6	>1	24	2	-	-
Insomnia	<1	-	<1	<1	<1	<1	3	<1	<1	<1
Malaise	-	-	-	-	<1	<1	<1	<1	<1	-
Nervousness	<1	✓	<1	<1	<1	<1	1	<1	<1	-
Paresthesias	<1	-	<1	<1	<1	-	2	<1	<1	<1
Peripheral edema	<1	-	-	-	-	>1	-	-	-	-
Somnolence/drowsiness	2	✓	<1	<1	<1	<1	1	<1	<1	<1
Vertigo	-	-	2	<1	<1	-	<1	<1	<1 to 2	<1
<b>Dermatologic</b>										
Alopecia	<1	-	<1	-	<1	<1	-	<1	-	-
Diaphoresis	<1	-	<1	<1	<1	<1	<1	<1	<1	-
Erythema multiforme	-	✓	<1	-	-	-	<1	-	<1	-
Exfoliative dermatitis	-	✓	<1	✓	-	-	✓	<1	-	-
Flushing	<1	<1	<1	<1	<1	2	-	-	-	<1
Pemphigus/pemphigoid	<1	✓	<1	-	<1	-	-	<1	-	<1
Photosensitivity	<1	✓	<1	<1	<1	<1	-	<1	-	-
Pruritus	<1	2	<1	<1	-	<1	<1	<1	<1	<1
Rash	<1	4 to 7	<1	<1	<1	2	2	1	<1	<1
Stevens-Johnson syndrome	<1	✓	<1	-	✓	-	-	-	<1	-
Toxic epidermal necrolysis	-	-	<1	-	✓	-	-	-	<1	-

Adverse Events	Benazepril	Captopril	Enalapril	Fosinopril	Lisinopril	Moexipril	Perindopril	Quinapril	Ramipril	Trandolapril
Urticaria	-	-	<1	<1	<1	<1	-	<1	<1	-
<b>Gastrointestinal</b>										
Abdominal pain	-	-	2	<1	2	<1	3	1	<1	<1
Anorexia	-	-	<1	-	-	-	-	-	<1	-
Constipation	<1	-	<1	<1	<1	<1	<1	<1	<1	<1
Diarrhea	-	-	1 to 2	>1	3 to 4	3	4	2	<1	<1
Dry mouth	-	-	<1	<1	<1	<1	<1	<1	<1	-
Dysgeusia	-	2 to 4	-	-	-	-	-	-	-	-
Dyspepsia	-	✓	<1	-	<1	>1	<1	<1	<1	<6
Hepatitis	-	✓	<1	<1	<1	<1	-	<1	<1	-
Nausea	1	-	1	1 to 2	2	>1	2	2	2	-
Pancreatitis	<1	✓	<1	<1	<1	<1	✓	<1	<1	<1
Vomiting	<1	-	1	1 to 2	<1	<1	2	2	2	<1
<b>Genitourinary</b>										
Decreased libido	<1	-	-	<1	<1	-	-	-	-	<1
Impotence	<1	✓	<1	-	1	-	-	<1	<1	<1
Oliguria	-	<1	<1	-	<1	<1	-	-	-	-
Urinary tract infection	<1	-	1	-	<1	-	3	<1	-	-
<b>Musculoskeletal</b>										
Arthralgia	<1	✓	✓	<1	<1	<1	<1	<1	<1	-
Arthritis	<1	-	✓	✓	<1	-	1	-	<1	-
Muscle cramps	-	-	<1	<1	<1	-	-	-	-	<1
Myalgia	<1	✓	✓	<1	<1	1	<1	-	<1	5
<b>Respiratory</b>										
Asthma	<1	✓	<1	-	<1	-	-	-	-	-
Bronchitis	<1	-	1	-	<1	-	<1	-	-	-
Bronchospasm	-	✓	<1	<1	<1	<1	-	2 to 4	-	-
Cough	1	<2	1 to 2	2 to 10	1 to 4	6	6 to 12	2 to 4	8	2 to 35
Dyspnea	<1	-	1	≥1	<1	<1	<1	<1	<1	<1
Pharyngitis	-	-	-	<1	<1	2	3	<1	-	-
Rhinitis	-	✓	-	<1	<1	>1	5	-	-	-
Sinusitis	<1	-	-	<1	<1	>1	<5	-	-	-
Upper respiratory tract infection	-	-	<1	2	2	>1	7	-	✓	<1
<b>Miscellaneous</b>										
Anemia	✓	<1	-	✓	<1	<1	-	<1	<1	-
Angioedema	<1	<1	✓	<1	<1	<1	<1	<1	<1	<1
Asthenia	<1	✓	1 to 2	-	1	-	8	-	2	3
Blurred vision	-	✓	<1	-	<1	-	-	-	-	-
Eosinophilia	-	✓	✓	✓	<1	-	-	-	<1	-
Fever	-	✓	<1	<1	<1	-	<1	-	<1	-
Syncope	<1	✓	1 to 2	<1	<2	<1	<1	<1	<2	6
Tinnitus	-	-	<1	<1	<1	<1	2	-	<1	-
Vasculitis	-	✓	✓	-	<1	-	✓	-	<1	-

✓ Percent not specified  
- Event not reported

**Table 8. Adverse Drug Events (%) Reported with the Angiotensin-Converting Enzyme Inhibitors-Combination Products<sup>3-19</sup>**

Adverse Event	Benazepril and HCTZ	Captopril and HCTZ	Enalapril and HCTZ	Fosinopril and HCTZ	Lisinopril and HCTZ	Quinapril and HCTZ
<b>Cardiovascular</b>						
Angina	-	0.2 to 0.3	-	-	-	-
Cardiac arrest	-	✓	-	-	-	-
Cerebrovascular accident	-	✓	-	-	-	-
Chest pain	-	1	-	0.5 to <2.0	-	1
Hypotension	0.6	✓	-	-	1.4	-
Myocardial infarction	-	0.2 to 0.3	-	-	-	-
Orthostatic hypotension	0.3 to 3.5	✓	2.3	1.8	0.5	≥0.5 to <1.0
Palpitations	-	1	0.5 to 2.0	-	-	≥0.5 to <1.0
Tachycardia	-	1	-	-	-	-
<b>Central Nervous System</b>						
Depression	-	✓	-	-	-	-
Dizziness	6.3	-	8.6	3.2	7.5	4.8
Fatigue	5.2	-	3.9	3.9	3.7	2.9
Headache	3.1	-	5.5	7	5.2	6.7
Insomnia	✓	-	0.5 to 2.0	-	-	1.2
Somnolence/drowsiness	1.2	✓	-	-	-	1.2
<b>Dermatologic</b>						
Flushing	0.3 to 1.0	0.2 to 0.5	✓	0.5 to <2.0	-	-
Pruritus	-	2	-	-	-	✓
Rash	-	4 to 7	✓	0.5 to <2.0	1.2	-
Stevens-Johnson syndrome	✓	✓	✓	✓	-	✓
<b>Gastrointestinal</b>						
Abdominal pain	-	-	-	-	-	1.7
Diarrhea	0.3 to 1.0	✓	2.1	0.5 to <2.0	2.5	1.4
Dysgeusia	-	2 to 4	-	-	-	-
Dyspepsia	-	✓	-	-	-	-
Hepatitis	-	✓	-	-	-	-
Jaundice	✓	✓	✓	✓	-	✓
Nausea	1.4	-	2.5	✓	2.2	✓
Pancreatitis	-	✓	-	-	-	-
<b>Genitourinary</b>						
Decreased libido	-	✓	-	✓	-	-
Impotence	1.2	-	2.2	-	1.2	≥0.5 to <1.0
Oliguria	-	0.1 to 0.2	-	-	-	-
<b>Musculoskeletal</b>						
Hypertonia	1.5	-	-	-	-	-
Muscle cramps	-	-	2.7	-	2	-
Musculoskeletal pain	-	-	-	2	-	-
Myalgia	-	✓	-	-	-	2.4
<b>Respiratory</b>						
Bronchitis	-	-	-	-	-	1.2
Cough	2.1	0.5 to 2.0	3.5	5.6	3.9	3.2
Rhinitis	-	✓	-	-	-	2
Upper respiratory tract infection	-	-	-	2.3	2.2	1.3

Adverse Event	Benazepril and HCTZ	Captopril and HCTZ	Enalapril and HCTZ	Fosinopril and HCTZ	Lisinopril and HCTZ	Quinapril and HCTZ
<b>Miscellaneous</b>						
Anemia	-	≤0.2	-	-	-	-
Angioedema	0.3	0.1	0.5 to 2.0	0.5 to <2.0	0.3 to 1.0	0.1
Asthenia	-	✓	2.4	-	1.8	≥0.5 to <1.0
Blurred vision	-	✓	-	-	-	-
Eosinophilia	-	✓	-	-	-	-
Fever	-	✓	-	-	-	-
Neutropenia	-	✓	-	0.5 to <2.0	-	-
Syncope	-	✓	-	-	-	-
Viral infection	-	-	-	-	-	1.9

HCTZ=hydrochlorothiazide

- ✓ Percent not specified
- Event not reported

**Table 9. Boxed Warning for the Angiotensin-Converting Enzyme Inhibitors<sup>18</sup>**

<b>WARNING</b>
When pregnancy is detected, discontinue therapy as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

## VII. Dosing and Administration

The usual dosing regimens for the angiotensin-converting enzyme inhibitors are listed in Table 10.

**Table 10. Usual Dosing Regimens for the Angiotensin-Converting Enzyme Inhibitors<sup>3-19</sup>**

<b>Generic Name(s)</b>	<b>Usual Adult Dose</b>	<b>Usual Pediatric Dose</b>	<b>Availability</b>
<b>Single Entity Agents</b>			
Benazepril	<u>Hypertension:</u> Tablet: initial, 5 to 10 mg once daily (for patients not receiving diuretics); maintenance, 20 to 40 mg/day as a single dose or in two equally divided doses; maximum, >80 mg/day has not been evaluated	<u>Hypertension for children ≥6 years of age:</u> Tablet: initial, 0.2 mg/kg once daily; maximum, >0.6 mg/kg (or in excess of 40 mg daily) has not been studied  Safety and efficacy in children <6 years of age have not been established.	Tablet: 5 mg 10 mg 20 mg 40 mg
Captopril	<u>Diabetic nephropathy:</u> Tablet: maintenance, 25 mg three times daily  <u>Heart failure:</u> Tablet: initial, 25 mg three times daily; maximum, 450 mg/day  <u>Hypertension:</u> Tablet: initial, 25 mg two to three times daily; maintenance, after one to two weeks can increase to 50 mg two to three times daily; maximum: 450 mg/day  <u>Myocardial infarction (left ventricular dysfunction after myocardial infarction):</u> Tablet: initial, 6.25 mg once, followed by 12.5 mg three times daily; target maintenance, 50 mg three times daily	Safety and efficacy in children have not been established.	Tablet: 12.5 mg 25 mg 50 mg 100 mg
Enalapril	<u>Heart failure:</u> Solution, tablet: initial, 2.5 mg/day; maintenance, 2.5 to 20 mg two times daily; maximum, 40 mg/day in divided doses  <u>Hypertension:</u> Solution, tablet: initial, 5 mg once daily; maintenance, 10 to 40 mg/day as a single dose or in two divided doses	<u>Hypertension in children 1 month to 16 years of age:</u> Solution, tablet: initial, 0.08 mg/kg (up to 5 mg) once daily; maximum, >0.58 mg/kg (or in excess of 40 mg) has not been studied  Safety and efficacy in children <1 month have not been established.	Solution: 1 mg/mL  Tablet: 2.5 mg 5 mg 10 mg 20 mg



Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<u>Left ventricular dysfunction:</u> Solution ,tablet: initial, 2.5 mg two times daily; target maintenance, 10 mg/day in divided doses		
Fosinopril	<u>Heart failure:</u> Tablet: initial, 10 mg once daily; maintenance, 20 to 40 mg/day; maximum, 40 mg once daily  <u>Hypertension:</u> Tablet: initial, 10 mg once daily; maintenance, 20 to 40 mg/day in a single or divided dose(s); maximum, 80 mg/day	<u>Hypertension in children 6 to 16 years of age:</u> Tablet (>50 kg): 5 to 10 mg once daily  Safety and efficacy in children <6 years of age have not been established.	Tablet: 10 mg 20 mg 40 mg
Lisinopril	<u>Heart failure:</u> Solution, tablet: initial, 5 mg once daily; maintenance, 5 to 20 mg once daily  <u>Hypertension:</u> Solution, tablet: initial, 10 mg once daily; maintenance, 20 to 40 mg once daily  <u>Post-myocardial infarction:</u> Solution, tablet: initial, 5 mg every 24 hours for two doses, followed by 10 mg every day for 6 weeks	<u>Hypertension in children 6 to 16 years of age:</u> Solution, tablet: initial, 0.07 mg/kg (up to 5 mg) once daily; doses >0.6 mg/kg (or in excess of 40 mg) have not been studied  Safety and efficacy in children <6 years of age have not been established.	Solution: 1 mg/ mL  Tablet: 2.5 mg 5 mg 10 mg 20 mg 30 mg 40 mg
Moexipril	<u>Hypertension:</u> Tablet: initial, 7.5 mg once daily; maintenance, 7.5 to 30 mg/day in a single or divided dose(s); maximum, 60 mg/day	Safety and efficacy in children have not been established.	Tablet: 7.5 mg 15 mg
Perindopril	<u>Cardiovascular risk reduction (coronary artery disease):</u> Tablet: initial: 4 mg once daily for 2 weeks; maintenance, increase as tolerated to 8 mg once daily  <u>Hypertension:</u> Tablet: initial, 4 mg once daily; maintenance, 4 to 8 mg/day in a single or divided dose(s); maximum, 16 mg/day	Safety and efficacy in children have not been established.	Tablet: 2 mg 4 mg 8 mg
Quinapril	<u>Heart failure:</u> Tablet: initial, 5 mg twice daily; maintenance, titrate at weekly intervals to 10 to 20 mg two times daily  <u>Hypertension:</u> Tablet: initial, 10 to 20 mg once daily; maintenance, 20 to 80 mg/day in a single or divided dose(s)	Safety and efficacy in children have not been established.	Tablet: 5 mg 10 mg 20 mg 40 mg
Ramipril	<u>Cardiovascular risk reduction:</u>	Safety and efficacy in	Capsule:

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<p>Capsule: initial, 2.5 mg once daily for one week, followed by 5 mg once daily for three weeks; maintenance, 10 mg once daily</p> <p><u>Hypertension:</u> Capsule: initial, 2.5 mg once daily; maintenance, 2.5 to 20 mg/day in single or divided dose(s)</p> <p><u>Post-myocardial infarction (heart failure after myocardial infarction):</u> Capsule: initial, 2.5 mg twice daily; target maintenance, 5 mg twice daily</p>	children have not been established.	1.25 mg 2.5 mg 5 mg 10 mg
Trandolapril	<p><u>Post-myocardial infarction (left ventricular dysfunction or heart failure after myocardial infarction):</u> Tablet: initial, 1 mg once daily; maintenance, titrate as tolerated to target of 4 mg once daily</p> <p><u>Hypertension:</u> Tablet: initial, 1 mg once daily in non-African American patients and 2 mg once daily in African American patients; maintenance, 2 to 4 mg once daily</p>	Safety and efficacy in children have not been established.	Tablet: 1 mg 2 mg 4 mg
<b>Combination Products</b>			
Benazepril and HCTZ	<p><u>Hypertension:</u> Tablet: initial, 10-12.5 or 20-12.5 mg/day if not adequately controlled on benazepril monotherapy; maintenance, titrate dose by clinical effect</p>	Safety and efficacy in children have not been established.	Tablet: 5-6.25 mg 10-12.5 mg 20-12.5 mg 20-25 mg
Captopril and HCTZ	<p><u>Hypertension:</u> Tablet: initial, 25-5 mg once daily; titrate dose by clinical effect; maximum, 150-50 mg/day</p>	Safety and efficacy in children have not been established.	Tablet: 25-15 mg 25-25 mg 50-15 mg 50-25 mg
Enalapril and HCTZ	<p><u>Hypertension:</u> Tablet: maximum, four tablets of 5-12.5 mg or two tablets of 10-25 mg</p>	Safety and efficacy in children have not been established.	Tablet: 5-12.5 mg 10-25 mg
Fosinopril and HCTZ	<p><u>Hypertension:</u> Tablet: titrate dose by clinical effect</p>	Safety and efficacy in children have not been established.	Tablet: 10-12.5 mg 20-12.5 mg
Lisinopril and HCTZ	<p><u>Hypertension:</u> Tablet: initial, 10-12.5 or 20-12.5 mg/day after failure on monotherapy; titrate dose by clinical effect; maximum, 80-50 mg/day</p>	Safety and efficacy in children have not been established.	Tablet: 10-12.5 mg 20-12.5 mg 20-25 mg
Quinapril and HCTZ	<p><u>Hypertension:</u> Tablet: initial, 10-12.5 or 20-12.5 mg/day; maintenance, titrate dose by clinical effect</p>	Safety and efficacy in children have not been established.	Tablet: 10-12.5 mg 20-12.5 mg 20-25 mg

HCTZ=hydrochlorothiazide

**VIII. Effectiveness**

Clinical studies evaluating the safety and efficacy of the angiotensin-converting enzyme inhibitors are summarized in Table 11.

**Table 11. Comparative Clinical Trials with the Angiotensin-Converting Enzyme Inhibitors**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Cardiovascular Disease</b>				
<p>Jamerson et al.<sup>39</sup> (2008) ACCOMPLISH</p> <p>Benazepril 20 to 40 mg QD and HCTZ 12.5 to 25 mg QD</p> <p>vs</p> <p>benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=11,506</p> <p>36 months (mean)</p>	<p>Primary: The composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for angina, resuscitation after sudden cardiac arrest, and coronary revascularization.</p> <p>Secondary: Death from cardiovascular causes, nonfatal MI, and nonfatal stroke</p>	<p>Primary: There were 552 primary-outcome events in the benazepril plus amlodipine group (9.6%) and 679 events in the benazepril plus HCTZ group (11.8%). The absolute risk reduction with benazepril plus amlodipine therapy was 2.2% and the relative risk reduction was 19.6% compared to benazepril plus HCTZ (HR, 0.80; 95% CI, 0.72 to 0.90; P&lt;0.001).</p> <p>Secondary: For the secondary end point of death from cardiovascular causes, nonfatal MI, and nonfatal stroke, there were 288 (5%) events in the benazepril plus amlodipine group compared to 364 (6.3%) events in the benazepril plus HCTZ group. The absolute risk reduction with benazepril plus amlodipine therapy was 1.3% and the RR reduction was 21.2% compared to benazepril plus HCTZ (HR, 0.79; 95% CI, 0.67 to 0.92; P=0.002).</p>
<p>Weber et al.<sup>40</sup> (2010) ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to 40-10 mg/day, followed by forced titration after 1 month on</p>	<p>Prespecified subanalysis of ACCOMPISH</p> <p>Men and women &gt;60 years of age with HTN and at high risk for cardiovascular events (history of</p>	<p>N=6,946</p> <p>Mean treatment duration 29.7 months for benazepril and amlodipine group and 29.5 months for</p>	<p>Primary: Primary: Time to first event (composite of cardiovascular event and death from cardiovascular causes)</p>	<p>Primary: The primary endpoint occurred in 8.8% of diabetic patients in the benazepril and amlodipine group and 11.0% in the benazepril and HCTZ group (HR, 0.79; P=0.003; NNT, 46). In high risk diabetic patients, 13.6% of patients in the benazepril and amlodipine group and 17.3% in the benazepril and HCTZ group (HR, 0.77, P=0.007; NNT, 28).</p> <p>Secondary: Due to early termination, the study had limited power to detect differences in the diabetic subgroups.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p> <p>benazepril and HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after one month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)</p>	<p>coronary events, MI, revascularization, or stroke; impaired renal function; peripheral arterial disease, left ventricular hypertrophy; or diabetes)</p> <p>(Subanalysis of patients with diabetes)</p>	<p>benazepril and HCTZ group</p>	<p>Secondary: Composite of cardiovascular events (the primary endpoint excluding fatal events) and composite of death from cardiovascular disease, nonfatal stroke and nonfatal MI</p>	<p>Peripheral edema was higher in the benazepril and amlodipine group compared to the benazepril and HCTZ group.</p>
<p>Weber et al.<sup>41</sup> (2013) ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to 40-10 mg/day, followed by forced titration after 1 month on benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p>	<p>Subanalysis of ACCOMPLISH based on body size</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=11,482</p> <p>Duration not specified</p>	<p>Primary: Composite of cardiovascular death or nonfatal MI or stroke</p> <p>Secondary: Cardiovascular death, total MI, total stroke</p>	<p>Primary: In patients receiving benazepril and HCTZ, the primary endpoint (per 1,000 patient-years) was 30.7 in normal weight (BMI &lt;25), 21.9 in overweight (BMI ≥25 to &lt;30), and 18.2 in obese patients (BMI ≥30) (overall P=0.0034). In patients receiving benazepril and amlodipine, the primary endpoint did not differ between the three BMI groups (18.2, 16.9, and 16.5, respectively; P=0.9721). In obese patients, primary event rates were similar between the two treatments, but rates were significantly lower with benazepril and amlodipine in overweight patients (HR, 0.76; 95% CI, 0.59 to 0.94; P=0.0369) and normal weight patients (HR, 0.57; 95% CI, 0.39 to 0.84; P=0.0037).</p> <p>Secondary: Comparing obese and overweight patients, event rates were all numerically lower, but not significantly lower, in obese patients. Cardiovascular deaths were significantly lower in overweight patients compared to normal weight patients (HR, 0.57; 95% CI, 0.37 to 0.89; P=0.0125). Cardiovascular death (HR, 0.40; 95% CI, 0.25 to 0.63;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
benazepril and HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after 1 month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)				P<0.0001) and total stroke (HR, 0.60; 95% CI, 0.37 to 0.96; P=0.0335) were significantly lower in obese patients compared to normal weight patients.
Swedberg et al. <sup>42</sup> (1992) CONSENSUS II  Enalapril 5 to 20 mg/day  vs  placebo  Treatment was started with an IV infusion of 1 mg of enalaprilat administered over 3 hours followed by oral enalapril 6 hours after the infusion was stopped.	DB, MC, PC, PG, RCT  Patients who presented within 24 hours of the onset of acute MI symptoms	N=6,090  180 days	Primary: Mortality rates within 6 months  Secondary: Mortality within 1 month, cause of death, re-infarction, or worsening heart failure	Primary: Mortality rates according to life-table analysis between the enalapril and placebo groups at six months were not significantly different (11 vs 10.2%; P=0.26). The RR associated with enalapril treatment and based on the mortality curves was 1.10 (95% CI, 0.93 to 1.29).  Secondary: Mortality rates between the enalapril and placebo groups at one month were not significantly different (7.2 vs 6.3%; P=0.26).  Death due to progressive heart failure occurred more frequently in patients treated with enalapril than placebo (4.3 vs 3.4%; P=0.06).  There were no significant differences in the rate of reinfarction between the enalapril or placebo groups (P value not significant).  Change in therapy because of heart failure occurred more in the placebo group (P<0.006) but there were no significant differences in hospitalization for heart failure (P value not significant).  Note: The first CONSENSUS trial excluded patients with a recent MI or unstable angina. The study was stopped early after recruiting 6,090 of the intended 9,000 patients since more patients had died on the drug than on placebo (although the difference was not statistically significant).
Wing et al. <sup>43</sup> (2003)	MC, OL, PRO, RCT	N=6,083	Primary: All cardiovascular	Primary: By the end of the study, blood pressure had decreased to a similar extent in

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ANBP2</p> <p>Enalapril</p> <p>vs</p> <p>HCTZ</p> <p>The choice of the specific agent and dose was made by the family practitioner.</p>	<p>Patients 65 to 84 years of age with average SBP while sitting of <math>\geq 160</math> mm Hg or an average DBP of <math>\geq 90</math> mm Hg (if the SBP was <math>\geq 140</math> mm Hg)</p>	<p>4.1 years (median)</p>	<p>events or death from any cause (both initial and subsequent fatal and nonfatal cardiovascular events)</p> <p>Secondary: Not reported</p>	<p>both groups (a decrease of 26/12 mm Hg).</p> <p>There were 695 cardiovascular events or deaths from any cause in the ACE inhibitor group (56.1 per 1,000 patient-years; HR, 0.89; 95% CI, 0.79 to 1.0; P=0.05) compared to 736 in the diuretic group (59.8 per 1,000 patient-years).</p> <p>The beneficial effects of ACE inhibitor treatment were more evident in male subjects (HR, 0.83; 95% CI, 0.71 to 0.97; P=0.02).</p> <p>The rates of nonfatal cardiovascular events and MI decreased with ACE inhibitor treatment, whereas a similar number of strokes occurred in each group (although there were more fatal strokes in the ACE inhibitor group).</p> <p>Secondary: Not reported</p>
<p>Nissen et al.<sup>44</sup> (2004)</p> <p>CAMELOT</p> <p>Enalapril 10 to 20 mg/day</p> <p>vs</p> <p>amlodipine 5 to 10 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients 30 to 79 years of age requiring coronary angiography for evaluation for chest pain or PCI and a diastolic pressure <math>&lt; 100</math> mm Hg, with or without treatment</p>	<p>N=1,991</p> <p>2 years</p>	<p>Primary: Composite of cardiovascular events (cardiovascular death, nonfatal MI, resuscitated cardiac arrest, coronary revascularization, hospitalization for angina pectoris, hospitalization for CHF, fatal or nonfatal stroke or TIA, and any new diagnosis of PVD), nominal change in percent atheroma volume (substudy)</p>	<p>Primary: Cardiovascular events occurred in 23.1% of placebo-treated patients, 16.6% amlodipine-treated patients (HR, 0.69; 95% CI, 0.54 to 0.88; P=0.003) and 20.2% enalapril-treated patients (HR, 0.85; 95% CI, 0.67 to 1.17; P=0.16).</p> <p>The primary end point comparison for enalapril vs amlodipine was not significant (HR, 0.81; 95% CI, 0.63 to 1.04; P=0.10).</p> <p>Secondary: Coronary revascularization was reduced in the amlodipine group from 15.7 to 11.8% (HR, 0.73; 95% CI, 0.54 to 0.98; P=0.03). Hospitalization for angina was reduced in the amlodipine group from 12.8 to 7.7% (HR, 0.58; 95% CI, 0.41 to 0.82; P=0.002).</p> <p>Individual components of the primary end point generally showed fewer events with enalapril treatment vs placebo, but none of the comparisons reached statistical significance.</p> <p>For components of the primary end point, only the rate of hospitalization for angina showed a statistically significant difference between amlodipine</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>Secondary: Incidence of adverse events; all-cause mortality, incidence of revascularization in vessels that had undergone previous stent placement</p>	<p>and enalapril (HR, 0.59; 95% CI, 0.42 to 0.84; P=0.003). A trend toward fewer episodes of revascularization in patients undergoing intervention at baseline was observed for amlodipine vs enalapril (HR, 0.66; 95% CI, 0.40 to 16; P=0.09).</p> <p>The mean change in percent atheroma volume was 0.5% for amlodipine (P=0.12 vs placebo), 0.8% for enalapril (P=0.32 vs placebo) and 1.3% for placebo. In patients with SBP greater than the mean, the amlodipine group showed a significantly slower progression (0.2%) compared to placebo (2.3%; P=0.02). Compared to baseline, intravascular ultrasound showed progression in patients receiving placebo (P&lt;0.001), a trend toward progression with enalapril (P=0.08) and no progression in patients receiving amlodipine (P=0.31). For the amlodipine group, correlation between blood pressure reduction and progression was r=0.19 (P=0.07).</p> <p>Discontinuation from the study for treatment-emergent adverse events was low, averaging 0.4% and not statistically significant between the three treatment groups.</p> <p>The only statistically significant difference in secondary end points was that amlodipine demonstrated a significant reduction in revascularization after previous stent placement compared to placebo (4.1 vs 7.9%; HR, 0.49; 95% CI, 0.31 to 0.78; P=0.002). The rate of revascularization was lower than enalapril (6.2%) but not statistically significant (HR 0.66, 95% CI, 0.40 to 16; P=0.09).</p>
<p>Pitt et al.<sup>45</sup> (2003) 4E-Left Ventricular Hypertrophy Study  Enalapril 40 mg QD  vs</p>	<p>AC, DB, PG, RCT  Patients with left ventricular hypertrophy, a history of HTN and predominantly in sinus rhythm</p>	<p>N=153  9 months</p>	<p>Primary: Change in left ventricular mass as assessed by MRI</p> <p>Secondary: Reduction in SBP and DBP, response rate (DBP &lt;90 mm Hg), change in urine albumin creatinine ratio</p>	<p>Primary: Both treatments were associated with a significant reduction in left ventricular mass from baseline (P&lt;0.001). The difference in left ventricular mass reduction from baseline between the two treatments was not significant (P=0.258).</p> <p>While enalapril plus eplerenone therapy demonstrated a significantly greater reduction in left ventricular mass from baseline compared to eplerenone therapy (P=0.007); the effect was not statistically different from that observed with enalapril therapy (P=0.107).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>eplerenone 200 mg QD</p> <p>vs</p> <p>enalapril 10 mg plus eplerenone 200 mg</p> <p>If the blood pressure was uncontrolled on study medication at week 8, OL HCTZ 12.5 to 25 mg/day and/or amlodipine 10 mg/day were allowed.</p>				<p>The SBP was reduced significantly more in enalapril plus eplerenone-treated patients compared to eplerenone-treated patients (P=0.048). The other treatment groups exhibited statistically comparable reductions from baseline in mean SBP and DBP (P value not reported).</p> <p>While 70.0% of eplerenone-treated patients responded to therapy, 40.7% of enalapril-treated patients responded (P=0.003). In addition, 79.6% of enalapril plus eplerenone-treated patients responded to therapy compared to 40.7% enalapril-treated patients (P=0.001).</p> <p>Enalapril plus eplerenone therapy was associated with a significant reduction in urine albumin creatinine ratio compared to either eplerenone or enalapril therapy (P&lt;0.05).</p> <p>Adverse events were reported with similar incidence among all treatment groups (P value not reported). Cough was significant in enalapril-treated patients compared to eplerenone-treated patients (P=0.033). Two cases of gynecomastia were reported (one eplerenone- and one enalapril plus eplerenone-treated patients). Four patients (three enalapril- and one enalapril plus eplerenone-treated patients) experienced impotence during the trial. Seven eplerenone-, two enalapril- and three enalapril plus eplerenone-treated patients experienced serious hyperkalemia (<math>\geq 6.0</math> mmol/L).</p>
<p>Hansson et al.<sup>46</sup> (1999) STOP-Hypertension</p> <p>Enalapril 10 mg or lisinopril 10 mg QD</p> <p>vs</p> <p>felodipine 2.5 mg or isradipine 2.5 mg QD</p>	<p>MC, OL, PRO, RCT</p> <p>Men and women, age 70 to 84 years with HTN (SBP <math>\geq 180</math>mm Hg or DBP <math>\geq 105</math> mm Hg or both)</p>	<p>N=6,614</p> <p>4 years</p>	<p>Primary: Fatal stroke, fatal MI, other fatal cardiovascular events</p> <p>Secondary: Blood pressure</p>	<p>Primary: The rate of prevention of cardiovascular deaths was similar in all groups (RR, 0.97 to 1.14; 95% CI, 0.86 to 1.26).</p> <p>Fatal cardiovascular events, including fatal stroke and fatal myocardial infarction MI, occurred in 19.8 per 1,000 patient-years in the <math>\beta</math>-blocker and/or HCTZ group, in the felodipine or isradipine group and in the enalapril or lisinopril group (RR, 0.99; 95% CI, 0.84 to 1.16).</p> <p>The RR of cardiovascular death in patients in the enalapril or lisinopril group as compared to the felodipine or isradipine group was 1.14 (95% CI, 0.86 to 1.26; P=0.67.)</p> <p>Secondary:</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  atenolol 50 mg or metoprolol 100 mg or pindolol 5 mg QD and/or HCTZ 25 mg with amiloride 2 to 5 mg QD				Decreases in blood pressure were similar among the groups.
ALLHAT <sup>47</sup> (2002) ALLHAT Lisinopril 10 to 40 mg/day  vs  amlodipine 2.5 to 10 mg/day  vs  chlorthalidone 12.5 to 25 mg/day  Doses were titrated to achieve a goal blood pressure of <140/90 mm Hg.	DB, MC, RCT  Patients ≥55 years with HTN and ≥1 additional CHD risk factor	N=33,357  4.9 years (mean)	Primary: Combined fatal CHD or nonfatal MI  Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization, heart failure, and PAD)	Primary: There were no significant differences in the primary outcome between lisinopril (11.4%), amlodipine (11.3%), and chlorthalidone (11.5%).  Secondary: All-cause mortality did not differ between groups.  Five year SBPs were significantly higher in the lisinopril (2 mm Hg; P<0.001) and amlodipine groups (0.8 mm Hg; P=0.03) compared to chlorthalidone, and five year DBPs were significantly lower with amlodipine (0.8 mm Hg; P<0.001).  Amlodipine had a higher six year rate of heart failure compared to chlorthalidone (10.2 vs 7.7%; RR, 1.38; 95% CI, 1.25 to 1.52).  Lisinopril had a higher six year rate of combined cardiovascular disease (33.3 vs 30.9%; RR, 1.10; 95% CI, 1.05 to 1.16); stroke (6.3 vs 5.6%; RR, 1.15; 95% CI, 1.02 to 1.30) and heart failure (8.7 vs 7.7%; RR, 1.19; 95% CI, 1.07 to 1.31).
Black et al. <sup>48</sup> (2008) ALLHAT Amlodipine 2.5 to	MC, RCT  Men and women, age 55 years old and older, with HTN and	N=17,515  4.9 years (mean)	Primary: Fatal coronary heart disease and nonfatal MI	Primary: For patients with metabolic syndrome, there was no significant difference in rates of coronary heart disease and nonfatal MI with amlodipine vs chlorthalidone (RR, 0.96; 95% CI, 0.79 to 1.16), or lisinopril vs chlorthalidone (RR, 1.15; 95% CI, 0.88 to 1.27).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>10 mg QD</p> <p>vs</p> <p>lisinopril 10 to 40 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg QD</p>	<p>metabolic syndrome</p>		<p>Secondary:</p> <p>All cause mortality, fatal and nonfatal stroke, combined coronary heart disease, combined cardiovascular disease</p>	<p>Secondary:</p> <p>For patients with metabolic syndrome, there were no significant differences found between amlodipine vs chlorthalidone in all secondary endpoints (P value not significant).</p> <p>For patients without metabolic syndrome, amlodipine treatment was associated with significantly more heart failure, but in patients with metabolic syndrome, there was no difference (P=0.03).</p> <p>Patients with metabolic syndrome who received lisinopril experienced more heart failure and cardiovascular disease than those who received chlorthalidone (RR, 1.31; 95% CI, 1.14 to 1.64 and RR, 1.19; 95% CI, 1.07 to 1.32).</p>
<p>Rahman et al.<sup>49</sup> (2012) ALLHAT</p> <p>Lisinopril 10 to 40 mg/day</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg/day</p> <p>vs</p> <p>chlorthalidone 12.5 to 25 mg/day</p>	<p>Long-term, post-trial, follow-up</p> <p>Patients in ALLHAT stratified based on eGFR</p>	<p>N=31,350</p> <p>4 to 8 years</p>	<p>Primary:</p> <p>Cardiovascular mortality</p> <p>Secondary:</p> <p>Total mortality, CHD, cardiovascular disease, stroke, heart failure, ESRD</p>	<p>Primary:</p> <p>After an average of 8.8 years of follow-up, total mortality was significantly higher in patients with moderate/severe eGFR reduction (eGFR &lt;60 mL/min/1.73 m<sup>2</sup>) compared to patients with normal/increased (eGFR ≥90 mL/min/1.73 m<sup>2</sup>) and mildly reduced eGFR (eGFR 60 to 89 mL/min/1.73 m<sup>2</sup>) (P&lt;0.001).</p> <p>In patients with moderate/severe eGFR reduction, there was no significant difference in cardiovascular mortality between chlorthalidone and amlodipine (P=0.64), or chlorthalidone and lisinopril (P=0.56).</p> <p>Secondary:</p> <p>No significant differences were observed for any of the secondary endpoints among eGFR reduction groups.</p>
<p>Muntner et al.<sup>50</sup> (2014) ALLHAT</p> <p>Chlorthalidone 12.5 to 25 mg/day</p>	<p>Post-hoc analysis of ALLHAT</p> <p>Patients in ALLHAT with 5, 6, or 7 visits in 6 to 28 months of follow-up</p>	<p>N=24,004</p> <p>6 to 28 months</p>	<p>Primary:</p> <p>Visit-to-visit variability (VVV) of blood pressure</p> <p>Secondary:</p> <p>Not reported</p>	<p>Primary:</p> <p>Each measure of VVV of SBP was lower among participants randomized to chlorthalidone and amlodipine compared with those randomized to lisinopril. All four VVV of SBP metrics were lower among participants randomized to amlodipine vs chlorthalidone after full multivariable adjustment.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs amlodipine 2.5 to 10 mg/day  vs lisinopril 10 to 40 mg/day				After multivariable adjustment including mean SBP across visits and compared with participants randomized to chlorthalidone, participants randomized to amlodipine had a 0.36 (standard error [SE]: 0.07) lower standard deviation (SD) of SBP and participants randomized to lisinopril had a 0.77 (SE=0.08) higher SD of SBP. Results were consistent using other VVV of SBP metrics. These data suggest chlorthalidone and amlodipine are associated with lower VVV of SBP than lisinopril.  Secondary: Not reported
Bangalore et al. <sup>51</sup> (2017) ALLHAT  Lisinopril 10 to 40 mg/day  vs amlodipine 2.5 to 10 mg/day  vs chlorthalidone 12.5 to 25 mg/day	Post-hoc analysis of ALLHAT  Patients in ALLHAT with average blood pressure $\geq$ 140 mmHg systolic or $\geq$ 90 mm Hg diastolic on $\geq$ 3 antihypertensive medications, or blood pressure $<$ 140/90 mmHg on $\geq$ 4 antihypertensive medications (i.e., identified as having apparent treatment-resistant hypertension) at 2-year follow up	N=14,684  4.9 years (mean)	Primary: Combined fatal CHD or nonfatal MI  Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization, heart failure, and PAD)	Primary: Of participants assigned to chlorthalidone, amlodipine, or lisinopril, 9.6%, 11.4%, and 19.7%, respectively, had treatment-resistant hypertension. During mean follow-up of 2.9 years, primary outcome incidence was similar for those assigned to chlorthalidone compared with amlodipine or lisinopril (amlodipine- vs chlorthalidone-adjusted HR, 0.86; 95% CI, 0.53 to 1.39; P=0.53; lisinopril- vs chlorthalidone-adjusted HR, 1.06; 95% CI, 0.70 to 1.60; P=0.78).  Secondary: Secondary outcome risks were similar for most comparisons except coronary revascularization, which was higher with amlodipine than with chlorthalidone (HR, 1.86; 95% CI, 1.11 to 3.11; P=0.02). An as-treated analysis based on diuretic use produced similar results.
Fox et al. <sup>52</sup> (2003) EUROPA  Perindopril 8 mg QD	DB, MC, PC, RCT  Patients $\geq$ 18 years of age with evidence of CHD (e.g., MI $>$ 3 months before screening,	N=12,218  4.2 years (mean)	Primary: Composite of cardiovascular death, MI, or cardiac arrest  Secondary:	Primary: Patients treated with perindopril had a significant reduction in the primary outcome compared to patients treated with placebo (8 vs 10%; RR reduction, 20%; 95% CI, 9 to 29; P=0.0003). The benefit began to appear at one year and gradually increased throughout the trial.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo	percutaneous or surgical coronary revascularization >6 months before screening, 70% narrowing of 1 or more major coronary arteries, history of chest pain) and without clinical heart failure or uncontrolled HTN		Composite of total mortality, nonfatal MI, hospital admission for unstable angina, and cardiac arrest with successful resuscitation; cardiovascular mortality and nonfatal MI; individual components of the secondary outcomes and revascularization, stroke, and admission for heart failure	<p>Compared to placebo, treatment with perindopril was associated with reductions in all secondary end points. However, not all changes were significant.</p> <p>There was a 14% reduction in total mortality, nonfatal MI, unstable angina, and cardiac arrest (P=0.0009).</p> <p>There was a 22% reduction in nonfatal MI with perindopril (P=0.001).</p> <p>Total mortality was 11% lower with perindopril but this finding was not significant (P=0.1).</p> <p>Hospital admission for heart failure was significantly reduced with perindopril by 39% (P=0.002).</p>
PREAMI Investigators <sup>53</sup> (2006)  Perindopril 8 mg/day  vs placebo	DB, MC, PC, PG, RCT  Patients ≥65 years with LVEF ≥40% and recent acute MI	N=1,252  12 months	<p>Primary: Composite of death, hospitalization for heart failure or left ventricular remodeling</p> <p>Secondary: Cardiovascular death, hospitalization for reinfarction or angina, revascularization</p>	<p>Primary: The primary end point occurred in 35% of patients taking perindopril and 57% of patients on placebo, with an absolute risk reduction of 0.22 (95% CI, 0.16 to 0.28; P&lt;0.001).</p> <p>A total of 126 patients (28%) and 226 patients (51%) in the perindopril and placebo groups, respectively, experienced remodeling (P&lt;0.001). The mean increase in left ventricular end-diastolic volume was 0.7 mL with perindopril compared to 4.0 mL with placebo (P&lt;0.001).</p> <p>Secondary: Cardiovascular death, hospitalization for subsequent acute MI or angina or revascularization was infrequent and not modified by treatment.</p> <p>Conclusion: Perindopril treatment for one year reduced progressive left ventricular remodeling but was not associated with better clinical outcomes.</p>
ADVANCE	DB, MC, PC, RCT	N=11,140	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Collaborative Group<sup>54</sup> (2007)</p> <p>Perindopril (2 to 4 mg) and indapamide (0.625 to 1.25 mg) QD</p> <p>vs</p> <p>placebo</p>	<p>Adults 55 years of age or older who were diagnosed with type 2 diabetes at age 30 or older, and a history of cardiovascular disease or <math>\geq 1</math> other risk factor for cardiovascular disease</p>	<p>Mean 4.3 years</p>	<p>Composites of major macrovascular and microvascular events (death from cardiovascular disease, nonfatal stroke, nonfatal MI, or new renal or diabetic eye disease)</p> <p>Secondary: Macrovascular and microvascular endpoints analyzed separately</p>	<p>The relative risk of a major macrovascular or microvascular event was reduced by 9% (861 [15.5%] active vs 938 [16.8%] placebo; HR, 0.91, 95% CI 0.83 to 1.0, P=0.04).</p> <p>Secondary: The RR of death from cardiovascular disease was reduced by 18% (211 [3.8%] active vs 257 [4.6%] placebo; 0.82, 0.68-0.98, p=0.03) and death from any cause was reduced by 14% (408 [7.3%] active vs 471 [8.5%] placebo; 0.86, 0.75-0.98, P=0.03).</p>
<p>HOPE Investigators<sup>55</sup> (2000)</p> <p>Ramipril 10 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, RCT, two-by-two factorial trial</p> <p>Men and women <math>\geq 55</math> years old with history of CAD, stroke, PVD, or diabetes and <math>\geq 1</math> other cardiovascular risk factor and who were not known to have a low ejection fraction (&lt;40%) or heart failure</p>	<p>N=9,297</p> <p>5 years (mean)</p>	<p>Primary: Composite of death from cardiovascular causes, MI, or stroke and each outcome separately</p> <p>Secondary: Death from any cause, revascularization, hospitalization for unstable angina or heart failure, and complications related to diabetes</p> <p>Other end points: Worsening angina,</p>	<p>Primary: Fewer patients on ramipril than placebo (14.0 vs 17.8%, respectively) died of cardiovascular causes or had a MI or stroke (RR, 0.78; 95% CI, 0.70 to 0.86; P&lt;0.001).</p> <p>Treatment with ramipril reduced the rates of death from cardiovascular causes (RR, 0.74; P&lt;0.001), MI (RR, 0.80; P&lt;0.001), and stroke (RR, 0.68; P&lt;0.001).</p> <p>Secondary: The risk of death from any cause was also significantly reduced by treatment with ramipril (RR, 0.84; P=0.005).</p> <p>Significantly fewer patients treated with ramipril underwent revascularization compared to placebo (RR, 0.85; P=0.002).</p> <p>Fewer hospitalizations for heart failure were reported with ramipril vs placebo but the risk reduction was not statistically significant (RR, 0.88; P=0.25).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			cardiac arrest, heart failure, unstable angina with ECG changes, and the development of diabetes	Fewer complications related to diabetes were reported in patients receiving ramipril (RR, 0.84; P=0.03).  Other end points: Significantly fewer patients treated with ramipril than placebo group had the following: worsening angina (RR, 0.89; P=0.004), cardiac arrest (RR, 0.62; P=0.02), heart failure (RR 0.77; P<0.001), and new diagnosis of diabetes (RR, 0.66; P<0.001). There was no difference between treatment groups for unstable angina with ECG changes (RR, 0.97; P=0.76).
<p>ONTARGET Investigators<sup>56</sup> (2008)</p> <p>Ramipril 10 mg/day</p> <p>vs</p> <p>telmisartan 80 mg/day</p> <p>vs</p> <p>ramipril 10 mg/day and telmisartan 80 mg/day</p>	<p>DB, MC, PC, RCT</p> <p>Patients with coronary, peripheral, or cerebrovascular disease or diabetes with end-organ damage</p>	<p>N=25,620</p> <p>56 months (median follow-up)</p>	<p>Primary: Death from cardiovascular causes, MI, stroke or hospitalization for heart failure</p> <p>Secondary: Composite of death from cardiovascular causes, MI or stroke; heart failure, worsening or new angina, new diagnosis diabetes mellitus, new atrial fibrillation, renal impairment, revascularization procedures</p>	<p>Primary: The primary outcome occurred in 16.5, 16.7, and 16.3% of patients receiving ramipril, telmisartan and combination therapy, respectively.</p> <p>Secondary: The composite of death from cardiovascular causes, MI or stroke occurred in 14.1% of patients in the ramipril group and 13.9% of patients in the telmisartan group (RR, 0.99; 95% CI, 0.91 to 1.07; P=0.001 for non-inferiority). Combination therapy was not significantly better than ramipril alone (RR, 0.99; 95% CI, 0.92 to 1.07).</p> <p>There were no significant differences in the rates of secondary outcomes, except for renal dysfunction, which occurred in 10.2% of patients receiving ramipril, 10.6% of patients receiving telmisartan and 13.5% of patients receiving combination therapy (P&lt;0.001 vs ramipril; P value not reported vs telmisartan).</p> <p>As compared to the ramipril group, the telmisartan group had lower rates of cough (1.1 vs 4.2%; P&lt;0.001) and angioedema (0.1 vs 0.3%; P=0.01) and a higher rate of hypotensive symptoms (2.6 vs 1.7%; P&lt;0.001); the rate of syncope was the same in the two groups (0.2%).</p> <p>As compared to the ramipril group, combination therapy had an increased risk of hypotensive symptoms (4.8 vs 1.7%; P&lt;0.001), syncope (0.3 vs 0.2%; P=0.03) and renal dysfunction (13.5 vs 10.2%; P&lt;0.001).</p>
<p>Redon et al.<sup>57</sup> (2012)</p> <p>ONTARGET</p>	<p>Post-hoc analysis</p> <p>Patients with</p>	<p>N=25,584</p> <p>56 months</p>	<p>Primary: Composite of cardiovascular</p>	<p>Primary: The primary outcome occurred in 20.2% (n=1,938) and 14.2% (n=2,276) of diabetic and nondiabetic patients. Compared to nondiabetic patients,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Ramipril 10 mg/day</p> <p>vs</p> <p>telmisartan 80 mg/day</p> <p>vs</p> <p>ramipril 10 mg/day and telmisartan 80 mg/day</p>	<p>coronary, peripheral, or cerebrovascular disease or diabetes with end-organ damage</p>	<p>(median follow-up)</p>	<p>death, nonfatal MI, nonfatal stroke, and hospitalized heart failure</p> <p>Secondary: Not reported</p>	<p>diabetic patients had a significantly higher risk for the primary endpoint (HR, 1.48; 95% CI, 1.38 to 1.57) and cardiovascular death (HR, 1.56; 95% CI, 1.42 to 1.71), MI (HR, 1.30; 95% CI, 1.17 to 1.46), stroke (HR, 1.39; 95% CI, 1.23 to 1.56), and CHF hospitalization (HR, 2.06; 95% CI, 1.82 to 2.32).</p> <p>Cardiovascular risk was significantly higher in diabetic patients compared to nondiabetic patients regardless of changes in SBP during treatment. In all patients, progressively greater SBP reductions were accompanied by reduced risk for the primary outcome only if baseline SBP levels ranged from 143 to 155 mm Hg; except for stroke, there was no benefit in fatal and nonfatal cardiovascular outcomes by reducing SBP &lt;130 mm Hg.</p> <p>Secondary: Not reported</p>
<p>Mann et al.<sup>58</sup> (2013) ONTARGET</p> <p>Ramipril with telmisartan</p>	<p>Subanalysis</p> <p>Patients in the ONTARGET trial with diabetes mellitus</p>	<p>N=3163 with CKD N=6465 no CKD</p> <p>56 months</p>	<p>Primary: Composite of death from cardiovascular cause, nonfatal MI, nonfatal stroke or hospitalization for CHF</p> <p>Secondary: composite renal outcome for this analysis was defined posthoc as chronic dialysis (&gt;2 months) or a doubling of baseline serum creatinine</p>	<p>Primary: The stroke rate in all participants with diabetes was not different between the treatment groups, 1.19 and 1.22 per 100 patient-years in those on dual and monotherapy, respectively (HR, 0.99; 95% CI, 0.82 to 1.20). The results were consistent in those with or without renal disease (P value for interaction =0.60; 1.59 vs 1.55 and 1.01 vs 1.08 strokes per 100 patient-years, respectively). Results for other major outcomes indicated no differences and no interaction of renal subgroups with treatment effects.</p> <p>Secondary: Dialysis-dependent acute kidney injury tended to occur more frequently in those allocated to dual than with monotherapy, 0.14 vs 0.08 cases per 100 patient-years, (HR, 1.55; 95% CI, 0.84 to 2.85), and hyperkalemia was more frequent, 1.82 vs 1.07 cases per 100 patient-years (HR, 1.71; 95% CI, 1.44 to 2.02). Both adverse outcomes were more frequent in those with renal disease; however, the excess due to dual therapy was similar in those with and without renal disease.</p>
<p>PEACE Trial Investigators<sup>59</sup></p>	<p>DB, PC, RCT</p>	<p>N=8,290</p>	<p>Primary: Combined rate of</p>	<p>Primary: No significant differences in the primary outcome measures between</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(2004) PEACE  Trandolapril 4 mg/day  vs  placebo	Patients $\geq 50$ years of age with stable CAD and normal or slightly reduced left ventricular function (LVEF $>40\%$ )	4.8 years (median)	nonfatal MI, death from cardiovascular causes, or coronary revascularization procedures  Secondary: Composite of death from cardiovascular causes, nonfatal MI, revascularization, unstable angina, new CHF, stroke, PVD, and cardiac arrhythmia	trandolapril and placebo were reported (21.9 vs 22.5%; HR, 0.96; 95% CI, 0.88 to 1.6; P=0.43).  Secondary: No significant differences in secondary outcome measures between trandolapril and placebo were reported (P $>0.05$ ).  Side effects leading to discontinuation of study medication occurred in 14.4% of patients receiving trandolapril and 6.5% of patients receiving placebo (P $<0.001$ ). The rates of cough (39.1 vs 27.5%; P $<0.01$ ) and syncope (4.8 vs 3.9%; P=0.04) were higher in patients receiving trandolapril vs placebo.  Note: This trial was conducted in low-risk patients with stable CAD and normal or slightly reduced left ventricular function. However, the HOPE trial was conducted in patients with coronary or other vascular disease or with diabetes and another cardiovascular risk factor and the EUROPA trial was conducted in patients with evidence of CHD.
Pilote et al. <sup>60</sup> (2004)  Captopril (50 mg), enalapril (10 mg), fosinopril (10 mg), lisinopril (10 mg), perindopril (4 mg), quinapril (20 mg), and ramipril (5 mg)	RETRO  Patients $\geq 65$ years who were hospitalized for acute myocardial infarction and filled a prescription for an ACE inhibitor within 30 days of discharge and who continued to receive the same drug for $\geq 1$ year	N=7,512  Average of 2.3 years since discharge	Primary: 1-year mortality following an acute MI  Secondary: Readmissions due to cardiac complications	Primary: Captopril (HR, 1.56; 95% CI, 1.13 to 2.15), enalapril (HR, 1.47; 95% CI, 1.14 to 1.89), fosinopril (HR, 1.71; 95% CI, 1.29 to 2.25), lisinopril (HR, 1.28; 95% CI, 0.98 to 1.67), and quinapril (HR, 1.58; 95% CI, 1.10 to 2.82) were associated with higher mortality than was ramipril.  No statistically significant difference was reported between perindopril and ramipril (HR, 0.98; 95% CI, 0.60 to 1.60).  Secondary: Enalapril (HR, 1.44; 95% CI, 1.13 to 2.01) and fosinopril (HR, 1.83; 95% CI, 1.27 to 2.62) were associated with higher readmission rates for CHF than ramipril. Readmissions for unstable angina and recurrent MI were similar across all prescription groups.
Dalhof et al. <sup>61</sup> (2005) ASCOT-BPLA  Amlodipine 5 to	MC, OL, RCT  Patients 40 to 79 years of age with HTN and $\geq 3$ other	N=19,257  5.5 years	Primary: Nonfatal MI (including silent MI) and fatal CHD	Primary: No statistically significant difference in nonfatal MI and fatal CHD was reported between the amlodipine plus perindopril group compared to the atenolol plus bendroflumethiazide groups (HR, 0.90; 95% CI, 0.79 to 1.2; P=0.1052).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>10 mg/day adding perindopril 4 to 8 mg/day as needed</p> <p>vs</p> <p>atenolol 50 to 100 mg/day adding bendroflumethiazide* 1.25 to 2.5 mg/day and potassium as needed</p> <p>If blood pressure was still not achieved, doxazosin 4 to 8 mg/day was added to the regimen.</p>	<p>cardiovascular risk factors (left ventricular hypertrophy, other specified abnormalities on ECG, type 2 diabetes, PAD, history of stroke or TIA, male, age <math>\geq 55</math> years, microalbuminuria or proteinuria, smoking, TC:HDL-C ratio <math>\geq 6</math>, or family history of CHD)</p>	<p>N=1,411</p> <p>1.3 years</p>	<p>Secondary: All-cause mortality, total stroke, primary end points minus silent MI, all coronary events, total cardiovascular events and procedures, cardiovascular mortality, nonfatal and fatal heart failure, effects on primary end point and on total cardiovascular events and procedures among prespecified subgroups</p> <p>Tertiary: Silent MI, unstable angina, chronic stable angina, PAD, life-threatening arrhythmias, development of diabetes, development of renal impairment</p>	<p>Secondary: Significantly greater reductions in the following secondary end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: all- cause mortality (P=0.0247), total stroke (P=0.0003), primary end points minus silent MI (P=0.0458), all coronary events (P=0.0070), total cardiovascular events and procedures (P&lt;0.0001), and cardiovascular mortality (P=0.0010).</p> <p>There were no significant differences in nonfatal and fatal heart failure between the two treatment groups (P=0.1257).</p> <p>The study was terminated early due to higher mortality and worse outcomes on several secondary end points observed in the atenolol study group.</p> <p>Tertiary: Significantly greater reductions in the following end points were observed with amlodipine plus perindopril compared to atenolol plus bendroflumethiazide: unstable angina (P=0.0115), PAD (P=0.0001), development of diabetes (P&lt;0.0001), and development of renal impairment (P=0.0187).</p> <p>There were no significant differences in the incidence of silent MI (P=0.3089), chronic stable angina (P=0.8323) or life-threatening arrhythmias (P=0.8009) between the two treatment groups.</p> <p>There was no significant difference in the percent of patients who stopped therapy because of an adverse event between the two treatment groups (overall 25%). There was, however, a significant difference in favor of amlodipine plus perindopril in the proportion of patients who stopped trial therapy because of a serious adverse events (2 vs 3%; P&lt;0.0001).</p>
<p>Chapman et al.<sup>62</sup> (2007) ASCOT-BPLA</p>	<p>Subanalysis of ASCOT-BPLA evaluating effects of spironolactone on</p>	<p>N=1,411</p> <p>1.3 years</p>	<p>Primary: Change in DBP and SBP, adverse effects</p>	<p>Primary: Spironolactone-treated patients lead to a significant 21.9 mm Hg reduction in SBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 20.8 to 23.0 mm</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Atenolol 50 to 100 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); bendroflumethiazide* plus potassium 1.25 to 2.5 mg plus doxazosin were added for additional blood pressure control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p> <p>vs</p> <p>amlodipine 5 to 10 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); perindopril 4 to 8 mg and doxazosin were added for additional control;</p>	<p>treatment-resistant HTN</p> <p>Patients 40 to 79 years of age with HTN and <math>\geq 3</math> cardiovascular risk factors, with SBP <math>\geq 160</math> mm Hg and/or DBP <math>\geq 100</math> mm Hg (not on antihypertensive therapy) or SBP <math>\geq 140</math> mm Hg and/or DBP <math>\geq 90</math> mm Hg (on antihypertensive therapy)</p>		<p>Secondary: Not reported</p>	<p>Hg; <math>P &lt; 0.001</math>).</p> <p>Spironolactone-treated patients lead to a significant 9.5 mm Hg reduction in DBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 9.0 to 10.1; <math>P &lt; 0.001</math>).</p> <p>Spironolactone-treated patients exhibited small but significant decreases in sodium, LDL-C and TC as well as increases in potassium, glucose, creatinine and HDL-C (<math>P &lt; 0.05</math>).</p> <p>The most common adverse effect reported in the trial was gynecomastia in men (<math>P</math> value not reported).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p>				
<p>Pepine et al.<sup>63</sup> (2003) INVEST</p> <p>Verapamil SR 240 mg/day (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p> <p>vs</p> <p>atenolol 50 mg/day (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p>	<p>MC, OL, RCT</p> <p>Patients with essential HTN</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: First occurrence of death (all cause), nonfatal MI or stroke</p> <p>Secondary: Cardiovascular death, angina, cardiovascular hospitalization, angina, blood pressure control (SBP/DBP &lt;140/90 mm Hg or &lt;130/85 mm Hg if diabetic or renal impairment), safety</p>	<p>Primary: At 24 months, in the calcium antagonist strategy subgroup, 81.5% of patients were taking verapamil SR, 62.9% trandolapril, and 43.7% HCTZ. In the non-calcium antagonist strategy, 77.5% of patients were taking atenolol, 60.3% HCTZ, and 52.4% trandolapril.</p> <p>After a follow-up of 61,835 patient-years (mean, 2.7 years per patient), 2,269 patients had a primary outcome event with no statistically significant difference between treatment strategies (9.93% in calcium antagonist strategy vs 10.17% in non-calcium antagonist strategy; RR, 0.98; 95% CI, 0.90 to 16; P=0.57).</p> <p>Secondary: There was no significant difference in the rate of cardiovascular death (P=0.94) or cardiovascular hospitalization (P=0.59) between the two treatment groups.</p> <p>At 24 months, angina episodes decreased in both groups, but the mean frequency was lower in the calcium antagonist strategy group (0.77 episodes/week) compared to the non-calcium antagonist strategy group (0.88 episodes/week; P=0.02).</p> <p>Two-year blood pressure control was similar between groups. The blood pressure goals were achieved by 65.0% (systolic) and 88.5% (diastolic) of calcium antagonist strategy patients and 64.0% (systolic) and 88.1% (diastolic) of non-calcium antagonist strategy patients. A total of 71.7% of calcium antagonist strategy patients and 70.7% of non-calcium antagonist strategy patients achieved an SBP &lt;140 mm Hg and DBP &lt;90 mm Hg.</p> <p>Both regimens were generally well tolerated. Patients in the calcium</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Trandolapril was recommended for all patients with heart failure, diabetes, or renal insufficiency.</p>				<p>antagonist strategy group reported constipation and cough more frequently than patients in the non-calcium antagonist strategy group, while non-calcium antagonist strategy patients experienced more dyspnea, lightheadedness, symptomatic bradycardia and wheezing (all were statistically significant with <math>P \leq 0.05</math>).</p>
<p>Lindholm et al.<sup>64</sup> (2005)</p> <p>Other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)</p> <p>or</p> <p>placebo</p> <p>vs</p> <p><math>\beta</math>-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol, or</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a <math>\beta</math>-blocker as first-line treatment (in <math>\geq 50\%</math> of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary: The RR of stroke was 16% higher with <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; <math>P=0.009</math>). The RR of stroke was the highest with atenolol (26% higher) compared to other non <math>\beta</math>-blockers (RR, 1.26%; 95% CI, 1.15 to 1.38; <math>P&lt;0.0001</math>).</p> <p>The relative risk of MI was 2% higher for <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (<math>P</math> value not reported).</p> <p>The RR of all-cause mortality was 3% higher for <math>\beta</math>-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; <math>P=0.14</math>).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>propranolol)</p> <p>Wiysonge et al.<sup>65</sup> (2007)</p> <p>Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)</p> <p>vs</p> <p>β-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)</p>	<p>MA</p> <p>13 RCTs evaluating patients ≥18 years of age with HTN</p>	<p>N=91,561</p> <p>Duration varied</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions</p>	<p>Primary: There was not a significant difference observed in all-cause mortality between β-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with β-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).</p> <p>Secondary: There was a significant decrease in stroke observed with β-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between β-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07)), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with β-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of β-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with β-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Blood Pressure Lowering Treatment Trialists' Collaboration<sup>66</sup> (2007)</p> <p>ACE inhibitors (17 trials)</p> <p>vs</p> <p>ARBs (9 trials)</p>	<p>MA</p> <p>Patients with high blood pressure, diabetes, history or CHD or cerebrovascular disease</p>	<p>N=146,838 (26 trials)</p> <p>Variable duration</p>	<p>Primary: Nonfatal myocardial infarction or death from CHD, including sudden death; heart failure causing death or requiring hospitalization; nonfatal stroke or death from cerebrovascular disease</p> <p>Secondary: Not reported</p>	<p>reported.</p> <p>Primary: From a total of 146,838 individuals with high blood pressure or an elevated risk of cardiovascular disease, major cardiovascular events were documented in 22,666 patients during follow-up. The analyses showed comparable blood pressure-dependent reductions in risk with ACE inhibitors and ARBs (<math>P \geq 0.3</math> for all three outcomes).</p> <p>ACE inhibitors produced a blood pressure-independent reduction in the relative risk of CHD of approximately 9% (95% CI, 3 to 14%). No similar effect was detected for ARBs, and there was some evidence of a difference between ACE inhibitors and ARBs in this regard (<math>P=0.002</math>).</p> <p>For both stroke and heart failure, there was no evidence of any blood pressure-independent effects of either ACE inhibitors or ARBs.</p> <p>Secondary: Not reported</p>
<b>Cerebrovascular Disease</b>				
<p>PROGRESS<sup>67</sup> (2001)</p> <p>Perindopril 4 mg/day</p> <p>vs</p> <p>perindopril 4 mg/day and indapamide 2 to 2.5 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients with a history of prior stroke or TIA within the previous 5 years</p>	<p>N=6,105</p> <p>4 years</p>	<p>Primary: Fatal or nonfatal stroke</p> <p>Secondary: Fatal or disabling stroke, total major vascular events comprising the composite of nonfatal stroke, nonfatal MI, or death due to any vascular cause (including unexplained sudden death); total and cause</p>	<p>Primary: Patients receiving active treatment experienced a 28% reduction in nonfatal or fatal stroke (95% CI, 17 to 38; <math>P &lt; 0.0001</math>).</p> <p>There were similar reductions in the risk of stroke in hypertensive and non-hypertensive subgroups (32 vs 27%; <math>P &lt; 0.01</math>)</p> <p>A trend towards a greater effect of active treatment among patients treated with combination therapy (43% risk reduction) than in those treated with single drug therapy (5% risk reduction) was reported.</p> <p>Secondary: There was a 33% reduction in fatal or disabling strokes in the active treatment group.</p> <p>Active treatment reduced the risk of total major vascular events by 26% (<math>P=0.02</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			specific deaths; hospital admissions	<p>There were no significant differences between active treatment and placebo in total deaths from vascular or nonvascular causes.</p> <p>Among those assigned active treatment, there was a 9% RR reduction in hospitalization, with a median reduction of 2.5 days in the time spent in the hospital during follow-up.</p> <p>Combination therapy with perindopril plus indapamide reduced blood pressure by 12/5 mm Hg and stroke risk by 43%. Single drug therapy reduced blood pressure by 5/3 mm Hg and produced no discernible reduction in the risk of stroke.</p>
<p>Arima et al.<sup>68</sup> (2011) PROGRESS</p> <p>Perindopril 4 mg/day</p> <p>vs</p> <p>perindopril 4 mg/day and indapamide 2 to 2.5 mg/day</p> <p>vs</p> <p>placebo</p>	<p>Post-hoc analysis</p> <p>Patients with a history of prior stroke or TIA within the previous 5 years</p>	<p>N=4,283</p> <p>4 years</p>	<p>Primary: Total major vascular events (nonfatal stroke, nonfatal MI, or vascular death)</p> <p>Secondary: Not reported</p>	<p>Primary: Among all patients, active treatment reduced the RR of major vascular events by 27% (95% CI, 10 to 41) in patients with isolated systolic HTN, by 28% (95% CI, -29 to 60) in patients with isolated diastolic HTN, and by 32% (95% CI, 17 to 45%) in patients with systolic-diastolic HTN. There was no evidence of differences in the magnitude of the effects of treatment among different types of HTN.</p> <p>Blood pressure reductions and RRs were consistently greater with combination therapy compared to single drug therapy (mean SBP difference, 12.3 vs 3.9 mm Hg, 7.7 vs 4.3 mm Hg, and 13.5 vs 5.2 mm Hg; RR reduction of major vascular events 34 vs 16%, 63 vs -78%, and 45 vs 10% for isolated systolic HTN, isolated diastolic HTN, and systolic-diastolic HTN).</p> <p>Secondary: Not reported</p>
<b>Heart Failure</b>				
<p>Pfeffer et al.<sup>69</sup> (1992) SAVE</p> <p>Captopril up to 50 mg TID</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Patients 21 to 80 years of age who had an acute MI within 3 to 16 days and left ventricular dysfunction with a</p>	<p>N=2,231</p> <p>42 months (average)</p>	<p>Primary: Mortality from all causes, mortality from cardiovascular causes, mortality combined with a decrease in</p>	<p>Primary: Mortality from all causes was significantly reduced in the captopril group (20%) vs placebo group (25%) for a 19% reduction in the risk of mortality from all causes (95% CI, 3 to 25; P=0.019).</p> <p>The incidence of fatal cardiovascular events was consistently reduced in the captopril group with a 21% reduced risk of mortality from cardiovascular causes (P=0.014).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo	LVEF $\leq$ 40%, but without overt heart failure or symptoms of myocardial ischemia		ejection fraction $\geq$ 9 units, cardiovascular morbidity, combination of cardiovascular mortality and morbidity  Secondary: Not reported	The incidence of nonfatal major cardiovascular events was consistently reduced in the captopril group with a 25% reduced risk of recurrent MI (P=0.015), 37% reduced risk for the development of severe heart failure (P<0.001), and 22% reduced risk of CHF requiring hospitalization (P=0.019).  Long-term captopril administration was associated with an improvement in survival and reduced morbidity and mortality due to major cardiovascular events.  Secondary: Not reported
Pitt et al. <sup>70</sup> (1997) ELITE  Captopril 50 mg TID  vs  losartan 50 mg QD	DB, MC, PG, RCT  Patients $\geq$ 65 years with symptomatic heart failure (NYHA class II to IV and LVEF $\leq$ 40%), and no history of prior ACE inhibitor therapy	N=722  1 year	Primary: Change in renal function  Secondary: Composite of death and/or hospital admission for heart failure, all-cause mortality, admission for heart failure, NYHA class, admission for MI or unstable angina	Primary: No difference between losartan and captopril was reported in the rate of persistent rise in serum creatinine concentrations (10.5% for both groups).  Secondary: Death and/or hospital admission for heart failure was recorded in 9.4% of patients receiving losartan and 13.2% for patients receiving captopril (risk reduction, 32%; 95% CI, -4 to 55; P=0.075). This risk reduction was primarily due to a decrease in all-cause mortality (4.8 vs 8.7%; risk reduction, 46%; 95% CI, 5 to 69; P=0.035).  Admissions with heart failure were the same in both groups (5.7%), as was improvement in NYHA functional class from baseline. Admission to hospital for any reason was less frequent with losartan than with captopril treatment (22.2 vs 29.7%; P=0.014).  More patients discontinued therapy due to adverse events with captopril (20.8%) than losartan (12.2%; P=0.002).
Pitt et al. <sup>71</sup> (2000) ELITE II  Captopril 50 mg TID	DB, MC, PG, RCT  Patients $\geq$ 60 years old with symptomatic heart failure (NYHA II to	N=3,152  555 days (mean follow-up)	Primary: All-cause mortality  Secondary: Composite of sudden cardiac	Primary: No significant difference in all-cause mortality was reported between losartan (17.7%) and captopril (15.9%; HR, 1.13; 95% CI, 0.95 to 1.35; P=0.16).  Secondary:



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs losartan 50 mg QD	IV and LVEF ≤40%), and no history of prior ACE inhibitor therapy		death or resuscitated cardiac arrest	Sudden death or resuscitated cardiac arrest was observed in 9.0% of patients receiving losartan and 7.3% of patients receiving captopril (HR, 1.25; 95% CI; 0.98 to 1.60; P=0.08).  Significantly fewer patients in the losartan group (excluding those who died) discontinued study treatment because of adverse events (9.7 vs 14.7%; P<0.001), including cough (0.3 vs 2.7%).  Note: ELITE II trial was a larger follow-up trial to the ELITE I trial to confirm the secondary end point from the ELITE I trial, which reported a greater reduction in all-cause mortality with losartan compared to captopril.
Dickstein et al. <sup>72</sup> (2002) OPTIMAAL  Captopril 50 mg TID  vs losartan 50 mg QD	DB, MC, PG, RCT  Patients ≥50 years with an acute MI and signs or symptoms of heart failure during the acute phase or a new Q-wave anterior infarction or reinfarction	N=5,477  2.7 years (mean)	Primary: All-cause mortality  Secondary: Composite of sudden cardiac death or resuscitated cardiac arrest	Primary: No significant difference in all-cause mortality was reported between patients receiving losartan and captopril (18 vs 16%, respectively; RR, 1.13; 95% CI, 0.99 to 1.28; P=0.07).  Secondary: No significant difference in sudden cardiac death or resuscitated cardiac arrest was reported between patients receiving losartan and captopril (9% vs 7%; RR, 1.19; 95% CI, 0.98 to 1.43; P=0.07).  Losartan was significantly better tolerated than captopril, with fewer patients discontinuing study medication (17 vs 23%; P<0.0001).
Pfeffer et al. <sup>73</sup> (2003) VALIANT  Captopril 50 mg TID  vs valsartan 160 mg BID  vs	DB, MC, RCT  Patients ≥18 years of age with an acute MI that was complicated by clinical or radiologic signs of heart failure and/or evidence of left ventricular systolic dysfunction	N=14,703  24.7 months	Primary: All-cause mortality  Secondary: Death from cardiovascular causes, recurrent MI, hospitalization for heart failure	Primary: No significant difference in all-cause mortality was reported between valsartan monotherapy and captopril monotherapy (P=0.98).  No significant difference in all-cause mortality was observed between valsartan plus captopril combination therapy and captopril monotherapy (P=0.73).  Secondary: The rate of death from cardiovascular causes, reinfarction, or hospitalization for heart failure was not significantly different between valsartan and captopril monotherapy (P=0.20).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
valsartan 80 mg BID and captopril 50 mg TID				<p>The rate of death from cardiovascular causes, reinfarction, or hospitalization for heart failure was not significantly different between valsartan and captopril combination therapy and captopril monotherapy (P=0.37).</p> <p>Combination therapy had the most drug-related adverse events. With monotherapy, hypotension and renal dysfunction were more common in the valsartan group and cough, rash, and taste disturbance were more common in the captopril group.</p>
<p>CONSENSUS Trial Study Group<sup>74</sup> (1987) CONSENSUS Enalapril 2.5 to 40 mg/day vs placebo</p>	<p>DB, MC, PC, PG, RCT  Patients with severe CHF (NYHA class IV symptoms), patients with recent MI and unstable angina were excluded</p>	<p>N=253  188 days (average)</p>	<p>Primary: 6-month mortality and the cause of death  Secondary: 12-month mortality and overall mortality</p>	<p>Primary: Mortality at six months was 26 and 44% for patients in the enalapril and placebo groups, respectively, for an overall reduction of 40% for enalapril (P=0.002).</p> <p>Secondary: At 12 months, enalapril reduced mortality by 31% compared to placebo (P=0.001).</p> <p>By the end of the study, there had been 50 deaths in the enalapril group and 68 deaths in the placebo group for a reduction of 27% (P=0.003). The entire reduction in total mortality was found to be among patients with progressive heart failure (a reduction of 50%), whereas no difference was seen in the incidence of sudden cardiac death.</p> <p>Note: The study was stopped early due to clear benefit with enalapril.</p>
<p>SOLVD Investigators<sup>75</sup> (1991) SOLVD Enalapril 2.5 to 20 mg/day vs placebo</p>	<p>DB, MC, PC, RCT  Patients with CHF and LVEF ≤35% receiving conventional therapy</p>	<p>N=2,569  41.4 months (average)</p>	<p>Primary: Mortality, rate of hospitalization for heart failure  Secondary: Not reported</p>	<p>Primary: Death was reported in 35.2 and 39.7% of patients receiving enalapril and placebo, respectively (risk reduction, 16%; 95% CI, 5 to 26; P=0.0036).</p> <p>Although reductions in mortality were observed in several categories of cardiac deaths, the largest reduction occurred among the deaths attributed to progressive heart failure (risk reduction, 22%; 95% CI, 6 to 35). There was little apparent effect of treatment on deaths classified as due to arrhythmia without pump failure.</p> <p>Fewer patients died or were hospitalized for worsening heart failure (risk reduction, 26%; 95% CI, 18 to 34; P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
SOLVD Investigators <sup>76</sup> (1992) SOLVD  Enalapril 2.5 mg to 20 mg/day  vs  placebo	DB, MC, PC, RCT  Patients 21 to 80 years of age with heart disease and an ejection fraction of ≤35% who were not receiving diuretics, digoxin or vasodilators for the treatment of heart failure	N=4,228  37.4 months (average)	Primary: All-cause mortality, incidence of heart failure, rate of hospitalization for heart failure  Secondary: Not reported	Primary: Enalapril resulted in an 8% reduction in risk for all-cause mortality (P=0.30). The difference was entirely due to a reduction in deaths due to cardiovascular causes, primarily progressive heart failure (risk reduction, 12%; P=0.12).  In the placebo group, 30.2% of patients developed heart failure compared to 20.7% for enalapril (risk reduction, 37%; P<0.001).  Rates of first hospitalization and multiple hospitalizations for CHF were higher with placebo (12.9 and 4.8%) than enalapril (8.7 and 2.7%; both P<0.001).  The total number of deaths and cases of heart failure were lower in the enalapril group than in the placebo group (risk reduction, 29%; P<0.001). In addition, fewer patients given enalapril died or were hospitalized for heart failure (risk reduction, 20%; P<0.001).  Secondary: Not reported
McMurray et al. <sup>77</sup> (2016) ATMOSPHERE  Enalapril 5 or 10 mg BID  vs  aliskiren 150 mg QD  vs	DB, DD, RCT  Patients with CHF (NYHA class II to IV) and EF ≤35% receiving stable doses of an ACE inhibitor (equivalent to at least 10 mg of enalapril daily) and of a β-blocker at the time of enrollment	N=7,016  Median of 36.6 months	Primary: Composite of death from cardiovascular causes or hospitalization for heart failure  Secondary: Change from baseline to 12 months in the Kansas City Cardiomyopathy	Primary: Overall, the primary outcome occurred in 770 patients (32.9%) in the combination-therapy group (11.7 events per 100 person-years), in 791 patients (33.8%) in the aliskiren group (12.1 events per 100 person-years), and in 808 patients (34.6%) in the enalapril group (12.4 events per 100 person-years). The HR in the combination-therapy group, as compared with the enalapril group, was 0.93 (95% CI, 0.85 to 1.03; P=0.17); the HR in the aliskiren group, as compared with the enalapril group, was 0.99 (95% CI, 0.90 to 1.10; P=0.91 for superiority). Although the noninferiority margin of 1.104 was met with the use of the 95% confidence interval, the one-sided P value of 0.0184 did not fulfill the prespecified requirement of a P value of 0.0123 or less. A sensitivity analysis that included only patients who received the assigned trial regimen gave consistent results, as did an analysis in which data that were collected after regulatory censoring

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
combination of aliskiren 150 mg QD and enalapril 5 or 10 mg BID			Questionnaire (KCCQ) clinical summary score	<p>were included.</p> <p>Secondary: There were no significant between-group differences in the secondary outcome. The exploratory composite renal outcome (the composite of death from renal causes, end-stage renal disease, or doubling of the serum creatinine level) occurred significantly more frequently in the combination-therapy group than in the enalapril group.</p>
<p>McKelvie et al.<sup>78</sup> (1999) RESOLVD</p> <p>Enalapril 10 mg BID</p> <p>vs</p> <p>candesartan 4 to 16 mg QD</p> <p>vs</p> <p>candesartan 4 to 8 mg QD and enalapril 10 mg BID</p>	<p>DB, PG, MC, RCT</p> <p>Patients with CHF (NYHA classes II to IV), a 6 minute walk distance of 500 meters or less, and an ejection fraction &lt;40%</p>	<p>N=768</p> <p>43 weeks</p>	<p>Primary: Change in 6-minute walk distance</p> <p>Secondary: Change in NYHA functional class, QOL, ejection fraction, ventricular volumes, neurohormone levels, safety</p>	<p>Primary: There were no significant differences among the groups with regards to the 6-minute walk distance over the 43 week study period.</p> <p>Secondary: There were no significant differences among the groups with regards to the NYHA functional class or QOL at 18 or 43 weeks.</p> <p>Ejection fraction increased more with candesartan plus enalapril than monotherapy with either agent; however, the difference was not statistically significant (P value not significant). End-diastolic volumes (P&lt;0.01) and end-systolic volumes (P&lt;0.05) increased less with combination therapy than with monotherapy with either agent.</p> <p>Aldosterone decreased with combination therapy at 17 but not 43 weeks compared to candesartan or enalapril (P&lt;0.05). Brain natriuretic peptide decreased with combination therapy compared to candesartan and enalapril alone (P&lt;0.01).</p> <p>Blood pressure decreased with combination therapy compared to candesartan or enalapril alone (P&lt;0.05).</p> <p>Compared to enalapril, potassium decreased with candesartan use (P&lt;0.05) and increased with candesartan plus enalapril (P&lt;0.05). The proportion of patients with potassium levels <math>\geq 5.5</math> mmol/L was not significantly different among the treatment groups. There were no significant differences in creatinine, mortality, or hospitalizations for CHF or any cause among the three groups.</p>
Willenheimer et	BE, MC, OL, PG,	N=1,010	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>al.<sup>79</sup> (2005) CIBIS-III</p> <p>Enalapril 2.5 to 10 mg BID</p> <p>vs</p> <p>bisoprolol 1.25 to 10 mg QD</p>	<p>RCT</p> <p>Patients ≥65 years with stable mild to moderate CHF (NYHA class II to III), LVEF of ≤35% ≥3 months prior to randomization, not on an ACE inhibitor, β-blocker or ARB therapy and no clinically relevant fluid retention of diuretic adjustment within the 7 days prior to randomization</p>	<p>1.22±0.42 years</p>	<p>Combined all-cause mortality or hospitalization</p> <p>Secondary: Combined end point at the end of the monotherapy phase and the individual components of the primary end point, cardiovascular death and cardiovascular hospitalization, permanent treatment cessation and the need for early introduction of the second drug as indicators of drug tolerability</p>	<p>There were 178 patients (35.2%) with a primary end point of combined all-cause mortality or all-cause hospitalization in the bisoprolol-first group, compared to 186 (36.8%) patients in the enalapril-first group (absolute difference, -1.6%; 95% CI, -7.6 to 4.4; HR, 0.94; 95% CI, 0.77 to 1.16; non-inferiority for bisoprolol-first vs enalapril-first treatment; P=0.019).</p> <p>Secondary: The combined endpoint at the end of the monotherapy phase occurred in 109 patients in the bisoprolol-first group compared to 108 patients in the enalapril-first group (HR, 1.02; 95% CI, 0.78 to 1.33; between-group difference P=0.90); 23 vs 32 patients died, respectively (HR, 0.72; 95% CI, 0.42 to 1.24; between-group difference P=0.24); and 99 vs 92 patients had been a hospitalization, respectively (HR, 1.08; 95% CI, 0.81 to 1.43; between-group difference P=0.59).</p> <p>There were 65 deaths in the bisoprolol-first group, as compared to 73 in the enalapril-first group (HR, 0.88; 95% CI, 0.63 to 1.22; between-group difference P=0.44).</p> <p>In the bisoprolol-first group, 151 patients were hospitalized, compared to 157 patients in the enalapril-first group (HR, 0.95; 95% CI, 0.76 to 1.19; between-group difference P=0.66).</p> <p>There was not a significant difference in cardiovascular death rate observed between the bisoprolol-first (55) and enalapril-first (56) treatment groups (HR, 0.97; 95% CI, 0.67 to 1.40; between-group difference P=0.86).</p> <p>During the monotherapy phase, 35 (6.9%) patients in the bisoprolol-first group permanently discontinued therapy, compared to 49 (9.7%) patients in the enalapril-first group. During the combined-therapy phase, 19 patients (4.2%) in the bisoprolol-first group permanently discontinued bisoprolol therapy and 47 (10.4%) discontinued enalapril therapy. In the enalapril-first group, 24 patients (5.5%) permanently discontinued bisoprolol and 16 (3.7%) discontinued enalapril.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				There was not a statistical significant difference observed in the early introduction of the second drug between the bisoprolol-first group (39 [7.7%] patients) compared to the enalapril-first group (37 [7.3%] patients; P=0.81).
<p>Cohn et al.<sup>80</sup> (1991) V-HEFT II</p> <p>Enalapril 20 mg/day</p> <p>vs</p> <p>hydralazine 300 mg plus isosorbide dinitrate 160 mg/day</p>	<p>AC, DB, MC, RCT</p> <p>Men between the ages of 18 and 75 years with chronic heart failure receiving digoxin and diuretic therapy</p>	<p>N=804</p> <p>2 years</p>	<p>Primary: Mortality</p> <p>Secondary: Peak oxygen consumption during exercise, LVEF</p>	<p>Primary: Mortality after two years was significantly lower in the group treated with enalapril (18%) than hydralazine plus isosorbide dinitrate (25%; P=0.016), and overall mortality tended to be lower (P=0.08).</p> <p>The lower mortality in the enalapril arm was attributable to a reduction in the incidence of sudden death, and this beneficial effect was more prominent in patients with less severe symptoms (NYHA class I or II).</p> <p>Secondary: Peak oxygen consumption during exercise was increased only by hydralazine plus isosorbide dinitrate (P&lt;0.05).</p> <p>While LVEF increased with both regimens during the two years after randomization, LVEF increased more (P&lt;0.05) during the first 13 weeks in the hydralazine plus isosorbide dinitrate group.</p>
<p>Tu et al.<sup>81</sup> (2005)</p> <p>Enalapril</p> <p>vs</p> <p>lisinopril, ramipril, and other ACE inhibitors (benazepril, captopril, cilazapril*, fosinopril, perindopril, quinapril, and trandolapril)</p>	<p>RETRO</p> <p>Patients &gt;65 years with newly diagnosed CHF initiated on ACE inhibitors who survived ≥30 days after hospital discharge</p>	<p>N=6,753</p> <p>≤2 years</p>	<p>Primary: Combined end point of readmission for CHF as a primary diagnosis or mortality</p> <p>Secondary: CHF readmission alone and mortality alone</p>	<p>Primary: Relative to enalapril users, there were no significant differences in combined end point of readmission for CHF or mortality with lisinopril (adjusted HR, 1.18; 95% CI, 0.94 to 1.23), ramipril (adjusted HR, 1.16; 95% CI, 0.92 to 1.24) or other ACE inhibitors (adjusted HR, 1.12; 95% CI, 0.90 to 1.17).</p> <p>Secondary: There were no significant differences among groups in readmission for CHF: enalapril 13% (adjusted HR, 1.0; 95% CI, 0.92 to 1.32), lisinopril 15% (adjusted HR, 1.11; 95% CI, 0.92 to 1.45), and other ACE inhibitors 15% (adjusted HR, 1.13; 95% CI, 0.96 to 1.34).</p> <p>There were no significant differences among groups in mortality: enalapril 12% (adjusted HR, 1.0; 95% CI, 0.90 to 1.31), lisinopril 13% (adjusted HR, 1.19; 95% CI, 0.78 to 1.20), and other</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				ACE inhibitors 11% (adjusted HR, 0.94; 95% CI, 0.78 to 1.13).
Packer et al. <sup>82</sup> (1999) ATLAS  Lisinopril 2.5 to 5 mg/day (low dose)  vs  lisinopril 32.5 to 35 mg/day (high dose)	DB, RCT  Patients with NYHA class II, III, or IV symptoms of heart failure associated with a LVEF $\leq$ 30% despite treatment with diuretics for $\geq$ 2 months	N=3,164  39 to 58 months	Primary: All-cause mortality  Secondary: cardiovascular mortality, hospitalizations (for any reason and for cardiovascular reasons), combinations of the primary and secondary end points	ACE inhibitors 11% (adjusted HR, 0.94; 95% CI, 0.78 to 1.13).  Primary: High-dose lisinopril was associated with a nonsignificant 8% lower risk of all-cause mortality compared to low-dose lisinopril (P=0.128).  Secondary: Cardiovascular mortality was reported in 40.2 and 37.2% of patients receiving low-dose and high-dose lisinopril, respectively (P=0.073).  High-dose lisinopril resulted in a 12% lower risk of death or hospitalizations for any reason (P=0.002), a 9% lower risk of cardiovascular mortality and hospitalization for cardiovascular reason (P=0.027) and 24% fewer hospitalizations for heart failure (P=0.002).  Dizziness and renal insufficiency were observed more frequently in the high-dose group, but the two groups were similar in the number of patients requiring discontinuation of the study medication.
AIRE Study Investigators <sup>83</sup> (1993) AIRE  Ramipril 2.5 to 5 mg BID  vs  placebo	DB, MC, PC, RCT  Patients $\geq$ 18 years of age with acute MI and clinical evidence of heart failure	N=2,006  15 months	Primary: All-cause mortality  Secondary: First event in an individual patient (death, progression to severe or resistant heart failure, reinfarction, or stroke)	ACE inhibitors 11% (adjusted HR, 0.94; 95% CI, 0.78 to 1.13).  Primary: On the intention-to-treat analysis, all-cause mortality was significantly lower for patients randomized to receive ramipril (17%) than placebo (23%). The observed risk reduction was 27% (95% CI, 11 to 40; P=0.002).  Secondary: Analysis of prespecified secondary outcomes revealed a 19% risk reduction in the ramipril group compared to placebo (95% CI, 5 to 31; P=0.008).
Wu et al. <sup>84</sup> (2021) AIRE-S (AIRE survival)  Ramipril target dose 10 mg/day	DB, MC, PC, PG, RCT  Patients $\geq$ 18 years of age with acute MI and clinical evidence of heart failure	N=603  29.6 years	Primary: All-cause mortality for UK participants  Secondary: Life expectancy and extensions of life (difference)	ACE inhibitors 11% (adjusted HR, 0.94; 95% CI, 0.78 to 1.13).  Primary: By April 9, 2019, death from all causes occurred in 266 (88.4%) patients in placebo arm and 275 (91.1%) patients in ramipril arm. Survival at the end of study follow-up was not different between the ramipril and placebo groups (HR, 0.96; 95% CI, 0.81 to 1.15) because more-or-less all patients had died at the end of follow-up.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			in median survival times)	Ramipril treatment improved life expectancy across all of the comorbidity groups studied, including diabetes (LED, 32.1 vs 5.0 months), previous MI (20.1 vs 4.9 months), history of heart failure (19.5 vs 4.9 months), hypertension (16.6 vs 8.3 months), angina (16.2 vs 5.0 months) and was greater for patients aged $\geq 65$ years than $< 65$ years (11.3 vs 5.7 months). The median survival time was 8.3 years (95% CI, 6.2 to 10.2 years) for placebo group and 9.6 years (95% CI, 8.3 to 10.9 years) for ramipril group. The extension of life between ramipril and placebo groups, measured by the difference in median survival time, was 14.5 (95% CI, 13.2 to 15.8) months.
Kober et al. <sup>85</sup> (1995) TRACE  Trandolapril 1 to 4 mg QD  vs  placebo  Medication was started between day 3 and 7 after the myocardial infarction.	DB, MC, PC, RCT  Men and women >18 years who were hospitalized with a recent MI and an LVEF $\leq 35\%$	N=1,749  24 to 50 months	Primary: Death from any cause  Secondary: Death from a cardiovascular cause, sudden death, progression to severe heart failure (defined as the first of the following events: hospital admission for heart failure, death due to progressive heart failure, or heart failure necessitating the administration of open-label ACE inhibition), recurrent infarction, change in the wall-motion index	Primary: During the study, 34.7% of patients in the trandolapril group died compared to 42.3% in the placebo group (P=0.001). The relative risk of death in the trandolapril group was 0.78 compared to placebo (95% CI, 0.67 to 0.91).  Secondary: Trandolapril reduced the risk of death from cardiovascular causes (RR, 0.75; 95% CI, 0.63 to 0.89; P=0.001) and sudden death (RR, 0.76; 95% CI, 0.59 to 0.98; P=0.03).  Progression to severe heart failure was less frequent in the trandolapril group (RR, 0.71; 95% CI, 0.56 to 0.89; P=0.003).  The risk of recurrent fatal or nonfatal MI was not significantly reduced (RR, 0.86; 95% CI, 0.66 to 1.13; P=0.29).  After three months, the mean change from the base-line index was 0.09 in the trandolapril group and 0.06 in the placebo group (P=0.03) but this statistically significant difference was absent at six and 12 months.



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Galløe et al.<sup>86</sup> (2006)</p> <p>Trandolapril 0.5 mg (0, 1, 2 or 4 tablets QD) plus bumetanide 0.5 mg (0, 1, 2 or 4 tablets BID)</p> <p>Treatment was combined to achieve 16 different dosage combinations.</p>	<p>DB, DD, RCT, multiple XO</p> <p>Patients with previous MI <math>\geq 3</math> years ago, had medical treatment for heart failure and ejection fraction between 0.36 and 0.54 estimated by echocardiography</p>	<p>N=16</p> <p>14 days</p>	<p>Primary: Patient reported QOL</p> <p>Secondary: Effects on kidney function, left ventricular function and blood pressure</p>	<p>Primary: Bumetanide 0.5 mg-treated patients experienced a 12% increase in well-being, but higher doses of bumetanide decreased patient's well-being by 12% compared to placebo (P&lt;0.002). Increasing doses of bumetanide tended to increase tiredness (P=0.072). There were no significant effects of bumetanide therapy on the patients' opinion of their health, degree of dyspnea, appetite or work capacity.</p> <p>Secondary: Bumetanide therapy increased 24 hour urine production in a straight dose-dependent manner (P&lt;0.0001), while trandolapril therapy had no effect (P=0.53). Bumetanide and trandolapril therapy did not alter the 24 hour creatinine excretion and creatinine clearance (P=0.33, P=0.11 and P=0.53, P=0.97, respectively).</p> <p>Bumetanide therapy decreased left ventricular function and increased heart rate in a dose-dependent manner (P&lt;0.001). Left ventricular function was also nonsignificantly decreased with trandolapril therapy (P&gt;0.062).</p> <p>Trandolapril therapy significantly reduced SBP by maximally of 7.6 mm Hg (5.8%) with the lowest dose of 0.5 mg/day (P=0.007). Bumetanide therapy had no significant effect on DBP (P=0.23).</p>
<p>Galloe et al.<sup>87</sup> (2006)</p> <p>Trandolapril 0.5 mg (0, 1, 2, or 4 tablets QD)</p> <p>vs</p> <p>bumetanide 0.5 mg (0, 1, 2, or 4 tablets BID)</p>	<p>DB, PC, RCT, XO</p> <p>Men and women with previous MI <math>\geq 3</math> years ago, had medical treatment for heart failure and ejection fraction between 0.36 and 0.54 estimated by echo-cardiography (wall motion index)</p>	<p>N=16</p> <p>14 days</p>	<p>Primary: Patient reported QOL</p> <p>Secondary: Effects on the involved organs: kidney function, left ventricular function, blood pressure</p>	<p>Primary: Patient's well-being increased 12% with 0.5 mg bumetanide BID but higher doses bumetanide decreased patient's well-being by 12% compared to placebo (P&lt;0.002). Increasing doses of bumetanide tended to increase tiredness (P=0.072). There were no statistically significant effects of bumetanide on the patient's opinion of their health, degree of dyspnea, appetite or work capacity.</p> <p>Secondary: Bumetanide increased 24-hour urine production in a straight dose-dependent manner (P&lt;0.0001) while trandolapril had no effect (P=0.53). Bumetanide and trandolapril did not alter the 24-hour creatinine excretion and creatinine clearance (P=0.33, P=0.11 and P=0.53, P=0.97, respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Bumetanide decreased left ventricular function and increased heart rate in a dose dependent manner (P&lt;0.001). Left ventricular function was also decreased with trandolapril but did not reach statistically significant. (P&gt;0.062).</p> <p>Trandolapril significantly reduced SBP by maximally of 7.6 mm Hg (5.8%) with the lowest dose of 0.5 mg/day (P=0.007). Bumetanide had no significant effect on DBP (P=0.23).</p>
<p>Fröhlich et al.<sup>88</sup> (2018)</p> <p>Enalapril vs lisinopril vs ramipril</p> <p>Medication used was at the discretion of the referring physician.</p>	<p>Cohort</p> <p>Outpatients with stable HFrEF (EF &lt;45%)</p>	<p>N=4,723</p> <p>≥6 months (Median follow-up of 50 months)</p>	<p>Primary: Mortality</p> <p>Secondary: Not reported</p>	<p>Baseline characteristics of HFrEF patients differed with respect to ACE inhibitor treatment for a number of variables. Overall, patients receiving ramipril were younger and more likely to have NYHA functional Class I or II symptoms than those on enalapril and lisinopril. NT-proBNP levels were lower in the ramipril group, whereas LVEF was similar in all three treatment groups. In patients using lisinopril, systolic BP was significantly higher when compared with patients on enalapril or ramipril.</p> <p>Primary: During a follow-up of 21,939 patient-years, 360 (49.5%), 337 (52.4%), and 1119 (33.4%) patients died among those prescribed enalapril, lisinopril, and ramipril, respectively. In univariable analysis of the general sample, enalapril and lisinopril were both associated with higher mortality when compared with ramipril treatment (HR, 1.46; 95% CI, 1.30 to 1.65; P&lt;0.001; and HR, 1.38; 95% CI, 1.22 to 1.56; P&lt;0.001, respectively). Patients prescribed enalapril or lisinopril had similar mortality (HR, 1.06; 95% CI, 0.92 to 1.24; P=0.41). However, there was no significant association between ACE inhibitor choice and all-cause mortality in any of the matched samples (HR, 1.07; 95% CI, 0.91 to 1.25; P=0.40; HR, 1.12; 95% CI, 0.96 to 1.32; P=0.16; and HR, 1.10; 95% CI, 0.93 to 1.31; P=0.25 for enalapril vs ramipril, lisinopril vs ramipril, and enalapril vs lisinopril, respectively). Results were confirmed in subgroup analyses with respect to age, sex, LVEF, New York Class Association functional class, cause of HFrEF, rhythm, and systolic BP.</p> <p>Secondary: Not reported</p>
<p>Lee et al.<sup>89</sup> (2004)</p>	<p>MA</p>	<p>N=38,080</p>	<p>Primary: All-cause mortality</p>	<p>Primary: ARBs were associated with reduced all-cause mortality (OR, 0.83) and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ARBs</p> <p>vs</p> <p>placebo (±ACE inhibitor)</p> <p>vs</p> <p>ACE inhibitor monotherapy</p>	<p>Patients with chronic heart failure and high-risk acute MI</p>	<p>Duration varied</p>	<p>and heart failure hospitalizations</p> <p>Secondary: Not reported</p>	<p>heart failure hospitalizations (OR, 0.64) vs placebo.</p> <p>There was no difference in all-cause mortality (OR, 1.06) and heart failure hospitalization (OR, 0.95) between ARBs and ACE inhibitors.</p> <p>When ARBs were combined with ACE inhibitors, all-cause mortality was not reduced (OR, 0.97) but heart failure hospitalizations were reduced (OR, 0.77) compared to treatment with ACE inhibitors alone.</p> <p>Two RCT comparing ARBs with ACE inhibitors in patients with high-risk acute MI did not reveal differences in all-cause mortality or heart failure hospitalization.</p> <p>Secondary: Not reported</p>
<b>Hypertension</b>				
<p>Kuschnir et al.<sup>90</sup> (1996)</p> <p>Benazepril 20 mg/day and amlodipine 5 mg/day</p> <p>vs</p> <p>amlodipine 5 mg/day</p> <p>vs</p> <p>benazepril 20 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PG, RCT</p> <p>Men and women 21 to 80 years of age with uncomplicated primary HTN</p>	<p>N=308</p> <p>8 weeks</p>	<p>Primary: Reduction in mean sitting DBP, SBP and percentage of patients with DBP &lt;90 mm Hg or a ≥10 mm Hg reduction</p> <p>Secondary: Not reported</p>	<p>Primary: All treatment groups significantly reduced mean sitting DBP compared to placebo (P&lt;0.001).</p> <p>Combination therapy had significantly greater reductions in DBP -13.2 mm Hg; P&lt;0.001) compared to amlodipine (-8.8 mm Hg) and benazepril (-6.7 mm Hg) monotherapy.</p> <p>Combination therapy had significantly greater reductions in SBP (-24.7 mm Hg; P&lt;0.001) compared to amlodipine (-16.2 mm Hg) and benazepril (-12.4 mm Hg).</p> <p>Significantly more patients on combination therapy reached DBP &lt;90 mm Hg or a ≥10 mm Hg reduction (87.0%; P≤0.005) compared to amlodipine (67.5%) and benazepril (53.3%) monotherapy.</p> <p>Adverse events considered to be drug related occurred in 15.6% of patients receiving combination therapy, 24.7% of patients receiving amlodipine monotherapy, 6.5% of patients on benazepril monotherapy and 11.7% of patients on placebo (P values not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Neutel et al. <sup>91</sup> (2005) SELECT  Benazepril and amlodipine 20-5 mg/day (fixed dose combination product)  vs  amlodipine 5 mg/day  vs  benazepril 20 mg/day	DB, RCT  Patients with stage 2 systolic HTN	N=443  8 weeks	Primary: Reduction in SBP, proportion of patients achieving blood pressure control  Secondary: Not reported	Secondary: Not reported  Primary: Significantly greater SBP reductions were achieved with combination therapy compared to amlodipine or benazepril monotherapy (P<0.0001).  Significantly more patients on combination therapy met blood pressure goals than on monotherapy (P<0.0001).  No significant difference was noted in the incidence of adverse events. Adverse events were low in all three treatment arms, with less peripheral edema in the combination group than in the amlodipine-treated group.  Secondary: Not reported
Chrysant <sup>92</sup> (2004)  Amlodipine and benazepril 5-40 mg QD for 4 weeks, followed by 10-40 mg QD for 4 weeks (fixed-dose combination product)  vs  benazepril 40	DB, RCT  Men and women (mean age 53 years) with mean sitting DBP ≥95 mm Hg not adequately controlled with benazepril 40 mg/day monotherapy	N=329  8 weeks	Primary: Reduction in mean sitting DBP and SBP, reduction in standing DBP and SBP, and change in heart rate, safety  Secondary: Not reported	Primary: Combination therapy had significantly greater reductions in sitting SBP (-17 mm Hg; P<0.0001) compared monotherapy (-5 mm Hg).  Combination therapy had significantly greater reductions in sitting DBP (-14 mm Hg; P<0.0001) compared to monotherapy (-7 mm Hg).  Combination therapy had significantly greater reductions in standing SBP (-17 mm Hg; P<0.0001) compared to monotherapy (-6 mm Hg).  Combination therapy had significantly greater reductions in standing DBP (-14 mm Hg; P<0.0001) compared to monotherapy (-7 mm Hg).  No significant differences in heart rate were observed (P>0.05).  No significant differences in adverse events were reported (P>0.05).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day for 8 weeks				Secondary: Not reported
Fogari et al. <sup>93</sup> (1997)  Benazepril 10 mg QD  vs  amlodipine and benazepril 2.5-10 to 5-10 mg QD (fixed-dose combination product)	DB, MC, PC, RCT  Men and women 24 to 73 years of age (mean 55 years) with HTN inadequately controlled with ACE inhibitor monotherapy	N=448  8 weeks	Primary: Reduction in mean sitting DBP  Secondary: Reduction in sitting SBP, standing DBP and SBP, and percentage of patients with DBP <90 mm Hg (deemed excellent response) or a ≥10 mm Hg reduction (deemed good response)	Primary: Significantly greater reductions in sitting DBP were observed with benazepril 10 mg and amlodipine 2.5 mg (-5.3 mm Hg, 97.5% CI, -8.3 to -2.4; P=0.0001) and benazepril 10 mg and amlodipine 5 mg (-4.5 mm Hg, 97.5% CI, -7.4 to -1.6; P=0.0006) compared to benazepril monotherapy.  Secondary: Significantly greater reductions in sitting SBP were seen with benazepril 10 mg and amlodipine 2.5 mg (-7.9 mm Hg, 97.5% CI, -12.3 to -3.5; P=0.0001) and benazepril 10 mg and amlodipine 5 mg (-7.9 mm Hg, 97.5% CI, -12.2 to -3.6; P=0.0000) compared to benazepril monotherapy.  Significantly greater reductions in standing DBP and SBP were also reported with the combination therapy compared to benazepril monotherapy (P≤0.001).  Significantly more patients had excellent or good response with benazepril 10 mg and amlodipine 2.5 mg (69.2%; P=0.0004) and 10-5 mg (65.8%; P=0.02) compared to benazepril monotherapy (40.5%).  Tolerability was good in the three treatment groups and no significant abnormal laboratory data was detected.
Chrysant et al. <sup>94</sup> (2012)  Study 1: Benazepril 40 mg/day (Group 1)  vs  amlodipine and benazepril 5-40 mg/day, up	Post-hoc analysis of 2 trials  Patients with HTN	N=1,013  14 weeks	Primary: Change in baseline mean sitting DBP and mean sitting SBP, rate of blood pressure control (<140/90 mm Hg), rate of blood pressure control (mean sitting DBP <90 mm Hg or ≥10 mm Hg decrease	Primary: Pooled results demonstrate that combination therapy resulted in significantly greater lowering of mean sitting DBP and mean seated SBP compared to benazepril or amlodipine (P<0.001). Amlodipine and benazepril 10-20 mg/day resulted in significantly greater blood pressure reductions in White patients (mean sitting DBP: 12.99 mm Hg; mean sitting SBP: 13.72 mm Hg) compared to Black patients (8.80 and 8.72 mm Hg) (P<0.004). Amlodipine and benazepril 10-40 mg/day resulted in similar reductions in blood pressure in both White and Black patients.  The proportion of patients who achieved blood pressure control with amlodipine and benazepril 10-40 mg/day was similar between White and

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>titrated to 10-40 mg/day after 4 weeks (fixed-dose combination product) (Group 2)</p> <p>Study 2: Amlodipine and benazepril 10-20 mg/day, uptitrated to 10-40 mg/day after 2 weeks (Group 3)</p> <p>vs</p> <p>amlodipine and benazepril 10-20 mg/day (fixed-dose combination product) (Group 4)</p> <p>vs</p> <p>amlodipine 10 mg/day (Group 5)</p>			<p>from baseline)</p> <p>Secondary: Safety</p>	<p>Black patients (60.7%), whereas with amlodipine and benazepril 10-20 mg/day the rate of control was higher with White patients (61.2 vs 39.4%; P&lt;0.023).</p> <p>There was no difference in the proportion of patients who responded to treatment between Black and White patients with amlodipine and benazepril 10-40 mg/day (74.8 vs 77%; P&lt;0.639). The proportion of patients who responded to amlodipine and benazepril 10-20 mg/day was significantly lower in Black patients (50.7 vs 73.5%; P&lt;0.007).</p> <p>Secondary: There were no serious clinical or metabolic side effects reported, with the exception of pedal edema which occurred more frequently with amlodipine monotherapy.</p>
<p>Messerli et al.<sup>95</sup> (2000)</p> <p><u>Study 1:</u> Amlodipine and benazepril 5-10 mg to 5-20 mg QD (fixed-dose combination</p>	<p>2 DB, MC, RCT</p> <p>Patients 18 to 80 years of age with uncomplicated essential HTN</p>	<p>N=1,079</p> <p>8 weeks</p>	<p>Primary: Change in DBP from baseline</p> <p>Secondary: Change from baseline in SBP and heart rate</p>	<p>Primary: Study 1 Significant reductions in DBP were observed with benazepril and amlodipine 10-5 and 20-5 mg (-9.4 and -9.7 mm Hg, respectively) compared to nifedipine 30 mg (-7.0 mm Hg; P&lt;0.05), but not nifedipine 60 mg (-8.5; P&gt;0.05).</p> <p>Study 2 Benazepril and amlodipine 10-5 (-8.9 mm Hg) and 20-5 mg (-9.1 mm Hg)</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>product)</p> <p>vs</p> <p>nifedipine 30 to 60 mg/day</p> <p><u>Study 2:</u> Amlodipine and benazepril 5-10 mg to 5-20 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>				<p>produced significantly greater reductions in DBP than amlodipine 5 mg (-6.8 mm Hg; P&lt;0.05), but not amlodipine 10 mg (-8.7 mm Hg; P&gt;0.05).</p> <p>Secondary: Study 1 Significant reductions in SBP were observed with benazepril and amlodipine 20-5 mg (-11.6 mm Hg) compared to nifedipine 30 mg (-7.9 mm Hg; P&lt;0.05).</p> <p>Significantly less edema was reported with combination therapies (3.1 to 3.8%; P≤0.001) compared to nifedipine 60 mg (15.5%; P=0.008) but not nifedipine 30 mg (5.4%).</p> <p>Study 2 Significant reductions in SBP were observed with benazepril and amlodipine 20-5 mg (-9.1 mm Hg) compared to amlodipine 5 mg (-5.3 mm Hg; P&lt;0.05). There were no significant difference in SBP between amlodipine 10 mg and the combination therapies.</p> <p>Significantly less edema (P&lt;0.001) was reported with amlodipine 5 mg (4.9%) and combination therapies (1.5 to 2.2%) compared to amlodipine 10 mg (23.6%).</p>
<p>Hilleman et al.<sup>96</sup> (1999)</p> <p>Benazepril and amlodipine (fixed-dose combination product)</p> <p>vs</p> <p>monotherapy (atenolol, HCTZ, captopril, enalapril,</p>	<p>MA</p> <p>Patients with mild-to-moderate essential HTN</p>	<p>82 trials</p> <p>≥4 weeks</p>	<p>Primary: Absolute change in supine DBP from baseline</p> <p>Secondary: Percent of patients who achieved blood pressure control, safety</p>	<p>Primary: The mean absolute decrease in supine DBP ranged from 9.7 to 13.3 mm Hg with verapamil showing the greatest effect and captopril the least. When studies were weighted by sample size, amlodipine and benazepril, atenolol, lisinopril, and verapamil showed the greatest blood pressure effect.</p> <p>Secondary: The average percentage of patients defined as controlled after treatment varied from 53.5 to 79.0%, with amlodipine and benazepril (74.3%) and lisinopril (79.0%) showing the highest percentage control (P=0.096).</p> <p>The incidence of adverse events ranged from 12.1 to 41.8%, with lisinopril and verapamil showing the lowest incidences (12.1% and 14.1%, respectively) and nifedipine the highest incidence. Lisinopril demonstrated</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
lisinopril, amlodipine, diltiazem, nifedipine, verapamil)				<p>significantly less overall side effects compared to nifedipine (P=0.030).</p> <p>Nifedipine demonstrated a higher withdrawal rate due to side effects compared to atenolol, HCTZ, enalapril, amlodipine, and diltiazem (P=0.002). Although amlodipine and benazepril had the lowest rate of withdrawals due to adverse events, lack of significant change was due to the low number of cohorts available for analysis.</p>
<p>Jamerson et al.<sup>97</sup> (2007) ACCOMPLISH</p> <p>Benazepril 20 to 40 mg QD and HCTZ 12.5 to 25 mg QD</p> <p>vs</p> <p>benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD</p>	<p>DB, MC, RCT</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=10,704</p> <p>Analysis performed at 6 months (complete trial duration 5 years)</p>	<p>Primary: Changes in mean SBP from baseline to 6 months, blood pressure control rates (SBP/DBP &lt;140/90 mm Hg or &lt;130/89 mm Hg for patients with diabetes and chronic kidney disease)</p> <p>Secondary: Not reported</p>	<p>Primary: At baseline, 97% of subjects were treated with antihypertensive medications at entry, but only 37% of participants had blood pressure control.</p> <p>Mean blood pressure fell from 145/80 to 132/74 mm Hg after six months of treatment with either combination regimen (P&lt;0.001).</p> <p>The six month blood pressure control rate was 73% in the overall trial (78% in the United States), 43% in diabetics, and 40% in patients with renal disease. Of the patients uncontrolled, 61% were not on maximal medications.</p> <p>Secondary: Not reported</p>
<p>Kereiakes et al.<sup>98</sup> (2007)</p> <p>Benazepril 10 mg/day for 2 weeks, then 20 mg/day for 2 weeks, then benazepril 20 mg/day plus amlodipine 5 mg/day for 4 weeks, then benazepril 20 mg/day plus</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients with stage 2 HTN</p>	<p>N=190</p> <p>12 weeks</p>	<p>Primary: Change in mean seated SBP at the end of week 12</p> <p>Secondary: DBP at the end of week 12, percent of patients attaining blood pressure goals of &lt;140/90, &lt;130/85, and &lt;130/80 mm Hg</p>	<p>Primary: Patients treated with olmesartan and HCTZ experienced significantly greater reductions in mean seated SBP at week 12 than patients treated with benazepril plus amlodipine (least square mean change, -32.5 vs -26.5 mm Hg; P=0.024; least square mean treatment difference, -6.0 mm Hg; 95% CI, -11.1 to -0.8).</p> <p>Secondary: The least square mean change for reduction in DBP approached statistical significance with olmesartan and HCTZ compared to benazepril plus amlodipine at week 12 (P=0.056).</p> <p>The percentage of patients achieving goal rates at the end of the study for olmesartan and HCTZ and benazepril plus amlodipine were 66.3 and 44.7% (P=0.006) for &lt;140/90 mm Hg, 44.9 vs 21.2% (P=0.001) for</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amlodipine 10 mg/day for 4 weeks</p> <p>vs</p> <p>olmesartan 20 mg/day for 2 weeks, then 40 mg/day for 2 weeks then olmesartan and HCTZ 40-12.5 mg/day for 4 weeks increased to 40-25 mg for 4 weeks</p>				<p>&lt;130/85 mm Hg, and 32.6 and 14.1% (P=0.006) for &lt;130/80 mm Hg.</p> <p>Both treatments were well tolerated.</p>
<p>Waeber et al.<sup>99</sup> (2001)</p> <p>Valsartan 80 mg QD, which was switched to valsartan 80 mg and HCTZ 12.5 mg QD or valsartan 80 mg and benazepril 10 mg QD</p>	<p>OL, RCT</p> <p>Patients with mild-to-moderate uncontrolled HTN (DBP ≥90) while on valsartan monotherapy</p>	<p>N=327</p> <p>4 weeks</p>	<p>Primary: Efficacy and safety</p> <p>Secondary: Not reported</p>	<p>Primary: The two combinations produced an additional blood pressure reduction compared to monotherapy (P&lt;0.001 for both), with similar DBP reductions reported for the two combination groups (-4.5 mm Hg with valsartan plus HCTZ and -3.3 mm Hg with valsartan plus benazepril).</p> <p>SBP reductions of -6.7 and -3.2 mm Hg with valsartan plus HCTZ and valsartan plus benazepril, respectively, were reported (P=0.1).</p> <p>At the end of the trial, the blood pressure of the responders to valsartan monotherapy was lower than that of patients requiring combination therapy.</p> <p>Valsartan given alone or in association with HCTZ or benazepril was well tolerated.</p> <p>Secondary: Not reported</p>
<p>Malacco et al.<sup>100</sup> (2002)</p>	<p>DB, MC, RCT</p>	<p>N=397</p>	<p>Primary: Reduction in</p>	<p>Primary: Significantly lower sitting DBP (-2.7 mm Hg; P&lt;0.001) and SBP (-3.7 mm</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Captopril and HCTZ 50-25 mg/day (fixed-dose combination)</p> <p>vs</p> <p>amlodipine and benazepril 5-10 mg/day (fixed-dose combination)</p>	<p>Patients with mild-to-moderate arterial HTN (sitting DBP &gt;95 mm Hg and/or SBP &gt;160 mm Hg) inadequately controlled by monotherapy with an ACE inhibitor, calcium-channel blocking agent or diuretic</p>	<p>12 weeks</p>	<p>sitting DBP and SBP</p> <p>Secondary: Percentage of patients responding to therapy (DBP&lt;90 mm Hg, reduction in DBP ≥10 mm Hg or SBP ≥20 mm Hg, or SBP &lt;150 mm Hg)</p>	<p>Hg; P&lt;0.001) were achieved with amlodipine and benazepril compared to captopril and HCTZ.</p> <p>Secondary: Significantly more amlodipine and benazepril patients responded to therapy (94.8%) compared to captopril and HCTZ (86.0%; P=0.004).</p> <p>No differences in adverse events were reported between the two treatment groups.</p>
<p>Elliot et al.<sup>101</sup> (1999)</p> <p>Enalapril 10 mg QD</p> <p>vs</p> <p>enalapril and felodipine ER 5-5 mg/day (fixed-dose combination)</p> <p>After 6 weeks, all patients received the fixed-dose combination for an additional 6 weeks.</p>	<p>DB, PG, PRO, RCT, XO</p> <p>Patients with sitting DBP &gt;95 mm Hg and &lt;115 mm Hg</p>	<p>N=217</p> <p>12 weeks</p>	<p>Primary: Change in sitting DBP, proportion of responders (DBP &lt;90 mm Hg or a reduction of &gt;10 mm Hg)</p> <p>Secondary: Not reported</p>	<p>Primary: Patients receiving combination therapy had significantly greater reductions in sitting SBP and DBP compared to baseline (P&lt;0.05 and P&lt;0.01, respectively).</p> <p>More patients receiving combination therapy were classified as responders than patients receiving enalapril monotherapy (59 vs 41%; P&lt;0.01).</p> <p>When patients originally taking 10 mg enalapril were crossed over to the combination therapy for an additional six weeks, there was a further blood pressure reduction and increase in response rate, with loss of significant differences compared to those treated continuously with the combination for the entire 12 weeks.</p> <p>There were no significant differences in tolerability between the regimens.</p> <p>Secondary: Not reported</p>
<p>Prisant et al.<sup>102</sup> (1995)</p> <p>Enalapril 5, 10, or</p>	<p>DB, MC, PG, RCT</p> <p>Patients ≥21 years with mild to</p>	<p>N=218</p> <p>17 weeks</p>	<p>Primary: Mean change from baseline in SBP and DBP, lab</p>	<p>Primary: Mean decreases in SBP and DBP from baseline were 13.4/10.7 mm Hg for bisoprolol and HCTZ patients, 12.8/10.2 mm Hg for amlodipine patients, and 7.3/6.6 mm Hg for enalapril patients. The hypotensive effects were</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>20 mg vs bisoprolol and HCTZ 2.5-6.25, 5-6.25, or 10-6.25 mg/day (fixed-dose combination product) vs amlodipine 2.5, 5, or 10 mg</p>	<p>moderate essential HTN, (average sitting DBP 95 to 114 mm Hg) each treatment was once daily and titrated to effect</p>		<p>measurements, adverse events, QOL questionnaire  Secondary: Not reported</p>	<p>significant for all three groups (P&lt;0.001).  SBP and DBP mean changes from baseline for the bisoprolol and HCTZ group and the amlodipine group were greater than the change from baseline for the enalapril group (P&lt;0.01).  Response rates (DBP ≤90 mm Hg or ≥10 mm Hg decrease from baseline) were 71% for the bisoprolol and HCTZ group, 69% for the amlodipine group, and 45% for the enalapril group. The response rates for the bisoprolol and HCTZ and the amlodipine groups differed significantly from the enalapril group (P&lt;0.01).  Twenty nine percent of bisoprolol patients had adverse experiences compared to 42% of amlodipine patients (P=0.12). Nearly 47% of enalapril patients had adverse experience compared to bisoprolol (P=0.04). Adverse events reported included headache, fatigue, peripheral edema, and dizziness.  Drug related adverse events were 16% for the bisoprolol and HCTZ patients, 21% for the amlodipine patients, and 23% for the enalapril patients. There was no significant difference between the groups.  Enalapril demonstrated a mean decrease from baseline of 7.9 mg/dL for TC (P=0.02 vs amlodipine) and 6.6 mg/dL for LDL-C (P=0.04 vs amlodipine) which were not significantly different from the increase from the bisoprolol and HCTZ group of 1.7 mg/dL (P=0.07 vs enalapril) for TC and +0.6 mg/dL in LDL-C. However, the increase in TGs was highest for bisoprolol and HCTZ-treated patients compared to amlodipine- and enalapril-treated patients (P=0.08, for bisoprolol and HCTZ vs enalapril).  There was not a significant difference from baseline or between treatment groups in QOL scores: 0.9 for the bisoprolol and HCTZ group, 0.5 for the amlodipine group, and 2.3 for the enalapril group.</p>
<p>Ruilope et al.<sup>103</sup> (2001)  Enalapril 5 mg</p>	<p>DB, MC, PG, RCT  Patients greater than 65 years of age with</p>	<p>N=334  12 weeks</p>	<p>Primary: Mean change from baseline in sitting SBP</p>	<p>Primary: No significant difference between groups in change from baseline in sitting SBP was observed (P=0.76).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD (titration to 10 mg followed by 20 mg was allowed every 3 weeks)</p> <p>vs</p> <p>eprosartan 600 mg QD (titration to 800 mg QD was allowed after 3 weeks)</p>	<p>essential HTN, either newly diagnosed or for whom a change in existing antihypertensive medication is indicated due to poor control</p>		<p>Secondary: Normalization rate for sitting SBP and DBP, response rate for sitting SBP and DBP, mean change from baseline in DBP</p>	<p>Secondary: No significant difference between groups in change from baseline in sitting DBP was observed (P=0.84).</p> <p>BP response rates for SBP and DBP were significantly greater for eprosartan at week three (P≤0.033) but the significant difference had disappeared by endpoint (P≥0.49).</p> <p>Normalization rates for SBP were low in both groups (P value not reported).</p> <p>Normalization rates for DBP were higher in both groups than SBP normalization rates (P value not reported).</p>
<p>Karlberg et al.<sup>104</sup> (1999) TEES</p> <p>Enalapril 5 to 20 mg QD</p> <p>vs</p> <p>telmisartan 20 to 80 mg QD</p> <p>HCTZ 12.5 or 25 mg QD could be added to either group as needed to reach DBP goal (≤90 mm Hg).</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients ≥65 years of age with mild- to moderate HTN</p>	<p>N=278</p> <p>26 weeks</p>	<p>Primary: Change from baseline in supine SBP and DBP</p> <p>Secondary: Proportion of responders, safety</p>	<p>Primary: Both treatments had similar rates of HCTZ use.</p> <p>Both treatments showed comparable decreases in blood pressure. Mean changes in DBP were -12.8 mm Hg for telmisartan and -11.4 mm Hg for enalapril (P=0.074). Mean changes in SBP were -22.1 mm Hg for telmisartan and -20.1 mm Hg for enalapril (P=0.350).</p> <p>Secondary: Overall, 63 and 62% of patients responded to telmisartan and enalapril, respectively, with a DBP of &lt;90 mm Hg. Both regimens provided effective blood pressure lowering over the 24-hour dosing interval, as determined by ambulatory blood pressure monitoring.</p> <p>Both regimens were well tolerated; however, the enalapril group had a higher incidence of cough than the telmisartan group (15.8 vs 6.5%; P value reported).</p>
<p>Estacio et al.<sup>105</sup> (1998) ABCD</p> <p>Enalapril 5 to 40 mg/day</p>	<p>DB, PRO, RCT</p> <p>Patients between the ages of 40 and 74 years with NIDDM, baseline DBP ≥90</p>	<p>N=470</p> <p>67 months</p>	<p>Primary: Effect of intensive (target DBP of 75 mm Hg) or moderate (target DBP between 80 to</p>	<p>Primary: Analysis of the 470 patients in the trial who had HTN (DBP ≥90 mm Hg) showed similar control of blood pressure, blood glucose and lipid concentrations between the two study medications throughout the five years of follow-up.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs nisoldipine 10 to 60 mg/day</p>	<p>mm Hg and receiving no antihypertensive medications at the time of randomization</p>		<p>89 mm Hg) blood pressure control on the incidence and progression of complications of diabetes; compare enalapril to nisoldipine as a first-line antihypertensive agent</p> <p>Secondary: Incidence of MI</p>	<p>Secondary: Nisoldipine was associated with a higher incidence of fatal and nonfatal MI than enalapril (RR, 7.0; 95% CI, 2.3 to 21.4).</p>
<p>Williams et al.<sup>106</sup> (2004)</p> <p>Enalapril 10 mg QD</p> <p>vs</p> <p>eplerenone 50 mg QD</p> <p>vs</p> <p>Both medications were titrated to 200 (eplerenone) or 40 (enalapril) mg/day if needed for optimal blood pressure control (DBP &lt; 90 mm Hg).</p>	<p>AC, DB, MC, PG, RCT</p> <p>Patients ≥18 years of age with stage 1 to 2 HTN (seated DBP ≥90 but &lt;110 mm Hg, with a seated SBP &lt;190 mm Hg)</p>	<p>N=499</p> <p>12 months</p>	<p>Primary: Change in seated trough DBP at 6 months</p> <p>Secondary: Change in seated trough SBP at 6 months, reduction in SBP and DBP at 12 months, reduction in urine albumin/ creatinine ratio, adverse events</p>	<p>Primary: At six months, both treatments exhibited comparable reductions in DBP from baseline (P=0.91).</p> <p>Secondary: At six months, both treatments exhibited comparable reductions in SBP from baseline (P=0.20).</p> <p>At 12 months, both treatments exhibited comparable reductions in SBP and DBP from baseline (P=0.25 and P=0.33).</p> <p>Eplerenone-treated patients exhibited a significant reduction from baseline in urine albumin/creatinine ratio compared to enalapril-treated patients (61.5 vs 25.7%; P=0.01).</p> <p>There were no significant differences in overall treatment-emergent adverse events between the two treatments (P value not reported). There were no sex hormone related adverse events in eplerenone-treated patients. There were no clinically significant differences between the two treatments in any of the laboratory tests assessed. There were two eplerenone- and enalapril-treated patients that experienced hyperkalemia of ≥5.5 mmol/L.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Tatti et al.<sup>107</sup> (1998) FACET</p> <p>Fosinopril 20 mg QD</p> <p>vs</p> <p>amlodipine 10 mg QD</p> <p>If blood pressure was not controlled on monotherapy, the other study drug was added.</p>	<p>OL, PRO, RCT</p> <p>Men and women, diagnosed with HTN (SBP &gt;140 mm Hg or DBP &gt;90 mm Hg) and non-insulin dependent diabetes</p>	<p>N=380</p> <p>Up to 3.5 years</p>	<p>Primary: Blood pressure</p> <p>Secondary: Fasting serum glucose, serum creatinine, plasma insulin, HbA<sub>1c</sub>, TC, HDL-C, TG, fibrinogen, microalbuminuria</p>	<p>Primary: Both treatment groups significantly lowered SBP and DBP from baseline (P&lt;0.05).</p> <p>SBP was lower in the amlodipine group by 4 mm Hg than in the fosinopril group (P&lt;0.01). There was no difference in DBP, both groups decreased by 8 mm Hg.</p> <p>Amlodipine was added by 30.7% of the fosinopril group and fosinopril was added by 26.2% of the amlodipine group (P&gt;0.1).</p> <p>Secondary: No difference between the groups was found for serum creatinine, HbA<sub>1c</sub>, and triglycerides at the endpoint (P&gt;0.05).</p> <p>Fasting serum glucose, serum insulin and microalbuminuria were significantly lower at endpoint for both groups but not significantly different from each other (P&gt;0.05).</p> <p>Total cholesterol increased in both groups, and high-density lipoprotein cholesterol increased significantly in the fosinopril group (P&lt;0.05).</p> <p>No difference in fibrinogen levels was observed between the groups at the end of the trial (P&gt;0.05).</p>
<p>Whelton et al.<sup>108</sup> (1990)</p> <p>Lisinopril 10 to 40 mg QD</p> <p>vs</p> <p>captopril 25 to 100 mg BID</p> <p>Doses were titrated until</p>	<p>DB, MC, PG, RCT</p> <p>Patients with mild-to-moderate essential HTN</p>	<p>N=70</p> <p>Up to 8 weeks</p>	<p>Primary: Reduction in blood pressure in both ambulatory and office settings</p> <p>Secondary: Not reported</p>	<p>Primary: Lisinopril-treated patients showed significantly greater reductions in SBP and DBP measured by 24-hour ambulatory blood pressure monitoring compared to captopril-treated patients (P=0.023 and P=0.007, respectively). Greater reductions (P&lt;0.05) were also noted in patients receiving lisinopril at hours 10 to 12, suggesting two blood pressure troughs for those receiving captopril.</p> <p>The difference in mean reductions between treatment groups from baseline to the final visit approached statistical significance for office SBP (P=0.06) and DBP (P=0.09) in favor of patients receiving lisinopril.</p> <p>Both drugs were well tolerated, and no patients withdrew from either</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
patients responded to treatment (defined by a decrease in office DBP to <90 mm Hg or ≥10 mm Hg decrease from baseline).				treatment group.  Secondary: Not reported
Strasser et al. <sup>109</sup> (2007)  Lisinopril 20 to 40 mg QD  vs  aliskiren 150 to 300 mg QD  HCTZ may be added if additional blood pressure control was required.	AC, DB, DD, MC, PG, RCT  Men and women with uncomplicated severe HTN (mean sitting DBP 105 to 119 mm Hg)	N=183  8 weeks	Primary: Safety  Secondary: Change in mean sitting DBP and SBP, percentage of responders	Primary: Both active treatments were well tolerated with an incidence of adverse events of 32.8% for aliskiren and 29.3% for lisinopril. The proportion of patients discontinuing treatment due to adverse events was 3.2% for aliskiren and 3.4% for lisinopril. The most frequently reported adverse events in both groups were headache, nasopharyngitis and dizziness.  Secondary: Aliskiren showed similar reductions from baseline to lisinopril in mean sitting DBP (-18.5 vs -20.1 mm Hg) and SBP (-20.0 and -22.3 mm Hg).  Responder rates were 81.5% with aliskiren and 87.9% with lisinopril. Approximately half of patients required the addition of HCTZ to achieve blood pressure control (53.6% for aliskiren and 44.8% for lisinopril).
Rosei et al. <sup>110</sup> (2003)  Lisinopril 20 mg QD  vs  nebivolol 5 mg QD	DB, MC, PG, RCT  Patients between 24 and 65 years with mild to moderate uncomplicated essential HTN that was newly diagnosed, or previous antihypertensive therapy was withdrawn at >1	N=65  12 weeks	Primary: Response rates, changes in sitting blood pressure  Secondary: Standing blood pressure, sitting and standing heart rate	Primary: There was not a significant difference in response rates observed between the two treatment groups.  Both treatment groups significantly reduced sitting SBP (P<0.0001) and DBP (P<0.0001) throughout the study compared to baseline but there were no significant differences observed between the treatment groups at most visits, but at week eight, DBP was significantly lower in the nebivolol group compared to the lisinopril group (P<0.05).  Secondary: There was not a significant difference observed between treatment groups in standing blood pressure measurements.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	month before active treatment, and had a sitting DBP of >95 and <114 mm Hg			Both treatment groups significantly reduced sitting heart rate (P<0.01) throughout the study compared to baseline but there were no significant differences observed between the treatment groups at most visits, but at week eight, heart rate were significantly lower in the nebivolol group compared to the lisinopril group (P<0.05).
Wald et al. <sup>111</sup> (2008)  Lisinopril 5mg QD  vs  atenolol 25 mg QD  vs  lisinopril 5 mg and atenolol 25 mg QD  vs  placebo	DB, DD, RCT, XO  Patients ≥ 40 years enrolled in a HTN or anticoagulation clinic	N=47  16 weeks	Primary: Reduction in blood pressure  Secondary: Not reported	Primary: The mean reductions in SBP in the atenolol alone, lisinopril alone and atenolol plus lisinopril groups were 16.1, 12.5, and 22.9 mm Hg, respectively. The mean reductions in DBP in the atenolol alone, lisinopril alone and atenolol plus lisinopril groups were 9.8, 6.8, and 13.9 mm Hg, respectively. The reductions with lisinopril plus atenolol group were significantly higher than either agent as monotherapy (P<0.001).  Secondary: Not reported
Karotsis et al. <sup>112</sup> (2006)  Lisinopril 10 mg QD  vs  chlorthalidone 12.5 mg QD	RCT  Patients 25 to 79 years of age with uncontrolled HTN (average office blood pressure >140/90 mm Hg for all or >153/85 mm Hg for diabetics or patients <65 years of	N=211  8 weeks	Primary: Blood pressure  Secondary: Not reported	Primary: There was a significant decline in both office and home SBP and DBP during the trial with all treatments. The antihypertensive effect was more pronounced and reached significance when home blood pressure monitoring was used in comparison to office blood pressure without the white-coat effect (P<0.001 for all blood pressure changes). With or without the white-coat effect, blood pressure still declined and the differences were significant (P<0.0001 for all blood pressure changes).  Secondary: Not reported



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs felodipine 5 mg QD vs valsartan 80 mg QD All patients also received diltiazem 240 mg QD.	age, confirmed on 2 office visits $\geq 1$ week apart) after $\geq 4$ weeks of OL monotherapy with diltiazem at 240 mg QD			
McInnes et al. <sup>113</sup> (2000) Lisinopril and HCTZ 10-12.5 mg/day (fixed-dose combination product) vs candesartan and HCTZ 8-12.5 mg/day (fixed-dose combination product)	DB, DD, MC, PG, RCT Patients 20 to 80 years of age with mild-to-moderate HTN on prior antihypertensive monotherapy	N=355 26 weeks	Primary: Mean changes in DBP Secondary: Mean changes in SBP and heart rate, proportion of responders and controlled patients, safety	Primary: Changes in mean sitting DBP did not differ significantly between the groups (mean difference, 0.5 mm Hg; P=0.20). Secondary: No significant differences between the groups were reported for mean sitting SBP, heart rate, proportion of responders and controlled patients. Both regimens were well tolerated but a greater percentage of those in the lisinopril based group (80 vs 69%) had a least one side effect (P=0.020). The proportion of patients spontaneously reporting cough (23.1 vs 4.6%) and discontinuing therapy due to adverse events (12.0 vs 5.9%) was also higher in the lisinopril based group compared to the candesartan based group.
Poldermans et al. <sup>114</sup> (2007) Lisinopril 10 to 20 mg QD and HCTZ 12.5 mg QD	AC, DB, MC, PG, RCT Males and females, ages 18 years and older with HTN (mean DBP $\geq 110$ mm Hg and $< 120$	N=130 6 weeks	Primary: Safety/adverse events, vital signs, hematology, biochemistry variables Secondary:	Primary: Both treatments were well tolerated, 26 (40.6%) of patients receiving amlodipine and valsartan and 21 (31.8%) of patients receiving lisinopril and HCTZ reported an adverse events and most were not considered drug related. Peripheral edema was reported more often in the amlodipine and valsartan group than the lisinopril and HCTZ group (7.7 vs 1.5%) and cough was

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>amlodipine 5 to 10 mg QD and valsartan 160 mg QD</p>	<p>mm Hg)</p>		<p>Efficacy (mean DBP, response rate, proportion of patients with mean DBP &lt;90 mm Hg or a <math>\geq</math>10 mm Hg reduction from baseline)</p>	<p>reported less often in the amlodipine and valsartan group than the receiving lisinopril and hydrochlorothiazide group (1.6 vs 3.0%).</p> <p>No difference was found between the treatments in changes in laboratory values or biochemistry variables.</p> <p>Secondary: Both treatments led to a reduction in mean SBP and DBP (P&lt;0.0001 for both from baseline) but were not significantly different from each other. Mean blood pressure for each group at study end: amlodipine and valsartan 135.0/83.6 mm Hg and lisinopril and HCTZ 138.7/85.2 mm Hg.</p> <p>The response rate was similar among the groups (100 vs 95.5%; P value not significant).</p>
<p>Duprez et al.<sup>115</sup> (2010) AGELESS</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>The addition of HCTZ was allowed at week 12 and amlodipine was allowed at week 22 in patients not achieving adequate blood pressure control.</p>	<p>AC, DB, MC, RCT</p> <p>Patients <math>\geq</math>65 years of age with essential HTN (mean sitting SBP <math>\geq</math>140 and &lt;180 mm Hg and mean sitting DBP &lt;110mm Hg)</p>	<p>N=901</p> <p>36 weeks</p>	<p>Primary: Change in mean seated SBP at week 12</p> <p>Secondary: Change in mean sitting SBP at week 36, change in mean sitting DBP at week 12 and week 36, percentage of patients who achieved blood pressure control (mean sitting SBP/DBP &lt;140/90 mm Hg in non-diabetic patients and &lt;130/80 mm Hg in diabetic patients)</p>	<p>Primary: At week 12, aliskiren lowered mean sitting SBP by 14 mm Hg and ramipril decreased mean sitting SBP by 11.6 mm Hg (difference, -2.3 mm Hg; 95% CI, -4.3 to -0.3). Aliskiren monotherapy showed statistically non-inferior (P&lt;0.001) and statistically superior (P=0.02) reductions in mean sitting SBP compared with ramipril monotherapy.</p> <p>Secondary: At week 22, aliskiren decreased mean sitting SBP by 19.6 mm Hg and ramipril decreased mean sitting SBP by 17 mm Hg (difference, -2.4 mm Hg; 95% CI, -4.5 to -0.3; P=0.03).</p> <p>At week 36, aliskiren decreased mean sitting SBP by 20 mm Hg and ramipril decreased mean sitting SBP by 18.1 mm Hg (difference, -1.9 mm Hg; 95% CI, -4.0 to 0.2; P=0.07).</p> <p>At week 12, aliskiren decreased mean sitting DBP by 5.1 mm Hg and ramipril decreased mean sitting DBP by 3.6 mm Hg (difference, -1.5 mm Hg; 95% CI, -2.6 to -0.5; P&lt;0.01).</p> <p>At week 22, aliskiren decreased mean sitting DBP by 8.2 mm Hg and ramipril decreased mean sitting DBP by 7.3 mm Hg (difference, -0.8 mm Hg; 95% CI, -2.0 to 0.3; P=0.14).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>at week 12 and week 36, percentage of patients who required add-on therapy</p>	<p>At week 36, aliskiren decreased mean sitting DBP by 8.2 mm Hg and ramipril decreased mean sitting DBP by 7.0 mm Hg (difference, -1.2 mm Hg; 95% CI, -2.3 to -0.1; P=0.03).</p> <p>The percentage of patients achieving blood pressure control was significantly greater with aliskiren (42%) compared to ramipril (33%) at week 12 (P&lt;0.01). At week 22, a significantly greater proportion of patients achieved blood pressure control with aliskiren (62%) compared to ramipril (50%; P&lt;0.001). At week 36, similar blood pressure control rates were achieved with aliskiren (59%) and ramipril (51%; P=0.01).</p> <p>By week 36, a significantly greater percentage of patients receiving ramipril compared to aliskiren required additional HCTZ (56 vs 46%; P&lt;0.01).</p> <p>By week 36, a greater percentage of patients receiving ramipril (16%) compared to aliskiren (12%) required add-on therapy with both HCTZ and amlodipine (P=0.048).</p> <p>More patients receiving aliskiren were receiving monotherapy (42%) than patients receiving ramipril (29%) at week 36.</p>
<p>Anderson et al.<sup>116</sup> (2008)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>The addition of HCTZ was allowed in patients not</p>	<p>AC, DB, MC, PC, RCT</p> <p>Men and women ≥18 years with essential HTN (mean sitting DBP 90 to 109 mm Hg)</p>	<p>N=842</p> <p>26 weeks</p>	<p>Primary: Change in mean sitting DBP at week 26</p> <p>Secondary: Change in mean sitting SBP at week 26, change in mean sitting SBP and DBP at week 6 and 12 (comparing aliskiren and ramipril monotherapy),</p>	<p>Primary: Reductions in mean sitting DBP at week 26 were significantly greater with aliskiren-based therapies (-13.2 mm Hg) compared to ramipril-based therapies (-12.0 mm Hg; P=0.0250).</p> <p>Secondary: Reductions in mean sitting SBP at week 26 were significantly greater with aliskiren-based therapies (-17.9 mm Hg) compared to ramipril-based therapies (-15.2 mm Hg; P=0.0036).</p> <p>Mean changes in sitting SBP were significantly greater with aliskiren (-12.9 and -14.0 mm Hg, respectively) compared to ramipril (-10.5 and -11.3, respectively) at weeks six and 12 (P=0.0041 and P=0.0027, respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>achieving adequate blood pressure control.</p> <p>The study did not specifically analyze the effects of HCTZ on either treatment regimen.</p>			<p>proportion achieving blood pressure control (&lt;140/90 mm Hg), proportion achieving SBP control (&lt;140 mm Hg), safety</p>	<p>Mean changes in sitting DBP were not significantly greater with aliskiren (-10.5 and -11.3 mm Hg, respectively) compared to ramipril (-9.5 and -9.7, respectively) at week six, but were significantly greater at week 12 (P=0.0689 and P=0.0056, respectively).</p> <p>The proportion of patients achieving overall blood pressure control (&lt;140/90 mm Hg) was significantly higher with aliskiren-based therapy (61.4%) compared to ramipril-based therapy (53.1%; P=0.0205) at week 26. Also, the proportion of patients achieving SBP control (&lt;140 mm Hg) was significantly higher with aliskiren-based therapy (72.5%) compared to ramipril-based therapy (64.1%; P=0.0075) at week 26.</p> <p>The majority of adverse events reported during the active treatment period were mild or moderate in intensity and transient. Most events occurred at a similar incidence in the two groups with the exception of cough which was considered treatment-related in 5.5% of patients receiving ramipril vs 2.1% of patients receiving aliskiren.</p>
<p>Miranda et al.<sup>117</sup> (2008)</p> <p>Ramipril 2.5 to 10 mg QD and amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Adults 40 to 79 years of age with stage 1 or 2 essential HTN</p>	<p>N=222</p> <p>18 weeks</p>	<p>Primary: Change in SBP and DBP</p> <p>Secondary: Safety and tolerability</p>	<p>Primary: The mean changes in ambulatory BP were greater with amlodipine and ramipril compared to amlodipine monotherapy (SBP, -20.21 vs -15.31 mm Hg and DBP, -11.61 vs -8.42 mm Hg, respectively; both, P=0.002]. There was no significant difference among the treatment groups in office BP (SBP, -26.60 vs -22.97 mm Hg and DBP, -16.48 vs -14.48 mm Hg; both, P value not significant).</p> <p>Secondary: Twenty-nine patients (22.1%) treated with combination therapy and 41 patients (30.6%) treated with monotherapy experienced ≥1 adverse event considered possibly related to study drug. The combination-therapy group had lower prevalence of edema (7.6 vs 18.7%; P=0.011) and a similar prevalence of dry cough (3.8 vs 0.8%; P value not significant).</p>
<p>Bönner et al.<sup>118</sup> (2013)</p> <p>Azilsartan (AZL) 20mg titrated to 40 mg</p>	<p>DB, RCT</p> <p>Patients ≥18 years of age with clinic systolic blood pressure (SBP) 150</p>	<p>N=884</p> <p>24 weeks</p>	<p>Primary: Change in trough, seated clinic SBP</p> <p>Secondary: Change from</p>	<p>Primary: After 24 weeks of treatment, trough, sitting, clinic SBP decreased significantly in all the groups. The changes from baseline were significantly greater for the AZL 40 and 80 mg treatment groups (-20.6±0.95 and -21.2±0.95 mm Hg, respectively) than for RAM 10 mg (-12.2±0.95 mm Hg). The differences between the AZL-treated subjects</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>azilsartan (AZL) 20mg titrated to 80 mg</p> <p>vs</p> <p>ramipril (RAM) 2.5 mg titrated to 10 mg</p>	<p>to 180 mm Hg</p>		<p>baseline to week 24 in trough, seated clinic DBP, measures of ambulatory BP, and BP response rates</p>	<p>and the RAM-treated subjects were -8.4 mm Hg for AZL 40 and -9.0 mm Hg for AZL 80 (P&lt;0.001 for both comparisons).</p> <p>Secondary: Change in trough, sitting, DBP was -10.2±0.55 mm Hg in the AZL 40 mg group, -10.5±0.55 mm Hg in the AZL 80 mg and -4.9±0.56 mm Hg in the RAM 10 mg group.</p> <p>AZL 40 and 80 mg reduced ambulatory SBP and DBP significantly more than RAM for all ABPM time intervals evaluated, including 24-hour mean, mean daytime, mean nighttime and mean trough pressure.</p> <p>The differences between the AZL and RAM groups proportion of subjects achieving SBP and DBP response criteria were highly significant (P&lt;0.001). More subjects achieved a reduction in clinic BP to &lt;140/90 mm Hg and/or a reduction in BP≥20/10 mm Hg at week 24 following treatment with AZL compared with RAM (54.0% and 53.6% for AZL 40 and 80 mg vs 33.8% with RAM 10 mg, respectively; P&lt;0.001).</p>
<p>Williams et al.<sup>119</sup> (2009) PRISMA I and PRISMA II</p> <p>Ramipril 2.5 mg QD for 2 weeks then force titration to 5 mg QD for 6 weeks then 10 mg QD for 6 weeks</p> <p>vs</p> <p>telmisartan 40 mg QD for 2 weeks then force titration to 80 mg QD for 12 weeks</p>	<p>Pooled analysis: blinded endpoint, OL, PRO, RCT</p> <p>Patients ≥18 years of age with mild- to moderate HTN</p>	<p>N=1,613</p> <p>14 weeks</p>	<p>Primary: Change from baseline in mean ambulatory BP during the final 6 hours of the 24- hour dosing interval</p> <p>Secondary: Change from baseline in mean ambulatory blood pressure during the 24-hour dosing interval, morning, daytime and nighttime ambulatory blood</p>	<p>Primary: A significantly greater reduction in mean ambulatory blood pressure during the last six hours of the 24-hour dosing interval was observed with telmisartan 80 mg group compared to ramipril 5 and 10 mg (P&lt;0.0001).</p> <p>Secondary: Significantly greater reductions in mean 24-hour, morning, daytime, nighttime and 24-hour blood pressure load were observed with telmisartan 80 mg compared to ramipril 5 and 10 mg (P&lt;0.0001).</p> <p>Significantly greater reductions in treatment response and blood pressure control rates were observed with telmisartan 80 mg compared to ramipril 5 and 10 mg (P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			pressure, 24-hour blood pressure load, treatment response, blood pressure control	
<p>O'Brien et al.<sup>120</sup> (2007)</p> <p>Aliskiren 150 mg QD for 3 weeks, then HCTZ 25 mg QD was added for an additional 3 weeks (if ABPM remained <math>\geq 135/85</math> mm Hg)</p> <p>vs</p> <p>irbesartan 150 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p> <p>vs</p> <p>ramipril 5 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p>	<p>3 OL studies</p> <p>Men and women 18 to 80 years with ambulatory SBP <math>\geq 140</math> and <math>\leq 180</math> mm Hg without treatment</p>	<p>N=67</p> <p>6 to 9 weeks</p>	<p>Primary: Change in daytime systolic ABPM with combination therapy compared with monotherapy</p> <p>Secondary: Change in daytime diastolic ABPM, nighttime systolic and diastolic ABPM, daytime and nighttime heart rates, plasma renin activity</p>	<p>Primary: Aliskiren coadministered with HCTZ (P=0.0007) or ramipril (P=0.03) led to significantly greater reductions in daytime systolic ABPM compared to monotherapy. There was a trend for a reduction in daytime systolic ABPM with the addition of aliskiren to irbesartan; however, this trend was not statistically significant.</p> <p>Secondary: Aliskiren plus HCTZ significantly lowered daytime diastolic ABPM compared to aliskiren monotherapy (P=0.0006). Changes in nighttime systolic and diastolic ABPM followed similar trends but did not achieve statistical significance (P=0.06 and P=0.09, respectively). No changes in heart rate were observed with either aliskiren regimen.</p> <p>Aliskiren added to irbesartan did not significantly change diastolic ABPM compared to irbesartan monotherapy; however, nighttime systolic and diastolic ABPM were significantly reduced (P&lt;0.05 for all). No changes in heart rate were observed with either irbesartan regimen.</p> <p>Mean diastolic ABPM was significantly decreased with the addition of aliskiren 150 mg (P&lt;0.05) but not aliskiren 75 mg to ramipril monotherapy. Both aliskiren doses significantly decreased nighttime systolic and diastolic ABPM (P&lt;0.05 for all). No changes in heart rate were observed with either ramipril regimen.</p> <p>Aliskiren alone significantly inhibited plasma renin activity by 65% (P&lt;0.0001), while ramipril and irbesartan monotherapy increased renin activity by 90 and 175%, respectively. When aliskiren was coadministered with HCTZ, ramipril or irbesartan, plasma renin activity remained similar to baseline levels or decreased.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
weeks				
Tytus et al. <sup>121</sup> (2007)  Trandolapril 1 to 4 mg/day  At 14 weeks after treatment initiation, subjects not achieving blood pressure targets could receive a combination of trandolapril 4 mg/day plus verapamil 240 mg/day with or without a diuretic.	MC, OL, PRO  Patients with stage 1 or 2 HTN who were treatment naïve (82%) or uncontrolled on a diuretic (11%) or calcium-channel blocker (7%); uncontrolled HTN was defined as $\geq 140/90$ mm Hg in subjects with no other risk factors or $\geq 130/80$ mm Hg in subjects with diabetes or kidney disease	N=1,683  26 weeks	Primary: Percentage of patients reaching target blood pressure at 14 weeks  Secondary: Percentages of subjects with stage 1 and 2 HTN who achieved target blood pressure, percentages of subjects who achieved a drop in SBP of $\geq 20$ mm Hg and/or DBP $\geq 10$ mm Hg, absolute changes in SBP and DBP, adverse events	Primary: At 14 weeks of treatment, 71.2% of patients who were treated with trandolapril monotherapy reached SBP/DBP $< 140/90$ mm Hg.  Secondary: At 26 weeks, 73.4% of patients achieved a target level of SBP/DBP $< 140/90$ mm Hg. Of the 683 subjects with stage 2 HTN, 64.6% achieved the target level after 14 weeks of trandolapril and 67.9% after 26 weeks.  At 14 weeks, 78.8% of subjects treated with a trandolapril regimen experienced a decrease in SBP of $\geq 20$ mm Hg or a decrease in DBP of $\geq 10$ mm Hg.  Statistically significant ( $P < 0.001$ ) and clinically relevant mean decreases in SBP of -16.1 mm Hg and in DBP of -8.8 mm Hg were observed from four weeks of treatment onward for the overall study population. The mean reductions in SBP and DBP were -21.5 and -11.9 mm Hg, respectively at 14 weeks ( $P < 0.001$ ), and -22.4 and -12.7 mm Hg, respectively, at 26 weeks ( $P < 0.001$ ).  A total of 343 predominantly mild, nonserious adverse events were attributed to the study drugs, reported by 15.3% of the 1,650 subjects. The most frequently reported nonserious adverse events were cough (6.3%); gastrointestinal disorders (2.3%), predominantly nausea; and headache (2.1%). No serious adverse events were attributed to the study treatment.
Tytus et al. <sup>122</sup> (2011) MAVIKtory  Trandolapril 1 to 2 mg/day  With or without existing antihypertensive therapy.	MC, OS  Patients with HTN	N=8,787  6 months	Primary: Proportion of patients reaching blood pressure targets, safety  Secondary: Not reported	Primary: The target of $< 140/90$ mm Hg was achieved by 67.3% of patients. The lower mean target of 133.4/83.3 mm Hg for nondiabetic patients and 128.6/79.3 mm Hg for diabetic patients were achieved by 52.2%. Mean reductions from baseline to trial end were 19.4 mm Hg (95% CI, -19.9 to -19.0) in SBP and 10.1 mm Hg (95% CI, -10.4 to -9.8) in DBP.  Cough was the most commonly reported adverse event (4.2%).  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Pauly et al.<sup>123</sup> (1994)</p> <p>Trandolapril 4 mg QD</p> <p>vs</p> <p>captopril 50 mg BID</p> <p>If blood pressure was not normalized at 8 weeks, HCTZ 25 mg was added.</p>	<p>DB, MC, RCT</p> <p>Patients between 21 to 65 years with mild-to-moderate essential HTN (DBP of 95 to 115 mm Hg)</p>	<p>N=180</p> <p>16 weeks</p>	<p>Primary: Morning pre-dosing supine DBP at 8 weeks of monotherapy</p> <p>Secondary: Supine SBP at 8 weeks of monotherapy, blood pressure at 16 weeks of therapy (including 8 weeks of monotherapy and 8 weeks of combination therapy with HCTZ)</p>	<p>Primary: Significantly greater mean reductions in supine DBP in the trandolapril group vs captopril group were observed after eight weeks of monotherapy (-13.5 vs -10.1 mm Hg; P=0.007).</p> <p>Secondary: Differences in supine SBP between treatment groups approached significance after eight weeks of monotherapy (P=0.06).</p> <p>Both SBP and DBP were significantly reduced at all time points compared to baseline for both treatment groups at the end of the study (P&lt;0.05).</p> <p>The proportion of patients whose blood pressure normalized (supine and standing blood pressure ≤160/90 mm Hg) at the end of the study was 61% for trandolapril and 44% for captopril (P=0.02).</p> <p>The overall proportion of responders (DBP fell by ≥10 or to &lt;90 mm Hg) was significantly greater in the trandolapril group (77%) than in the captopril group (58%; P&lt;0.007).</p>
<p>Vaur et al.<sup>124</sup> (1995)</p> <p>Trandolapril 2 mg QD in the morning</p> <p>vs</p> <p>enalapril 20 mg QD in the morning</p>	<p>DB, RCT</p> <p>Patients between 18 to 70 years with mild-to-moderate primary HTN</p>	<p>N=88</p> <p>3 weeks</p>	<p>Primary: 24-hour ambulatory SBP and DBP over an active 24-hour period and subsequent 24-hour period (to mimic a missed dose)</p> <p>Secondary: Not reported</p>	<p>Primary: Both trandolapril and enalapril showed similar reductions in SBP and DBP over the 24-hour period. In the trandolapril group, SBP and DBP decreased from 148/92 to 135/83 mm Hg (P&lt;0.001). In the enalapril group, SBP and DBP decreased from 143/91 to 133/83 mm Hg (P&lt;0.001).</p> <p>The trough/peak ratio on active treatment was 90% (SBP) and 54% (DBP) in the trandolapril group and 49% (SBP and DBP) in the enalapril group. Following the missed dose, trough/peak ratio decreased to 58% (SBP)/36% (DBP) for trandolapril and 10% (SBP)/19% (DBP) for enalapril. The blood pressure control was better sustained with trandolapril, such that significant falls in blood pressure were observed during the daytime, nighttime and early morning periods after a missed dose, whereas during the same periods, enalapril only significantly reduced blood pressure in the daytime period.</p> <p>Secondary: Not reported</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Karlberg et al.<sup>125</sup> (2000)</p> <p>Trandolapril 2 mg/day</p> <p>vs</p> <p>verapamil 240 mg/day</p> <p>vs</p> <p>trandolapril and verapamil 2-180 mg/day (fixed-dose combination)</p>	<p>DB, MC, PRO, RCT, XO</p> <p>Patients with uncomplicated primary HTN (sitting DBP between 95 and 115 mm Hg) between the ages of 20 to 80 years</p>	<p>N=226</p> <p>2 months</p>	<p>Primary: Change in blood pressure and rate pressure product</p> <p>Secondary: Predictive value of plasma concentrations of active renin regarding the blood pressure response to the different treatment regimens, safety</p>	<p>Primary: The mean fall in blood pressure was significantly greater with the combination (20/15 mm Hg; P&lt;0.00054), as compared to trandolapril (14/11 mm Hg) or verapamil (13/11) mm Hg. The difference between verapamil and trandolapril was not significant.</p> <p>Rate pressure product decreased significantly more on the combination (P&lt;0.001) than on trandolapril or verapamil alone.</p> <p>Secondary: There was a significant positive correlation between blood pressure fall and plasma concentrations of active renin (e.g., the higher the initial active renin, the better the blood pressure response to trandolapril [P&lt;0.045 for SBP and P&lt;0.004 for DBP]). No relationships were found for either verapamil or the combination.</p> <p>All treatments were well tolerated and safe.</p>
<p>Pepine et al.<sup>126</sup> (2006)</p> <p>INVEST</p> <p>Verapamil SR (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p> <p>vs</p> <p>atenolol (step 1), then add HCTZ if</p>	<p>Post hoc analysis of INVEST</p> <p>Patients with essential HTN</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Risk for adverse outcome associated with baseline factors, follow-up blood pressure and drug treatments</p> <p>Secondary: Not reported</p>	<p>Primary: Previous heart failure (adjusted HR, 1.96), as well as diabetes (HR, 1.77), increased age (HR, 1.63), United States residency (HR, 1.61), renal impairment (HR, 1.50), stroke/TIA (HR, 1.43), smoking (HR, 1.41), MI (HR, 1.34), PVD (HR, 1.27), and revascularization (HR, 1.15) predicted increased risk.</p> <p>Follow-up SBP &lt;140 mm Hg (HR, 0.82) or DBP &lt;90 mm Hg (HR, 0.70) and trandolapril with verapamil SR (HR, 0.78 and 0.79) were associated with reduced risk.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)				
<p>Brunner et al.<sup>127</sup> (2007) INVEST</p> <p>Verapamil SR 240 mg and trandolapril 1 to 4 mg</p>	<p>Post hoc analysis of INVEST</p> <p>Patients with essential HTN</p>	<p>N=1,832</p> <p>24 months</p>	<p>Primary: Factors influencing blood pressure response to trandolapril add-on therapy</p> <p>Secondary: Not reported</p>	<p>Primary: Trandolapril decreased mean unadjusted SBP and DBP by -9.1 and -4.1 mm Hg, respectively. The percentage of patients with blood pressure under control (&lt;140/90 mm Hg) increased from 6.7 to 41.3% (P&lt;0.0001).</p> <p>Adjusted blood pressure response was significantly associated with age and baseline SBP and DBP (P&lt;0.0001). Whereas the decrease in SBP was more pronounced in younger patients, the opposite was observed for DBP decrease.</p> <p>DBP response was significantly associated with race. Specifically, the adjusted DBP decrease was significantly smaller in Hispanics and African Americans than whites (P=0.0032 and P=0.0069, respectively). However, Hispanics achieved a decrease in SBP and an increase in blood pressure control similar to the other ethnic groups.</p> <p>Secondary: Not reported</p>
<p>Cifkova et al.<sup>128</sup> (2000)</p> <p>Verapamil and trandolapril 180-2 mg QD (fixed-dose combination) (VT)</p> <p>vs</p>	<p>AC, OL, RCT, XO</p> <p>Caucasian patients aged 18 to 75 years with mild-to-moderate essential HTN (SBP 140 to 209 mm Hg and DBP 90 to 119 mm Hg)</p>	<p>N=100</p> <p>8 months</p>	<p>Primary: LDL-C</p> <p>Secondary: Other lipid parameters (HDL-C, TC, TG, apolipoproteins AI and B, lipoprotein(a)), blood pressure</p>	<p>Primary: LDL-C was not significantly different between the two treatment groups (P=0.909).</p> <p>Secondary: All secondary lipid parameters remained unaltered except for HDL-C which was significantly higher with VT (1.39 vs 1.35 mmol/L; P&lt;0.03).</p> <p>Serum potassium declined while uric acid and glucose increased on CH (P&lt;0.001 for all).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>captopril and HCTZ 50-25 mg QD (fixed-dose combination) (CH)</p> <p>After 16 weeks, patients were switched to the other fixed combination for an additional 16 weeks.</p>			parameters	<p>While there were no significant differences with respect to adjusted mean DBP, adjusted mean SBP was slightly higher on treatment with VT than with CH. These differences reached statistical significance for the 24-hour and night-time means, although the absolute adjusted mean treatment differences were only 2.3 mm Hg (P=0.02) and 3.5 mm Hg (P=0.01), respectively. The number of patients who achieved DBP &lt;90 mm Hg at the end of each treatment did not differ (56% VT vs 46% CH; P value not significant). Heart rate was significantly lower in the VT group than the CH group (treatment differences ranged from 2.8 to 4.5 bpm; P≤0.001 for all).</p>
<p>de Leeuw et al.<sup>129</sup> (1997)</p> <p>Verapamil SR and trandolapril 180-2 mg/day, atenolol and chlorthalidone 100-25 mg/day, or lisinopril and HCTZ 20-12.5 mg/day (fixed-dose combination products)</p> <p>vs</p> <p>placebo</p> <p>All patients entered a SB, placebo 4 week run in period.</p>	<p>DB, MC, PC, RCT</p> <p>Patients 18 to 70 years of age with essential HTN (WHO I or II) newly or unsuccessfully treated, with supine DBP 101 to 114 mm Hg in week 4 of the run in period</p>	<p>N=205</p> <p>12 weeks</p>	<p>Primary: Changes in supine blood pressure, standing blood pressure response rates, normalization rates</p> <p>Secondary: Not reported</p>	<p>Primary: Each of the three treatments was significantly more effective than placebo in reducing seated DBP. Changes in DBP were as follows: verapamil SR and trandolapril, -13 (95% CI, -16 to -9); atenolol and chlorthalidone, -13 (95% CI, -16 to -9); lisinopril and HCTZ, -12 (95% CI, -15 to -9) and placebo, -3 (95% CI, -7 to 0) (P=0.0001 for all vs placebo), but there was not a significance among the treatments (P values not reported).</p> <p>Each of the three treatments was significantly more effective than placebo in reducing seated SBP. Changes in SBP were as follows: verapamil SR and trandolapril, -27 (95% CI, -33 to -21); atenolol and chlorthalidone, -28 (95% CI, -34 to -22); lisinopril and HCTZ, -23 (95% CI, -29 to -17) and placebo, -3 (95% CI, -9 to 3) (P=0.0001 for all vs placebo), but there was not a significance among the treatments (P values not reported).</p> <p>Effects on standing blood pressure demonstrated similar results as the effects on sitting blood pressure (P values not reported).</p> <p>Normalization of DBP (&lt;90 mm Hg), corrected for placebo, were significantly higher with all treatments compared to placebo (verapamil SR and trandolapril, 33% [95% CI, 16 to 50; P&lt;0.0005]; atenolol and chlorthalidone, 31% [95% CI, 14 to 48; P&lt;0.002] and lisinopril and HCTZ, 25% [95% CI, 9 to 42; P&lt;0.005]).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Response rates (normalization of DBP or a reduction in DBP &gt;10 mm Hg), corrected for placebo, were significantly higher with all treatments compared to placebo (verapamil SR and trandolapril, 40% [95% CI, 22 to 58; P&lt;0.0001], atenolol and chlorthalidone, 44% [95% CI, 27 to 61; P&lt;0.0001] and lisinopril and HCTZ, 37% [95% CI, 19 to 55; P&lt;0.0002]).</p> <p>Secondary: Not reported</p>
<p>Stanton et al.<sup>130</sup> (2010)</p> <p>Aliskiren 300 mg QD</p> <p>vs</p> <p>irbesartan, losartan, valsartan, ramipril, HCTZ, placebo</p>	<p>MA</p> <p>Adults with mild to moderate essential HTN</p>	<p>N=4,877 (8 trials)</p> <p>4 to 12 weeks</p>	<p>Primary: Paradoxical blood pressure rises, as well as the percentage of patients with SBP increases (&gt;10 or &gt;20 mm Hg) or DBP increases (&gt;5 or &gt;10 mm Hg) from baseline</p> <p>Secondary: Not reported</p>	<p>Primary: There were no significant differences among the pooled aliskiren, irbesartan, losartan, valsartan, ramipril, and HCTZ groups in the incidence of SBP increases &gt;10 mm Hg (P=0.30) and &gt;20 mm Hg (P=0.28) or DBP increases &gt;5 mm Hg (P=0.65) and &gt;10 mm Hg (P=0.5).</p> <p>Increases in SBP and DBP occurred significantly more frequently in the pooled placebo group than the aliskiren group (P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Van Bortel et al.<sup>131</sup> (2008)</p> <p>ACE inhibitor, ARB, β-blocker, calcium channel blocker, or placebo</p> <p>vs</p> <p>nebivolol</p>	<p>MA</p> <p>12 RCTs involving &gt;25 patients with essential HTN where nebivolol 5 mg QD was compared to placebo or other active drugs for &gt;1 month</p>	<p>N=2,653</p> <p>Duration varied</p>	<p>Primary: Antihypertensive effect and tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: Overall, higher response rates were observed with nebivolol than all other antihypertensive agents combined (OR, 1.41; 95% CI, 1.15 to 1.73; P=0.001) and compared to the ACE inhibitors (OR, 1.92; 1.30 to 2.85; P=0.001), but response rates to nebivolol were similar to β-blockers (OR, 1.29; 95% CI, 0.81 to 2.04; P=0.283), calcium channel blockers (OR, 1.19; 95% CI, 0.83 to 1.70; P=0.350) and losartan (OR, 1.35; 95% CI, 0.84 to 2.15; P=0.212).</p> <p>Overall, a higher percentage of patients obtained normalized blood pressure with nebivolol compared to the other antihypertensive agents combined (OR, 1.35; 95% CI, 1.07 to 1.72; P=0.012). A higher percentage of patient receiving nebivolol obtained normalized blood pressure compared to losartan (OR, 1.98; 95% CI, 1.24 to 3.15; P=0.004) and calcium channel blockers (OR, 1.96; 95% CI, 1.05 to 1.96; P=0.024), but</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>not when compared to other <math>\beta</math>-blockers (OR, 1.29; 95% CI, 0.81 to 1.65; P=0.473).</p> <p>Overall, the percentage of adverse events was significantly lower with nebivolol compared to the other antihypertensive agents combined (OR, 0.59; 95% CI, 0.48 to 0.72; P&lt;0.001) and similar to placebo (OR, 1.16; 95% CI, 0.76 to 1.67; P=0.482). In comparing nebivolol to the individual treatments, nebivolol had a lower percentage of adverse events compared to losartan (OR, 0.52; 95% CI, 0.30 to 0.89; P=0.016), the other <math>\beta</math>-blockers (OR, 0.56; 95% CI, 0.36 to 0.85; P=0.007) and calcium channel blockers (OR, 0.49; 95% CI 0.33 to 0.72; P&lt;0.001), but was similar to ACE inhibitors (OR, 0.75; 95% CI 0.52 to 1.08).</p> <p>Secondary: Not reported</p>
<p>Baguet et al.<sup>132</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the <math>\beta</math>-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).</p> <p>The weighted average reduction of SBP and DBP for each drug class were as follows: Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively. <math>\beta</math>-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively. Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.            ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively.            Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary:            Not reported</p>
<b>Diabetes/Diabetic Nephropathy/Renal Dysfunction</b>				
<p>Bakris et al.<sup>133</sup> (2010)            ACCOMPLISH</p> <p>Benazepril and amlodipine 40-5 to 40-10 mg/day, followed by forced titration after 1 month on benazepril and amlodipine 20-5 mg (fixed-dose combination product)</p> <p>vs</p> <p>benazepril and</p>	<p>Prespecified subanalysis of ACCOMPISH</p> <p>Men and women &gt;60 years of age with HTN and at high risk for cardiovascular events (history of coronary events, MI, revascularization, or stroke; impaired renal function; PAD, left ventricular hypertrophy; or diabetes)</p>	<p>N=11,482</p> <p>2.9 years (mean duration)</p>	<p>Primary:            Time to first event of doubling of serum creatinine concentration or end stage renal disease (defined as eGFR &lt;15 mL/min/1.73 m<sup>2</sup> or need for chronic dialysis)</p> <p>Secondary:            Progression of chronic kidney disease plus death, change in albuminuria, and change in eGFR</p>	<p>Primary:            There were fewer chronic kidney disease events in the benazepril and amlodipine group (2.0% of patients) compared to the benazepril and HCTZ group (3.7%; HR, 0.52; 95% CI, 0.41 to 0.65; P&lt;0.0001).</p> <p>Secondary:            The composite endpoint of progression of chronic kidney disease and all-cause mortality was lower in the benazepril and amlodipine group (6.0%) compared to the benazepril and HCTZ group (8.1%; HR, 0.73; 95% CI, 0.64 to 0.84; P&lt;0.0001). There was a slower decline in eGFR in the benazepril and amlodipine group compared to the benazepril and HCTZ group (-0.88 vs -4.22 mL/min/1.73 m<sup>2</sup>; P=0.01). Of the patients with baseline microalbuminuria, there was a reduction in the urinary albumin:creatinine in the benazepril and HCTZ group of -63.8% (median change) compared to a median change of -29.0% in the benazepril and amlodipine group (P&lt;0.0001).</p> <p>There was a higher percentage of patients reporting peripheral edema in the benazepril and amlodipine group compared to the benazepril and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 40-12.5 to 40-25 mg/day, followed by forced titration after 1 month on benazepril and HCTZ 20-12.5 mg (fixed-dose combination product)</p>				<p>HCTZ group (P&lt;0.0001).</p>
<p>Hou et al.<sup>134</sup> (2007) ROAD</p> <p>Benazepril 10 mg/day vs individual up-titration (10 to 40 mg/day with median dose of 20 mg/day)</p> <p>or</p> <p>losartan 50 mg/day vs individual up-titration (50 to 200 mg/day with median dose of 100 mg/day)</p> <p>Up-titration was performed to optimal antiproteinuric and tolerated</p>	<p>OL, PRO, RCT</p> <p>Patients aged 18 to 70 years with proteinuria and chronic renal insufficiency who did not have diabetes</p>	<p>N=360</p> <p>3.7 years (median follow-up)</p>	<p>Primary: Time to composite of doubling of serum creatinine, ESRD or death</p> <p>Secondary: Changes in level of proteinuria, rate of progression of renal disease</p>	<p>Primary: Compared to the conventional dosages, optimal antiproteinuric dosages of benazepril and losartan that were achieved through up-titration were associated with a 51 and 53% reduction in the risk for the primary end point (P=0.028 and P=0.022, respectively).</p> <p>There was no statistically significant difference between benazepril and losartan in the overall relative risk reduction at their respective optimal antiproteinuric dosages or at conventional dosages.</p> <p>Secondary: Optimal antiproteinuric dosages of benazepril and losartan at comparable blood pressure control, achieved a greater reduction in both proteinuria and the rate of decline in renal function compared to their conventional dosages.</p> <p>There was no significant difference in proteinuria reduction between benazepril and losartan at both conventional and optimal antiproteinuric dosages. Changes in renal function were similar between benazepril and losartan arms at both conventional and optimal antiproteinuric doses (P&gt;0.05).</p> <p>There was no significant difference for the overall incidence of major adverse events between groups that were given conventional and optimal dosages in any of the treatment arms.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
dosages, and then these dosages were maintained.				
<p>Bakris et al.<sup>135</sup> (2008) GUARD</p> <p>Benazepril and HCTZ (fixed-dose combination)</p> <p>vs</p> <p>amlodipine and benazepril (fixed-dose combination)</p>	<p>DB, RCT</p> <p>Hypertensive, albuminuric type 2 diabetic patients, mean age 58 years were randomized to receive either initial fixed-dose combination product</p>	<p>N=322</p> <p>52 weeks</p>	<p>Primary: Change in urinary albumin to creatinine ratio after 1 year of initial treatment with either fixed-dose combination, blood pressure reductions</p> <p>Secondary: Proportion who progressed to overt diabetic nephropathy, safety</p>	<p>Primary: Both combinations significantly reduced the urinary albumin to creatinine ratio compared to baseline (P&lt;0.0001). The median percent change was -72.1% for benazepril and HCTZ and -40.5% for amlodipine and benazepril (P&lt;0.0001).</p> <p>Both regimens significantly reduced SBP and DBP compared to baseline (P&lt;0.0001). The mean reduction in both SBP and DBP was greater in the amlodipine-based arm than in the HCTZ-based arm; however, significance in favor of the amlodipine regimen was observed only for DBP (SBP, -20.5 vs -18.8; P=0.19; DPB, -13.1 vs -9.97; P=0.02).</p> <p>A greater proportion of patients who had microalbuminuria at baseline and treated with benazepril and HCTZ compared to amlodipine and benazepril attained normalization of the urinary albumin to creatinine ratio, defined as &lt;30 mg/g (69.2 vs 47.8%; P=0.0004).</p> <p>Secondary: The percentage of patients progressing to overt proteinuria was similar for both groups.</p> <p>Overall, both study drugs were well tolerated. Adverse reactions possibly related to the study medications occurred in 11.4 and 3.6% of patients receiving amlodipine and benazepril and benazepril and HCTZ, respectively. They included peripheral edema (7.8 vs 2.4%, respectively), fatigue (1.2% in each group), pitting edema (1.2 vs 0.0%), face edema (0.6 vs 0.0%) and thirst (0.6 vs 0.0%). More patients receiving the HCTZ-based regimen (10.8%) discontinued study drug than with the amlodipine-based regimen due to side effects (5.4%).</p>
<p>Esnault et al.<sup>136</sup> (2008)</p> <p>Enalapril 5 to 20 mg/day</p>	<p>MC, DB, PC, RCT</p> <p>Nondiabetic, adult patients with estimated creatinine</p>	<p>N=263</p> <p>3 years</p>	<p>Primary: Change in GFR measured yearly by blood clearance</p>	<p>Primary: No statistically significant difference was found between amlodipine and enalapril in GFR decline (-4.92 and -3.98 mL/min., respectively, at last observation).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs amlodipine 5 to 10 mg QD	clearance of 20 to 60 ml/min		Secondary: Composite of renal events and tolerability	Secondary: No statistically significant difference was found between amlodipine and enalapril in the composite secondary end point after a median follow-up of 2.9 years, including in the subgroup of patients with proteinuria >1 g/d at baseline.
Barnett et al. <sup>137</sup> (2004) DETAIL Enalapril 20 mg/day vs telmisartan 80 mg/day	DB, MC, PG, RCT  Patients aged 35 to 80 years with type 2 diabetes and HTN	N=250  5 years	Primary: Change in the GFR  Secondary: Annual changes in GFR, serum creatinine level, urinary albumin excretion, and blood pressure; rates of ESRD and cardiovascular events; all-cause mortality	Primary: After five years, GFR decreased by 17.9 mL/minute/1.73 m <sup>2</sup> with telmisartan compared to 14.9 mL/min/1.73 m <sup>2</sup> with enalapril (mean difference, -3.0 mL/min/1.73 m <sup>2</sup> ; 95% CI, -7.6 to 1.6). Therefore, the changes in GFR were comparable between the groups.  Secondary: The effects of the two agents on the secondary end points were not significantly different after five years.
Mogensen et al. <sup>138</sup> (2000) CALM Lisinopril 20 mg QD vs candesartan 16 mg QD vs lisinopril 20 mg QD plus candesartan 16 mg QD	DB, DD, MC, PG, RCT  Patients 30 to 75 years old with HTN, type 2 diabetes, and microalbuminuria	N=199  24 weeks	Primary: Blood pressure and urinary albumin:creatinine ratio  Secondary: Not reported	Primary: At 12 weeks, mean reductions in DBP were 9.7 mm Hg (P<0.001) and 9.5 mm Hg (P<0.001), respectively, and in urinary albumin:creatinine ratio were 46% (P<0.001) and 30% (P<0.001) for lisinopril and candesartan, respectively.  Compared to either agent alone, at 24 weeks the combination of lisinopril plus candesartan resulted in 16.3 mm Hg reduction in mean DBP vs 10.4 mm Hg for candesartan alone (P<0.001) and 10.7 mm Hg for lisinopril alone (P<0.001).  The reduction in urinary albumin:creatinine ratio with combination treatment (50%) was greater than with lisinopril alone (39%; P<0.001) and candesartan alone (24%; P=0.05).  All treatments were generally well tolerated.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Patients received 12 weeks monotherapy followed by an additional 12 weeks of monotherapy or combination therapy.</p>				<p>Not reported</p>
<p>Fried et al.<sup>139</sup> (2013) VA NEPHRON-D</p> <p>Losartan with lisinopril</p> <p>vs</p> <p>losartan alone</p>	<p>DB, MA, RCT</p> <p>Veterans with proteinuric diabetic kidney disease, an estimated GFR of 30.0 to 89.9 ml/minute/1.73 m<sup>2</sup>, and a urinary albumin-to-creatinine ratio of ≥300</p>	<p>N=1448</p> <p>Median follow-up 2.2 years</p>	<p>Primary: First occurrence of a decline in the eGFR (an absolute decrease of ≥30 ml/minute/1.73 m<sup>2</sup> if the eGFR was ≥60 ml/minute/1.73 m<sup>2</sup> at randomization or a relative decrease of ≥50% if the eGFR was &lt;60 ml/minute/1.73 m<sup>2</sup>), ESRD, or death</p> <p>Secondary: First occurrence of a decline in the eGFR or ESRD</p> <p>Tertiary: CV events, slope of change in eGFR, and change in albuminuria at 1</p>	<p>The trial was stopped early because the absolute risk of serious adverse events appeared to be greater than the potential benefit of reducing primary end-point events.</p> <p>Primary: There were 152 primary end-point events in the monotherapy group (21.0%) and 132 in the combination-therapy group (18.2%).The risk of the primary end point did not differ significantly between the two groups.</p> <p>Secondary: There were 101 secondary end-point events (a decline in the estimated GFR or ESRD) in the monotherapy group (14.0%) and 77 events in the combination-therapy group (10.6%).There was no significant between-group difference in mortality or ESRD (Table 2), though the number of ESRD events was small.</p> <p>Tertiary: There was no significant difference in the rate of cardiovascular events between the two groups. There was no significant difference in treatment effect on the decline in the estimated GFR (P=0.17). During adjustment of the losartan dose, the median urinary albumin-to-creatinine ratio declined from 959 to 807 (P=0.001). There was a further decline from randomization to 1 year, with a greater decline in the combination-therapy group (from 786 to 517) than in the monotherapy group (from 829 to 701) (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			year	
DREAM Trial Investigators <sup>140</sup> (2006) DREAM  Ramipril up to 15 mg/day  vs  placebo	DB, MC, PC, PRO, RCT, 2-by-2 factorial design  Adults aged 30 years or more with impaired fasting glucose and/or impaired glucose tolerance and no previous cardiovascular disease	N=5,269  3 years (median)	Primary: Composite of newly diagnosed diabetes or death  Secondary: Regression to normoglycemia, glucose levels, composite of cardiac and renal events (were not yet analyzed at the time of this publication)	Primary: The composite primary outcome did not differ significantly between the ramipril group (18.1%; HR, 0.91; 95% CI, 0.81 to 1.3; P=0.15) and the placebo group (19.5%).  Secondary: Participants receiving ramipril were more likely to have regression to normoglycemia than those receiving placebo (HR, 1.16; 95% CI, 1.07 to 1.27; P=0.001).  At the end of the study, the median fasting plasma glucose level was not significantly lower in the ramipril group than in the placebo group (P=0.07), though plasma glucose levels two hours after an oral glucose load were significantly lower in the ramipril group (P=0.01).
GISEN Group <sup>141</sup> (1997) REIN  Ramipril 1.25 mg/day  vs  placebo	DB, PC, RCT  Patients between 18 and 70 years who were either normotensive (<140/90 mm Hg) or hypertensive with chronic nephropathy and persistent proteinuria, who had not received ACE inhibition therapy for ≥2 months	N=166  16 months	Primary: Rate of GFR decline, extent to which this effect was dependent on the drug's antiproteinuric effect  Secondary: Blood pressure control, time to doubling of baseline serum creatinine or progression to end-stage renal failure, cardiovascular complications, total and cardiovascular	Primary: Mean rate of GFR decline per month was significantly lower in the ramipril group than in the placebo group (0.53 mL/min vs 0.88 mL/min; P=0.03).  Among the ramipril-assigned patients, percentage reduction in proteinuria was inversely correlated with decline in GFR (P=0.035) and predicted the reduction in risk of doubling of baseline creatinine or end-stage renal failure (18 ramipril vs 40 placebo; P=0.04).  Secondary: Blood pressure control and the overall number of cardiovascular events were similar in the two treatment groups.  Fifty-eight patients (18 in the ramipril group and 40 in the placebo group) reached the combined end point of doubling of baseline serum creatinine concentration or end-stage renal failure (P=0.02). The risk of progression was still significantly reduced after adjustment for changes in SBP (P=0.04) and DBP (P=0.04) with ramipril, but not after adjustment for changes in proteinuria.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			mortality	Note: Originally, 352 patients were placed into stratum 1 (urinary protein excretion exceeding 1 g/24 hours) or stratum 2 (urinary protein excretion exceeding 3.0 g/24 hours). At the second planned interim analysis, the difference in decline in GFR between the ramipril and placebo groups in stratum 2 was highly significant (P=0.001). The Independent Adjudicating Panel therefore decided to open the randomization code and do the final analysis in this stratum while stratum 1 continued in the trial.
<p>Uresin et al.<sup>142</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and ramipril 5 to 10 mg QD</p>	<p>DB, MC, RCT</p> <p>Patients ≥18 years of age with type 1 or type 2 diabetes mellitus and stage 1 to 2 HTN (mean sitting DBP) &gt;95 and &lt;110 mm Hg)</p>	<p>N=837</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, proportion of patients with a successful response to treatment (trough mean sitting DBP &lt;90 mm Hg and/or ≥10 mm Hg reduction from baseline), rates of blood pressure control (blood pressure &lt;130/80 mm Hg), changes from baseline in 24-hour ABPM measurements, and changes in biomarkers (plasma renin concentration, plasma renin</p>	<p>Primary: Aliskiren monotherapy, ramipril monotherapy, and aliskiren and ramipril combination therapy lowered mean sitting DBP by 11.3, 10.7, and 12.8 mm Hg, respectively. Treatment with aliskiren and ramipril combination therapy produced significantly greater reductions from baseline in mean sitting DBP compared to either aliskiren monotherapy (P=0.043) or ramipril monotherapy (P=0.004). Aliskiren 300 mg was statistically non-inferior (P=0.0002) to ramipril 10 mg for the change in mean sitting DBP.</p> <p>Secondary: Aliskiren monotherapy, ramipril monotherapy, and aliskiren and ramipril combination therapy lowered mean sitting SBP by 14.7, 12.0, and 16.6 mm Hg, respectively. Treatment with aliskiren and ramipril combination therapy produced significantly greater reductions from baseline in mean sitting SBP compared to ramipril monotherapy (P&lt;0.0001), but not aliskiren monotherapy (P=0.088). Aliskiren monotherapy was statistically superior to ramipril for the change in mean sitting SBP (P=0.021).</p> <p>The proportion of patients with a successful response to therapy was similar for aliskiren and ramipril combination therapy (74.1%) and aliskiren monotherapy (73.1%). The responder rates in both groups were significantly higher (P&lt;0.05) compared to ramipril monotherapy (65.8%).</p> <p>Rates of blood pressure control with aliskiren and ramipril combination pressure (13.1%) were not significantly different compared to aliskiren monotherapy (8.2%) or ramipril monotherapy (8.4%).</p> <p>All treatments significantly lowered mean 24-hour ambulatory blood pressure. Aliskiren and ramipril combination therapy was significantly more effective compared to ramipril monotherapy in lowering 24-hour</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			activity, aldosterone)	mean ambulatory DBP (P=0.034). There was no significant difference in 24-hour ambulatory SBP compared to ramipril monotherapy.  Aliskiren significantly reduced plasma renin activity from baseline as monotherapy (by 66%, P<0.0001) or in combination with ramipril (by 48%, P<0.0001).
Agodoa et al. <sup>143</sup> (2001) AASK  Ramipril 2.5 to 10 mg QD  vs  amlodipine 5 to 10 mg QD	DB, MC, RCT  African American patients, age 18 to 70 years old, with hypertensive renal disease (GFR 20 to 65 mL/min)	N=1,094  4 years	Primary: Rate of change in GFR (GFR slope)  Secondary: Composite of: confirmed reduction GFR by 50% or by 25 mL/min for baseline, ESRD	Primary: The average decline in GFR was slower, by 36% in the ramipril group as compared to the amlodipine group (P=0.002). However, during the first three months, GFR increased more in the amlodipine group than the ramipril group (P<0.001). The mean total slope did not differ between the groups (P=0.38).  Secondary: The risk reduction for the composite secondary outcome was significantly greater for the ramipril group than the amlodipine group (P=0.005). The rate of ESRD was significantly lower in the ramipril group (P=0.01).
Wright et al. <sup>144</sup> (2002) AASK  Ramipril 2.5 to 10 mg/day  vs  amlodipine 5 to 10 mg/day  vs  metoprolol 50 to 200 mg/day	DB, MC, RCT  Patients were self-identified African Americans aged 18 to 70 years with HTN and a GFR between 20 and 65 mL/min/ 1.73 m <sup>2</sup> and no other identified cause of renal insufficiency	N=1,094  3-6.4 years	Primary: Rate of change in GFR (grouped by usual blood pressure [MAP goal 102 to 107 mm Hg] vs lower blood pressure [ $\leq$ 92 mm Hg])  Secondary: Clinical composite outcome (reduction in GFR by 50% or more, ESRD, or death)	Primary: No significant difference in primary outcome was reported between the usual blood pressure group compared to the lower blood pressure group (P=0.24).  None of the drug group comparisons showed consistently significant differences in the GFR slope.  Secondary: The lower blood pressure goal did not significantly reduce the rate of the clinical composite outcome (risk reduction for lower blood pressure group, 2%; 95% CI, -22 to 21; P=0.85).  Ramipril resulted in significant risk reductions in the clinical composite outcomes compared to amlodipine (38%; 95% CI, 14 to 56; P=0.004) and metoprolol (22%; 95% CI, 1 to 38; P=0.04).  There was no significant difference in the clinical composite outcome between the amlodipine and metoprolol groups.
Bianchi et al. <sup>145</sup>	RCT, OL	N=128	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2010)</p> <p>Ramipril 10 mg and atorvastatin 10 mg QD (conventional therapy)</p> <p>vs</p> <p>spironolactone 25 mg, ramipril 10 mg, irbesartan 300 mg, and atorvastatin 10 mg QD (intensive therapy)</p> <p>The addition of diuretics, calcium antagonists, <math>\beta</math>-blockers or <math>\alpha</math>1-receptor antagonists were added to achieve blood pressure &lt;130/80 mm Hg</p>	<p>Patients with a clinical diagnosis of idiopathic chronic glomerulonephritis and urine protein-creatinine ratio &gt;1 g/g</p>	<p>36 months</p>	<p>Changes over time in proteinuria and eGFR</p> <p>Secondary: Adverse events, drop outs</p>	<p>SBP decreased more in the intensive-therapy group (from 156.6 to 113.5 mm Hg) than in the conventional therapy group (from 155.7 to 122.7 mm Hg; <math>P&lt;0.01</math>).</p> <p>Urine protein excretion decreased from 2.65 to 0.45 g/g creatinine with intensive therapy (<math>P&lt;0.001</math>). With conventional therapy, urine protein excretion decreased from 2.60 to 1.23 g/g creatinine (<math>P&lt;0.001</math>).</p> <p>With intensive therapy, eGFR did not significantly change over time (64.6 vs 62.9 mL/min/1.73 m<sup>2</sup>). With conventional therapy, eGFR decreased from 62.5 to 55.8 mL/min/1.73 m<sup>2</sup> (<math>P&lt;0.01</math>).</p> <p>Secondary: In the conventional therapy group, eight patients discontinued the study due to hyperkalemia, cough, and rapid deterioration in kidney function. In the intensive therapy group, 15 dropped out due to hyperkalemia, cough, and hypotension. Nine patients in the intensive therapy group developed gynecomastia. Twelve patients on conventional and 31 on intensive therapy had to interrupt the study temporarily because of low blood pressure. No patient developed an increase in creatine kinase, alanine aminotransferase, and alkaline phosphatase levels during the study.</p>
<p>Chrysostomou et al.<sup>146</sup> (2006)</p> <p>Ramipril 5 mg/day plus spironolactone 25 mg/day and placebo</p>	<p>DB, PC, RCT</p> <p>Patients 18 to 75 years of age, with a 24 hour urinary protein excretion &gt;1.5 g/24 hours on <math>\geq 2</math> occasions <math>\geq 3</math> months apart, serum creatinine level <math>\leq 200</math></p>	<p>N=41</p> <p>6 months</p>	<p>Primary: Change in 24 hour urinary protein excretion at three months</p> <p>Secondary: Change in 24 hour urinary protein excretion at six</p>	<p>Primary: Compared to ramipril-treated patients, the 24 hour urinary protein excretion reduction at three months was significantly greater in ramipril plus spironolactone-treated patients (<math>P=0.004</math>).</p> <p>Ramipril-, irbesartan- and spironolactone-treated patients exhibited a significant reduction in 24 hour urinary protein excretion compared to ramipril-treated patients (<math>P&lt;0.001</math>).</p> <p>There was no significant difference in 24 hour urinary protein excretion</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>ramipril 5 mg/day plus irbesartan 150 mg/day and placebo</p> <p>vs</p> <p>ramipril 5 mg/day plus placebo and placebo</p> <p>vs</p> <p>spironolactone 25 mg/day plus irbesartan 150 mg/day and ramipril 5 mg/day</p>	<p>μmol/L with &lt;20% variability in the preceding 3 months and treatment with an ACE inhibitor ≥6 months</p>		<p>months, change in blood pressure and creatinine clearance, adverse effects</p>	<p>with ramipril- and ramipril plus irbesartan-treated patients (P=1.00).</p> <p>At three months, spironolactone-treated patients exhibited a significant reduction in proteinuria from baseline (P≤0.001). In contrast, non-spironolactone-treated patients did not experience a significant reduction in proteinuria from baseline (P=0.840).</p> <p>Secondary: At six months, spironolactone-treated patients exhibited the greatest reduction in proteinuria compared to the other treatments (P&lt;0.05).</p> <p>At six months, DBP was higher among ramipril monotherapy-treated patients compared to the other treatments (P=0.046). There was no difference in SBP among the treatments (P value not reported).</p> <p>There were no differences in creatinine clearance among the treatments (P&gt;0.05).</p> <p>Gynecomastia was not observed with any of the treatments.</p>
<p>Nakao et al.<sup>147</sup> (2003) COOPERATE</p> <p>Trandolapril 3 mg/day</p> <p>vs</p> <p>losartan 100 mg/day</p> <p>vs</p> <p>trandolapril and losartan at equivalent doses</p>	<p>DB, MC, PC, RCT</p> <p>Patients aged 18 to 70 years with chronic nephropathy (nondiabetic renal disease)</p>	<p>N=263</p> <p>3 years</p>	<p>Primary: Composite of time to doubling of serum creatinine or ESRD</p> <p>Secondary: Changes in blood pressure, daily urinary protein excretion, adverse effects</p>	<p>Primary: The combined end point was reached in 11% of patients in the combination trandolapril and losartan group compared to 23% of patients in the trandolapril (P=0.018) and 23% of patients in the losartan group (P=0.016).</p> <p>Secondary: Mean SBP and DBP reductions were similar among the three treatment groups (P=0.109).</p> <p>All patients receiving active treatment had significant decreases in urinary protein excretion, but the greatest difference was seen with the combination trandolapril and losartan group compared to trandolapril or losartan (-75.6, -44.3, and -42.1%, respectively; P=0.01).</p> <p>The frequency of adverse events did not differ between groups, although a slightly higher occurrence of hyperkalemia and dry cough was recorded in</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Ruggenenti et al.<sup>148</sup> (2004) BENEDICT</p> <p>Trandolapril 2 mg/day</p> <p>vs</p> <p>verapamil SR 240 mg/day</p> <p>vs</p> <p>trandolapril and verapamil SR 2-180 mg/day (fixed-dose combination)</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, RCT</p> <p>Patients ≥40 years with type 2 diabetes (not exceeding 25 years) and HTN (SBP ≥130 mm Hg and/or DBP ≥85 mm Hg ) but with normoalbuminuria (urinary albumin excretion rate of &lt;20 mcg/minute)</p>	<p>N=1,204</p> <p>3.6 years (median)</p>	<p>Primary: Development of persistent microalbuminuria comparing combination therapy to placebo, acceleration factor</p> <p>Secondary: Primary end point comparing trandolapril and verapamil monotherapy to placebo, blood pressure, adverse events</p>	<p>the trandolapril and combination groups than in the losartan group.</p> <p>Primary: The primary outcome was reached in 5.7% of patients receiving combination therapy vs 10.0% for patients receiving placebo. The estimated acceleration factor (which quantifies the effect of one treatment relative to another in accelerating or slowing disease progression) adjusted for predefined baseline characteristics was 0.39 for the comparison between verapamil plus trandolapril and placebo (P=0.01).</p> <p>Secondary: The primary outcome was reached in 6.0% of patients receiving trandolapril, 11.9% receiving verapamil, and 10.0% receiving placebo. The estimated acceleration factor was 0.47 for trandolapril vs placebo (P=0.01) and 0.83 for verapamil vs placebo (P=0.54).</p> <p>Trandolapril plus verapamil and trandolapril alone delayed the onset of microalbuminuria by factors of 2.6 and 2.1, respectively.</p> <p>Throughout the study the average trough SBP/DBP was 139/80 mm Hg for patients receiving trandolapril plus verapamil, 139/81 mm Hg for trandolapril, 141/82 mm Hg for verapamil and 142/83 mm Hg for placebo. The comparison was significant (P≤0.002) between trandolapril plus verapamil or trandolapril alone vs placebo, but not for verapamil vs placebo.</p> <p>Serious adverse events were similar in all treatment groups.</p>
<p>Casas et al.<sup>149</sup> (2005)</p> <p>ACE inhibitor or ARBs compared to placebo</p> <p>vs</p> <p>ACE inhibitor or ARBs compared</p>	<p>MA (127 trials)</p> <p>Studies in adults that examined the effect of any drug treatment with a blood pressure lowering action on progression of renal disease</p>	<p>N=not reported</p> <p>4.2 years (mean)</p>	<p>Primary: Doubling of serum creatinine, and ESRD</p> <p>Secondary: Serum creatinine, urine albumin excretion and GFR</p>	<p>Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>Secondary:</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>to other antihypertensive drugs (β-adrenergic blocking agents, α-adrenergic blocking agents, calcium-channel blocking agents, or combinations)</p> <p>Specific agents and doses were not specified.</p>				<p>Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).</p> <p>Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).</p> <p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>
<p>Strippoli et al.<sup>150</sup> (2004)</p> <p>ACE inhibitors</p> <p>vs</p> <p>placebo</p> <p>or</p> <p>ARBs</p> <p>vs</p> <p>placebo</p> <p>or</p> <p>ACE inhibitors</p> <p>vs</p>	<p>MA</p> <p>Patients with diabetic nephropathy</p>	<p>43 trials</p> <p>≥6 months (range 6 to 63.6 months)</p>	<p>Primary: All-cause mortality, renal outcomes (ESRD, doubling of serum creatinine, microalbuminuria to macroalbuminuria)</p> <p>Secondary: Not reported</p>	<p>Primary: ACE inhibitors significantly reduced all-cause mortality compared to placebo or no treatment (RR, 0.79; 95% CI, 0.63 to 0.99; P=0.04). There was a nonsignificant trend for reduction in ESRD (P=0.07) and doubling of serum creatinine (P=0.08) with ACE inhibitors compared to placebo or no treatment. ACE inhibitors significantly reduced the risk of progression from microalbuminuria to macroalbuminuria (P=0.0007) and increased regression back to normoalbuminuria (P&lt;0.0001) compared to placebo or no treatment.</p> <p>ARBs did not significantly reduce all-cause mortality compared to placebo or no treatment (RR, 0.99; 95% CI, 0.85 to 1.17; P=0.95). ARBs significantly reduced the risk of ESRD (P=0.001) and doubling of serum creatinine (P=0.004). ARBs significantly decreased the risk of progression to macroalbuminuria (P=0.001) and increased regression to normoalbuminuria (P=0.02) compared to placebo or no treatment.</p> <p>The three trials that compared ACE inhibitors to ARBs did not report on all-cause mortality, ESRD or doubling of serum creatinine. Progression from microalbuminuria to macroalbuminuria was reported in one trial (N=92) and there was no significant difference in risk, with the point estimate favoring ACE inhibitors (RR, 0.16; 95% CI, 0.02 to 1.44). Regression from microalbuminuria to normoalbuminuria in 1 trial showed</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
ARBs				<p>a nonsignificant difference in the risk.</p> <p>Secondary: Not reported</p>
<p>Strippoli et al.<sup>151</sup> (2006)</p> <p>ACE inhibitors vs placebo or ARBs vs placebo or ACE inhibitors vs ARBs</p>	<p>MA</p> <p>Patients with diabetic kidney disease</p>	<p>N=12,067 (49 trials)</p> <p>≥6 months</p>	<p>Primary: All-cause mortality, ESRD, doubling of serum creatinine concentration, progression from micro- to macroalbuminuria, regression from micro- to normoalbuminuria, drug-related toxicity (including cough, headache, hyperkalemia, impotence and pedal edema)</p> <p>Secondary: Not reported</p>	<p>Primary: There was no significant difference in the risk of all-cause mortality for ACE inhibitors vs placebo or no treatment (RR, 0.91; 95% CI, 0.71 to 1.17) and ARBs vs placebo or no treatment (RR, 0.99; 95% CI, 0.85 to 1.17). No statistically significant reduction in the risk of all-cause mortality was found in the three studies that compared ACE inhibitors with ARBs (RR, 0.92; 95% CI, 0.31 to 2.78).</p> <p>A subgroup analysis of studies showed a significant reduction in the risk of all-cause mortality with the use of full-dose ACE inhibitors (RR, 0.78; 95% CI, 0.61 to 0.98) but not when using half or less than half the maximum tolerable dose of ACE inhibitors (RR, 1.18; 95% CI, 0.41 to 3.44).</p> <p>There was a significant reduction in the risk of ESRD with ACE inhibitors and ARBS compared to placebo or no treatment (RR, 0.60; 95% CI, 0.39 to 0.93 and RR, 0.78; 95% CI, 0.67 to 0.91, respectively). There was a significant reduction in the risk of doubling of serum creatinine concentration with ACE inhibitors and ARBS (RR, 0.68; 95% CI, 0.47 to 1.0 and RR, 0.79; 95% CI, 0.67 to 0.93, respectively).</p> <p>ACE inhibitors and ARBS significantly reduced the risk of progression from micro- to macroalbuminuria (RR, 0.45; 95% CI, 0.29 to 0.69 and RR, 0.49; 95% CI, 0.32 to 0.75, respectively). ACE inhibitors and ARBS significantly increased the regression from micro- to normoalbuminuria compared to placebo or no treatment (RR, 3.06; 95% CI, 1.76 to 5.35 and RR, 1.42; 95% CI, 1.15 to 1.93, respectively).</p> <p>The seven studies that compared ACE inhibitors to ARBS did not report the outcome of ESRD or doubling of serum creatinine. Progression from micro- to macroalbuminuria and from micro- to normoalbuminuria were evaluated each in one trial and showed a nonsignificant difference in the risk between ACE inhibitors and ARBS.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>ACE inhibitors were associated with a significant increase in the risk of cough but not hyperkalemia, headache or impotence when compared to placebo or no treatment. ARBS were associated with a significant increase in the risk of hyperkalemia but not cough or headache compared to placebo or no treatment.</p> <p>Secondary: Not reported</p>
<b>Miscellaneous</b>				
<p>Montalescot et al.<sup>152</sup> (2009) ARCHIPELAGO</p> <p>Enalapril 10 mg QD, followed by 20 mg QD on day 15</p> <p>vs</p> <p>irbesartan 150 mg QD, followed by 300 mg QD on day 15</p>	<p>AC, DB, MC, RCT</p> <p>Adults with non-ST elevation ACS</p>	<p>N=429</p> <p>60 days</p>	<p>Primary: Change from baseline in high-sensitivity C-reactive protein at day 60</p> <p>Secondary: Changes in other inflammatory markers such as troponin I</p>	<p>Primary: High-sensitivity C-reactive protein levels were comparable in both treatment groups (irbesartan: 15.2 mg/L at baseline, 6.5 mg/L at day 60; absolute change of -8.7 mg/L; enalapril: 12.6 mg/L at baseline, 5.5 mg/L at day 60; absolute change of -7.1 mg/L, P value not significant).</p> <p>Secondary: Similarly, mean levels of markers of myocardial injury (troponin I) and endothelial dysfunction (microalbuminuria) also decreased from baseline to day 60, with no significant differences between treatment groups.</p>
<p>Dagenais et al.<sup>153</sup> (2008)</p> <p>Ramipril 15 mg or rosiglitazone 8 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, RCT</p> <p>Adults &gt;30 years with impaired fasting glucose or impaired glucose tolerance without known cardiovascular disease or renal insufficiency</p>	<p>N=5,269</p> <p>3 years</p>	<p>Primary: Composite cardiorenal outcome (first occurrence of any cardiovascular death, nonfatal MI, stroke, new heart failure, progression to microalbuminuria</p>	<p>Primary: Compared to placebo, neither ramipril (15.7 vs 16.0%; HR, 0.98; P=0.75) nor rosiglitazone (15.0 vs 16.8%; HR, 0.87; P=0.07) reduced the risk of the cardiorenal composite outcome.</p> <p>Secondary: Ramipril had no impact on the cardiovascular disease and renal components. Rosiglitazone increased heart failure (0.53 vs 0.08%; HR, 7.04; P=0.01), but reduced the risk of the renal component (HR, 0.80; P=0.005).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>or proteinuria, renal insufficiency requiring dialysis or transplantation)</p> <p>Secondary: Subcomponents of the primary analysis</p>	
<p>Belluzzi et al.<sup>154</sup> (2009)</p> <p>Ramipril 5mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, RCT</p> <p>Adults with lone atrial fibrillation without heart disease or HTN</p>	<p>N=62</p> <p>3 years</p>	<p>Primary: Relapse of atrial fibrillation as determined by clinical assessment, ECG, 24 hour Holter monitor, and questionnaire collection.</p> <p>Secondary: Not reported</p>	<p>Primary: At the end of the study, atrial fibrillation relapses were observed in three ramipril-treated patients and in 10 control patients (P&lt;0.03).</p> <p>Secondary: Not reported</p>
<p>Hansson et al.<sup>155</sup> (1998)</p> <p>HOT</p> <p>Aspirin 75 mg QD</p> <p>vs</p> <p>placebo</p> <p>A 5 step antihypertensive treatment regimen: 1) felodipine 5 mg</p>	<p>MC, RCT, OL</p> <p>Adults with HTN and a DBP between 100 and 115 mm Hg</p>	<p>N=18,790</p> <p>3.8 years</p>	<p>Primary: Major cardiovascular events (fatal and nonfatal, fatal and nonfatal stroke, and all other cardiovascular deaths)</p> <p>Secondary: Not reported</p>	<p>Primary: There were 9.9, 10.0, and 9.3 major cardiovascular events per 1,000-patients years, respectively, in the DBP ≤90, DBP ≤85, and DBP ≤80 treatment groups (P=0.50), thus suggesting that the reduction of DBP below 90 mm Hg does not provide any mortality or morbidity advantage.</p> <p>Aspirin reduced major cardiovascular events by 15% (P=0.03) and all MI by 36% (P=0.002), with no effect on stroke. There were seven fatal bleeds in the aspirin group and eight in the placebo group, and 129 versus 70 non-fatal major bleeds in the two groups, respectively (P&lt;0.001).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
QD, 2) ACE inhibitor or $\beta$ -blocker, 3) dose titrations, 4) dose titrations, 5) diuretic.				

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, ER=extended-release, QD=once daily, SR=sustained-release, TID=three times daily

Study design abbreviations: AC=active-controlled, BE=blinded endpoint, DB=double blind, DD=double dummy, MA=meta-analysis, MC=multicenter, OL=open label, OS=observational, PC=placebo controlled, PG=parallel group, PRO=prospective, RETRO=retrospective, RCT=randomized controlled trial, SB=single-blind, XO=crossover

Miscellaneous abbreviations: ABPM=ambulatory blood pressure monitoring, ACE inhibitor=angiotensin converting enzyme inhibitor, ACS=acute coronary syndrome, ARB=angiotensin II receptor blocker, BMI=body mass index, CAD=coronary artery disease, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure,

EKG=electrocardiogram, eGFR=estimated glomerular filtration rate, ESRD=end stage renal disease, GFR=glomerular filtration rate, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HR=hazard ratio, HTN=hypertension, IV=intravenous, LDL-C=low-density lipoprotein cholesterol, LVEF=left ventricular ejection fraction, MAP=mean arterial pressure, MI=myocardial infarction, MMSE=Mini Mental State Examination, MRI=magnetic resonance imaging, NIDDM=non-insulin dependent diabetes mellitus, NNT=number needed to treat, NYHA=New York Heart Association, OR=odds ratio, PAD=peripheral arterial disease, PCI=percutaneous coronary intervention, PVD=peripheral vascular disease, QOL=quality of life, RR=relative risk, SBP=systolic blood pressure, TC=total cholesterol, TG=triglyceride, TIA=transient ischemic attack, WHO=World Health Organization

**Additional Evidence**

Dose Simplification

Taylor et al. evaluated adherence rates with amlodipine and benazepril fixed-dose combination compared to an ACE inhibitor plus a long-acting dihydropyridine administered as separate formulations. There was no significant difference in adherence in younger subjects (18 to 39 year olds); however, in all age group combined, adherence rates were higher with amlodipine and benazepril compared to the use of an ACE inhibitor plus a long-acting dihydropyridine (80.8 vs 73.8%; P<0.001).<sup>156</sup> Dickson et al. evaluated adherence rates with amlodipine and benazepril fixed-dose combination compared to an ACE inhibitor plus a long-acting dihydropyridine administered as separate formulations in an elderly Medicaid population. Over a 12 month period, adherence rates were reported to be significantly higher with fixed-dose combination product compared to the administration of an ACE inhibitor and dihydropyridine as separate formulations (63.4 vs 49.0%; P<0.0001).<sup>157</sup> Dezzi et al. also reported significantly higher compliance rates at 12 months in patients receiving fixed-dose lisinopril and hydrochlorothiazide (68.7%) or enalapril and hydrochlorothiazide (70.0%) vs administration of the components as separate formulations (57.8 and 57.5%, respectively; P<0.05 for both comparisons).<sup>158</sup>

Stable Therapy

Sapienza et al. evaluated the impact of converting long-term care patients from high-dose calcium-channel blockers or ACE inhibitor plus calcium-channel blockers to a fixed-dose combination of amlodipine/benazepril. There was no significant change in blood pressure from baseline following the conversion; however, there was a significant reduction (81.8%) in the number of patients reporting ≥1 drug-related adverse event (22 vs 4; P<0.05), particularly edema (75% reduction).<sup>159</sup>

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 12. Relative Cost of the Angiotensin-Converting Enzyme Inhibitors**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Benazepril	tablet	Lotensin®*	\$\$\$	\$
Captopril	tablet	N/A	N/A	\$\$\$
Enalapril	solution, tablet	Epaned®*, Vasotec®*	\$\$\$\$	\$\$\$
Fosinopril	tablet	N/A	N/A	\$
Lisinopril	solution, tablet	Prinivil®*, Qbrelis®, Zestril®*	\$\$\$\$	\$
Moexipril	tablet	N/A	N/A	\$\$\$\$
Perindopril	tablet	N/A	N/A	\$
Quinapril	tablet	Accupril®*	\$\$\$\$	\$
Ramipril	capsule	Altace®*	\$\$\$\$	\$
Trandolapril	tablet	N/A	N/A	\$
<b>Combination Products</b>				
Benazepril and HCTZ	tablet	Lotensin HCT®*	\$\$\$	\$
Captopril and HCTZ	tablet	N/A	N/A	\$\$\$\$
Enalapril and HCTZ	tablet	Vaseretic®*	\$\$\$\$	\$\$
Fosinopril and HCTZ	tablet	N/A	N/A	\$\$\$
Lisinopril and HCTZ	tablet	Prinzide®*, Zestoretic®*	\$\$\$\$	\$
Quinapril and HCTZ	tablet	Accuretic®*	\$\$\$\$	\$\$

\*Generic is available in at least one dosage form or strength.  
HCTZ=hydrochlorothiazide, N/A=not available

## X. Conclusions

All of the angiotensin-converting enzyme (ACE) inhibitors are approved for the treatment of hypertension. Some of the products are also approved for the treatment of diabetic nephropathy, heart failure, and post-myocardial infarction.<sup>3-19</sup> The ACE inhibitors are available as single entity products, as well as in combination with hydrochlorothiazide or verapamil. All of the products are available in a generic formulation.

There are numerous national and international guidelines that recommend the use of ACE inhibitors in patients with the following conditions: acute coronary syndrome, cerebrovascular disease, coronary artery disease, diabetes, diabetic nephropathy, heart failure, hypertension, left ventricular dysfunction, left ventricular hypertrophy, previous myocardial infarction, and renal disease. In general, guidelines do not give preference to one ACE inhibitor over another.<sup>20-38</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>30-35</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>30</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>30-38</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>30-37</sup>

In clinical trials, the ACE inhibitors have been shown to reduce cardiovascular morbidity and mortality, preserve renal function in patients with nephropathy, and effectively lower blood pressure when administered as monotherapy or in combination with other antihypertensive agents.<sup>39-150</sup> Most patients will need more than one antihypertensive agent to achieve blood pressure goals. The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>30-35,156-158</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations.

There is insufficient evidence to support that one brand angiotensin-converting enzyme inhibitor is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand angiotensin-converting enzyme inhibitors within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand angiotensin-converting enzyme inhibitor is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.



## XII. References

1. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
2. Givertz MM, Colucci WS. Pharmacologic therapy of heart failure with reduced ejection fraction: Mechanisms of action. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2022 [cited 2022 Jan]. Available from: <http://www.uptodate.com/utd/index.do>.
3. Lotensin<sup>®</sup> [package insert]. Parsippany (NJ): Validus Pharmaceuticals LLC; 2019 Jan.
4. Epaned<sup>®</sup> [package insert]. Wilmington (MA): Azurity Pharmaceuticals, Inc; 2020 Mar.
5. Vasotec<sup>®</sup> [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Dec.
6. Prinivil<sup>®</sup> [package insert]. Whitehouse Station (NJ): Merck & Co., Inc.; 2021 Nov.
7. Qbrelis<sup>®</sup> [package insert]. Wilmington (MA): Azurity Pharmaceuticals, Inc; 2020 Mar.
8. Zestril<sup>®</sup> [package insert]. Pine Brook (NJ): Almatica Pharma; 2020 Mar.
9. Accupril<sup>®</sup> [package insert]. New York (NY): Pfizer Inc; 2017 Apr.
10. Altace<sup>®</sup> [package insert]. New York (NY): Pfizer Inc; 2021 Aug.
11. Lotensin HCT<sup>®</sup> [package insert]. Parsippany (NJ): Validus Pharmaceuticals LLC; 2020 Oct.
12. Vaseretic<sup>®</sup> [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Sep.
13. Prinzide<sup>®</sup> [package insert]. Whitehouse Station (NJ). Merck & Co, Inc; 2013 Feb.
14. Zestoretic<sup>®</sup> [package insert]. Pine Brook (NJ): Almatica Pharma, Inc.; 2021 Jul.
15. Accuretic<sup>®</sup> [package insert]. New York (NY): Pfizer; 2021 Feb.
16. Tarka<sup>®</sup> [package insert]. North Chicago (IL): AbbVie Inc.; 2019 Sep.
17. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
18. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
19. Micromedex<sup>®</sup> Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
20. Fraker T, Fihn S, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of chronic stable angina: a report of the American College of Cardiology/American Heart Association task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *Circulation*. 2007 Dec 4;116(23):2762-72.
21. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425.
22. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline From the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
23. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
24. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>
25. O’Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.

26. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
27. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].
28. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
29. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
30. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
31. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
32. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
33. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
34. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
35. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
36. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
37. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
38. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
39. Jamerson K, Weber MA, Bakris GL, et al. Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high risk patients. *N Engl J Med* 2008;359:2417-28.
40. Weber MA, Bakris GL, Jamerson K, Weir M, Kjeldsen SE, Devereux RB, et al.; ACCOMPLISH Investigators. Cardiovascular events during differing hypertension therapies in patients with diabetes. *J Am Coll Cardiol*. 2010 Jun 29;56(1):77-85.
41. Weber MA, Jamerson K, Bakris G, Weir MR, Zappe D, Zhang Y, et al. Effects of body size and hypertension treatments on cardiovascular event rates: subanalysis of the ACCOMPLISH randomised controlled trial. *Lancet*. 2013;381:537-45.
42. Swedberg K, Held P, Kjeksus J, et al. Effects of the early administration of enalapril on mortality in patients with acute myocardial infarction results of the cooperative New Scandinavian enalapril survival study II (CONSENSUS II). *N Engl J Med*. 1992 Sep 3;327(10):678-84.

43. Wing LMH, Reid CM, Ryan P, et al; for the Second Australian National Blood Pressure Study Group. A comparison of outcomes with angiotensin-converting enzyme inhibitors and diuretics for hypertension in the elderly. *N Engl J Med.* 2003 Feb 13;348(7):583-92.
44. Nissen SE, Tuzcu EM, Libby P, et al; for the CAMELOT Investigators. Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: a randomized controlled trial. *JAMA.* 2004 Nov 10;292(18):2217-26.
45. Pitt B, Reichek N, Willenbrock R, Zannad F, Philips RA, Roniker B, et al. Effects of eplerenone, enalapril, and eplerenone/enalapril in patients with essential hypertension and left ventricular hypertrophy: the 4E-left ventricular hypertrophy study. *Circulation.* 2003;108(15):1831-8.
46. Hansson L, Lindholm LH, Ekblom T, et al. Randomized trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet.* 1999 Nov 20;354(9192):1751-6.
47. ALLHAT Officers and Coordinators for ALLHAT Collaborative Research Group. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium-channel blocker vs diuretic: the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA.* 2002 Dec 18;288(23):2981-97.
48. Black HR, Davis B, Barzilay J, et al; Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Metabolic and clinical outcomes in nondiabetic individuals with the metabolic syndrome assigned to chlorthalidone, amlodipine, or lisinopril as initial treatment for hypertension: a report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Diabetes Care.* 2008 Feb;31(2):353-60.
49. Rahman M, Ford CE, Cutler JA, Davis BR, Piller LB, Whelton PK, et al. Long-term renal and cardiovascular outcomes in antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT) participants by baseline estimated GFR. *Clin J Am Soc Nephrol.* 2012;7:989-1002.
50. Muntner P, Levitan EB, Lynch AI, et al. Effect of chlorthalidone, amlodipine, and lisinopril on visit-to-visit variability of blood pressure: results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. *J Clin Hypertens (Greenwich).* 2014 May;16(5):323-330.
51. Bangalore S, Davis BR, Cushman WC, Pressel SL, Muntner PM, Calhoun DA, et al. Treatment-Resistant Hypertension and Outcomes Based on Randomized Treatment Group in ALLHAT. *Am J Med.* 2017 Apr;130(4):439-448.e9.
52. Fox KM, Bertrand M, Ferrari R, et al. EUROPA Investigators. Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomized, double-blind, placebo-controlled, multicentre trial (the EUROPA study). *Lancet.* 2003 Sep 6;362:782-8.
53. PREAMI Investigators. Effects of angiotensin-converting enzyme inhibition with perindopril on left ventricular remodeling and clinical outcome: results of the randomized perindopril and remodeling in elderly with acute myocardial infarction (PREAMI) study. *Arch Intern Med.* 2006 Mar 27;166:659-66.
54. ADVANCE Collaborative Group. Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): A randomized controlled trial. *Lancet* 2007;370:829-40.
55. The Heart Outcomes Prevention Evaluation Study Investigators. Effects of an angiotensin-converting enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med.* 2000 Jan 20;342:145-53.
56. ONTARGET Investigators. Telmisartan, ramipril, or both in patients at high risk for vascular events. *N Engl J Med.* 2008 Apr 10;358(15):1547-59.
57. Redon J, Mancia G, Sleight P, Schumacher H, Gao P, Pogue J, et al. Safety and efficacy of low blood pressures among patients with diabetes. *J Am Coll Cardiol.* 2012;59(1):78-83.
58. Mann JF, Anderson C, Gao P, et al. Dual inhibition of the renin-angiotensin system in high-risk diabetes and risk for stroke and other outcomes: results of the ONTARGET trial. *J Hypertens.* 2013 Feb;31(2):414-21.
59. The PEACE Trial Investigators. Angiotensin-converting enzyme inhibition in stable coronary artery disease. *N Engl J Med.* 2004 Nov 11;351(20):2058-68.
60. Pilote L, Abrahamowicz M, Rodrigues E, et al. Mortality rates in elderly patients who take different angiotensin-converting enzyme inhibitors after acute myocardial infarction: a class effect? *Ann Intern Med.* 2004 Jul 20;141(2):102-12.
61. Dahlof B, Sever PS, Poulter NR, et al; for the ASCOT Investigators. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure

- Lowering Arm (ASCOT-BPLA): a multicentre randomized controlled trial. *Lancet*. 2005 Sep 10;366(9489):895-906.
62. Chapman N, Dobson J, Wilson S, Dahlöf B, Sever PS, Wedel H, Poulter NR; Anglo-Scandinavian Cardiac Outcomes Trial Investigators. Effect of spironolactone on blood pressure in subjects with resistant hypertension. *Hypertension*. 2007 Apr;49(4):839-45.
  63. Pepine CJ, Handberg EM, Cooper-DeHoff RM, et al. A calcium antagonist vs a non-calcium antagonist hypertension treatment strategy for patients with coronary artery disease: the international verapamil-trandolapril study (INVEST): a randomized controlled trial. *JAMA*. 2003 Dec 3;290(21):2805-16.
  64. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*. 2005 Oct 29-Nov 4;366(9496):1545-53.
  65. Wiyosong CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
  66. Blood Pressure Lowering Treatment Trialists' Collaboration. Blood pressure-dependent and independent effects of agents that inhibit the renin-angiotensin system. *J Hypertens*. 2007 Jun;25(5):951-8.
  67. PROGRESS Collaborative group. Randomized trial of a perindopril-based blood-pressure-lowering regimen among 6105 individuals with previous stroke or transient ischemic attack. *Lancet*. 2001 Sep 29;358:1033-41.
  68. Arima H, Anderson C, Omae T, Woodward M, Hata J, Murakami Y, et al. Effects of blood pressure lowering on major vascular events among patients with isolated diastolic hypertension: the perindopril protection against recurrent stroke study (PROGRESS) trial. *Stroke*. 2011;42:2339-2341.
  69. Pfeffer MA, Braunwald E, Moye LA, et al; on behalf of the SAVE Investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: results of the Survival and Ventricular Enlargement Trial. *N Engl J Med*. 1992 Sep 3;327(10):669-77.
  70. Pitt B, Segal R, Martinez FA, et al; on behalf of ELITE Study Investigators. Randomized trial of losartan versus captopril in patients over 65 with heart failure (Evaluation of Losartan in the Elderly Study, ELITE). *Lancet*. 1997 Mar 15;349:747-52.
  71. Pitt B, Poole-Wilson PA, Segal R, et al; on behalf of the ELITE II Investigators. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomized trial—the Losartan Heart Failure Survival Study, ELITE II. *Lancet*. 2000 May 6;355:1582-7.
  72. Dickstein K, Kjekshus J, and the OPTIMAAL Steering Committee, for the OPTIMAAL Study Group. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomized trial. *Lancet*. 2002 Sep 7;360:752-60.
  73. Pfeffer MA, McMurray JJV, Velazquez EJ, et al; for the Valsartan in Acute Myocardial Infarction Trial Investigators. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med*. 2003 Nov 13;349:1893-906.
  74. The CONSENSUS Trial Study Group. Effects of enalapril on mortality in severe congestive heart failure: results of the cooperative North Scandinavian enalapril survival study. *N Engl J Med*. 1987 Jun 4;316(23):1429-35.
  75. The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med*. 1991 Aug 1; 325(5):293-302.
  76. The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med*. 1992 Sep 3;327(10):685-91.
  77. McMurray JJ, Krum H, Abraham WT, Dickstein K, Køber LV, Desai AS, et al. Aliskiren, Enalapril, or Aliskiren and Enalapril in Heart Failure. *N Engl J Med*. 2016 Apr 21;374(16):1521-32.
  78. McKelvie RS, Yusuf S, Pericak D, et al. Comparison of candesartan, enalapril, and their combination in congestive heart failure: randomized evaluation of strategies for left ventricular dysfunction (RESOLVD) pilot study: the RESOLVD Pilot Study Investigators. *Circulation*. 1999 Sep 7;100:1056-64.
  79. Willenheimer R, van Veldhuisen DJ, Silke B, Erdmann E, Follath F, Krum H, et al; CIBIS III Investigators. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite sequence: results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation*. 2005 Oct 18;112(16):2426-35.
  80. Cohn JN, Johnson G, Ziesche S, et al. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med*. 1991 Aug 1;325:303-10.
  81. Tu K, Mamdani M, Kopp A, Lee D. Comparison of angiotensin-converting enzyme inhibitors in the treatment of congestive heart failure. *Am J Cardiology*. 2005 Jan 15;95:283-6.

82. Packer M, Poole-Wilson PA, Armstrong PW, et al; on behalf of the ATLAS Study Group. Comparative effects of low and high doses of angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. *Circulation*. 1999 Dec 7;100:2312-18.
83. The Acute Infarction Ramipril Efficacy (AIRE) Study Investigators. Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet*. 1993;342:821-28.
84. Wu J, Hall AS, Gale CP, AIREX Study Investigators. Long-term survival benefit of ramipril in patients with acute myocardial infarction complicated by heart failure. *Heart*. 2021 Mar;107(5):389-395.
85. Kober L, Torp-Pedersen C, Carlsen JE, et al; for the Trandolapril Cardiac Evaluation (TRACE) Study Group. A clinical trial of the angiotensin-converting enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med*. 1995 Dec 21;333(25):1670-6.
86. Galløe AM, Skagen K, Christensen NJ, Nielsen SL, Frandsen EK, Bie P, et al. Dosage dependent hormonal counter regulation to combination therapy in patients with left ventricular dysfunction. *J Clin Pharm Ther*. 2006 Apr;31(2):139-47.
87. Galløe AM, Skagen K, Christensen NJ, Nielsen SL, Frandsen EK, Bie P, et al. Dosage dependent hormonal counter regulation to combination therapy in patients with left ventricular dysfunction. *J Clin Pharm Ther*. 2006 Apr;31(2):139-47.
88. Fröhlich H, Henning F, Täger T, Schellberg D, Grundtvig M, Goode K, et al. Comparative effectiveness of enalapril, lisinopril, and ramipril in the treatment of patients with chronic heart failure: a propensity score-matched cohort study. *Eur Heart J Cardiovasc Pharmacother*. 2018 Apr 1;4(2):82-92.
89. Lee VC, Rhew DC, Dylan M, Badamgarav E, Braunstein GD, Weingarten SR. Meta-analysis: angiotensin-receptor blockers in chronic heart failure and high-risk acute myocardial infarction. *Ann Intern Med*. 2004 Nov 2;141(9):693-704.
90. Kuschnir E, Acuna E, Sevilla D, Vasquez J, Bendersky M, Resk J, et al. Treatment of patients with essential hypertension: amlodipine 5 mg/benazepril 20 mg compared with amlodipine 5 mg, benazepril 20 mg, and placebo. *Clin Ther*. 1996;18(6):1213-24.
91. Neutel JM, Smith DH, Weber MA, Schofield L, Purkayastha D, Gatlin M. Efficacy of combination therapy for systolic blood pressure in patients with severe systolic hypertension: the Systolic Evaluation of Lotrel Efficacy and Comparative Therapies (SELECT) study. *J Clin Hypertens*. 2005 Nov;7(11):641-6.
92. Chrysant SG, Bakris GL. Amlodipine/benazepril combination therapy for hypertensive patients nonresponsive to benazepril monotherapy. *Am J Hypertens*. 2004 Jul;17(7):590-6.
93. Fogari R, Corea L, Cardoni O, et al. Combined therapy with benazepril and amlodipine in the treatment of hypertension inadequately controlled by an ACE inhibitor alone. *J Cardiovasc Pharmacol*. 1997 Oct;30(4):497-503.
94. Chrysant SG. Blood pressure effects of high-dose amlodipine-benazepril combination in black and white hypertensive patients not controlled on monotherapy. *Drugs RD*. 2012;12(2):57-64.
95. Messerli FH, Oparil S, Feng Z. Comparison of efficacy and side effects of combination therapy of angiotensin-converting enzyme inhibitor (benazepril) with calcium antagonist (either nifedipine or amlodipine) versus high-dose calcium antagonist monotherapy for systemic hypertension. *Am J Cardiol*. 2000 Dec 1;86:1182-7.
96. Hilleman DE, Ryschon KL, Mohiuddin SM, Wurdeman RL. Fixed-dose combination vs monotherapy in hypertension: a meta-analysis evaluation. *J Hum Hypertens*. 1999;13:477-83.
97. Jamerson K, Bakris GL, Dahlof B, et al; for the ACCOMPLISH Investigators. Exceptional early blood pressure control rates: the ACCOMPLISH trial. *Blood Press*. 2007;16(2):80-6.
98. Kereiakes DJ, Neutel JM, Punzi HA, et al. Efficacy and safety of olmesartan medoxomil and hydrochlorothiazide compared with benazepril and amlodipine besylate. *Am J Cardiovasc Drugs*. 2007;7(5):36-72.
99. Waeber B, Aschwanden R, Sadecky L, Ferber P. Combination of hydrochlorothiazide or benazepril with valsartan in hypertensive patients unresponsive to valsartan alone. *J Hypertens*. 2001 Nov;19(11):2097-104.
100. Malacco E, Piazza S, Carretta R, et al; Italian Blood Pressure Study Group. Comparison of benazepril-amlodipine and captopril-thiazide combinations in the management of mild-to-moderate hypertension. *Int J Clin Pharmacol Ther*. 2002 Jun;40(6):263-9.
101. Elliott WJ, Montoro R, Smith D, et al. Comparison of two strategies for intensifying antihypertensive treatment: low-dose combination (enalapril plus felodipine ER) versus increased dose of monotherapy (enalapril). *Am J Hypertens*. 1999 Jul;12:691-6.
102. Prisant LM, Weir MR, Papademetriou V, et al. Low-dose combination therapy: an alternative first-line approach to hypertension treatment. *Am Heart J*. 1995 Aug;130(2):359-66.

103. Ruilope L, Jager B, Prichard B. Eprosartan versus enalapril in elderly patients with hypertension: a double-blind, randomized trial. *Blood Pressure*. 2001;10:223-9.
104. Karlberg BE, Lins LE, Hermansson K. Efficacy and safety of telmisartan, a selective AT1 receptor antagonist, compared with enalapril in elderly patients with primary hypertension. TEES Study Group. *J Hypertens*. 1999 Feb;17(2):293-302.
105. Estacio RO, Jeffers BW, Hiatt WR, et al. The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med*. 1998 Mar 5;338(10):645-52.
106. Williams GH, Burgess E, Kolloch RE et al. Efficacy of eplerenone versus enalapril as monotherapy in systemic hypertension. *Am J of Cardiol*. 2004;93(8):990-6.
107. Tatti P, Pahor M, Byington RP, et al. Outcome results of the Fosinopril versus Amlodipine Cardiovascular Events Trial (FACET) in patients with hypertension and non-insulin dependent diabetes mellitus. *Diabetes Care*. 1998; 21:597-603.
108. Whelton A, Miller WE, Dunne, Jr B, et al. Once-daily lisinopril compared with twice-daily captopril in the treatment of mild-to-moderate hypertension: assessment of office and ambulatory blood pressure. *J Clin Pharmacol*. 1990 Dec;30(12):1074-80.
109. Strasser RH, Puig JG, Farsang C, et al. A comparison of the tolerability of the direct renin inhibitor aliskiren and lisinopril in patients with severe hypertension. *J Hum Hypertens*. 2007 Oct;21(10):780-7.
110. Rosei EA, Rizzoni D, Comini S, Boari G; Nebivolol-Lisinopril Study Group. Evaluation of the efficacy and tolerability of nebivolol versus lisinopril in the treatment of essential arterial hypertension: a randomized, multicentre, double-blind study. *Blood Press Suppl*. 2003 May;1:30-5.
111. Wald DS, Law M, Mills S, Bestwick JP, Morris JK, Wald NJ. A 16-week, randomized, double-blind, placebo-controlled, crossover trial to quantify the combined effect of an angiotensin-converting enzyme inhibitor and a beta-blocker on blood pressure reduction. *Clin Ther*. 2008 Nov;30(11):2030-9.
112. Karotsis AK, Symeonidis A, Mastorantonakis SE, Stergiou GS. Additional antihypertensive effect of drugs in hypertensive subjects uncontrolled on diltiazem monotherapy: a randomized controlled trial using office and home blood pressure monitoring. *Clin Exp Hypertens*. 2006;28(7):655-62.
113. McInnes GT, O'Kane KP, Istad H, Keinänen-Kiukaanniemi S, Van Mierlo HF. Comparison of the AT1-receptor blocker, candesartan cilexetil, and the ACE inhibitor, lisinopril, in fixed combination with low dose hydrochlorothiazide in hypertensive patients. *J Hum Hypertens*. 2000 Apr;14(4):263-9.
114. Poldermans D, Glazer R, Karagiannis S, et al. Tolerability and blood pressure-lowering efficacy of the combination of amlodipine plus valsartan compared with lisinopril plus hydrochlorothiazide in adult patients with stage 2 hypertension. *Clin Ther*. 2007 Feb;29(2):279-89.
115. Duprez D, Munger M, Botha J, et al. Aliskiren for geriatric lowering of systolic hypertension: a randomized controlled trial. *J Hum Hypertens* 2010;24:600-608.
116. Andersen K, Weinberger MH, Egan B, et al. Comparative efficacy and safety of aliskiren, an oral direct renin inhibitor, and ramipril in hypertension: a 6-month, randomized, double-blind trial. *J Hypertens*. 2008;26(3):589-99.
117. Miranda RD, Mion D, Rocha JC, et al. An 18-week, prospective, randomized, double-blind, multicenter study of amlodipine/ramipril combination versus amlodipine monotherapy in the treatment of hypertension: The assessment of combination therapy of amlodipine/ramipril (ATAR) study. *Clin Ther* 2008;30:1618-28.
118. Bönner G, Bakris GL, Sica D, et al. Antihypertensive efficacy of the angiotensin receptor blocker azilsartan medoxomil compared with the angiotensin-converting enzyme inhibitor ramipril. *J Hum Hypertens*. 2013 Aug;27(8):479-86.
119. Williams B, Lacourcière Y, Schumacher H, Gosse P, Neutel JM. Antihypertensive efficacy of telmisartan vs ramipril over the 24-h dosing period, including the critical early morning hours: a pooled analysis of the PRISMA I and II randomized trials. *J Hum Hypertens*. 2009 Sep;23(9):610-9.
120. O'Brien E, Barton J, Nussberger J, et al. Aliskiren reduces blood pressure and suppresses plasma renin activity in combination with a thiazide diuretic, an angiotensin-converting enzyme inhibitor, or an angiotensin receptor blocker. *Hypertension*. 2007 Feb;49(2):276-84.
121. Tytus RH, Burgess ED, Assouline L, Vanjaka A. A 26-week, prospective, open-label, uncontrolled, multicenter study to evaluate the effect of an escalating-dose regimen of trandolapril on change in blood pressure in treatment-naïve and concurrently treated adult hypertensive subjects (TRAIL). *Clin Ther*. 2007 Feb;29(2):305-15.
122. Tytus RH, Assouline L, Vanjaka A. Blood pressure control rates with an antihypertensive regimen including trandolapril in a Canadian usual-care setting. *Adv Ther*. 2011;28(9):789-798.

123. Pauly NC, Safar ME; for the Investigator Study Group. Comparison of the efficacy and safety of trandolapril and captopril for 16 weeks in mild-to-moderate essential hypertension. *J Cardiovasc Pharmacol.* 1994;23 Suppl 4:s73-6.
124. Vaur L, Dutrey-Dupagne C, Boussac J, et al. Differential effects of a missed dose of trandolapril and enalapril on blood pressure control in hypertensive patients. *J Cardiovasc Pharmacol.* 1995;Jul;26(1):127-31.
125. Karlberg BE, Andrup M, Oden A; on behalf of the Swedish Tarka Trialists. Efficacy and safety of a new long-acting drug combination, trandolapril/verapamil as compared to monotherapy in primary hypertension. *Blood Pressure* 2000;9:140-5.
126. Pepine CJ, Kowey PR, Kupfer S, et al. Predictors of adverse outcome among patients with hypertension and coronary artery disease. *J Am Coll Cardiol.* 2006;47(3):547-51.
127. Brunner M, Cooper-DeHoff RM, Gong Y, et al; for the INVEST Investigators. Factors influencing blood pressure response to trandolapril add-on therapy in patients taking verapamil SR (from the International Verapamil SR/Trandolapril [INVEST] Study). *Am J Cardiol.* 2007;99:1549-54.
128. Cifkova R, Nakov R, Novozamska E, et al. Evaluation of the effects of fixed combinations of sustained-release verapamil/trandolapril versus captopril/hydrochlorothiazide on metabolic and electrolyte parameters in patients with essential hypertension. *J Hum Hypertens* 2000;14:347-54.
129. de Leeuw PW, Notter T, Zilles P. Comparison of different fixed antihypertensive combination drugs: a double-blind, placebo-controlled parallel group study. *J Hypertens.* 1997 Jan;15(1):87-91.
130. Stanton AV, Gradman AH, Schmieder RE, et al. Aliskiren monotherapy does not cause paradoxical blood pressure rises. Meta-analysis of data from 8 clinical trials. *Hypertension* 2010;55:54-60.
131. Van Bortel LM, Fici F, Mascagni F. Efficacy and tolerability of nebivolol compared with other antihypertensive drugs: a meta-analysis. *Am J Cardiovasc Drugs.* 2008;8(1):35-44.
132. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig.* 2007;27(11):735-53.
133. Bakris GL, Sarafidis PA, Weir MR, Dahlöf B, Pitt B, Jamerson K, et al.; ACCOMPLISH Trial investigators. Renal outcomes with different fixed-dose combination therapies in patients with hypertension at high risk for cardiovascular events (ACCOMPLISH): a prespecified secondary analysis of a randomized controlled trial. *Lancet.* 2010 Apr 3;375(9721):1173-81.
134. Hou FF, Xie D, Zhang X, et al. Renoprotection of optimal antiproteinuric doses (ROAD) study: a randomized controlled study of benazepril and losartan in chronic renal insufficiency. *J Am Soc Nephrol.* 2007;18:1889-98.
135. Bakris GL, Toto RD, McCullough PA, et al; on behalf of GUARD (Gauging Albuminuria Reduction With Lotrel in Diabetic Patients With Hypertension) Study Investigators. Effects of different ACE inhibitor combinations on albuminuria: results of the GUARD study. *Kidney Int.* 2008 Jun;73(11):1303-9.
136. Esnault VL, Brown EA, Apetrei E, et al. The effects of amlodipine and enalapril on renal function in adults with hypertension and nondiabetic nephropathies: A 3-year, randomized, multicenter, double-blind, placebo-controlled study. *Clin Ther* 2008;30:482-98.
137. Barnett AH, Bain SC, Bouter P, et al. Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N Engl J Med.* 2004 Nov 4;351:1953-61.
138. Morgensen CE, Neldam S, Tikkanen I, et al. Randomized controlled trial of dual blockade of renin-angiotensin system in patients with hypertension, microalbuminuria, and non-insulin dependent diabetes: the candesartan and lisinopril microalbuminuria (CALM) study. *BMJ.* 2000 Dec 9;321:1440-4.
139. Fried LF, Emanuele N, Zhang JH, et al. Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med.* 2013 Nov 14;369(20):1892-903.
140. DREAM Trial Investigators. Effect of ramipril on the incidence of diabetes. *N Engl J Med.* 2006 Oct 12;355(15):1551-62.
141. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). Randomized placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. *Lancet.* 1997 Jun 28;349:1857-63.
142. Uresin Y, Taylor AA, Kilo C, et al. Efficacy and safety of the direct renin inhibitor aliskiren and ramipril alone or in combination in patients with diabetes and hypertension. *J Renin Angiotensin Aldosterone Syst* 2007;8:190-8.
143. Agodoa LY, Appel L, Bakris GL, et al; African American Study of Kidney Disease and Hypertension (AASK) Study Group. Effect of ramipril vs amlodipine on renal outcomes in hypertensive nephrosclerosis: a randomized controlled trial. *JAMA.* 2001 Jun 6;285(21):2719-28.

144. Wright JT Jr, Bakris G, Greene T, et al. Effects of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA*. 2002 Nov 20;288(19):2421-31.
145. Bianchi S, Bigazzi R, Campese VM. Intensive versus conventional therapy to slow the progression of idiopathic glomerular disease. *Am J Kidney Dis* 2010;55:671-81.
146. Chrysostomou A, Pedagogos E, MacGregor L, Becker GJ. Double-blind, placebo-controlled study on the effect of the aldosterone receptor antagonist spironolactone in patients who have persistent proteinuria and are on long-term angiotensin-converting enzyme inhibitor therapy, with or without an angiotensin II receptor blocker. *Clin J Am Soc Nephrol*. 2006 Mar;1(2):256-62.
147. Nakao N, Yoshimura A, Morita H, et al. Combination treatment of angiotensin-II receptor blocker and angiotensin-converting enzyme inhibitor in non-diabetic renal disease (COOPERATE): a randomized controlled trial. *Lancet*. 2003 Jan 11;361:117-24.
148. Ruggenenti P, Fassi A, Ilieva AP, et al; Bergamo Nephrologic Diabetes Complications Trial (BENEDICT) Investigators. *N Engl J Med*. 2004 Nov 4;351(19):1941-51.
149. Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet*. 2005 Dec 10;366:2026-33.
150. Strippoli GFM, Craig M, Deeks JJ, et al. Effects of angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists on mortality and renal outcomes in diabetic nephropathy: systematic review. *BMJ*. 2004 Sep 30; 329:828-38.
151. Strippoli GFM, Bonifati C, Craig M, Navaneethan DS, Craig JC. Angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists for preventing the progression of diabetic kidney disease (review). *Cochrane Database Syst Rev*. 2006 Oct 18;(4):CD006257.
152. Montalescot G, Drexler H, Gallo R, et al. Effect of irbesartan and enalapril in non-ST elevation acute coronary syndrome: results of the randomized, double-blind ARCHPELAGO study. *Euro Heart J* 2009;30:2733-41.
153. The Dream Trial Investigators. Results of the diabetes reduction assessment with ramipril and rosiglitazone medication (DREAM) trial: Effects of ramipril and rosiglitazone on cardiovascular and renal outcomes in people with impaired glucose tolerance or impaired fasting glucose. *Diabetes Care* 2008;31:1007-1014.
154. Belluzzi F, Sernesi L, Preti P, et al. Prevention of recurrent lone atrial fibrillation by the angiotensin II converting enzyme inhibitor ramipril in normotensive patients. *J Am Coll Cardiol* 2009;53:24-9.
155. Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomized trial. *Lancet* 1998;351:1755-62.
156. Taylor AA, Shoheiber O. Adherence to antihypertensive therapy with fixed-dose amlodipine besylate/benazepril HCl versus comparable component-based therapy. *Congest Heart Fail* 2003;9:324-32.
157. Dickson M, Plauschinat CA. Compliance with antihypertensive therapy in the elderly: a comparison of fixed-dose combination amlodipine/benazepril versus component-based free-combination therapy. *Am J Cardiovasc Drugs* 2008;1:45-50.
158. Dezii CM. A retrospective study of persistence with single-pill combination therapy vs concurrent two-pill therapy in patients with hypertension. *Managed Care* 2000;9:S2-6.
159. Sapienza S, Sacco P, Floyd K, et al. Results of a pilot pharmacotherapy quality improvement program using fixed-dose, combination amlodipine/benazepril antihypertensive therapy in a long-term care setting. *Clin Ther* 2003;25:1872-87.



**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Angiotensin II Receptor Antagonists  
AHFS Class 243208  
May 4, 2022**

**I. Overview**

The renin-angiotensin-aldosterone system (RAAS) is the most important component in the homeostatic regulation of blood pressure. Excessive activity of the RAAS may lead to hypertension, as well as fluid and electrolyte disorders.<sup>1,2</sup> Renin catalyzes the conversion of angiotensinogen to angiotensin I. Angiotensin I is then cleaved to angiotensin II by angiotensin-converting enzyme (ACE). Angiotensin II may also be generated through other pathways (angiotensin I convertase).<sup>1</sup> Angiotensin II can increase blood pressure by direct vasoconstriction, as well as through actions on the brain and autonomic nervous system.<sup>1,2</sup> In addition, angiotensin II stimulates aldosterone synthesis from the adrenal cortex, leading to sodium and water reabsorption. Angiotensin II exerts other detrimental effects, including ventricular hypertrophy, remodeling, and myocyte apoptosis.<sup>1,2</sup>

The angiotensin II receptor antagonists are approved for the treatment of diabetic nephropathy, heart failure, hypertension and post-myocardial infarction.<sup>3-20</sup> Since angiotensin II may be generated through other pathways that do not depend upon ACE, blockade of angiotensin II by ACE inhibitors is incomplete. Angiotensin II receptor antagonists block the angiotensin II receptor subtype AT<sub>1</sub>, preventing the negative effects of angiotensin II, regardless of its origin. They do not appear to affect bradykinin and may be an option for patients who cannot tolerate ACE inhibitors.<sup>20,21</sup> The angiotensin II receptor antagonists are available as single entity products, as well as in combination with hydrochlorothiazide or chlorthalidone. Hydrochlorothiazide inhibits the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride. Telmisartan and olmesartan are also available in combination with amlodipine, a nondihydropyridine calcium-channel blocking agent, which is a potent vasodilator.<sup>20,21</sup>

The angiotensin II receptor antagonists that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All single entity products with the exception of azilsartan are available generically. Fixed-dose combination products are available in a generic formulation with the exception of azilsartan-chlorthalidone. Byvalson<sup>®</sup> (neбиволол/valsartan) was discontinued in 2018. This class was last reviewed in February 2020.

**Table 1. Angiotensin II Receptor Antagonists Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Azilsartan	tablet	Edarbi <sup>®</sup>	none
Candesartan	tablet	Atacand <sup>®*</sup>	candesartan
Irbesartan	tablet	Avapro <sup>®*</sup>	irbesartan
Losartan	tablet	Cozaar <sup>®*</sup>	losartan
Olmesartan	tablet	Benicar <sup>®*</sup>	olmesartan
Telmisartan	tablet	Micardis <sup>®*</sup>	telmisartan
Valsartan	tablet	Diovan <sup>®*</sup>	valsartan
<b>Combination Products</b>			
Azilsartan and chlorthalidone	tablet	Edarbyclor <sup>®</sup>	none
Candesartan and hydrochlorothiazide	tablet	Atacand HCT <sup>®*</sup>	candesartan and hydrochlorothiazide
Irbesartan and hydrochlorothiazide	tablet	Avalide <sup>®*</sup>	irbesartan and hydrochlorothiazide
Losartan and hydrochlorothiazide	tablet	Hyzaar <sup>®*</sup>	losartan and hydrochlorothiazide
Olmesartan and amlodipine and	tablet	Tribenzor <sup>®*</sup>	olmesartan and amlodipine

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
hydrochlorothiazide			and hydrochlorothiazide
Olmesartan and hydrochlorothiazide	tablet	Benicar HCT®*	olmesartan and hydrochlorothiazide
Telmisartan and amlodipine	tablet	N/A	telmisartan and amlodipine
Telmisartan and hydrochlorothiazide	tablet	Micardis HCT®*	telmisartan and hydrochlorothiazide
Valsartan and hydrochlorothiazide	tablet	Diovan HCT®*	valsartan and hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the angiotensin II receptor antagonists are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Angiotensin II Receptor Antagonists**

Clinical Guideline	Recommendations
American College of Cardiology/American Heart Association: <b>2007 Chronic Angina Focused Update of the 2002 Guidelines for the Management of Patients With Chronic Stable Angina (2007)</b> <sup>22</sup>	<ul style="list-style-type: none"> <li>Aspirin should be started at 75 to 162 mg/day and continued indefinitely in all patients, unless contraindicated.</li> <li>Use of warfarin in conjunction with aspirin and/or clopidogrel is associated with an increased risk of bleeding and should be monitored closely.</li> <li>Patients with hypertension and established coronary artery disease (CAD) should be treated with blood pressure medication(s) as tolerated, including angiotensin-converting enzyme inhibitors (ACE inhibitors) and/or <math>\beta</math>-adrenergic antagonists (<math>\beta</math>-blockers) with the addition of other medications as needed to achieve blood pressure goals of &lt;140/90 or &lt;130/80 mm Hg for patients with chronic kidney disease or diabetes.</li> <li>Long-acting calcium-channel blocking agents or long-acting nitrates may be used if <math>\beta</math>-blockers are contraindicated. Immediate-release and short-acting dihydropyridine calcium channel blockers can increase adverse cardiac events and should not be used.</li> <li>Long-acting calcium channel blockers or long-acting nitrates may be used with <math>\beta</math>-blockers if initial treatment is not successful.</li> <li>ACE inhibitors should be used indefinitely in patients with a left ventricular ejection fraction (LVEF) of <math>\leq 40\%</math> and in those with hypertension, diabetes or chronic kidney disease, unless contraindicated.</li> <li>ACE inhibitors should also be used indefinitely in patients at lower risk (mildly reduced or normal LVEF in whom cardiovascular risk factors remain well controlled and revascularization has been performed), unless contraindicated.</li> <li>Angiotensin II receptor blockers (ARBs) are recommended in patients with hypertension, those who have an indication for an ACE inhibitor and are intolerant to them, who have heart failure, or who have had a myocardial infarction (MI) and have a LVEF of <math>\leq 40\%</math>.</li> <li>ARBs may be considered in combination with an ACE inhibitor for heart failure due to left ventricular systolic dysfunction.</li> <li>Aldosterone blockade is recommended in patients post-MI without significant renal dysfunction or hyperkalemia who are already receiving therapeutic doses of an ACE inhibitor and a <math>\beta</math>-blocker, have a LVEF <math>\leq 40\%</math> and have either diabetes or heart failure.</li> <li>It is beneficial to start and continue <math>\beta</math>-blocker therapy indefinitely in all patients who have had a MI, acute coronary syndrome or left ventricular dysfunction with or without heart failure symptoms, unless contraindicated.</li> </ul>

Clinical Guideline	Recommendations
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)</b><sup>23</sup></p>	<ul style="list-style-type: none"> <li>• Annual influenza vaccination is recommended in patients with cardiovascular disease.</li> </ul> <p><b>Pharmacological management of stable coronary artery disease (CAD) patients</b></p> <ul style="list-style-type: none"> <li>• The two aims of the pharmacological management of stable CAD patients are to obtain relief of symptoms and to prevent CV events.</li> <li>• Optimal medical treatment indicates at least one drug for angina/ischemia relief plus drugs for event prevention.</li> <li>• It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>• It is indicated to review the patient's response soon after starting therapy.</li> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin, consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</li> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief:             <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a beta-blocker and/or a non-DHP-calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</li> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</li> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischaemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and beta-blockers avoided.</li> </ul> </li> <li>• Event prevention:             <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><b>Treatment in patients with microvascular angina</b></p> <ul style="list-style-type: none"> <li>• It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>• <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>• Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>• ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>• Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>• Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy (DAPT).</li> <li>• Aspirin is recommended for elective stenting.</li> <li>• Clopidogrel is recommended for elective stenting.</li> <li>• Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>• GP IIb/IIIa antagonists should be considered for bailout situation only.</li> <li>• Platelet function testing or genetic testing may be considered in specific or high risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>• Prasugrel or ticagrelor may be considered in specific high risk situations of elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</li> <li>• Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>• Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>• Prasugrel or ticagrelor is not recommended in low risk elective stenting.</li> <li>• After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low</li> <li>• Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months</li> </ul> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all revascularized patients receive a secondary prevention and be scheduled for follow-up visit.</li> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after second generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/ American College of Cardiology Foundation/ American Heart Association/ American Association</p>	<p><u>Medical therapy to prevent MI and death in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> </ul>

Clinical Guideline	Recommendations
<p>for Thoracic Surgery/ Preventive Cardiovascular Nurses Association/ Society of Thoracic Surgeons: <b>Management of Stable Ischemic Heart Disease (2012)</b><sup>24</sup></p>	<ul style="list-style-type: none"> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</li> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</li> <li>• Patients should receive an annual influenza vaccine.</li> </ul> <p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> <li>• Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes (2014)</b><sup>25</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>• Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation <math>&lt; 90\%</math>, respiratory distress, or other high risk features of hypoxemia.</li> <li>• Anti-ischemic and analgesic medications             <ul style="list-style-type: none"> <li>○ Nitrates                 <ul style="list-style-type: none"> <li>▪ Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>▪ Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>▪ Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>○ Analgesic therapy                 <ul style="list-style-type: none"> <li>▪ In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>▪ Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>○ Beta-adrenergic blockers                 <ul style="list-style-type: none"> <li>▪ Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock,</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval <math>&gt;0.24</math> second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</p> <ul style="list-style-type: none"> <li>▪ In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>▪ Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> <p>○ Calcium channel blockers (CCBs)</p> <ul style="list-style-type: none"> <li>▪ In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval <math>&gt;0.24</math> seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</li> <li>▪ Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>▪ CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>▪ Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>▪ Immediate-release nifedipine should not be administered to patients with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</li> </ul> <p>○ Other anti-ischemic interventions</p> <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> <p>○ Cholesterol management</p> <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> <p>● Inhibitors of renin-angiotensin-aldosterone system</p> <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF <math>&lt;0.40</math> and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF <math>&lt;0.40</math> who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine <math>&gt;2.5</math> mg/dL in men or <math>&gt;2.0</math> mg/dL in women) or hyperkalemia (<math>K &gt;5.0</math> mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF <math>&lt;0.40</math>, diabetes mellitus, or heart failure.</li> </ul> <p>● Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy</p> <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by</li> </ul>

Clinical Guideline	Recommendations
	<p>a daily maintenance dose should be administered.</p> <ul style="list-style-type: none"> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include: <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatid or tirofiban.</li> </ul> </li> </ul> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>● Antiplatelet agents <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg, prasugrel 60 mg, or ticagrelor 180 mg.</li> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatid, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> </li> <li>● Anticoagulant therapy <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>● Timing of CABG in relation to use of antiplatelet agents <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</p> <ul style="list-style-type: none"> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>● Medications at discharge <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</li> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication type, purpose, dose, frequency, side effects, and duration of use.</li> <li>○ For patients who are post-NSTEMI-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> </li> <li>● Late hospital and post-hospital oral antiplatelet therapy <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTEMI-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTEMI-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTEMI-ACS <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTEMI-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTEMI-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the</b></p>	<p><b>Pharmacological treatment of ischemia</b></p> <ul style="list-style-type: none"> <li>● Sublingual or intravenous nitrates and early initiation of β-blocker treatment is recommended in patients with ongoing ischemic symptoms and without</li> </ul>



Clinical Guideline	Recommendations
<p><b>Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation (2020)</b><sup>26</sup></p>	<p>contraindications.</p> <ul style="list-style-type: none"> <li>• Continuation of chronic <math>\beta</math>-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>• Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>• In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and <math>\beta</math>-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> <li>• A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds. <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTEMI-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</li> </ul> </li> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH</li> </ul>

Clinical Guideline	Recommendations
	<p>is recommended during the procedure.</p> <ul style="list-style-type: none"> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul> <p><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score <math>\geq 2</math>, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is <math>&lt; 2.5</math> in VKA-treated patients.</li> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li>• If at low bleeding risk (HAS-BLED <math>\leq 2</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED <math>\geq 3</math>), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED <math>\geq 3</math> and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are</li> </ul>

Clinical Guideline	Recommendations
	<p>contraindications such as excessive risk of bleeding.</p> <ul style="list-style-type: none"> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>
<p>American College of Cardiology/American Heart Association: <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>27</sup></p>	<p><u>Routine medical therapies: <math>\beta</math>-blockers</u></p> <ul style="list-style-type: none"> <li>• Oral <math>\beta</math>-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral <math>\beta</math>-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li>• <math>\beta</math>-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>• Patients with initial contraindications to the use of <math>\beta</math>-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>• It is reasonable to administer intravenous <math>\beta</math>-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia.</li> </ul> <p><u>Routine medical therapies: renin-angiotensin-aldosterone system inhibitors</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor should be administered within the first 24 hours to all patients with ST-segment elevation myocardial infarction with anterior location, heart failure, or ejection fraction <math>\leq 40\%</math>, unless contraindicated.</li> <li>• An ARB should be given to patients who have indications for but are intolerant of ACE inhibitors.</li> <li>• ACE inhibitors are reasonable for all patients with no contraindications to their use.</li> <li>• An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq 40\%</math> and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>• High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>• It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>28</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of ST-elevation myocardial infarction (STEMI)</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>• Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>• A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>• In patients with an indication for oral anticoagulation, oral anticoagulants are</li> </ul>

Clinical Guideline	Recommendations
	<p>indicated in addition to antiplatelet therapy.</p> <ul style="list-style-type: none"> <li>• In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the estimated risk of recurrent coronary events and bleeding).</li> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age ≥50 years, and at least one of the following risk factors: age ≥65 years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR &lt;60 mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</li> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with β-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with β-blockers is indicated in patients with heart failure or left ventricular dysfunction, LVEF ≤40% unless contraindicated.</li> <li>• Intravenous β-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> <li>• Intravenous β-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of &lt;1.8 mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C &gt;1.8 mmol/L (&gt;70 mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction ≤40% and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
American College of	<u>Top 10 messages for the primary prevention of cardiovascular disease</u>

Clinical Guideline	Recommendations
<p>Cardiology/ American Heart Association: <b>Guideline on the Primary Prevention of Cardiovascular Disease (2019)</b><sup>29</sup></p>	<ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</li> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</li> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</li> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be <math>&lt;130/80</math> mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a decision is made for statin therapy, a moderate-intensity statin should be recommended.</li> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) patients, LDL-C levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</li> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt; 20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> </li> <li>• In patients at borderline risk (5% to <math>&lt; 7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul> <p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>• In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include: <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>• In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>• In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>• In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>• In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than</li> </ul>

Clinical Guideline	Recommendations
	<p>130/80 mm Hg.</p> <ul style="list-style-type: none"> <li>In adults with an estimated 10-year ASCVD risk &lt;10% and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering medication are recommended.</li> <li>In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul> <p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> <li>To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</li> <li>Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)<sup>30</sup></p>	<p><u>Treatment of Stage A heart failure (HF)</u></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>In patients with a recent or remote history of MI or ACS and LVEF≤40%,</li> </ul>

Clinical Guideline	Recommendations
	<p>evidence-based beta blockers should be used to reduce mortality. (LoE: B)</p> <ul style="list-style-type: none"> <li>• In patients who are at least 40 days post-MI with LVEF <math>\leq 30\%</math> and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF <math>\leq 40\%</math>, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF <math>\leq 50\%</math>, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF <math>\leq 50\%</math>, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to</li> </ul>



Clinical Guideline	Recommendations
	<p>improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to</li> </ul>

Clinical Guideline	Recommendations
	<p>prevent morbidity. (LoE: C)</p> <ul style="list-style-type: none"> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>31</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults</b> (2014)<sup>32</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type</li> </ul>

Clinical Guideline	Recommendations
	<p>diuretic, CCB, ACE inhibitor, or ARB classes should be added.</p> <ul style="list-style-type: none"> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>33</sup></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumption is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week. Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><u>Pharmacological Treatment</u></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> <li>● Core drug-treatment strategy               <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>34</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>● Initial therapy should be with either monotherapy or single pill combination (SPC).           <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are:               <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt;</math>60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>like diuretic monotherapy.</p> <ul style="list-style-type: none"> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial</u></p>

Clinical Guideline	Recommendations
	<p><b>infarction</b></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><b>Treatment of hypertension in association with heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction &lt;40%), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><b>Treatment of hypertension in association with stroke</b></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><b>Treatment of hypertension in association with LVH</b></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or</li> </ul>



Clinical Guideline	Recommendations
	<p><b>minoxidil should not be used.</b></p> <p><b>Treatment of hypertension in association with nondiabetic chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><b>Treatment of hypertension in association with renovascular disease</b></p> <ul style="list-style-type: none"> <li>Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amenable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><b>Treatment of hypertension in association with diabetes mellitus</b></p> <ul style="list-style-type: none"> <li>Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><b>Resistant hypertension</b></p> <ul style="list-style-type: none"> <li>Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>Accurate office and out-of-office BP measurement is essential.</li> <li>Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen</li> </ul>

Clinical Guideline	Recommendations
	<p>decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</p> <ul style="list-style-type: none"> <li>Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90th percentile in children with risk factors or target organ damage.</li> <li>Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>35</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater</li> </ul>

Clinical Guideline	Recommendations
	<p>effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</p> <ul style="list-style-type: none"> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (≥65 years) are treated with the same recommendations in non-older patient population. In very old patients (≥80 years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should be avoided.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq</math>220 mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq</math>140/90 mmHg.</li> <li>• In patients with HF<sub>r</sub>EF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HF<sub>p</sub>EF, BP treatment threshold and target values should be the same as for HF<sub>r</sub>EF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq</math>130 mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
National Institute for Health and Clinical	<u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u>

Clinical Guideline	Recommendations
<p>Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>36</sup></p>	<ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to</li> </ul>

Clinical Guideline	Recommendations
	<p>ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</p> <ul style="list-style-type: none"> <li>If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>37</sup></p>	<ul style="list-style-type: none"> <li>To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p><b>Kidney Disease Improving Clinical Outcomes Group:</b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the</li> </ul>

Clinical Guideline	Recommendations
<p><b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>38</sup></p>	<p>management of high BP in adults.</p> <ul style="list-style-type: none"> <li>The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults</b> (2017)<sup>39</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor,</li> </ul>



Clinical Guideline	Recommendations
	<p>angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</p> <ul style="list-style-type: none"> <li>• For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>• Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be <math>&lt; 130</math> mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP <math>&lt; 130/80</math> mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP <math>&lt; 130</math> mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal <math>&lt; 130/80</math> mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to</li> </ul>

Clinical Guideline	Recommendations
	<p>a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</p> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq</math>220/120 mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq</math>140/90 mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP <math>\geq</math>180 mmHg or DBP <math>\geq</math>110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>40</sup></p>	<p><b>Hypertension/blood pressure control</b></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq</math>140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq</math>180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq</math>15%), a blood pressure target of <math>&lt;</math>130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;</math>15%), treat to blood pressure target of <math>&lt;</math>140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;</math>120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><u>Chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"><li data-bbox="500 205 1414 268">• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate &lt;30 mL/min/1.73 m<sup>2</sup>.</li><li data-bbox="500 268 1414 321">• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li></ul>

\*Agent not available in the United States.

### III. Indications

The FDA-approved indications for the angiotensin II receptor antagonists are noted in Tables 3 and 4. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively on the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Angiotensin II Receptor Antagonists-Single Entity Agents<sup>3-20</sup>**

Indication(s)	Single Entity Agents							
	Azil-sartan	Cande-sartan		Irbe-sartan	Lo-sartan	Olme-sartan	Telmi-sartan	Val-sartan
<b>Cardiovascular Risk Reduction</b>								
Reduce the risk of cardiovascular mortality in clinically stable patients with left ventricular failure or left ventricular dysfunction following myocardial infarction								✓
Reduce the risk of myocardial infarction, stroke, or death from cardiovascular causes in patients ≥55 years of age at high risk of developing major cardiovascular events who are unable to take an angiotensin converting enzyme inhibitor							✓	
Reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy					✓ *			
<b>Heart Failure</b>								
Heart failure (New York Heart Association functional class II to IV)								✓
Heart failure (New York Heart Association functional class II to IV) in adults with left ventricular systolic dysfunction (ejection fraction ≤40%) to reduce cardiovascular death and to reduce heart failure hospitalizations		✓						
<b>Hypertension</b>								
Hypertension, alone or in combination with other antihypertensive agents	✓	✓	✓	✓	✓	✓	✓	✓
<b>Nephropathy in Type 2 Diabetic Patients</b>								
Diabetic nephropathy with an elevated serum creatinine and proteinuria (>300 mg/day) in patients with type 2 diabetes and hypertension				✓	✓			

\*There is evidence that this benefit does not apply to Black patients.

**Table 4. FDA-Approved Indications for the Angiotensin II Receptor Antagonists-Combination Products<sup>3-20</sup>**

Indication(s)	Combination Products								
	Azilsartan and chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmesartan and Amlodipine and HCTZ	Olmesartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
<b>Cardiovascular Risk</b>									

Indication(s)	Combination Products								
	Azilsartan and chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmесartan and Amlodipine and HCTZ	Olmесartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
<b>Reduction</b>									
Reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy				✓ *					
<b>Hypertension</b>									
Hypertension	✓	✓ †	✓	✓ §	✓ †	✓ †	✓ ‡	✓ †	✓

\*There is evidence that this benefit does not apply to Black patients.

†This fixed dose combination is not indicated for initial therapy.

‡May be used alone or in combination with other antihypertensive agents.

§This fixed dose combination is not indicated for initial therapy, except when the hypertension is severe enough that the value of achieving prompt blood pressure control exceeds the risk of initiating combination therapy in these patients.

HCTZ=hydrochlorothiazide



#### IV. Pharmacokinetics

The pharmacokinetic parameters of the angiotensin II receptor antagonists are listed in Table 5.

**Table 5. Pharmacokinetic Parameters of the Angiotensin II Receptor Antagonists<sup>21</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Azilsartan	60	>99	Liver (% not reported)	Feces (55) Renal (42)	11
Candesartan	15	>99	Intestinal wall (>99)	Feces (67) Renal (33)	9
Irbesartan	60 to 80	90	Liver (50 to 70)	Feces (65) Renal (20)	11 to 15
Losartan	25 to 35	99	Liver (14)	Feces (50 to 60) Renal (13 to 35)	2
Olmesartan	26	99	Intestinal wall (100)	Feces (50 to 65) Renal (35 to 50)	13
Telmisartan	42 to 58	>99	Liver (<3)	Feces (97)	24
Valsartan	25	95	Liver, minimal (% not reported)	Feces (83) Renal (13)	6 to 9
<b>Combination Products</b>					
Azilsartan and Chlorthalidone	60/not reported	>99/75	Liver (% not reported)/ Not reported	Feces (55) Renal (42)/ Renal, major (% not reported)	12/45
Candesartan and HCTZ	15/70	>99/40	Liver, minimal (% not reported)/ Not metabolized	Feces (67) Renal (26)/ Renal (61)	5.1 to 10.5/ 5.6 to 14.8
Irbesartan and HCTZ	60 to 80/not reported	90/40	Liver (% not reported)/ Not metabolized	Feces, majority (% not reported) Renal (20)/ Renal (61)	10 to 12/ 11 to 15
Losartan and HCTZ	33/not reported	Not reported/not reported	Systemic (% not reported)/ Not metabolized	Feces (60) Renal (35)/ Renal (% not reported)	2/ 5.6 to 14.8
Olmesartan and amlodipine and HCTZ	26/ 64 to 90/ Not reported	99/ 93/ Not reported	Intestinal wall, extensive/ Liver (90)/ Not reported	Feces (50 to 65) Renal (35 to 50)/ Renal (10)/ Renal (61)	13/ 30 to 50/ 5.6 to 14.8
Olmesartan and HCTZ	26/Not reported	99/Not reported	Hydrolysis (complete)/Not metabolized	Feces (% not reported) Renal (35 to 50)/ Renal (61)	13/ 5.6 to 14.8
Telmisartan and amlodipine	64 to 90/ 42 to 58	99.5/93	Hepatic, minimal (% not reported)/ Hepatic (90)	Feces (>97) Renal (<1)/ Feces (20 to 25) Renal (10)	24/ 30 to 50
Telmisartan and HCTZ	42 to 58/Not reported	Not reported/Not reported	Not reported/Not reported	Feces (97)/ Renal (61)	24/ 5.6 to 14.8
Valsartan and HCTZ	25/70	95/40 to 70	Liver, minimal (% not reported)/ Not reported	Feces (83) Renal (13)/ Renal (70)	6 to 9/ 10 to 12

HCTZ=hydrochlorothiazide

## V. Drug Interactions

Major drug interactions with the angiotensin II receptor antagonists are listed in Table 6.

**Table 6. Major Drug Interactions with the Angiotensin II Receptor Antagonists<sup>21</sup>**

Generic Name(s)	Interaction	Mechanism
ARBs (azilsartan, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan)	ACE inhibitors	Concurrent use of angiotensin converting enzyme inhibitors and ARBs may result in increased risk of adverse events (i.e., hypotension, syncope, hyperkalemia, changes in renal function, acute renal failure).
ARBs (azilsartan, irbesartan)	Fluconazole	Concurrent use of fluconazole and selected ARBs may result in increased exposure of ARB and increased risk of toxicity.
ARBs (azilsartan, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan)	Lithium	Angiotensin II receptor antagonists may decrease lithium renal excretion by enhancing its reabsorption. Lithium levels may increase, resulting in an increase in pharmacologic and toxic effects of lithium.
ARBs (azilsartan, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan)	Trimethoprim	Angiotensin II receptor antagonists and trimethoprim may act additively or synergistically to inhibit renal excretion of potassium, increasing the risk of hyperkalemia.
ARBs (azilsartan, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan)	Aliskiren	Concurrent use of aliskiren and ARBs may result in an increased risk of hyperkalemia, renal impairment, and hypotension.
Telmisartan	Digoxin	Concurrent use of digoxin and telmisartan may result in an increased risk of digoxin toxicity (nausea, vomiting, arrhythmias).
Dihydropyridines (amlodipine)	HIV protease inhibitors	Pharmacologic effects of amlodipine may be enhanced by protease inhibitors.
Dihydropyridines (amlodipine)	Imidazoles	Imidazoles may increase the plasma concentrations and pharmacologic effects of amlodipine.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics may induce hypokalemia which may increase the risk of torsades de pointes.
Thiazide diuretics (HCTZ)	Lithium	Thiazide diuretics decrease the renal clearance of lithium which leads to increased serum lithium levels. Lithium toxicity has occurred.
Thiazide diuretics (HCTZ)	Diazoxide	The combination of diazoxide with a thiazide diuretic may lead to hyperglycemia through an unknown mechanism; therefore the combination should be avoided.
Thiazide diuretics (HCTZ)	Digitalis glycosides	Diuretic-induced electrolyte disturbances may predispose the patient to digitalis-induced cardiac arrhythmias.

ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor antagonist, HCTZ=hydrochlorothiazide, HIV=human immunodeficiency virus

## VI. Adverse Drug Events

The most common adverse drug events reported with the angiotensin II receptor antagonists are listed in Table 7. The most common adverse drug events reported with amlodipine and hydrochlorothiazide are listed in Table 8. The boxed warning for the angiotensin II receptor antagonists is listed in Table 9.

**Table 7. Adverse Drug Events (%) Reported with the Combination Angiotensin II Receptor Antagonists-Single Entity Agents<sup>3-20</sup>**

Adverse Events	Single Entity Agents						
	Azilsartan	Candesartan	Irbesartan	Losartan	Olmesartan	Telmisartan	Valsartan
<b>Cardiovascular</b>							
Chest pain	-	>1	≥1	≥1	>0.5	1	-
Hypertension	-	-	<1	-	-	-	-
Hypotension	-	-	<1	<1	-	-	<1
Orthostatic hypotension	-	-	✓	-	-	-	-
Tachycardia	-	≥0.5	≥1	<1	>0.5	>0.3	-
<b>Central Nervous System</b>							
Anxiety/nervousness	-	≥0.5	≥1	<1	-	>0.3	>0.2
Depression	-	≥0.5	<1	<1	-	>0.3	-
Dizziness	≥0.3	4	≥1	4	3	1	>1
Dizziness, postural	≥0.3	-	-	-	-	-	-
Fatigue	-	>1	4	-	>0.5	1	2
Headache	-	≥1	≥1	≥1	>1	1	>1
Insomnia	-	-	-	1	>0.5	>0.3	>0.2
<b>Dermatological</b>							
Rash	-	≥0.5	≥1	<1	>0.5	>0.3	>0.2
<b>Gastrointestinal</b>							
Abdominal pain	-	>1	≥1	≥1	>0.5	1	2
Diarrhea	2	>1	3	2	>1	3	>1
Dyspepsia/heartburn	-	≥0.5	2	1	>0.5	1	>0.2
Nausea/vomiting	≥0.3	>1	≥1	≥1	>0.5	1	>1
<b>Genitourinary</b>							
Albuminuria	-	>1	-	-	-	-	-
Hematuria	-	≥0.5	-	-	>1	-	-
Urinary tract infection	-	-	≥1	<1	>0.5	1	-
<b>Laboratory Test Abnormalities</b>							
Creatine phosphokinase increased	-	≥0.5	-	-	>1	-	-
Decreased hematocrit	0.4	-	-	-	-	-	-
Decreased hemoglobin	0.2	-	-	-	-	-	-
Decreased red blood counts	0.3	-	-	-	-	-	-
Hyperglycemia	-	≥0.5	-	-	>1	-	-
Hyperkalemia	-	✓	✓	✓	-	-	✓
Hypertriglyceridemia	-	≥0.5	-	-	>1	-	-
<b>Musculoskeletal</b>							
Arthralgia	-	>1	-	<1	>0.5	>0.3	>1
Muscle cramp	-	-	-	1.1	-	-	>0.2
Muscle spasm	≥0.3	-	-	-	-	-	-

Adverse Events	Single Entity Agents						
	Azilsartan	Candesartan	Irbesartan	Losartan	Olmesartan	Telmisartan	Valsartan
Myalgia	-	≥0.5	-	1	>0.5	1	>0.2
Pain (includes back and leg)	-	3	≥1	1 to 2	>1	1 to 3	>0.2
Trauma	-	-	2	-	-	-	-
<b>Respiratory</b>							
Bronchitis	-	>1	-	<1	>1	>0.3	-
Cough	≥0.3	>1	3	3	-	1	>1
Influenza/influenza-like symptoms	-	-	≥1	<1	>1	1	-
Nasal congestion	-	-	-	2	-	-	-
Pharyngitis	-	2	≥1	≥1	>1	1	>1
Rhinitis	-	2	≥1	<1	>1	>0.3	>1
Sinus disorder	-	-	≥1	2	-	-	-
Sinusitis	-	>1	-	1	>1	3	>1
Upper respiratory tract infection	-	6	9	8	>1	7	>1
<b>Miscellaneous</b>							
Allergic reactions	-	✓	✓	✓	✓	✓	✓
Angioedema	-	✓	✓	✓	✓	✓	✓
Asthenia	≥0.3	-	-	-	-	-	-
Edema	-	>1	≥1	≥1	>0.5	1	>1
Fatigue	≥0.3	-	-	-	-	-	-
Inflicted injury	-	-	-	-	>1	-	-
Viral infection	-	-	-	-	-	-	3

✓ Percent not specified  
- Event not reported

**Table 8. Adverse Drug Events (%) Reported with the Combination Angiotensin II Receptor Antagonists-Combination Products<sup>3-20</sup>**

Adverse Event	Azilsartan and Chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmesartan and Amlodipine and HCTZ	Olmesartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
<b>Cardiovascular</b>									
Abnormal electrocardiogram	-	≥0.5	-	-	-	-	-	-	-
Angina	-	<0.5	-	-	-	-	-	-	-
Bradycardia	-	≥0.5	-	-	-	-	-	-	-
Chest pain	-	≥0.5	2	-	-	>1	-	-	>0.2
Extrasystoles	-	≥0.5	-	-	-	-	-	-	-
Hypotension	1.7	-	0.6 to 0.9	0.6	-	-	<2.0	<2	>0.2 to 1.0
Myocardial infarction	-	<0.5	-	-	-	-	-	-	-
Palpitations	-	≥0.5	-	1.4	-	-	-	-	>0.2
Syncope	0.3	-	-	-	1	-	<2.0	-	✓
Tachycardia	-	≥0.5	1	-	-	-	-	<2	>0.2
<b>Central Nervous System</b>									
Anxiety	-	≥0.5	≥1	-	-	-	-	-	>0.2
Asthenia	-	≥0.5	-	≥1	-	-	-	-	>0.2
Depression	-	≥0.5	-	-	-	-	-	-	✓
Dizziness	8.9	2.9	1 to 8	5.7	-	9	3.0	1 to 7	>0.2 to 6.0

Adverse Event	Azilsartan and Chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmesartan and Amlodipine and HCTZ	Olmesartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
Headache	-	2.9	1.0 to 5.5	≥1	6.4	>2	-	≥2	-
Hypesthesia	-	≥0.5	-	-	-	-	-	-	-
Insomnia	-	≥0.5	-	-	-	-	-	-	>0.2
Nervousness	-	-	≥1	-	-	-	-	-	-
Paresthesia	-	≥0.5	-	-	-	-	-	-	>0.2
Somnolence	-	-	-	-	-	-	-	-	>0.2
Vertigo	-	≥0.5	-	-	-	>1	-	-	>0.2
<b>Dermatological</b>									
Alopecia	-	-	-	-	-	✓	-	-	✓
Dermatitis	-	≥0.5	-	-	-	-	-	-	-
Eczema	-	≥0.5	-	-	-	-	-	-	-
Pruritus	-	≥0.5	-	-	-	✓	-	-	✓
Rash	-	≥0.5	≥1	1.4	-	>1	-	<2	>0.2
Sweating	-	≥0.5	-	-	-	-	-	-	>0.2
Urticaria	-	-	✓	-	-	✓	-	-	-
<b>Gastrointestinal</b>									
Abdominal pain	-	≥0.5	2	1.2	-	>1	-	<2	>0.2
Constipation	-	-	-	-	-	-	-	-	✓
Diarrhea	-	≥0.5	≥1	≥1	2.6	>1	-	3	>0.2
Dry mouth	-	-	-	-	-	-	-	-	>0.2
Dyspepsia	-	≥0.5	2	-	-	>1	-	<2	>0.2
Flatulence	-	-	-	-	-	-	-	-	>0.2
Gastritis	-	≥0.5	-	-	-	-	-	-	-
Gastroenteritis	-	≥0.5	-	-	-	>1	-	-	>0.2
Hepatic function abnormal	-	≥0.5	-	-	-	-	-	-	-
Hepatitis	-	-	✓	-	-	-	-	-	✓
Nausea	-	≥0.5	3	≥1	3.0	3	-	2	>0.2
Vomiting	-	≥0.5	3	-	-	✓	-	<2	>0.2
<b>Laboratory Test Abnormalities</b>									
Bilirubin increased	-	✓	-	✓	-	-	-	✓	-
Blood urea nitrogen increased	-	≥0.5	-	0.6	-	1.3	-	2.8	>0.2
Creatine phosphokinase increased	-	≥0.5	-	-	-	>1	-	-	-
Hematocrit decreased	-	✓	-	✓	-	0.4	-	0.6	-
Hemoglobin decreased	-	✓	-	✓	-	-	-	1.2	-
Hyperglycemia	-	≥0.5	-	-	-	>1	-	-	-
Hyperkalemia	-	-	0.2 to 1.2	-	-	✓	-	-	✓
Hyperlipidemia	-	-	-	-	-	>1	-	-	-
Hyperuricemia	-	≥0.5	-	-	-	4	-	-	-
Hypokalemia	-	≥0.5	0.6 to 0.9	-	-	-	-	<2	-
Serum creatinine increased	-	✓	-	0.8	-	-	-	1.4	-
Thrombocytopenia	-	-	-	✓	-	-	-	-	-
Transaminase levels increased	-	≥0.5	-	✓	-	>1	-	✓	✓
<b>Musculoskeletal</b>									
Arthralgia	-	≥0.5	-	-	-	>1	-	-	>0.2

Adverse Event	Azilsartan and Chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmesartan and Amlodipine and HCTZ	Olmesartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
Arthritis	-	≥0.5	-	-	-	>1	-	-	-
Arthrosis	-	≥0.5	-	-	-	-	-	-	-
Back pain	-	3.3	-	2.1	-	>1	2.2	<2	>0.2
Joint swelling	-	-	-	-	2.1	-	-	-	-
Leg cramps	-	≥0.5	-	-	-	-	-	-	-
Muscle cramps	-	-	>1	-	-	-	-	-	>0.2
Muscle spasms	-	-	-	-	3.1	-	-	-	-
Muscle weakness	-	-	-	-	-	-	-	-	✓
Musculoskeletal pain	-	-	6	-	-	-	-	-	-
Myalgia	-	≥0.5	-	-	-	>1	-	-	>0.2
Pain in extremity	-	-	-	-	-	-	-	-	>0.2
Rhabdomyolysis	-	-	✓	-	-	✓	-	-	✓
Sciatica	-	≥0.5	-	-	-	-	-	-	-
<b>Respiratory</b>									
Bronchitis	-	≥0.5	-	≥1	-	-	-	<2	>0.2
Bronchospasm	-	-	-	-	-	-	-	-	✓
Cough	-	≥0.5	≥1	2.6	-	>1	-	≥2	>0.2
Dyspnea	-	≥0.5	-	-	-	-	-	-	>0.2
Epistaxis	-	≥0.5	-	-	-	-	-	-	✓
Nasal congestion	-	-	-	-	-	-	-	-	>0.2
Nasopharyngitis	-	-	-	-	3.5	-	-	2.4	-
Pharyngitis	-	≥0.5	≥1	≥1	-	-	-	<2	✓
Pharyngolaryngeal pain	-	-	-	-	-	-	-	-	>0.2
Rhinitis	-	≥0.5	≥1	-	-	-	-	-	-
Sinus abnormality	-	-	≥1	-	-	-	-	-	-
Sinus congestion	-	-	-	-	-	-	-	-	>0.2
Sinusitis	-	≥0.5	-	1.2	-	-	-	4	>0.2
Upper respiratory tract infection	-	3.6	≥1	6.1	2.8	7	-	8	>0.2
<b>Miscellaneous</b>									
Abnormal vision	-	-	-	-	-	-	-	-	✓
Acute renal failure	-	-	-	-	-	✓	-	-	-
Allergy	-	-	1	-	-	-	-	-	-
Anaphylaxis	-	-	-	-	-	-	-	-	✓
Angioedema	-	<0.5	✓	✓	-	✓	-	-	✓
Appetite increased	-	-	-	-	-	-	-	-	✓
Conjunctivitis	-	≥0.5	-	-	-	-	-	-	-
Cystitis	-	≥0.5	-	-	-	-	-	-	-
Dehydration	-	-	-	-	-	-	-	-	✓
Dysuria	-	-	-	-	-	-	-	-	✓
Edema	-	-	3	1.3	-	-	<2.0	-	-
Erectile dysfunction	-	-	-	-	-	-	-	-	>0.2
Facial edema	-	-	-	-	-	✓	-	-	-
Fatigue	2.0	≥0.5	6	>1	4.2	-	-	3	>0.2
Fever	-	-	-	-	-	-	-	-	>0.2

Adverse Event	Azilsartan and Chlorthalidone	Candesartan and HCTZ	Irbesartan and HCTZ	Losartan and HCTZ	Olmesartan and Amlodipine and HCTZ	Olmesartan and HCTZ	Telmisartan and Amlodipine	Telmisartan and HCTZ	Valsartan and HCTZ
Flushing	-	-	-	-	-	-	-	-	✓
Gout	-	-	-	-	-	-	-	-	✓
Hematuria	-	≥0.5	-	-	-	>1	-	-	-
Inflicted injury	-	≥0.5	-	-	-	-	-	-	-
Influenza-like symptoms	-	2.5	3	-	-	-	-	2	>0.2
Infection	-	≥0.5	-	-	-	-	-	-	-
Libido decreased	-	-	-	-	-	-	-	-	✓
Pain	-	≥0.5	-	-	-	-	-	≥2	-
Peripheral edema	-	≥0.5	-	-	7.7	>1	4.8	-	>0.2
Pollakiuria	-	-	-	-	-	-	-	-	>0.2
Renal impairment	-	-	-	-	-	-	-	-	✓
Sunburn	-	-	-	-	-	-	-	-	✓
Tinnitus	-	≥0.5	-	-	-	-	-	-	>0.2
Urinary tract infection	-	≥0.5	≥1	-	2.4	>2	-	≥2	-
Urination abnormal	-	-	2	-	-	-	-	-	>0.2
Vasculitis	-	-	-	-	-	-	-	-	✓
Viral infection	-	≥0.5	-	-	-	-	-	-	✓

✓ Percent not specified

- Event not reported

HCTZ=hydrochlorothiazide

**Table 9. Boxed Warning for the Angiotensin II Receptor Antagonists<sup>20,21</sup>**

<b>WARNING</b>
When pregnancy is detected, discontinue therapy as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and even death to the developing fetus.

## VII. Dosing and Administration

The usual dosing regimens for the angiotensin II receptor antagonists are listed in Table 10.

**Table 10. Usual Dosing Regimens for the Angiotensin II Receptor Antagonists<sup>3-21</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Azilsartan	<u>Hypertension:</u> Tablet: initial, 40 or 80 mg once daily; maintenance, 80 mg once daily	Safety and efficacy in children have not been established.	Tablet: 40 mg 80 mg
Candesartan	<u>Heart Failure:</u> Tablet: initial, 4 mg once daily; maintenance, 32 mg once daily  <u>Hypertension:</u> Tablet: initial, 16 mg once daily when used as monotherapy in patients who are not volume-depleted; maintenance: 8 to 32 mg/day in a single or divided dose(s)	<u>Hypertension in children 1 to 6 years of age:</u> Tablet: initial, 0.2 mg/kg/day; maintenance, 0.05 to 0.4 mg/kg/day  <u>Hypertension in children 7 to 17 years of age and &lt;50 kg:</u> Tablet: initial, 4 to 8 mg/day; maintenance, 2 to 16 mg/day  <u>Hypertension in children 7 to 17 years of age and &gt;50 kg:</u> Tablet: initial, 8 to 16 mg/day; maintenance, 4 to 32 mg/day  Safety and efficacy in children with heart failure have not been established.	Tablet: 4 mg 8 mg 16 mg 32 mg
Irbesartan	<u>Diabetic nephropathy:</u> Tablet: 300 mg once daily  <u>Hypertension:</u> Tablet: initial, 150 mg once daily in patients who are not volume-depleted; maximum, 300 mg once daily	Safety and efficacy in children have not been established.	Tablet: 75 mg 150 mg 300 mg
Losartan	<u>Cerebrovascular risk reduction (hypertension and left ventricular hypertrophy):</u> Tablet: initial, 50 mg once daily; maintenance, 100 mg once daily  <u>Diabetic nephropathy:</u> Tablet: initial, 50 mg once daily; maintenance, dose should be increased to 100 mg once daily based on blood pressure response	<u>Hypertension in children ≥6 years of age:</u> Tablet: initial, 0.7 mg/kg once daily (up to 50 mg total); maximum, >1.4 mg/kg/day (or in excess of 100 mg) have not been studied  Safety and efficacy in children <6 years of age have not been established.	Tablet: 25 mg 50 mg 100 mg



Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<u>Hypertension:</u> Tablet: initial, 50 mg once daily in patients who are not volume-depleted; maintenance, 25 to 100 mg/day in a single or divided dose(s)		
Olmesartan	<u>Hypertension:</u> Tablet: initial, 20 mg once daily when used as monotherapy in patients who are not volume depleted; maximum, 40 mg once daily	<u>Hypertension in children 6 to 16 years of age and 20 to &lt;35 kg:</u> Tablet: initial, 10 mg once daily; maximum, 20 mg once daily  <u>Hypertension in children 6 to 16 years of age &gt;35 kg:</u> Tablet: initial, 20 mg once daily; maximum, 40 mg once daily  Safety and efficacy in children <6 years of age have not been established.	Tablet: 5 mg 20 mg 40 mg
Telmisartan	<u>Cardiovascular risk reduction:</u> Tablet: 80 mg once daily  <u>Hypertension:</u> Tablet: initial, 40 mg once daily; maximum: 80 mg per day	Safety and efficacy in children have not been established.	Tablet: 20 mg 40 mg 80 mg
Valsartan	<u>Cardiovascular risk reduction (post-myocardial infarction):</u> Tablet: initial, 20 mg twice daily; maintenance: 160 mg twice daily  <u>Heart Failure:</u> Tablet: Initial, 40 mg twice daily; maintenance, up titrate to 80 to 160 mg twice daily; maximum, 320 mg in divided doses  <u>Hypertension:</u> Tablet: initial, 80 to 160 mg once daily when used as monotherapy in patients who are not volume depleted; maintenance, 80 to 320 mg once daily	<u>Hypertension in children 1 to 16 years of age:</u> Tablet: initial, 1 mg/kg once daily (up to 40 mg total) administered as a tablet or suspension; maximum, >4 mg/kg/day (or in excess of 160 mg) have not been studied  <u>Safety and efficacy in children with hypertension &lt;1 years of age, or with heart failure, or for cardiovascular risk reduction have not been established.</u>	Tablet: 40 mg 80 mg 160 mg 320 mg
<b>Combination Products</b>			
Azilsartan and chlorthalidone	<u>Hypertension:</u> Tablet: initial, 40-12.5 mg once daily; maintenance, 40-25 mg once daily; maximum, 40-25 mg /day	Safety and efficacy in children have not been established.	Tablet: 40-12.5 mg 40-25 mg
Candesartan and HCTZ	<u>Hypertension:</u> Tablet: 16-12.5 to 32-25 mg/day	Safety and efficacy in children have not been established.	Tablet: 16-12.5 mg 32-12.5 mg 32-25 mg

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Irbesartan and HCTZ	<u>Hypertension:</u> Tablet: initial, 150-12.5 mg once daily; maximum, 300-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 150-12.5 mg 300-12.5 mg
Losartan and HCTZ	<u>Cerebrovascular risk reduction (hypertension and left ventricular hypertrophy):</u> Tablet: initial, 50-12.5 mg once daily; maintenance, 100-12.5 mg once daily; maximum, 100-25 mg once daily  <u>Hypertension:</u> Tablet: initial, 50-12.5 mg once daily; maintenance, 100-12.5 mg once daily; maximum, 100-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 50-12.5 mg 100-12.5 mg 100-25 mg
Olmesartan and amlodipine and HCTZ	<u>Hypertension:</u> Tablet: maximum, 40-10-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 20-5-12.5 mg 40-5-12.5 mg 40-5-25 mg 40-10-12.5 mg 40-20-25 mg
Olmesartan and HCTZ	<u>Hypertension:</u> Tablet: 20-12.5 to 40-25 mg/day	Safety and efficacy in children have not been established.	Tablet: 20-12.5 mg 40-12.5 mg 40-25 mg
Telmisartan and amlodipine	<u>Hypertension:</u> Tablet: initial, 40-5 or 80-5 mg once daily; maintenance, titrate as needed; maximum , 80-10 mg once daily	Safety and efficacy in children have not been established.	Tablet: 40-5 mg 40-10 mg 80-5 mg 80-10 mg
Telmisartan and HCTZ	<u>Hypertension:</u> Tablet: 40-12.5 to 80-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 40-12.5 mg 80-12.5 mg 80-25 mg
Valsartan and HCTZ	<u>Hypertension:</u> Tablet: 80-12.5 to 320-25 mg once daily	Safety and efficacy in children have not been established.	Tablet: 80-12.5 mg 160-12.5 mg 160-25 mg 320-12.5 mg 320-25 mg

HCTZ=hydrochlorothiazide

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the angiotensin II receptor antagonists are summarized in Table 11.

**Table 11. Comparative Clinical Trials with the Angiotensin II Receptor Antagonists**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Cardiovascular Risk Reduction</b>				
Pfeffer et al. <sup>41</sup> (2003) VALIANT  Captopril 50 mg TID  vs  valsartan 160 mg BID  vs  valsartan 80 mg BID and captopril 50 mg TID	DB, MC, RCT  Patients ≥18 years of age with an acute MI that was complicated by clinical or radiologic signs of heart failure and/or evidence of left ventricular systolic dysfunction	N=14,703  24.7 months	Primary: All-cause mortality  Secondary: Death from cardiovascular causes, recurrent MI, hospitalization for heart failure	Primary: No significant difference in all-cause mortality was reported between valsartan monotherapy and captopril monotherapy (P=0.98).  No significant difference in all-cause mortality was observed between valsartan plus captopril combination therapy and captopril monotherapy (P=0.73).  Secondary: The rate of death from cardiovascular causes, reinfarction, or hospitalization for heart failure was not significantly different between valsartan and captopril monotherapy (P=0.20).  The rate of death from cardiovascular causes, reinfarction, or hospitalization for heart failure was not significantly different between valsartan and captopril combination therapy and captopril monotherapy (P=0.37).  Combination therapy had the most drug-related adverse events. With monotherapy, hypotension and renal dysfunction were more common in the valsartan group and cough, rash, and taste disturbance were more common in the captopril group.
Dickstein et al. <sup>42</sup> (2002) OPTIMAAL  Captopril 50 mg TID  vs  losartan 50 mg QD	DB, MC, PG, RCT  Patients ≥50 years with an acute MI and signs or symptoms of heart failure during the acute phase or a new Q-wave anterior infarction or	N=5,477  2.7 years (mean)	Primary: All-cause mortality  Secondary: Composite of sudden cardiac death or resuscitated cardiac arrest	Primary: No significant difference in all-cause mortality was reported between patients receiving losartan and captopril (18 vs 16%, respectively; RR, 1.13; 95% CI, 0.99 to 1.28; P=0.07).  Secondary: No significant difference in sudden cardiac death or resuscitated cardiac arrest was reported between patients receiving losartan and captopril (9 vs 7%; RR, 1.19; 95% CI, 0.98 to 1.43; P=0.07).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	reinfarction			Losartan was significantly better tolerated than captopril, with fewer patients discontinuing study medication (17 vs 23%; P<0.0001).
Graham et al. <sup>43</sup> (2014)  Olmesartan  vs  other ARBs	Cohort, RETRO  Medicare patients (age 65 years or older) who filled at least one ARB prescription and had no recorded prescription for an ACE inhibitor or ARB 6 months prior to initiating a study drug. Results also stratified by diabetes or no diabetes	N=882,727  Mean duration of study drug use was 130 days	Primary: Acute MI, stroke, and all-cause mortality enriched for acute cardiovascular death  Secondary: Not reported	Primary: In the combined study population, there was no difference in the HRs for acute MI or stroke. The HR for death was reduced for olmesartan compared with other ARB users (HR, 0.82; 95% CI, 0.73 to 0.93; P=0.002).  In strata defined by presence or absence of diabetes, there was no difference in risk between users of olmesartan and other ARBs for any study endpoint at lower doses of therapy, regardless of duration of use. With high-dose therapy, the risk of acute MI was nonsignificantly increased in diabetic patients treated for 6 months or longer with olmesartan. For nondiabetic patients, the risk of acute MI was statistically significantly reduced with high-dose olmesartan over all durations combined. There was no effect of dose or duration on stroke risk with olmesartan in diabetic or nondiabetic patients.  Mortality risk was increased in diabetic patients treated with high-dose olmesartan for 6 months or longer (HR, 2.03; 95% CI, 1.09 to 3.75; P=0.02) and was reduced in nondiabetic patients during the first 6 months of olmesartan use (HR, 0.72; 95% CI, 0.55 to 0.96; P=0.02), and with use of 6 months or longer (HR, 0.46; 95% CI, 0.24 to 0.86; P=0.01).  Secondary: Not reported
<b>Diabetes/Diabetic Nephropathy/Renal Disease</b>				
White et al. <sup>44</sup> (2016)  Azilsartan medoxomil 40 or 80 mg  vs  olmesartan 40 mg  vs	Pooled analysis of 3 DB, PC or AC, RCTs  Patients with impaired fasting glucose (prediabetes mellitus) and T2DM	N=3821  6 to 8 weeks	Primary: Changes from baseline in both 24-h and clinic SBP  Secondary: Safety and tolerability	Primary: Baseline 24-h mean SBPs were approximately 145 and 146mmHg in the prediabetes mellitus and T2DM subgroups, respectively; corresponding clinic SBPs were approximately 158 and 159mmHg. Baseline HbA1c values for each subgroup were normoglycemic, 5.3%; prediabetes mellitus, 6.0%; and T2DM, 6.9%. Changes from baseline in 24-h or clinic SBP were significantly greater with azilsartan, 80mg compared with either olmesartan 40mg or valsartan 320mg in all subgroups in each pool.  Secondary: Safety and tolerability were similar among the active treatment and placebo subgroups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>valsartan 320 mg</p> <p>Mogensen et al.<sup>45</sup> (2000) CALM</p> <p>Lisinopril 20 mg QD</p> <p>vs</p> <p>candesartan 16 mg QD</p> <p>vs</p> <p>lisinopril 20 mg QD plus candesartan 16 mg QD</p> <p>Patients received 12 weeks monotherapy followed by an additional 12 weeks of monotherapy or combination therapy.</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients 30 to 75 years old with HTN, type 2 diabetes, and microalbuminuria</p>	<p>N=199</p> <p>24 weeks</p>	<p>Primary: Blood pressure and urinary albumin:creatinine ratio</p> <p>Secondary: Not reported</p>	<p>Primary: At 12 weeks, mean reductions in DBP were 9.7 mm Hg (P&lt;0.001) and 9.5 mm Hg (P&lt;0.001), respectively, and in urinary albumin:creatinine ratio were 46% (P&lt;0.001) and 30% (P&lt;0.001) for lisinopril and candesartan, respectively.</p> <p>Compared to either agent alone, at 24 weeks the combination of lisinopril plus candesartan resulted in 16.3 mm Hg reduction in mean DBP vs 10.4 mm Hg for candesartan alone (P&lt;0.001) and 10.7 mm Hg for lisinopril alone (P&lt;0.001).</p> <p>The reduction in urinary albumin:creatinine ratio with combination treatment (50%) was greater than with lisinopril alone (39%; P&lt;0.001) and candesartan alone (24%; P=0.05).</p> <p>All treatments were generally well tolerated.</p> <p>Secondary: Not reported</p>
<p>Lewis et al.<sup>46</sup> (2001) IDNT</p> <p>Irbesartan 300 mg/day</p>	<p>DB, MC, PC, PRO, RCT</p> <p>Patients 30 to 70 years old, with type 2 diabetes mellitus, HTN, and</p>	<p>N=1,715</p> <p>2.6 years</p>	<p>Primary: Composite of risk of doubling serum creatinine, ESRD, or death from any cause</p>	<p>Primary: Compared to placebo, irbesartan 300 mg/day resulted in a 20% lower relative risk of the composite primary outcome (P=0.02). Irbesartan treatment was associated with a 33% lower risk of doubling serum creatinine (P=0.003) and 23% trend towards lower risk of ESRD (P=0.07) compared to placebo. There was no significant difference in risk of death from any cause for irbesartan compared to placebo (P=0.57).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs amlodipine 10 mg/day vs placebo	nephropathy		Secondary: Composite of death from cardiovascular causes, nonfatal MI, heart failure requiring hospitalization, permanent neurologic deficit caused by a cerebrovascular event, or lower limb amputation	Compared to amlodipine, irbesartan treatment resulted in a 23% lower risk of composite primary outcome (P=0.006). Irbesartan treatment was associated with a 37% lower risk of doubling serum creatinine vs amlodipine (P<0.001) and 23% trend towards lower risk of ESRD vs amlodipine (P=0.07). There was no significant difference in risk of death from any cause (P=0.80).  Secondary: There were no significant differences in the secondary cardiovascular composite end point (P=0.40 and P=0.79 for irbesartan vs placebo and amlodipine, respectively).
Parving et al. <sup>47</sup> (2001) IRMA2  Irbesartan 150 or 300 mg/day  vs placebo	DB, MC, PC, RCT  Patients with HTN, type 2 diabetes mellitus and microalbuminuria	N=590  2 years	Primary: Time to onset of diabetic nephropathy  Secondary: Changes in level of albuminuria and creatinine clearance and restoration of normoalbuminuria	Primary: The primary end point was reached in 5.2% of patients in the irbesartan 300 mg group (P<0.001) and 9.7% of patients in the irbesartan 150 mg group (P=0.08) compared to 14.9% of patients receiving placebo.  Secondary: Irbesartan reduced the level of urinary albumin excretion by 38% in patients receiving the 300 mg dose and 24% in patients receiving the 150 mg dose vs 2% for placebo (P<0.001 for the combined irbesartan groups vs placebo and P<0.001 for the 300 vs 150 mg doses).  There was no significant difference in the decline in creatinine clearance among the 3 groups.  Restoration of normoalbuminuria was observed in 34% of patients receiving irbesartan 300 mg (P=0.006), 24% of patients receiving irbesartan 150 mg and 21% with placebo.
Persson et al. <sup>48</sup> (2009)  Irbesartan 300 mg QD  vs	DB, RCT, XO  Adults with type 2 diabetes, HTN, and albuminuria	N=26  Four 2-month treatment periods	Primary: Albuminuria (urinary albumin excretion rate)  Secondary: 24-hour blood	Primary: Treatment with aliskiren led to a significant reduction in albuminuria by 48% compared to placebo (P<0.001). Treatment with irbesartan led to a significant reduction in albuminuria by 58% compared to placebo (P<0.001). There was no significant difference in albuminuria between aliskiren and irbesartan (P value not reported). The combination of aliskiren and irbesartan significantly reduced albuminuria by 71% compared to placebo (P<0.001), which was also

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>aliskiren 300 mg QD</p> <p>vs</p> <p>aliskiren 300 mg QD and irbesartan 300 mg QD</p> <p>vs</p> <p>placebo</p>			<p>pressure and GFR</p>	<p>significantly better than with monotherapy (P&lt;0.001 for aliskiren and P=0.028 for irbesartan).</p> <p>Secondary: SBP and DBP 24-hr blood pressure were reduced by 3 and 4 mm Hg, respectively by aliskiren (P value not significant and P=0.009, respectively), 12 and 5 mm Hg, respectively by irbesartan (P&lt;0.001 and P=0.002, respectively), and 10 and 6 mm Hg, respectively with the combination (P=0.001 and P &lt;0.001, respectively) compared to placebo. There was no significant change in 24-hr blood pressure with irbesartan compared to combination therapy.</p> <p>GFR was significantly reduced 4.6 mL/min/1.73 m<sup>2</sup> with aliskiren (P=0.037), 8.0 mL/min/1.73 m<sup>2</sup> with irbesartan (P&lt;0.001), and 11.7 mL/min/1.73 m<sup>2</sup> with the combination (P&lt;0.001) compared to placebo.</p>
<p>Chrysostomou et al.<sup>49</sup> (2006)</p> <p>Ramipril 5 mg/day plus spironolactone 25 mg/day and placebo</p> <p>vs</p> <p>ramipril 5 mg/day plus irbesartan 150 mg/day and placebo</p> <p>vs</p> <p>ramipril 5 mg/day plus placebo and placebo</p>	<p>DB, PC, RCT</p> <p>Patients 18 to 75 years of age, with a 24 hour urinary protein excretion &gt;1.5 g/24 hours on ≥2 occasions ≥3 months apart, serum creatinine level ≤200 μmol/L with &lt;20% variability in the preceding 3 months and treatment with an ACE inhibitor ≥6 months</p>	<p>N=41</p> <p>6 months</p>	<p>Primary: Change in 24 hour urinary protein excretion at three months</p> <p>Secondary: Change in 24 hour urinary protein excretion at six months, change in blood pressure and creatinine clearance, adverse effects</p>	<p>Primary: Compared to ramipril-treated patients, the 24 hour urinary protein excretion reduction at three months was significantly greater in ramipril plus spironolactone-treated patients (P=0.004).</p> <p>Ramipril-, irbesartan- and spironolactone-treated patients exhibited a significant reduction in 24 hour urinary protein excretion compared to ramipril-treated patients (P&lt;0.001).</p> <p>There was no significant difference in 24 hour urinary protein excretion with ramipril- and ramipril plus irbesartan-treated patients (P=1.00).</p> <p>At three months, spironolactone-treated patients exhibited a significant reduction in proteinuria from baseline (P≤0.001). In contrast, non-spironolactone-treated patients did not experience a significant reduction in proteinuria from baseline (P=0.840).</p> <p>Secondary: At six months, spironolactone-treated patients exhibited the greatest reduction in proteinuria compared to the other treatments (P&lt;0.05).</p> <p>At six months, DBP was higher among ramipril monotherapy-treated patients</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  spironolactone 25 mg/day plus irbesartan 150 mg/day and ramipril 5 mg/day				compared to the other treatments (P=0.046). There was no difference in SBP among the treatments (P value not reported).  There were no differences in creatinine clearance among the treatments (P>0.05).  Gynecomastia was not observed with any of the treatments.
Bianchi et al. <sup>50</sup> (2010)  Ramipril 10 mg and atorvastatin 10 mg QD (conventional therapy)  vs  spironolactone 25 mg, ramipril 10 mg, irbesartan 300 mg, and atorvastatin 10 mg QD (intensive therapy)  The addition of diuretics, calcium antagonists, $\beta$ -blockers or $\alpha$ 1-receptor antagonists were added to achieve blood pressure <130/80 mm Hg	RCT, OL  Patients with a clinical diagnosis of idiopathic chronic glomerulonephritis and urine protein-creatinine ratio >1 g/g	N=128  36 months	Primary: Changes over time in proteinuria and eGFR  Secondary: Adverse events, drop outs	Primary: SBP decreased more in the intensive-therapy group (from 156.6 to 113.5 mm Hg) than in the conventional therapy group (from 155.7 to 122.7 mm Hg; P<0.01).  Urine protein excretion decreased from 2.65 to 0.45 g/g creatinine with intensive therapy (P<0.001). With conventional therapy, urine protein excretion decreased from 2.60 to 1.23 g/g creatinine (P<0.001).  With intensive therapy, eGFR did not significantly change over time (64.6 vs 62.9 mL/min/1.73 m <sup>2</sup> ). With conventional therapy, eGFR decreased from 62.5 to 55.8 mL/min/1.73 m <sup>2</sup> (P<0.01).  Secondary: In the conventional therapy group, eight patients discontinued the study due to hyperkalemia, cough, and rapid deterioration in kidney function. In the intensive therapy group, 15 dropped out due to hyperkalemia, cough, and hypotension. Nine patients in the intensive therapy group developed gynecomastia. Twelve patients on conventional and 31 on intensive therapy had to interrupt the study temporarily because of low blood pressure. No patient developed an increase in creatine kinase, alanine aminotransferase, and alkaline phosphatase levels during the study.
Brenner et al. <sup>51</sup> (2001)	DB, PC, RCT	N=1,513	Primary: Composite of risk of	Primary: Compared to placebo, losartan resulted in a 16% reduction of composite



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
RENAAL  Losartan 50 to 100 mg QD  vs  placebo	Patients 31 to 70 years of age with HTN, type 2 diabetes mellitus and nephropathy on conventional antihypertensive therapy	3.4 years	doubling of serum creatinine, ESRD, or death from any cause  Secondary: Composite of morbidity and mortality from cardiovascular causes, proteinuria, rate of progression of renal disease	primary end point (P=0.02).  Losartan treatment produced a 25% reduction of doubling serum creatinine vs placebo (P=0.006) and 28% reduction in ESRD vs placebo (P=0.002).  No differences in mortality were reported (P=0.88).  Secondary: There was no significant difference between the losartan and placebo groups in the composite end point of morbidity and mortality from cardiovascular causes.  Losartan treatment led to an average reduction in the level of proteinuria by 35% (P<0.001 vs placebo).  Losartan reduced the rate of decline in renal function by 18% (P=0.01 vs placebo).
Kiernan et al. <sup>52</sup> (2015) HEAAL  Losartan 50 mg  vs  losartan 150 mg	DB, MC, RCT  Patients with HF <sub>r</sub> EF, NYHA functional class II to IV; LVEF ≤40%; stable cardiovascular medical therapy for at least two weeks; and known intolerance to ACEIs	N=3,843	Primary: eGFR levels, SCr, renal function and association with clinical outcomes  Secondary: Not reported	Primary: Compared with 50 mg, 150 mg losartan led to a greater reduction in eGFR across time (mean difference, -3.76 ml/min/1.73 m <sup>2</sup> ; P<0.0001). This difference was driven by early changes, and differences in eGFR after four months were not significant (mean difference, 0.42 ml/min/1.73 m <sup>2</sup> ; P=0.15). Although an increase in SCr >0.3 mg/dL from baseline was associated with increased risk of death or hospitalization for HF (HR, 1.36; P<0.0001), the relationship was not significant if the change occurred before four months (HR, 1.09; P=0.20). Despite increased risk of worsening renal function, 150 mg losartan was associated with reduced risk of death or hospitalization for HF compared with 50 mg (HR, 0.85; P<0.0001).  Secondary: Not reported
Hou et al. <sup>53</sup> (2007) ROAD  Benazepril 10 mg/day vs	OL, PRO, RCT  Patients aged 18 to 70 years with proteinuria and chronic renal	N=360  3.7 years (median follow-up)	Primary: Time to composite of doubling of serum creatinine, ESRD or death	Primary: Compared to the conventional dosages, optimal antiproteinuric dosages of benazepril and losartan that were achieved through up-titration were associated with a 51 and 53% reduction in the risk for the primary end point (P=0.028 and P=0.022, respectively).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>individual up-titration (10 to 40 mg/day with median dose of 20 mg/day)</p> <p>or</p> <p>losartan 50 mg/day vs individual up-titration (50 to 200 mg/day with median dose of 100 mg/day)</p> <p>Up-titration was performed to optimal antiproteinuric and tolerated dosages, and then these dosages were maintained.</p>	<p>insufficiency who did not have diabetes</p>		<p>Secondary: Changes in level of proteinuria, rate of progression of renal disease</p>	<p>There was no statistically significant difference between benazepril and losartan in the overall relative risk reduction at their respective optimal antiproteinuric dosages or at conventional dosages.</p> <p>Secondary: Optimal antiproteinuric dosages of benazepril and losartan at comparable blood pressure control, achieved a greater reduction in both proteinuria and the rate of decline in renal function compared to their conventional dosages.</p> <p>There was no significant difference in proteinuria reduction between benazepril and losartan at both conventional and optimal antiproteinuric dosages. Changes in renal function were similar between benazepril and losartan arms at both conventional and optimal antiproteinuric doses (P&gt;0.05).</p> <p>There was no significant difference for the overall incidence of major adverse events between groups that were given conventional and optimal dosages in any of the treatment arms.</p>
<p>Fried et al.<sup>54</sup> (2013) VA NEPHRON-D</p> <p>Losartan with lisinopril</p> <p>vs</p> <p>losartan alone</p>	<p>DB, MA, RCT</p> <p>Veterans with proteinuric diabetic kidney disease, an estimated GFR of 30.0 to 89.9 ml/minute/1.73 m<sup>2</sup>, and a urinary albumin-to-creatinine ratio of ≥300</p>	<p>N=1448</p> <p>Median follow-up 2.2 years</p>	<p>Primary: First occurrence of a decline in the eGFR (an absolute decrease of ≥30 ml/minute/1.73 m<sup>2</sup> if the eGFR was ≥60 ml/minute/1.73 m<sup>2</sup> at randomization or a relative decrease of ≥50% if the eGFR was &lt;60 ml/minute/1.73 m<sup>2</sup>), ESRD, or death</p>	<p>The trial was stopped early because the absolute risk of serious adverse events appeared to be greater than the potential benefit of reducing primary end-point events.</p> <p>Primary: There were 152 primary end-point events in the monotherapy group (21.0%) and 132 in the combination-therapy group (18.2%). The risk of the primary end point did not differ significantly between the two groups.</p> <p>Secondary: There were 101 secondary end-point events (a decline in the estimated GFR or ESRD) in the monotherapy group (14.0%) and 77 events in the combination-therapy group (10.6%). There was no significant between-group difference in mortality or ESRD (Table 2), though the number of ESRD events was small.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>Secondary: First occurrence of a decline in the eGFR or ESRD</p> <p>Tertiary: CV events, slope of change in eGFR, and change in albuminuria at 1 year</p>	<p>Tertiary: There was no significant difference in the rate of cardiovascular events between the two groups. There was no significant difference in treatment effect on the decline in the estimated GFR (P=0.17). During adjustment of the losartan dose, the median urinary albumin-to-creatinine ratio declined from 959 to 807 (P=0.001). There was a further decline from randomization to 1 year, with a greater decline in the combination-therapy group (from 786 to 517) than in the monotherapy group (from 829 to 701) (P&lt;0.001).</p>
<p>Nakao et al.<sup>55</sup> (2003) COOPERATE</p> <p>Trandolapril 3 mg/day</p> <p>vs</p> <p>losartan 100 mg/day</p> <p>vs</p> <p>trandolapril and losartan at equivalent doses</p>	<p>DB, MC, PC, RCT</p> <p>Patients aged 18 to 70 years with chronic nephropathy (nondiabetic renal disease)</p>	<p>N=263</p> <p>3 years</p>	<p>Primary: Composite of time to doubling of serum creatinine or ESRD</p> <p>Secondary: Changes in blood pressure, daily urinary protein excretion, adverse effects</p>	<p>Primary: The combined end point was reached in 11% of patients in the combination trandolapril and losartan group compared to 23% of patients in the trandolapril (P=0.018) and 23% of patients in the losartan group (P=0.016).</p> <p>Secondary: Mean SBP and DBP reductions were similar among the three treatment groups (P=0.109).</p> <p>All patients receiving active treatment had significant decreases in urinary protein excretion, but the greatest difference was seen with the combination trandolapril and losartan group compared to trandolapril or losartan (-75.6, -44.3, and -42.1%, respectively; P=0.01).</p> <p>The frequency of adverse events did not differ between groups, although a slightly higher occurrence of hyperkalemia and dry cough was recorded in the trandolapril and combination groups than in the losartan group.</p>
<p>Mann et al.<sup>56</sup> (2009) TRANSCEND</p> <p>Telmisartan 80 mg QD</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Adults with known cardiovascular disease or diabetes with end-organ damage but without macroalbuminuria or heart failure who</p>	<p>N=5927</p> <p>56 months</p>	<p>Primary: Composite outcome: first occurrence of dialysis, renal transplant, doubling of serum creatinine, or death</p>	<p>Primary: The composite outcome of dialysis, doubling of serum creatinine, or death did not significantly differ between the telmisartan and placebo groups (412 patients [14.0%] vs 381 patients [12.8%]; HR, 1.10 [CI, 0.95 to 1.26]; P=0.193).</p> <p>The incidence of the composite outcome of dialysis or doubling of serum creatinine was similar with telmisartan and placebo (58 patients [1.96%] vs 46 patients [1.55%]; HR, 1.29 [95% CI, 0.87 to 1.89]; P=0.20).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo	cannot tolerate ACE inhibitors		Secondary: Changes in the eGFR, progression of proteinuria, and individual components of the primary outcome	Secondary: Doubling of serum creatinine was more frequent with telmisartan than with placebo (56 vs 36 patients; P=0.031).  Decreases in eGFR were greater with telmisartan than with placebo (mean change in eGFR, -3.2 mL/min per 1.73 m <sup>2</sup> [SD, 18.3] vs -0.26 mL/min per 1.73 m <sup>2</sup> [SD, 18.0]; P <0.001).
Foulquier et al. <sup>57</sup> (2014) TRANSCEND  Telmisartan 80 mg QD  vs  placebo	Post-hoc analysis  Patients in the TRANSCEND trial stratified by hypertensive and nonhypertensive	N=5927  56 months	Primary: Composite outcome: first occurrence of dialysis, renal transplant, doubling of serum creatinine, or death  Secondary: Changes in the eGFR, progression of proteinuria, and individual components of the primary outcome	Primary: For the primary four-fold endpoint, No difference in the effect of treatment between hypertensive and nonhypertensive patients was found. No significant improvement with telmisartan over placebo in both hypertensive and nonhypertensive patients was seen.  Secondary: New onset of LVH, evaluated by ECG, was significantly less in hypertensive and nonhypertensive patients treated with telmisartan (hypertensive patients: -36%; P=0.0002; nonhypertensive patients: -58%; P=0.027).  Albuminuria increased less with telmisartan than with placebo in the hypertensive population, as the risks for new microalbuminuria and macroalbuminuria were lower than with placebo (P=0.0004 and P=0.009, respectively). In the nonhypertensive population, the risks were not modified by the treatment. However, according to the interaction tests, there is no difference in the effect of telmisartan in hypertensive and nonhypertensive patients, suggesting that telmisartan might also reduce the new onset of microalbuminuria and macroalbuminuria in nonhypertensive patients.
Barnett et al. <sup>58</sup> (2004) DETAIL  Enalapril 20 mg/day  vs  telmisartan 80	DB, MC, PG, RCT  Patients aged 35 to 80 years with type 2 diabetes and HTN	N=250  5 years	Primary: Change in the GFR  Secondary: Annual changes in GFR, serum creatinine level, urinary albumin excretion, and blood pressure; rates of	Primary: After five years, GFR decreased by 17.9 mL/minute/1.73 m <sup>2</sup> with telmisartan compared to 14.9 mL/min/1.73 m <sup>2</sup> with enalapril (mean difference, -3.0 mL/min/1.73 m <sup>2</sup> ; 95% CI, -7.6 to 1.6). Therefore, the changes in GFR were comparable between the groups.  Secondary: The effects of the two agents on the secondary end points were not significantly different after five years.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day			ESRD and cardiovascular events; all-cause mortality	
Galle et al. <sup>59</sup> (2008)  Telmisartan 80 mg QD  vs  valsartan 160 mg QD  Additional antihypertensive therapy was allowed.	DB, MC, PG, PRO, RCT, non-inferiority study  Hypertensive patients (SBP/DBP >130/80 mm Hg) with type 2 diabetes, proteinuria and serum creatinine ≤3.0 mg/dL	N=885  12 months	Primary: Change from baseline in the 24-hour proteinuria  Secondary: Changes in 24-hour albuminuria, eGFR and inflammatory parameters	Primary: Telmisartan and valsartan produced comparable reductions in 24-hour urinary protein excretion rates: geometric mean reduction was 33% for both telmisartan and valsartan.  Secondary: No significant differences between treatments were seen in changes from baseline in 24-hour urinary albumin excretion rate and GFR at 12 months.  With both treatments, greater renoprotection was seen among patients with better blood pressure control.  No significant changes in C-reactive protein were noted for either group at 12 months.
Fogari et al. <sup>60</sup> (2007)  Telmisartan and amlodipine 40 to 160-2.5 QD (fixed-dose combination)  vs  telmisartan and amlodipine 40-2.5 mg QD (fixed-dose combination)	DB, MC, RCT  Patients 35 to 70 years of age with essential HTN, type 2 diabetes mellitus and microalbuminuria (UAER >30 and <300 mg/24 hr)	N=210  64 weeks	Primary: Blood pressure, UAER, creatinine clearance, plasma potassium, fasting glycemia, and HbA <sub>1c</sub>  Secondary: Not reported	Primary: High-dose telmisartan/low-dose amlodipine and low-dose telmisartan/high-dose amlodipine combination produced a similar reduction in SBP and DBP with no significant difference between the two regimens at any time of the study.  With increasing doses of telmisartan (40, 80, 120, and 160 mg), SBP and DBP values were reduced from baseline by 16 and 10 mm Hg, respectively (P<0.01), 24 and 21 mm Hg, respectively (P<0.001), 23 and 21 mm Hg, respectively (P<0.001), and 24 and 21 mm Hg, respectively (P<0.001).  With increasing dose of amlodipine (2.5, 5, 7.5, and 10 mg) SBP and DBP values were reduced from baseline by 16 and 10 mm Hg, respectively (P<0.01), 25 and 22 mm Hg, respectively (P<0.001), 25 and 21 mm Hg, respectively (P<0.001), and 25 and 22 mm Hg, respectively (P<0.001).  Reductions of UAER from baseline were of 34.6 mg/24 hr (P<0.05 vs baseline), 62.9 mg/24 hr (P<0.01 vs baseline and P<0.05 vs A group), 86.5

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>mg/24 hr (P&lt;0.001 vs baseline and P&lt;0.01 vs A group) and 102 mg/24 hr (P&lt;0.0001 vs baseline and P&lt;0.001 vs A group) for telmisartan 40, 80, 120, and 160 mg/amlodipine 2.5 mg daily, respectively.</p> <p>Reductions of UAER from baseline were of 35.1 mg/24 hr (P&lt;0.05 vs baseline), 46.2 mg/24 hr (P&lt;0.03 vs baseline), 50.3 mg/24 hr (P&lt;0.03 vs baseline), and 45 mg/24 hr (P&lt;0.03 vs baseline) for amlodipine-telmisartan 2.5-40, 5-40, 7.5-40, and 10-40 mg/day, respectively.</p> <p>Creatinine clearance did not significantly change with either treatment. Neither combination affected levels of plasma potassium or fasting glucose. The HbA<sub>1c</sub> levels were not significantly influenced by either treatment.</p>
<p>Viberti et al.<sup>61</sup> (2002) MARVAL</p> <p>Valsartan 80 mg QD</p> <p>vs</p> <p>amlodipine 5 mg QD</p> <p>A target blood pressure of 135/85 mm Hg was aimed for by dose-doubling followed by the addition of bendrofluazide* and doxazosin whenever needed.</p>	<p>AC, DB, RCT</p> <p>Patients 35-75 years old with type 2 diabetes mellitus and microalbuminuria, with or without HTN</p>	<p>N=332</p> <p>24 weeks</p>	<p>Primary: Change in UAER; proportion of patients who returned to normal albuminuria</p> <p>Secondary: Proportion of patients returning to normoalbuminuria</p>	<p>Primary: Valsartan resulted in a UAER reduction of 44% at 24 weeks compared to baseline vs an 8% reduction with amlodipine (P&lt;0.001). Valsartan lowered UAER similarly in both the hypertensive and normotensive groups.</p> <p>Over the study period, blood pressure reductions were similar between the two treatments and at no time point was there a between-group significant difference in blood pressure values in either the hypertensive or the normotensive subgroup.</p> <p>Secondary: The proportion of patients returning to normal albuminuria was greater with valsartan (29.9%) vs amlodipine (14.5%; P=0.001).</p>
<p>Casas et al.<sup>62</sup> (2005)</p> <p>ACE inhibitor or</p>	<p>MA (127 trials)</p> <p>Studies in adults that examined the effect</p>	<p>N=not reported</p> <p>4.2 years</p>	<p>Primary: Doubling of serum creatinine, and ESRD</p>	<p>Primary: Treatment with ACE inhibitors or ARBs resulted in a nonsignificant reduction in the risk of doubling of creatinine vs other antihypertensives (P=0.07) with no differences in the degree of change of SBP or DBP between the groups.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ARBs compared to placebo</p> <p>vs</p> <p>ACE inhibitor or ARBs compared to other antihypertensive drugs (<math>\beta</math>-adrenergic blocking agents, <math>\alpha</math>-adrenergic blocking agents, calcium-channel blocking agents, or combinations)</p> <p>Specific agents and doses were not specified.</p>	<p>of any drug treatment with a blood pressure lowering action on progression of renal disease</p>	<p>(mean)</p>	<p>Secondary: Serum creatinine, urine albumin excretion and GFR</p>	<p>A small reduction in ESRD was observed in patients receiving ACE inhibitors or ARBs compared to other antihypertensives (P=0.04) with no differences in the degree of change of SBP or DBP between the groups.</p> <p>Secondary: Small reductions in serum creatinine and in SBP were noted when ACE inhibitors or ARBs were compared to other antihypertensives (P=0.01).</p> <p>Small reduction in daily urinary albumin excretion in favor of ACE inhibitor or ARBs were reported when these agents were compared to other antihypertensives (P=0.001).</p> <p>Compared to other drugs, ACE inhibitors or ARBs had no effect on the GFR.</p>
<p>Strippoli et al.<sup>63</sup> (2004)</p> <p>ACE inhibitors</p> <p>vs</p> <p>placebo</p> <p>or</p> <p>ARBs</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients with diabetic nephropathy</p>	<p>43 trials</p> <p><math>\geq 6</math> months (range 6 to 63.6 months)</p>	<p>Primary: All-cause mortality, renal outcomes (ESRD, doubling of serum creatinine, microalbuminuria to macroalbuminuria)</p> <p>Secondary: Not reported</p>	<p>Primary: ACE inhibitors significantly reduced all-cause mortality compared to placebo or no treatment (RR, 0.79; 95% CI, 0.63 to 0.99; P=0.04). There was a nonsignificant trend for reduction in ESRD (P=0.07) and doubling of serum creatinine (P=0.08) with ACE inhibitors compared to placebo or no treatment. ACE inhibitors significantly reduced the risk of progression from microalbuminuria to macroalbuminuria (P=0.0007) and increased regression back to normoalbuminuria (P&lt;0.0001) compared to placebo or no treatment.</p> <p>ARBs did not significantly reduce all-cause mortality compared to placebo or no treatment (RR, 0.99; 95% CI, 0.85 to 1.17; P=0.95). ARBs significantly reduced the risk of ESRD (P=0.001) and doubling of serum creatinine (P=0.004). ARBs significantly decreased the risk of progression to macroalbuminuria (P=0.001) and increased regression to normoalbuminuria (P=0.02) compared to placebo or no treatment.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>or ACE inhibitors vs ARBs</p>				<p>The three trials that compared ACE inhibitors to ARBs did not report on all-cause mortality, ESRD or doubling of serum creatinine. Progression from microalbuminuria to macroalbuminuria was reported in one trial (N=92) and there was no significant difference in risk, with the point estimate favoring ACE inhibitors (RR, 0.16; 95% CI, 0.02 to 1.44). Regression from microalbuminuria to normoalbuminuria in 1 trial showed a nonsignificant difference in the risk.</p> <p>Secondary: Not reported</p>
<p>Strippoli et al.<sup>64</sup> (2006) ACE inhibitors vs placebo or ARBs vs placebo or ACE inhibitors vs ARBs</p>	<p>MA  Patients with diabetic kidney disease</p>	<p>N=12,067 (49 trials)  ≥6 months</p>	<p>Primary: All-cause mortality, ESRD, doubling of serum creatinine concentration, progression from micro- to macroalbuminuria, regression from micro- to normoalbuminuria, drug-related toxicity (including cough, headache, hyperkalemia, impotence and pedal edema)</p> <p>Secondary: Not reported</p>	<p>Primary: There was no significant difference in the risk of all-cause mortality for ACE inhibitors vs placebo or no treatment (RR, 0.91; 95% CI, 0.71 to 1.17) and ARBs vs placebo or no treatment (RR, 0.99; 95% CI, 0.85 to 1.17). No statistically significant reduction in the risk of all-cause mortality was found in the three studies that compared ACE inhibitors with ARBs (RR, 0.92; 95% CI, 0.31 to 2.78).</p> <p>A subgroup analysis of studies showed a significant reduction in the risk of all-cause mortality with the use of full-dose ACE inhibitors (RR, 0.78; 95% CI, 0.61 to 0.98) but not when using half or less than half the maximum tolerable dose of ACE inhibitors (RR, 1.18; 95% CI, 0.41 to 3.44).</p> <p>There was a significant reduction in the risk of ESRD with ACE inhibitors and ARBS compared to placebo or no treatment (RR, 0.60; 95% CI, 0.39 to 0.93 and RR, 0.78; 95% CI, 0.67 to 0.91, respectively). There was a significant reduction in the risk of doubling of serum creatinine concentration with ACE inhibitors and ARBS (RR, 0.68; 95% CI, 0.47 to 10 and RR, 0.79; 95% CI, 0.67 to 0.93, respectively).</p> <p>ACE inhibitors and ARBS significantly reduced the risk of progression from micro- to macroalbuminuria (RR, 0.45; 95% CI, 0.29 to 0.69 and RR, 0.49; 95% CI, 0.32 to 0.75, respectively). ACE inhibitors and ARBS significantly increased the regression from micro- to normoalbuminuria compared to placebo or no treatment (RR, 3.06; 95% CI, 1.76 to 5.35 and RR, 1.42; 95% CI, 1.15 to 1.93, respectively).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The seven studies that compared ACE inhibitors to ARBS did not report the outcome of ESRD or doubling of serum creatinine. Progression from micro- to macroalbuminuria and from micro- to normoalbuminuria were evaluated each in one trial and showed a nonsignificant difference in the risk between ACE inhibitors and ARBS.</p> <p>ACE inhibitors were associated with a significant increase in the risk of cough but not hyperkalemia, headache or impotence when compared to placebo or no treatment. ARBS were associated with a significant increase in the risk of hyperkalemia but not cough or headache compared to placebo or no treatment.</p> <p>Secondary: Not reported</p>
<b>Heart Failure</b>				
<p>Cohn et al.<sup>65</sup> (2001) Val-HeFT  Valsartan 160 mg BID  vs  placebo</p>	<p>DB, PC, RCT  Patients ≥18 years old with a cardiovascular history and NYHA II to IV heart failure</p>	<p>N=5,010  2 years</p>	<p>Primary: Mortality and composite end point of morbidity and mortality  Secondary: Change in NYHA class, ejection fraction, signs and symptoms of heart failure, QOL</p>	<p>Primary: Compared to placebo, valsartan resulted in no significant differences in all-cause mortality.  Patients treated with valsartan experienced a 13% decrease in the composite end point (P=0.009) and 27% decrease in heart failure hospitalizations (P&lt;0.001).  Secondary: Treatment with valsartan resulted in significant improvements in NYHA class, ejection fraction, signs and symptoms of heart failure and QOL as compared to placebo (P&lt;0.01).  In a post hoc analysis of the combined end point and mortality in subgroups defined according to baseline treatments with ACE inhibitors or β-blockers, valsartan had a favorable effect in patients receiving neither or one of these types of drugs but an adverse effect in patients receiving both types of drugs.</p>
<p>Pfeffer et al.<sup>66</sup> (2003) CHARM Overall Programme  Candesartan 32</p>	<p>DB, PC, PG, RCT  Summary of all CHARM sub-studies</p>	<p>N=7,599  37.7 months</p>	<p>Primary: All-cause mortality (Overall Programme) and cardiovascular death or hospital</p>	<p>Primary: In the overall analysis, candesartan 32 mg daily resulted in an 18% decreased risk of all-cause mortality compared to placebo (23 vs 25%; unadjusted HR, 0.91; 95% CI, 0.83 to 1.0; P=0.055; covariate adjusted HR, 0.90; 95% CI, 0.82 to 0.99; P=0.032).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day ( $\pm$ ACE inhibitor)  vs  placebo ( $\pm$ ACE inhibitor)			admission for CHF (all of the component trials)  Secondary: Not reported	Annual mortality rates were 8.1 and 8.8% for patients treated with candesartan and placebo, respectively.  The lower mortality in patients treated with candesartan vs placebo was attributed to fewer cardiovascular deaths (18 vs 20%; unadjusted HR, 0.88; 95% CI, 0.79 to 0.97; P=0.012).  Hospital admissions for CHF were significantly fewer in patients treated with candesartan than placebo (20 vs 24%; P<0.0001).  Secondary: Not reported
McMurray et al. <sup>67</sup> (2003) CHARM-Added  Candesartan 32 mg/day in patients already taking ACE inhibitors  vs  placebo in patients already taking ACE inhibitors	DB, MC, PC, RCT  Patients $\geq$ 18 years old with LVEF $\leq$ 40%, NYHA II to IV heart failure and treatment with an ACE inhibitor at a constant dose for 30 days or longer	N=2,548  41 months	Primary: Composite of cardiovascular death and hospitalization for heart failure  Secondary: Composites of primary end point and MI, nonfatal stroke and coronary revascularization	Primary: Compared to placebo, candesartan 32 mg/day when added to ACE inhibitors resulted in a 15% reduction in the primary end point (P=0.011), 16% decrease in cardiovascular deaths (P=0.029) and 17% reduction in heart failure hospitalizations (P=0.014).  Secondary: Fewer patients experienced cardiovascular death, hospital admission for CHF, MI, stroke, or coronary revascularization in the candesartan group (42.9%) compared to placebo (46.9%; P=0.015).
Granger et al. <sup>68</sup> (2003) CHARM-Alternative  Candesartan 32 mg/day  vs  placebo	DB, PC, RCT  Patients $\geq$ 18 years old with LVEF $\leq$ 40%, NYHA II to IV heart failure and intolerance to ACE inhibitors	N=2,028  33.7 months	Primary: Composite of cardiovascular death and hospitalization for heart failure  Secondary: Composites of primary end point and MI, nonfatal stroke and coronary	Primary: Compared to placebo, candesartan 32 mg/day resulted in a 30% reduction of the composite end point (P<0.0001).  A 20% decrease in cardiovascular death (P=0.02) and 39% reduction in heart failure hospitalizations (P<0.0001) were noted in patients treated with candesartan compared to placebo.  Study drug discontinuation rates were similar in the candesartan (30%) and placebo (29%) groups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			revascularization	Secondary: Fewer patients experienced cardiovascular death, hospital admission for CHF, MI, stroke, or coronary revascularization in the candesartan group (39.1%) compared to placebo (44.9%; P<0.0001).
Yusuf et al. <sup>69</sup> (2003) CHARM-Preserved  Candesartan 32 mg/day  vs  placebo	DB, PC, RCT  Patients ≥18 years old with preserved ejection fraction (>40%) and symptomatic heart failure	N=3,025  36.6 months	Primary: Composite of cardiovascular death and hospitalization for heart failure  Secondary: Composites of primary end point and MI, nonfatal stroke and coronary revascularization	Primary: Compared to placebo, candesartan 32 mg/day resulted in an insignificant 14% trend towards lower incidence of the primary end point (P=0.051).  Candesartan significantly reduced the risk of heart failure hospitalization (16%; P=0.047) but did not significantly decrease the risk of cardiovascular death (P=0.635).  Secondary: The composite of cardiovascular death, hospitalization for CHF, MI, and stroke was significantly lower in the candesartan group compared to placebo (25.6 vs 28.4%; P=0.037).  There was no significant difference in the composite of cardiovascular death, hospital admission for CHF, MI, stroke, or coronary revascularization in the candesartan group (30.4%) compared to placebo (32.9%; P=0.130).
Castagno et al. <sup>70</sup> (2012) CHARM  Candesartan 32 mg/day (±ACE inhibitor)  vs  placebo (±ACE inhibitor)	Subgroup analysis according to baseline heart rate and LVEF  Patients with chronic heart failure	N=7,597  Duration varied	Primary: Composite of cardiovascular death or heart failure hospital stay  Secondary: Not reported	Primary: Patients with the highest heart rate tertile had worse outcomes when compared to patients in the lowest heart rate group (HR, 1.23; 95% CI, 1.11 to 1.36; P<0.001). The relationship between heart rate and outcomes was similar across LVEF categories, and was not influenced by use of β-blockers (P>0.10 for both endpoints).  Secondary: Not reported
Pitt et al. <sup>71</sup> (1997) ELITE  Captopril 50 mg	DB, MC, PG, RCT  Patients ≥65 years with symptomatic heart failure (NYHA	N=722  1 year	Primary: Change in renal function  Secondary:	Primary: No difference between losartan and captopril was reported in the rate of persistent rise in serum creatinine concentrations (10.5% for both groups).  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
TID vs losartan 50 mg QD	class II to IV and LVEF $\leq$ 40%), and no history of prior ACE inhibitor therapy		Composite of death and/or hospital admission for heart failure, all-cause mortality, admission for heart failure, NYHA class, admission for MI or unstable angina	Death and/or hospital admission for heart failure was recorded in 9.4% of patients receiving losartan and 13.2% for patients receiving captopril (risk reduction, 32%; 95% CI, -4 to 55; P=0.075). This risk reduction was primarily due to a decrease in all-cause mortality (4.8 vs 8.7%; risk reduction, 46%; 95% CI, 5 to 69; P=0.035).  Admissions with heart failure were the same in both groups (5.7%), as was improvement in NYHA functional class from baseline. Admission to hospital for any reason was less frequent with losartan than with captopril treatment (22.2 vs 29.7%; P=0.014).  More patients discontinued therapy due to adverse events with captopril (20.8%) than losartan (12.2%; P=0.002).
Pitt et al. <sup>72</sup> (2000) ELITE II  Captopril 50 mg TID vs losartan 50 mg QD	DB, MC, PG, RCT  Patients $\geq$ 60 years old with symptomatic heart failure (NYHA II to IV and LVEF $\leq$ 40%), and no history of prior ACE inhibitor therapy	N=3,152  555 days (mean follow-up)	Primary: All-cause mortality  Secondary: Composite of sudden cardiac death or resuscitated cardiac arrest	Primary: No significant difference in all-cause mortality was reported between losartan (17.7%) and captopril (15.9%; HR, 1.13; 95% CI, 0.95 to 1.35; P=0.16).  Secondary: Sudden death or resuscitated cardiac arrest was observed in 9.0% of patients receiving losartan and 7.3% of patients receiving captopril (HR, 1.25; 95% CI, 0.98 to 1.60; P=0.08).  Significantly fewer patients in the losartan group (excluding those who died) discontinued study treatment because of adverse events (9.7 vs 14.7%; P<0.001), including cough (0.3 vs 2.7%).  Note: ELITE II trial was a larger follow-up trial to the ELITE I trial to confirm the secondary end point from the ELITE I trial, which reported a greater reduction in all-cause mortality with losartan compared to captopril.
McKelvie et al. <sup>73</sup> (1999) RESOLVD  Enalapril 10 mg BID vs	DB, MC, PG, RCT  Patients with CHF (NYHA classes II to IV), a 6 minute walk distance of 500 meters or less, and an ejection fraction	N=768  43 weeks	Primary: Change in 6-minute walk distance  Secondary: Change in NYHA functional class, QOL, ejection	Primary: There were no significant differences among the groups with regards to the 6-minute walk distance over the 43 week study period.  Secondary: There were no significant differences among the groups with regards to the NYHA functional class or QOL at 18 or 43 weeks.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>candesartan 4 to 16 mg QD</p> <p>vs</p> <p>candesartan 4 to 8 mg QD and enalapril 10 mg BID</p>	<p>&lt;40%</p>		<p>fraction, ventricular volumes, neurohormone levels, safety</p>	<p>Ejection fraction increased more with candesartan plus enalapril than monotherapy with either agent; however, the difference was not statistically significant (P value not significant). End-diastolic volumes (P&lt;0.01) and end-systolic volumes (P&lt;0.05) increased less with combination therapy than with monotherapy with either agent.</p> <p>Aldosterone decreased with combination therapy at 17 but not 43 weeks compared to candesartan or enalapril (P&lt;0.05). Brain natriuretic peptide decreased with combination therapy compared to candesartan and enalapril alone (P&lt;0.01).</p> <p>Blood pressure decreased with combination therapy compared to candesartan or enalapril alone (P&lt;0.05).</p> <p>Compared to enalapril, potassium decreased with candesartan use (P&lt;0.05) and increased with candesartan plus enalapril (P&lt;0.05). The proportion of patients with potassium levels <math>\geq 5.5</math> mmol/L was not significantly different among the treatment groups. There were no significant differences in creatinine, mortality, or hospitalizations for CHF or any cause among the three groups.</p>
<p>Lee et al.<sup>74</sup> (2004)</p> <p>ARBs</p> <p>vs</p> <p>placebo (<math>\pm</math>ACE inhibitor)</p> <p>vs</p> <p>ACE inhibitor monotherapy</p>	<p>MA</p> <p>Patients with chronic heart failure and high-risk acute MI</p>	<p>N=38,080</p> <p>Duration varied</p>	<p>Primary: All-cause mortality and heart failure hospitalizations</p> <p>Secondary: Not reported</p>	<p>Primary: ARBs were associated with reduced all-cause mortality (OR, 0.83) and heart failure hospitalizations (OR, 0.64) vs placebo.</p> <p>There was no difference in all-cause mortality (OR, 1.06) and heart failure hospitalization (OR, 0.95) between ARBs and ACE inhibitors.</p> <p>When ARBs were combined with ACE inhibitors, all-cause mortality was not reduced (OR, 0.97) but heart failure hospitalizations were reduced (OR, 0.77) compared to treatment with ACE inhibitors alone.</p> <p>Two RCT comparing ARBs with ACE inhibitors in patients with high-risk acute MI did not reveal differences in all-cause mortality or heart failure hospitalization.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Hypertension</b>				
Rakugi et al. <sup>75</sup> (2012)  Azilsartan 20 to 40 mg QD  vs  candesartan 8 to 12 mg QD	DB, MC, RCT  Japanese patients with grade I or II essential HTN	N=622  16 weeks	Primary: Change in baseline mean sitting DBP at week 16  Secondary: Change in baseline mean sitting SBP at week 16	Primary: After 16 weeks, the mean baseline change in sitting DBP was -12.4 and -9.8 mm Hg with azilsartan and candesartan (difference, -2.6; 95% CI, -4.08 to -1.22; P=0.0003).  Secondary: After 16 weeks, the mean baseline change in sitting SBP was -21.8 and -17.5 mm Hg with azilsartan and candesartan (difference, -4.4 mm Hg; 95% CI, -6.53 to -2.20; P<0.0001).
Sica et al. <sup>76</sup> (2001)  Azilsartan 40 or 80 mg QD  vs  valsartan 320 mg QD	DB, MC, RCT  Patients with primary HTN	N=984  24 weeks	Primary: Change in baseline 24 hour mean ambulatory and clinic SBP  Secondary: Change in baseline 24 hour mean ambulatory and clinic DBP	Primary: Azilsartan 40 and 80 mg/day significantly lowered 24 hour mean ambulatory systolic blood pressure (-14.9 and -15.3 mm Hg) compared to valsartan 320 mg/day (-11.3 mm Hg; P<0.001). Clinic SBP reductions were consistent with ambulatory blood pressure results. (-14.9 and -16.9 vs -11.6 mm Hg; P=0.015 and P<0.001).  Secondary: Reductions in 24 hour mean and clinic DBP were significantly greater with azilsartan compared to valsartan (P≤0.001 for all comparisons).
Cushman et al. <sup>77</sup> (2012)  Azilsartan and chlorthalidone 40-25 or 80-25 mg QD (fixed-dose combination product)  vs  olmesartan and HCTZ 40-25 mg QD (fixed-dose	DB, RCT  Patients with clinic SBP 160 to 190 mm Hg and DBP ≤119 mm Hg	N=1,071  12 weeks	Primary: Change in baseline clinical SBP  Secondary: Change in baseline ambulatory SBP, safety	Primary: Changes in clinic SBP were significantly greater with azilsartan and chlorthalidone (-42.5±0.8 and -44.0±0.8 mm Hg) compared to olmesartan and HCTZ (-37.1±0.8 mm Hg; P<0.0001).  Secondary: Changes in ambulatory SBP were significantly greater with azilsartan and chlorthalidone (-33.9±0.8 and -36.3±0.8 mm Hg) compared to olmesartan and HCTZ (-27.5±0.8 mm Hg; P<0.0001).  Adverse events leading to discontinuation of study medications were 7.9, 14.5, and 7.1% of patients receiving azilsartan and chlorthalidone 40-25 mg/day, azilsartan and chlorthalidone 80-25 mg/day, and olmesartan and HCTZ.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
combination product)				
Bönnner et al. <sup>78</sup> (2013)  Azilsartan (AZL) 20mg titrated to 40 mg  vs  azilsartan (AZL) 20mg titrated to 80 mg  vs  ramipril (RAM) 2.5 mg titrated to 10 mg	DB, RCT  Patients ≥18 years of age with clinic systolic blood pressure (SBP) 150 to 180 mm Hg	N=884  24 weeks	Primary: Change in trough, seated clinic SBP  Secondary: Change from baseline to week 24 in trough, seated clinic DBP, measures of ambulatory BP, and BP response rates	Primary: After 24 weeks of treatment, trough, sitting, clinic SBP decreased significantly in all the groups. The changes from baseline were significantly greater for the AZL 40 and 80 mg treatment groups (-20.6±0.95 and -21.2±0.95 mm Hg, respectively) than for RAM 10 mg (-12.2±0.95 mm Hg). The differences between the AZL-treated subjects and the RAM-treated subjects were -8.4 mm Hg for AZL 40 and -9.0 mm Hg for AZL 80 (P<0.001 for both comparisons).  Secondary: Change in trough, sitting, DBP was -10.2±0.55 mm Hg in the AZL 40 mg group, -10.5±0.55 mm Hg in the AZL 80 mg and -4.9±0.56 mm Hg in the RAM 10 mg group.  AZL 40 and 80 mg reduced ambulatory SBP and DBP significantly more than RAM for all ABPM time intervals evaluated, including 24-hour mean, mean daytime, mean nighttime and mean trough pressure.  The differences between the AZL and RAM groups proportion of subjects achieving SBP and DBP response criteria were highly significant (P<0.001). More subjects achieved a reduction in clinic BP to <140/90 mm Hg and/or a reduction in BP≥20/10 mm Hg at week 24 following treatment with AZL compared with RAM (54.0% and 53.6% for AZL 40 and 80 mg vs 33.8% with RAM 10 mg, respectively; P<0.001).
Handley et al. <sup>79</sup> (2016)  All subjects initiated treatment with azilsartan 40 mg QD on day one, which was added to existing treatments (a maximum of two	MC, OL, cohort  Patients >18 years of age with essential HTN	N=669  56 weeks	Primary: Safety and tolerability  Secondary: Efficacy	Primary: Approximately 76% of subjects overall in the two cohorts experienced an adverse event. Within each cohort, more events were reported among subjects who received add-on therapy with chlorthalidone or HCTZ. The most commonly reported adverse events (≥5% of subjects) in both cohorts combined, regardless of add-on diuretic therapy, were dizziness (14.3%), headache (9.9%), fatigue (7.2%), upper respiratory tract infection (6.7%) and urinary tract infection (5.7%). Transient serum creatinine elevations were more frequent with add-on chlorthalidone.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>other antihypertensive agents), if applicable; at week four, azilsartan was force-titrated to 80 mg QD, if tolerated</p> <p>In Cohort 1, chlorthalidone 25 mg QD was the initial add-on agent for subjects who did not achieve target BP on azilsartan alone, In Cohort 2, HCTZ 12.5 mg QD was the initial add-on agent</p>				<p>By week 56 in Cohort 1, the overall change from baseline in clinic SBP (observed cases) was <math>-25.2 \pm 18.1</math> mmHg (<math>n = 259</math>; <math>21.1 \pm 15.2</math> mmHg for subjects receiving azilsartan alone [<math>n = 93</math>] and <math>-27.4 \pm 19.2</math> mmHg for those requiring add-on chlorthalidone [<math>n = 166</math>]). In Cohort 2, the overall change from baseline in clinic SBP was <math>-24.2 \pm 16.0</math> mmHg (<math>n = 201</math>; <math>-21.6 \pm 14.2</math> for mmHg azilsartan alone [<math>n = 68</math>] and <math>-25.6 \pm 16.7</math> mmHg for add-on HCTZ [<math>n = 133</math>]).</p> <p>By week 56 in Cohort 1, the overall change from baseline in clinic DBP (observed cases) was <math>-18.4 \pm 9.5</math> mmHg (<math>-18.0 \pm 8.8</math> mmHg for azilsartan alone and <math>-18.6 \pm 9.9</math> mmHg with add-on CLD). By week 56 in Cohort 2, the change from baseline in clinic DBP was <math>-17.9 \pm 10.9</math> mmHg (<math>-17.9 \pm 9.4</math> mmHg for subjects azilsartan alone and <math>-18.0 \pm 11.6</math> mmHg with add-on HCTZ).</p>
<p>Kipnes et al.<sup>80</sup> (2015)</p> <p>For the 26-week, OL phase, patients received an initial dose of azilsartan 40 mg QD. At week four, the dose was force-titrated to 80 mg QD, and from week eight to week 22, chlorthalidone 25 mg QD could be added to</p>	<p>7-day screening phase; 26-week OL phase; 6-week randomized, DB, reversal phase; and a 7-day post-treatment AE follow-up phase</p> <p>Patients &gt;18 years of age with essential HTN</p>	<p>26 weeks OL (N=418) followed by 6 weeks DB (N=299)</p>	<p>Primary: Change in trough clinic sitting DBP measured during the DB reversal phase</p> <p>Secondary: Change in trough clinic sitting SBP during the DB reversal phase; safety and tolerability</p>	<p>Primary: At the DB phase baseline (week 26), the mean clinic DBP was similar in the azilsartan and placebo groups (83.5 mm Hg and 82.3 mm Hg, respectively). This DBP level was maintained to the final visit (week 32) in patients who received azilsartan. In contrast, DBP increased among patients who received placebo, demonstrating a loss of efficacy after discontinuation of azilsartan. The least-squares mean difference between azilsartan and placebo was <math>-7.8</math> mm Hg (95% CI, <math>-9.8</math> to <math>-5.8</math>; <math>P &lt; 0.001</math>) at final visit.</p> <p>Secondary: At the DB phase baseline (week 26), the mean clinic SBP was also similar in the azilsartan and placebo groups (129.8 mm Hg and 128.2 mm Hg, respectively). As with DBP, this SBP level was maintained from week 26 to week 32 in patients who received azilsartan, whereas it increased in patients receiving placebo. The least-squares mean difference between azilsartan and placebo was <math>-12.4</math> mm Hg (95% CI, <math>-15.5</math> to <math>-9.3</math>; <math>P &lt; 0.001</math>) at final visit, and</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>achieve target BP</p> <p>At week 26 (end of OL phase), patients were randomized into a 6-week, DB reversal phase in which they continued to receive azilsartan at their final dose level or were switched to placebo</p>				<p>the LS mean difference was statistically significant at each scheduled double-blind dosing visit.</p> <p>During the OL phase, approximately half (54.1%) of patients overall experienced an adverse event and these were predominantly (&gt;90%) mild to moderate in severity. The most commonly reported adverse events overall were dizziness (8.9%) and headache (7.2%).</p>
<p>Neutel et al.<sup>81</sup> (2017)</p> <p>Azilsartan-chlorthalidone 20/40/12.5mg once daily titrated to 80/25mg if needed</p> <p>vs</p> <p>olmesartan-HCTZ 20/12.5mg FDC once daily titrated to 40/25mg if needed</p>	<p>MC, OL, RCT</p> <p>Patients with hypertension <math>\geq 18</math> years of age with clinic SBP 160 to 190 mmHg and DBP 119 mmHg or less</p>	<p>N=837</p> <p>52 weeks</p>	<p>Primary: Percentage of patients with one or more treatment-emergent adverse event</p> <p>Secondary: Clinical laboratory tests, vital signs, BP</p>	<p>Primary: The percentage of patients who reported one or more treatment-emergent adverse event was 78.5% in the azilsartan-chlorthalidone group and 76.4% in the olmesartan-HCTZ group. The most commonly reported adverse events (azilsartan-chlorthalidone vs olmesartan-HCTZ) were dizziness (16.3% vs 12.6%), blood creatinine increase (21.5% vs 8.6%), headache (7.4% vs 11.0%), and nasopharyngitis (12.2% vs 11.5%). Events of hypokalemia were uncommon in both treatment groups (1.0% vs 0.7%).</p> <p>Secondary: Mean changes in SBP from baseline to the final visit (last observation carried forward) were <math>-42.3 \pm 14.2</math> mm Hg (azilsartan-chlorthalidone) vs <math>-38.0 \pm 14.1</math> mm Hg (olmesartan-HCTZ), and mean changes in DBP were <math>-18.4 \pm 9.0</math> mm Hg (azilsartan-chlorthalidone) vs <math>-15.6 \pm 9.8</math> mm Hg (olmesartan-HCTZ), respectively. No clinically relevant differences in urinalysis parameters, electrocardiographic parameters, or vital signs (including heart rate, body weight, or orthostatic BP) were observed between azilsartan-chlorthalidone and olmesartan-HCTZ.</p>
<p>Cushman et al.<sup>82</sup> (2018)</p> <p>Azilsartan-</p>	<p>DB, RCT</p> <p>Patients with primary</p>	<p>N=1,085</p> <p>8 weeks</p>	<p>Primary: Change from baseline in clinic SBP</p>	<p>Primary: Greater reductions in clinic SBP from a baseline of 165 mmHg were observed (<math>P &lt; 0.001</math>) in both azilsartan-chlorthalidone arms (<math>-37.6</math> and <math>-38.2</math> mmHg) versus olmesartan-HCTZ (<math>-31.5</math> mmHg), despite greater dose titration in the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
chlorthalidone 20/12.5mg once daily titrated to 40/25mg if needed or 40/12.5mg once daily titrated to 80/25mg if needed  vs  olmesartan-HCTZ 20/12.5mg FDC once daily titrated to 40/25mg if needed	hypertension $\geq 18$ years of age with clinic SBP 160 to 190 mmHg and DBP 119 mmHg or less		Secondary: Changes in clinic DBP, changes in ambulatory BP monitoring parameters, proportion of patients achieving BP target	olmesartan-HCTZ group.  Secondary: Greater reductions in clinic DBP in both azilsartan-chlorthalidone arms (-16.1 and -16.5 mmHg; $P < 0.001$ vs olmesartan-HCTZ for both) versus olmesartan-HCTZ (-12.8 mmHg). At eight weeks, both azilsartan-chlorthalidone doses reduced 24-hour SBP more than olmesartan-HCTZ (-26.4 and -27.9 versus -20.7 mmHg; both $P < 0.001$ ), and higher proportions in both azilsartan-chlorthalidone groups achieved target BP compared with the olmesartan-HCTZ group (69.4 and 68.9 versus 54.7%, both $P < 0.001$ ). Adverse events leading to drug discontinuation occurred in 6.2, 9.5, and 3.1% with the azilsartan-chlorthalidone lower and higher doses, and olmesartan-HCTZ, respectively.
Lithell et al. <sup>83</sup> (2003) SCOPE  Candesartan 16 mg/day  vs  placebo  Patients also received conventional therapy with diuretics, ACE inhibitors, $\beta$ -blockers, and calcium-channel blocking agents	DB, MC, PC, PG, RCT  Patients 70 to 89 years of age with mild-to-moderate HTN (SBP 160 to 179 mm Hg and/or DBP 90 to 99 mm Hg) and MMSE scores $\geq 24$	N=4,964  3.7 years	Primary: First major coronary event including cardiovascular death, nonfatal MI, or nonfatal stroke  Secondary: cardiovascular death, nonfatal and fatal stroke and MI, cognitive function	Primary: Results showed no significant difference in the primary end point between candesartan and placebo ( $P=0.19$ ).  Secondary: Candesartan treatment reduced nonfatal stroke by 27.8% ( $P=0.04$ ) and all stroke by 23.6% ( $P=0.056$ ) compared to placebo.  There were no significant differences in MI and cardiovascular mortality.  Mean MMSE score fell from 28.5 to 28.0 in the candesartan group and from 28.5 to 27.9 in the control group ( $P=0.20$ ). The proportion of patients who had a significant cognitive decline or developed dementia was not different in the 2 groups.
Baguet et al. <sup>84</sup> (2006)	DB, RCT	N=256	Primary: Change in mean	Primary: At the end of the six weeks, the mean change in DBP between the baseline and

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Candesartan 8 mg QD</p> <p>vs</p> <p>losartan 50 mg QD</p> <p>vs</p> <p>placebo</p>	<p>Patients with mild-to-moderate essential HTN (DBP 95 to 115 mm Hg)</p>	<p>6 weeks</p>	<p>ambulatory DBP from baseline to the 0-24 hour period after the last dose of study medication</p> <p>Secondary: Change in mean ambulatory SBP from baseline to the 0-24 hour period after the last dose of study medication, change in DBP and SBP during the daytime and nighttime, change in DBP and SBP between 12 and 24 hours after dosing</p>	<p>the 0-24 hour period after the last dose of study medication was greater in patients receiving candesartan 8 mg compared to losartan (-7.3 vs -5.1 mm Hg; P&lt;0.05) or placebo (0.3 mm Hg; P&lt;0.001).</p> <p>Secondary: The mean change in SBP between the baseline and the 0-24 hour period after the last dose of study medication was greater in patients receiving candesartan (-10.8 mm Hg) or losartan (-8.8 mm Hg) than placebo (1.2 mm Hg; P&lt;0.001).</p> <p>Candesartan was associated with a greater reduction in DBP and SBP relative to placebo, when compared to losartan during both the daytime and nighttime, and between 12 and 24 hours after dosing (P&lt;0.001).</p> <p>Both active treatments were well tolerated.</p>
<p>Ohma et al.<sup>85</sup> (2000)</p> <p>Candesartan 16 mg</p> <p>vs</p> <p>losartan 50 mg</p> <p>All patients received HCTZ 12.5 mg QD.</p>	<p>DB, MC, RCT</p> <p>Patients aged 20 to 80 years with mild-to-moderate uncontrolled HTN while on monotherapy (any kind of medication)</p>	<p>N=340</p> <p>12 weeks</p>	<p>Primary: Change in sitting DBP</p> <p>Secondary: SBP, proportion of responders, safety and tolerability</p>	<p>Primary: Greater reductions in DBP were reported with candesartan and HCTZ vs losartan and HCTZ (-10.4 vs -7.8 mm Hg; P=0.016).</p> <p>Secondary: Greater decreases in SBP were reported with candesartan and HCTZ (-19.4 mm Hg) vs losartan and HCTZ (-13.7 mm Hg; P=0.004).</p> <p>The proportion of patients achieving a DBP ≤90 mm Hg was greater with candesartan and HCTZ (60.9 vs 49.3%; P=0.044).</p> <p>There were eight withdrawals due to adverse effects in the candesartan and HCTZ group and 12 in the losartan and HCTZ group. The most common adverse effects were headache, tachycardia/palpitations, dizziness, and fatigue.</p>
<p>Mengden et al.<sup>86</sup> (2011)</p> <p>CHILI CU Soon</p>	<p>MC, OL, PRO</p> <p>High risk patients</p>	<p>N=4,131</p> <p>10 weeks</p>	<p>Primary: Change in baseline office blood</p>	<p>Primary: Baseline office blood pressure was 162.1±14.8/94.7±9.2 mm Hg, and after ten weeks, a reduction to 131.7±10.5/80.0±6.6 mm Hg was achieved (P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Candesartan and HCTZ 32-12.5 or 32-25 mg QD (fixed-dose combination)	≥18 years of age with uncontrolled HTN, on prior antihypertensive agents, and presence of additional cardiovascular risk factors		pressure and ambulatory blood pressure, safety  Secondary: Not reported	Reductions in blood pressure were comparable irrespective of prior or concurrent medications.  Baseline ambulatory blood pressure was 158.2/93.7 mm Hg during the day and 141.8/85.2 mm Hg during the night. After ten weeks, ambulatory blood pressure reduced to 133.6/80.0 and 121.0/72.3 mm Hg, respectively.  During the trial, 49 adverse events were reported in 1.19% of patients receiving combination therapy. Of these events, seven were regarded as serious, and most of the events were related to the nervous system or cardiac disorders.  Secondary: Not reported
McInnes et al. <sup>87</sup> (2000)  Candesartan and HCTZ 8-12.5 mg/day (fixed-dose combination product)  vs  lisinopril and HCTZ 10-12.5 mg/day (fixed-dose combination product)	DB, DD, MC, PG, RCT  Patients 20 to 80 years of age with mild-to-moderate HTN on prior antihypertensive monotherapy	N=355  26 weeks	Primary: Mean changes in DBP  Secondary: Mean changes in SBP and heart rate, proportion of responders and controlled patients, safety	Primary: Changes in mean sitting DBP did not differ significantly between the groups (mean difference, 0.5 mm Hg; P=0.20).  Secondary: No significant differences between the groups were reported for mean sitting SBP, heart rate, proportion of responders and controlled patients.  Both regimens were well tolerated but a greater percentage of those in the lisinopril based group (80 vs 69%) had a least one side effect (P=0.020). The proportion of patients spontaneously reporting cough (23.1 vs 4.6%) and discontinuing therapy due to adverse events (12.0 vs 5.9%) was also higher in the lisinopril based group compared to the candesartan based group.
Hosaka et al. <sup>88</sup> (2015)  Candesartan 8 mg/HCTZ 6.25 mg single pill combination QD	MC, OL, PRO, RCT  Patients, 20 to 80 years old, with newly diagnosed, untreated hypertensive	N=206  8 weeks	Primary: Difference in home morning SBP reduction  Secondary: Time-dependent	Primary: The home BP reduction at eight weeks after randomization was 11.4±1.3 mm Hg for SBP and 5.3±0.7 mm Hg for DBP in the combination group and 7.8±1.2 mm Hg for SBP and 3.6±0.6 mm Hg for DBP in the maximum dose group. Analyses using analysis of covariance adjusted by baseline home BP values as covariates showed that the combination regimen provided additional reductions of 4.0 mm Hg (95% CI, 0.8 to 7.2 mm Hg) for SBP and 1.8 mm Hg

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs candesartan 12 mg</p>	<p>patients or those who were on monotherapy with any antihypertensive drug alone</p>		<p>changes in the antihypertensive effects and nocturnal BP reduction</p>	<p>(95% CI, -0.02 to 3.7 mm Hg) for DBP over the maximum dose regimen at four weeks after randomization, whereas at eight weeks after randomization, these reductions were 4.0 mm Hg (95% CI, 0.9 to 7.2 mm Hg) and 1.7 mm Hg (95% CI, -0.05 to 3.4 mm Hg), respectively.</p> <p>Secondary: For home nocturnal BP, the analysis included the 53 patients who measured nocturnal BP at baseline, four and eight weeks after randomization. The reduction in home nocturnal BP at eight weeks after randomization was not different between the two regimen groups.</p> <p>The maximal antihypertensive effect and stabilization time for home SBP were 9.4 mm Hg and 37.1 days (P&lt;0.0001), respectively, with the combination regimen. The maximum dose regimen decreased home SBP with a very gentle slope, and estimated maximal effect and estimated stabilization time were not significant (P&gt;0.2). The rate of achieving target BP (home morning SBP &lt;135 mm Hg) was significantly higher with the combination regimen than with the maximum dose regimen (52.4 vs 30.1%, P=0.002).</p>
<p>Fogari et al.<sup>89</sup> (2007) CANDIA  Candesartan 16 mg and HCTZ 12.5 mg QD  vs  amlodipine 10 mg QD</p>	<p>DB, MC, RCT  Patients, 20 to 80 years old, with mild to moderate uncomplicated HTN not controlled on monotherapy with an antihypertensive (SBP &lt;180 mg Hg and DBP 90 to 110 mg Hg)</p>	<p>N=203  8 weeks</p>	<p>Primary: Decrease in DBP  Secondary: Sitting SBP, reduction of the orthostatic blood pressure at least two minutes after standing, change in heart rate, percentage of patients normalized (DBP &lt;90 mm Hg and SBP &lt;140 mm Hg), percentage of responders (reduction in DBP ≥5 mm Hg)</p>	<p>Primary: There was no significant difference in the mean decrease in DBP between treatment groups; the difference in final DBP was -0.02 mm Hg (95% CI, -1.48 to 1.52 mm Hg; P=0.979).</p> <p>Secondary: There was no significant difference between the groups at week eight for the following: sitting SBP (P=0.835), heart rate (P&lt;0.500), orthostatic SBP (P=0.883), orthostatic DBP (P=0.264), percentage of patients normalized (P=10), percentage of responders (P=0.900).</p> <p>The number of patients reporting an adverse event was greater in the amlodipine group (P=0.001).</p> <p>The number of patients reporting an adverse drug-related event was greater in the amlodipine group (P&lt;0.001).</p> <p>Changes in blood chemistry and other secondary measurements were not significantly different between the treatment groups.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Gradman et al.<sup>90</sup> (2005)</p> <p>Irbesartan 150 mg QD</p> <p>vs</p> <p>aliskiren 150 to 600 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PG, RCT</p> <p>Men and women, age 18 years or older, with mild-to-moderate essential HTN (mean sitting DBP <math>\geq</math>95 mm Hg and <math>&lt;</math>110 mm Hg)</p>	<p>N=652</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP and SBP</p> <p>Secondary: Proportion of patients achieving blood pressure control (<math>&lt;</math>140/90 mm Hg), safety</p>	<p>Primary: Decreases in mean sitting DBP at eight weeks were significantly greater with all doses of aliskiren compared to placebo (<math>P&lt;</math>0.001). The least-squares mean reductions in trough DBP for aliskiren 150, 300, and 600 mg were 9.3, 11.8, and 11.5 mm Hg, respectively, vs 6.3 mm Hg for placebo.</p> <p>Decreases in mean sitting SBP at eight weeks were significantly greater with all doses of aliskiren compared to placebo (<math>P&lt;</math>0.001). The least-squares mean reductions in trough SBP for aliskiren 150, 300, and 600 mg were 11.4, 15.8, and 15.7 mm Hg, respectively, vs 5.3 mm Hg for placebo.</p> <p>The antihypertensive effect of aliskiren 150 mg was comparable to irbesartan 150 mg with reductions of 8.9 and 12.5 mm Hg for mean sitting DBP and SBP, respectively. Aliskiren 300 and 600 mg produced significantly greater mean sitting DBP reductions than irbesartan 150 mg (<math>P&lt;</math>0.05). While the reductions in mean sitting SBP were greater with aliskiren 300 and 600 mg than irbesartan 150 mg, these differences were not statistically significant).</p> <p>Secondary: The percentage of patients achieving blood pressure control was significantly greater with all doses of aliskiren (37.8%-150 mg, 50.0%-300 mg, 45.7%-600 mg) and irbesartan (33.8%) compared to placebo (20.8%; <math>P&lt;</math>0.05). More patients on aliskiren 300 and 600 mg achieved blood pressure control compared to irbesartan (<math>P&lt;</math>0.05).</p> <p>Drug-related adverse events for both aliskiren and irbesartan were comparable to placebo and the most commonly reported adverse events were headache, dizziness, and diarrhea. The number of patients discontinuing therapy was similar in all groups.</p>
<p>Jordan et al.<sup>91</sup> (2007)</p> <p>Irbesartan 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p>	<p>DB, DD, MC, PG, RCT</p> <p>Obese men and women (BMI <math>\geq</math>30 kg/m<sup>2</sup>) <math>\geq</math>18 years with essential HTN (mean sitting DBP</p>	<p>N=489</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting DBP with aliskiren 300 mg plus HCTZ vs HCTZ alone at 8 weeks</p>	<p>Primary: Aliskiren 300 mg added to HCTZ 25 mg significantly reduced mean sitting DBP compared with HCTZ alone at week eight (mean difference, -4.0; <math>P&lt;</math>0.0001).</p> <p>Secondary: Aliskiren 300 mg added to HCTZ caused numerically larger reductions in mean sitting DBP and SBP compared with amlodipine 10 mg plus HCTZ and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>aliskiren 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>HCTZ 25 mg QD (existing therapy)</p>	<p>95 to 109 mm Hg and SBP &lt;180 mm Hg) who had not responded to 4 weeks of treatment with HCTZ 25 mg</p>		<p>Secondary:</p> <p>Comparisons of mean sitting DBP and SBP with aliskiren plus HCTZ vs the other treatment groups, percentage of responders (mean sitting DBP &lt;90 mm Hg or <math>\geq</math>10 mm Hg reduction from baseline), proportion of patients achieving blood pressure control (mean sitting blood pressure &lt;140/90 mm Hg), plasma renin activity, safety and tolerability</p>	<p>irbesartan 300 mg plus HCTZ at week eight, but there were no statistically significant differences between treatment groups (P&gt;0.05).</p> <p>Responder rates were significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0193) and week 12 (P=0.004) but comparable to responder rates observed with amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>The proportion of patients achieving blood pressure control was significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0005) and week 12 (P=0.0001) but not statistically different than amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>Plasma renin activity significantly increased (P&lt;0.05) during four weeks of HCTZ monotherapy. Combination with aliskiren neutralized this increase and led to an overall significant reduction in plasma renin activity compared with pretreatment baseline (P&lt;0.05) whereas amlodipine and irbesartan led to further significant increases (P&lt;0.05).</p> <p>All of the study treatments were generally well tolerated. Amlodipine plus HCTZ (45.2%) was associated with a higher incidence of adverse events than the other treatment groups (36.1 to 39.3%), largely due to a higher rate of peripheral edema (11.1 vs 0.8 to 1.6%).</p>
<p>O'Brien et al.<sup>92</sup> (2007)</p> <p>Irbesartan 150 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p> <p>vs</p>	<p>3 OL studies</p> <p>Men and women 18 to 80 years with ambulatory SBP <math>\geq</math>140 and <math>\leq</math>180 mm Hg without treatment</p>	<p>N=67</p> <p>6 to 9 weeks</p>	<p>Primary:</p> <p>Change in daytime systolic ABPM with combination therapy compared with monotherapy</p> <p>Secondary:</p> <p>Change in daytime diastolic ABPM, nighttime systolic and diastolic ABPM, daytime and nighttime heart</p>	<p>Primary:</p> <p>Aliskiren coadministered with HCTZ (P=0.0007) or ramipril (P=0.03) led to significantly greater reductions in daytime systolic ABPM compared to monotherapy. There was a trend for a reduction in daytime systolic ABPM with the addition of aliskiren to irbesartan; however, this trend was not statistically significant.</p> <p>Secondary:</p> <p>Aliskiren plus HCTZ significantly lowered daytime diastolic ABPM compared to aliskiren monotherapy (P=0.0006). Changes in nighttime systolic and diastolic ABPM followed similar trends but did not achieve statistical significance (P=0.06 and P=0.09, respectively). No changes in heart rate were observed with either aliskiren regimen.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>aliskiren 150 mg QD for 3 weeks, then HCTZ 25 mg QD was added for an additional 3 weeks (if ABPM remained <math>\geq</math>135/85 mm Hg)</p> <p>vs</p> <p>ramipril 5 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p>			<p>rates, plasma renin activity</p>	<p>Aliskiren added to irbesartan did not significantly change diastolic ABPM compared to irbesartan monotherapy; however, nighttime systolic and diastolic ABPM were significantly reduced (<math>P&lt;0.05</math> for all). No changes in heart rate were observed with either irbesartan regimen.</p> <p>Mean diastolic ABPM was significantly decreased with the addition of aliskiren 150 mg (<math>P&lt;0.05</math>) but not aliskiren 75 mg to ramipril monotherapy. Both aliskiren doses significantly decreased nighttime systolic and diastolic ABPM (<math>P&lt;0.05</math> for all). No changes in heart rate were observed with either ramipril regimen.</p> <p>Aliskiren alone significantly inhibited plasma renin activity by 65% (<math>P&lt;0.0001</math>), while ramipril and irbesartan monotherapy increased renin activity by 90 and 175%, respectively. When aliskiren was coadministered with HCTZ, ramipril or irbesartan, plasma renin activity remained similar to baseline levels or decreased.</p>
<p>Derosa et al.<sup>93</sup> (2005)</p> <p>Irbesartan 300 mg QD</p> <p>vs</p> <p>doxazosin 4 mg QD</p>	<p>DB, PG, RCT</p> <p>Patients with type 2 diabetes and mild HTN</p>	<p>N=96</p> <p>1 year</p>	<p>Primary: Blood pressure, glucose metabolism and lipid parameters</p> <p>Secondary: Not reported</p>	<p>Primary: Blood pressure was significantly reduced in both treatment groups compared to baseline (<math>P&lt;0.01</math>).</p> <p>Irbesartan was significantly better in lowering blood pressure compared to doxazosin (<math>P&lt;0.05</math>).</p> <p>Doxazosin significantly reduced glycosylated hemoglobin, fasting plasma glucose, fasting plasma insulin, TC, LDL-C, HDL-C, and TG (<math>P\leq 0.05</math> for all parameters).</p> <p>As monotherapy, neither of the drugs achieved adequate blood pressure control.</p> <p>Secondary: Not reported</p>
<p>Neutel et al.<sup>94</sup> (2006)</p>	<p>AC, DB, MC, RCT</p>	<p>N=737</p>	<p>Primary: Proportion of</p>	<p>Primary: Significantly more patients on combination therapy achieved seated DBP <math>&lt;90</math></p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Irbesartan 150 to 300 mg QD</p> <p>vs</p> <p>irbesartan and HCTZ 150 to 300-12.5 to 25 mg QD (fixed-dose combination)</p>	<p>Patients <math>\geq 18</math> years with severe HTN who were untreated (seated DBP <math>\geq 110</math> mm Hg) or currently receiving antihypertensive monotherapy with DBP <math>\geq 100</math> mm Hg</p>	<p>7 weeks</p>	<p>patients with DBP <math>&lt; 90</math> mm Hg at week 5</p> <p>Secondary: Proportion of patients who achieved seated SBP/DBP <math>&lt; 140/90</math> mm Hg</p>	<p>mm Hg at week five compared to monotherapy (47.2 vs 33.2%; <math>P=0.0005</math>).</p> <p>Secondary: Significantly more patients attained SBP/DBP <math>&lt; 140/90</math> mm Hg at week five (34.6 vs 19.2%, respectively; <math>P&lt;0.0001</math>), while the mean difference between combination and monotherapy in seated DBP and SBP was 4.7 and 9.7 mm Hg, respectively (<math>P&lt;0.0001</math>).</p> <p>Greater and more rapid blood pressure reduction with irbesartan and HCTZ was achieved without additional side effects.</p>
<p>Neutel (abstract).<sup>95</sup> (2011)</p> <p>Irbesartan and HCTZ 150-12.5 mg QD, up titrated to 300-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>irbesartan 150 mg QD, up titrated to 300 mg QD</p>	<p>Post-hoc analysis of 2 PRO, RCT</p> <p>Patients with uncontrolled or untreated moderate to severe HTN who are obese or who have diabetes</p>	<p>N=1,268</p> <p>7 weeks (severe HTN)</p> <p>12 weeks (moderate HTN)</p>	<p>Primary: Changes in baseline blood pressure, blood pressure goal rate</p> <p>Secondary: Safety</p>	<p>Primary: After seven to eight weeks of treatment, SBP/DBP decreased in patients with diabetes by 26.9/17.8 and 21.8/15.8 mm Hg with combination irbesartan and HCTZ and irbesartan treatment, respectively (<math>P=0.09/P=0.27</math>). In obese patients, SBP/DBP decreased by 29.4/20.2 and 20.1/15.9 mm Hg with combination irbesartan and HCTZ and irbesartan treatment, respectively (<math>P&lt;0.0001</math>).</p> <p>More patients with type 2 diabetes achieved a blood pressure goal of <math>&lt; 130/80</math> mm Hg at week seven to eight with combination irbesartan and HCTZ treatment compared to irbesartan (12 vs 5%; <math>P=0.22</math>). Significantly more obese patients achieved blood pressure goals with combination irbesartan and HCTZ treatment compared to irbesartan (48 vs 23%; <math>P&lt;0.0001</math>).</p> <p>Secondary: Treatment emergent adverse event rates were similar between treatment groups regardless of the presence of diabetes or BMI status. In patients with moderate or severe HTN and with a BMI <math>\geq 30</math> kg/m<sup>2</sup>, initial treatment with combination irbesartan and HCTZ was more effective compared to irbesartan.</p>
<p>Neutel et al.<sup>96</sup> (2008)</p> <p>Irbesartan and HCTZ 300-25 mg QD (fixed-dose combination)</p>	<p>AC, DB, RCT</p> <p>Patients with moderate HTN (seated SBP 160 to 179 mm Hg when DBP <math>&lt; 110</math> mm Hg;</p>	<p>N=538</p> <p>12 weeks</p>	<p>Primary: Change in SBP after week 8</p> <p>Secondary: Change from baseline in DBP at</p>	<p>Primary: At week eight, there was a reduction in SBP of 27.1 mm Hg with irbesartan and HCTZ compared to 22.1 mm Hg with irbesartan monotherapy (<math>P=0.0016</math>) and 15.7 mm Hg with HCTZ (<math>P&lt;0.0001</math>).</p> <p>Secondary: At week eight, there was a reduction in DBP of 14.6 mm Hg with irbesartan</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
product) vs irbesartan 300 mg QD vs HCTZ 25 mg QD	or DBP 100 to 109 mm Hg when SBP <180 mm Hg)		weeks 8 and 12, SBP at week 12, proportion of responders (SBP <140 mm Hg and DBP <90 m Hg) at weeks 8 and 12	and HCTZ compared to 11.6 mm Hg with irbesartan monotherapy (P=0.0013) and 7.3 mm Hg with HCTZ (P<0.0001).  A significantly greater percentage of patients reached a treatment goal of SBP <140 mm Hg and DBP <90 mm Hg by week eight with irbesartan and HCTZ (53.4%) compared to irbesartan (40.6%; P=0.0254) and HCTZ (20.2%; P<0.0001) alone.  Treatment was well tolerated in all three treatment groups with a slight increase in adverse events in the combination therapy group.
Weir et al. <sup>97</sup> (2007)  Irbesartan and HCTZ 300-25 mg QD (fixed-dose combination)	Pooled analysis of 2 DB, MC, RCT  Patients with stage 1 or 2 HTN evaluated according to age	N=796  7 to 8 weeks	Primary: Antihypertensive efficacy, tolerability  Secondary: Not reported	Primary: SBP/DBP reductions (27 to 31/16 to 22 mm Hg) were similar regardless of age, obesity and type 2 diabetes status and were greater in high- vs low-risk patients.  Dizziness (2.0 to 3.7%), hypotension (0 to 0.7%), and syncope (0%) were rare and not centered in any subgroup. There was no hypotension in the elderly or in patients with type 2 diabetes.  Secondary: Not reported
Bobrie et al. <sup>98</sup> (2005)  Irbesartan and HCTZ 150-12.5 mg QD (fixed-dose combination product) vs valsartan and HCTZ 80-12.5 mg QD (fixed-dose combination product)	OL, RCT  Patients whose blood pressure remained uncontrolled after 5 weeks of HCTZ 12.5 mg QD	N=464  8 weeks	Primary: Blood pressure reductions, safety  Secondary: Not reported	Primary: Irbesartan and HCTZ produced greater reductions in average SBP and DBP measured by home blood pressure monitoring than valsartan and HCTZ (SBP, -13.0 vs -10.6 mm Hg; P=0.0094; DBP, -9.5 vs -7.4 mm Hg; P=0.0007). These differences were more pronounced in the morning than in the evening.  Normalization rates observed with home blood pressure monitoring (SBP <135 mm Hg and DBP <85 mm Hg) were significantly greater with irbesartan and HCTZ than with valsartan and HCTZ (50.2 vs 33.2%; P=0.0003).  The overall safety was similar in the two groups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Stanton et al.<sup>99</sup> (2003)</p> <p>Losartan 100 mg QD</p> <p>vs</p> <p>aliskiren 37.5 to 300 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Men and women 21 to 70 years of age with mild-to-moderate HTN (SBP <math>\geq</math>140 mm Hg)</p>	<p>N=226</p> <p>4 weeks</p>	<p>Primary: Change in daytime ambulatory SBP</p> <p>Secondary: Changes in clinic SBP and DBP, plasma renin activity, plasma aliskiren levels, adverse events</p>	<p>Primary: A dose-dependent reduction in daytime ambulatory SBP was observed with increasing aliskiren doses (with mean changes of -0.40 mm Hg with aliskiren 37.5 mg, -5.3 mm Hg with aliskiren 75 mg, -8.0 mm Hg with aliskiren 150 mg, and -11 mm Hg with aliskiren 300 mg; P=0.0002). The change in daytime SBP with losartan 100 mg (-10.9 mm Hg) was significantly different than aliskiren 37.5 mg, but not the other higher aliskiren dosages).</p> <p>Secondary: Clinic SBP and DBP, both in the sitting and standing positions, decreased with aliskiren in a dose-dependent manner, whereas heart rate was unaltered. The decreases in clinic blood pressures were similar for losartan 100 mg and aliskiren 150 and 300 mg.</p> <p>Dose-dependent reductions in plasma renin activity were also observed (median change -55, -60, -77, and -83% with 37.5, 75, 150, and 300 mg aliskiren, respectively; P=0.0008). By contrast, plasma renin activity increased by 110% with losartan 100 mg.</p> <p>Rate of adverse events was 22% with aliskiren 37.5 mg, 35% with aliskiren 75 mg, 25% with aliskiren 150 mg, 23% with aliskiren 300 mg, and 32% with losartan 100 mg. There was no increase in the number of adverse events when increasing the dose of aliskiren.</p>
<p>Ribeiro et al.<sup>100</sup> (2007)</p> <p>LAMHYST</p> <p>Losartan 50 to 100 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>DB, DD, RCT</p> <p>Males and females, age 18 to 79 years old, with diagnosis of mild (&gt;95 mm Hg but &lt;115 mm Hg) to moderate essential HTN and not taking an antihypertensive medication (within last 4 weeks)</p>	<p>N=194</p> <p>12 weeks</p>	<p>Primary: Difference between treatment groups in mean change in ABPM for last 9 hours of treatment and during drug holiday</p> <p>Secondary: Not reported</p>	<p>Primary: After 12 weeks, mean reductions in SBP were significantly larger in the amlodipine group than the losartan group (-18.1 vs -10.1 mm Hg; P&lt;0.001). Mean reductions in DBP were significantly larger in the amlodipine group than the losartan group (-18.1 vs -10.1 mm Hg; P&lt;0.05).</p> <p>Mean increases in SBP were similar between the groups during the two day drug holiday (P&gt;0.05).</p> <p>After the two day drug holiday, SBP was lower than baseline in both groups (P&lt;0.001), with the amlodipine group SBP remaining significantly lower (P&lt;0.01).</p> <p>Mean increases in DBP were similar between the groups during the two day</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>drug holiday (P&gt;0.05). After the two day drug holiday, DBP was lower than baseline in both groups (P=0.0001), with the amlodipine group DBP remaining significantly lower (P&lt;0.05).</p> <p>Secondary: Not reported</p>
<p>Oparil et al.<sup>101</sup> (1996)</p> <p>Losartan 50 to 100 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p> <p>If goal DBP (<math>\leq 90</math> mm Hg) was not attained, drug doses could be doubled and/or HCTZ mg was added.</p>	<p>DB, DD, MC, RCT</p> <p>Patients with HTN</p>	<p>N=900</p> <p>12 weeks</p>	<p>Primary: Efficacy, tolerability, effects on QOL</p> <p>Secondary: Not reported</p>	<p>Primary: DBP reductions after 4, 8, and 12 weeks of therapy were clinically comparable (losartan group: 7.3, 10.4, and 11.1 mm Hg, respectively; amlodipine group: 7.9, 11.2, and 11.8 mm Hg, respectively; P value not significant).</p> <p>Similar reductions in SBP were seen for both treatment groups (P value not significant).</p> <p>The percentage of patients reaching goal DBP (<math>\leq 90</math> mm Hg) or DBP <math>\geq 90</math> mm Hg with a <math>\geq 10</math> mm Hg decrease from baseline) was comparable for the two groups, with 68% of patients in the losartan group and 71% of patients in the amlodipine group reaching goal.</p> <p>Significantly more patients in the amlodipine group had drug-related adverse experiences (27 vs 13%; P=0.029). Edema was more common in patients receiving the amlodipine regimen than in those receiving the losartan regimen (11 vs 1%; P=0.004).</p> <p>Overall QOL was not different in the two treatment groups.</p> <p>Secondary: Not reported</p>
<p>Dahlöf et al.<sup>102</sup> (2002)</p> <p>LIFE</p> <p>Losartan 50 to 100 mg QD</p> <p>vs</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200 to 95 to 115 mm Hg) and left ventricular</p>	<p>N=9,193</p> <p><math>\geq 4</math> years</p>	<p>Primary: Composite of cardiovascular death, MI and stroke</p> <p>Secondary: All-cause mortality, hospitalization for angina or heart</p>	<p>Primary: SBP fell by 30.2 and 29.1 mm Hg in the losartan and atenolol groups, respectively (treatment difference, P=0.017) and DBP fell by 16.6 and 16.8 mm Hg, respectively (treatment difference, P=0.37). MAP was 102.2 and 102.4 mm Hg, respectively (P value not significant). Heart rate decreased more in patients assigned to atenolol than losartan (-7.7 vs -1.8 beats/minute, respectively; P&lt;0.0001).</p> <p>Compared to atenolol, the primary composite occurred in 13.0% fewer patients</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
atenolol 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	hypertrophy		failure, revascularization procedures, resuscitated cardiac arrest, new-onset diabetes	receiving losartan (RR, 0.87; 95% CI, 0.77 to 0.98; P=0.021).  While there was no difference in the incidence cardiovascular mortality (P=0.206) and MI (P=0.491), losartan treatment resulted in a 24.9% relative risk reduction in stroke compared to atenolol (P=0.001).  Secondary: A 25% lower incidence of new-onset diabetes was reported with losartan compared to atenolol (P=0.001). There was no significant difference among the other secondary end points between the two treatment groups.  Note: At end point or end of follow-up, 18 and 26% of patients on losartan were receiving HCTZ alone or with other drugs, respectively. In the atenolol group, 16 and 22% of patients were receiving HCTZ alone or with other drugs, respectively.
Julius et al. <sup>103</sup> (2004) LIFE Black Subset  Losartan 50 to 100 mg QD  vs  atenolol 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	Post hoc analysis  Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy	N=523  ≥4 years	Primary: Composite of cardiovascular death, MI and stroke  Secondary: Not reported	Primary: Compared to atenolol (11.2%), losartan in the United States African American population resulted in a greater incidence of the composite end point (17.4%; P=0.033).  HRs favored atenolol across all parameters (P=0.246 for cardiovascular mortality, P=0.140 for MI, and P=0.030 for stroke).  In African American patients, blood pressure reduction was similar in both groups, and regression of electrocardiographic-left ventricular hypertrophy was greater with losartan.  Secondary: Not reported
Lindholm et al. <sup>104</sup> (2002) LIFE Diabetic Subset  Losartan 50 to 100	Post hoc analysis  Patients 55 to 80 years old with essential HTN (sitting SBP/DBP	N=1,195  ≥4 years	Primary: Composite of cardiovascular death, MI and stroke  Secondary:	Primary: Compared to atenolol, losartan resulted in a 24% decrease in the primary composite end point (P=0.031).  Losartan treatment resulted in a 37% risk reduction in cardiovascular deaths vs atenolol (P=0.028).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg QD vs atenolol 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy		All-cause mortality	Losartan treatment resulted in a 39% risk reduction in all-cause mortality vs atenolol (P=0.002).  Mean blood pressure fell to 146/79 mm Hg in losartan patients and 148/79 mm Hg in atenolol patients.  Secondary: Mortality from all causes was 63 and 104 in the losartan and atenolol groups, respectively (RR, 0.61; P=0.002).
Kjeldsen et al. <sup>105</sup> (2002) LIFE Isolated Systolic Hypertension Subset  Losartan 50 to 100 mg QD vs atenolol 50 to 100 mg QD  HCTZ 12.5 to 25 mg QD was added if needed for blood pressure control.	Post hoc analysis  Patients 55 to 80 years old with isolated systolic HTN (SBP of 160 to 200 mm Hg and DBP <90 mm Hg) and left ventricular hypertrophy	N=1,326  ≥4 years	Primary: Composite of cardiovascular death, MI, or stroke  Secondary: All-cause mortality	Primary: Compared to atenolol, losartan resulted in a trend towards a 25% reduction in the primary end point (P=0.06).  Losartan treatment resulted in a 46% risk reduction in cardiovascular mortality (P=0.01) and 40% risk reduction in stroke compared to atenolol (P=0.02). There was no difference in the incidence of MI.  Blood pressure was reduced by 28/9 and 28/9 mm Hg in the losartan and atenolol arms.  Secondary: Patients receiving losartan also had reductions in all-cause mortality (28%; P<0.046).
Fossum et al. <sup>106</sup> (2006) ICARUS, a LIFE substudy  Losartan 50 to 100 mg/day	DB, DD, PG, RCT  Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115	N=81  3 years	Primary: Amount and density of atherosclerotic lesions in the common carotid arteries and carotid bulb	Primary: The amount of plaque decreased in the losartan group and increased in the atenolol group, though the difference between groups was not statistically significant (P=0.471).  Patients in the atenolol group had a greater increase in plaque index compared to the losartan group, though the difference between groups was not

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>atenolol 50 to 100 mg/day</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>mm Hg) and left ventricular hypertrophy</p>		<p>Secondary: Not reported</p>	<p>statistically significant (P=0.742)</p> <p>Secondary: Not reported</p>
<p>Kizer et al.<sup>107</sup> (2005) LIFE substudy</p> <p>Losartan 50 to 100 mg/day</p> <p>vs</p> <p>atenolol 50 to 100 mg/day</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy</p>	<p>N=9,193</p> <p>≥4 years</p>	<p>Primary: Reduction in the risk of different stroke subtypes and neurological deficits</p> <p>Secondary: Not reported</p>	<p>Primary: The risk of fatal stroke was significantly decreased in the losartan group compared to the atenolol group (P=0.032).</p> <p>The risk of atherothrombotic stroke was significantly decreased in the losartan group compared to the atenolol group (P=0.001).</p> <p>Comparable risk reductions were observed for hemorrhagic and embolic stroke but did not reach statistical significance.</p> <p>The risk of recurrent stroke was significantly reduced in the losartan arm compared to the atenolol arm (P=0.017).</p> <p>The number of neurological deficits per stroke was similar (P=0.68), but there were fewer strokes in the losartan group for nearly every level of stroke severity.</p> <p>Secondary: Not reported</p>
<p>Wachtell et al.<sup>108</sup> (2005) LIFE substudy</p> <p>Losartan 50 to 100 mg/day</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115</p>	<p>N=8,851 (patients in LIFE with no baseline history of atrial fibrillation</p>	<p>Primary: Incidence of new-onset atrial fibrillation and outcome</p> <p>Secondary:</p>	<p>Primary: Significantly fewer patients in the losartan group experienced new-onset atrial fibrillation compared to the atenolol group (P&lt;0.001).</p> <p>Randomization to losartan treatment was associated with a 33% lower rate of new onset atrial fibrillation independent of other risk factors (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs atenolol 50 to 100 mg/day</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>mm Hg) and left ventricular hypertrophy</p>	<p>but at risk for atrial fibrillation)</p> <p>≥4 years</p>	<p>Not reported</p>	<p>Patients in the losartan group had a 40% lower rate of composite events consisting of cardiovascular death, fatal or non-fatal stroke, and fatal or non-fatal MI (P=0.03).</p> <p>Significantly fewer strokes occurred in the losartan group compared to the atenolol group (P=0.01), and there was a trend toward fewer MIs in the losartan group (P=0.16).</p> <p>There was no significant difference in cardiovascular mortality between groups.</p> <p>In contrast, the atenolol group experienced significantly fewer hospitalizations for heart failure (P=0.004) and a trend toward fewer sudden cardiac deaths (P=0.07).</p> <p>Secondary: Not reported</p>
<p>Wachtell et al.<sup>109</sup> (2005) LIFE substudy</p> <p>Losartan 50 to 100 mg/day</p> <p>vs atenolol 50 to 100 mg/day</p> <p>All patients received HCTZ 12.5 to 25 mg/day if need for blood pressure control.</p>	<p>DB, DD, PG, RCT</p> <p>Patients 55 to 80 years old with essential HTN (sitting SBP/DBP 160 to 200/95 to 115 mm Hg) and left ventricular hypertrophy</p>	<p>N=342 (LIFE patients with AF at the start of the LIFE study)</p> <p>≥4 years</p>	<p>Primary: Cardiovascular morbidity and mortality</p> <p>Secondary: Not reported</p>	<p>Primary: Patients with a history of atrial fibrillation had significantly higher rates of cardiovascular and all-cause mortality, fatal and non-fatal stroke, heart failure, revascularization and sudden cardiac death compared to patients without atrial fibrillation (P&lt;0.001).</p> <p>Patients with a history of atrial fibrillation had similar rates of MI and hospitalization for angina pectoris (P≥0.209).</p> <p>The primary composite endpoint of cardiovascular mortality, stroke and MI occurred in significantly fewer patients in the losartan group compared to the atenolol group (P=0.009).</p> <p>The difference in MI between groups was not significant.</p> <p>Treatment with losartan trended toward lower all-cause mortality (P=0.09) and fewer pacemaker implantations (P=0.065).</p> <p>Secondary: Not reported</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Van Bortel et al.<sup>110</sup> (2005)</p> <p>Losartan 50 mg QD</p> <p>vs</p> <p>nebivolol 5 mg QD</p> <p>If after 6 weeks, DBP was not normalized, then HCTZ 12.5 mg QD was added to therapy</p>	<p>DB, MC, PG, RCT</p> <p>Patients &lt;70 years of age with DBP at randomization between 95 and 114 mm Hg</p>	<p>N=314</p> <p>12 weeks</p>	<p>Primary: Effects on blood pressure, overall QOL</p> <p>Secondary: Comparison of different aspects of QOL</p>	<p>Primary: At the end of 12 weeks, both nebivolol and losartan significantly reduced SBP compared to baseline (P&lt;0.0001 for both), but the agents were not significantly different from each other.</p> <p>Both agents also significantly decreased DBP compared to baseline (P&lt;0.0001), but nebivolol significantly reduced DBP compared to losartan (P&lt;0.02).</p> <p>At the end of 12 weeks, both nebivolol and losartan significantly improved QOL scores compared to baseline (P&lt;0.007), but the agents were not significantly different from each other.</p> <p>Secondary: At week 12 there was not a significant difference observed in the individual questions of the QOL questionnaire between the groups. Questions inquired about headaches, lightheadedness, sleepiness, flushing, and sexual function.</p>
<p>Flack et al.<sup>111</sup> (2003)</p> <p>Losartan 50 mg QD</p> <p>vs</p> <p>eplerenone 50 mg QD</p> <p>vs</p> <p>placebo</p> <p>Doses were increased if blood pressure remained uncontrolled.</p>	<p>DB, MC, PG, RCT</p> <p>Men and women ≥18 years old, with mild to moderate HTN, with SBP &lt;180 mm Hg and DBP 95 to 109 mm Hg (off medication) or if patients were receiving antihypertensive therapy their blood pressure was &lt;140/90 mm Hg</p>	<p>N=551</p> <p>16 weeks</p>	<p>Primary: Mean change from baseline in DBP at 16 weeks</p> <p>Secondary: Mean change from baseline at 16 weeks in SBP, SBP and DBP within and between racial groups, response rate (defined as the percentage of patients with DBP &lt;90 mm Hg or DBP ≥90 mm Hg but ≥10 mm Hg below baseline), urinary albumin/creatinine ratio, effect of</p>	<p>Primary: At 16 weeks, patients randomized to eplerenone exhibited significantly greater mean changes in DBP from baseline compared to either losartan- or placebo-treated groups (P&lt;0.001).</p> <p>Secondary: At 16 weeks, patients randomized to eplerenone exhibited significantly greater mean changes in SBP from baseline compared to either losartan- or placebo-treated groups (P&lt;0.001).</p> <p>At 16 weeks, African American patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the placebo-treated African American patients (P&lt;0.001).</p> <p>At 16 weeks, African American patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the losartan-treated African American patients (P≤0.001).</p> <p>At 16 weeks, white patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the placebo-treated white patients (P=0.001). However, the difference in SBP- and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>eplerenone in patients with various baseline renin and aldosterone levels, adverse effects</p>	<p>DBP-lowering effects was not significant different between the eplerenone and losartan groups (P=0.126, P=0.068, respectively).</p> <p>Significantly greater percentage of patients randomized to eplerenone exhibited a positive response to therapy compared to either placebo (64.5 vs 41.2%; P&lt;0.001) or losartan group (64.5 vs 48.3%; P=0.003).</p> <p>The eplerenone group (regardless of race) exhibited statistically significant improvement in urinary albumin/creatinine ratio from baseline compared to placebo (P=0.003). However, the difference in urinary albumin/creatinine ratio change from baseline was not significantly different between the eplerenone and losartan groups (P=0.652).</p> <p>Compared to losartan, eplerenone was more effective in lowering SBP and DBP in patients with low-moderate baseline renin levels (P&lt;0.05). However, the difference was not statistically significant in patients with high baseline renin levels.</p> <p>Compared to losartan, eplerenone was more effective in lowering SBP in patients with low or high baseline aldosterone levels (P&lt;0.05). However, the difference was not statistically significant in patients with moderate baseline aldosterone levels.</p> <p>Compared to losartan, eplerenone was more effective in lowering DBP in patients with low baseline aldosterone levels (P&lt;0.05). However, the difference was not statistically significant in patients with moderate-high baseline aldosterone levels.</p> <p>There were no significant differences in the incidence of adverse events noted in eplerenone, placebo or losartan groups. The reported incidence of gynecomastia, breast pain, menstrual abnormalities, impotence, hyperkalemia and decreased libido with eplerenone was low and comparable to losartan and placebo.</p>
<p>Hood et al.<sup>112</sup> (2007) SALT</p>	<p>DB, RCT, XO  Adult patients with seated blood</p>	<p>N=57  42 weeks</p>	<p>Primary: Change in blood pressure and plasma renin from baseline</p>	<p>Primary: Spironolactone 100 mg/day- and bendroflumethiazide 5 mg/day-treated patients did not exhibit a significant difference in BP reduction from baseline (P value not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Losartan 100 mg/day</p> <p>vs</p> <p>spironolactone 50 mg/day</p> <p>vs</p> <p>spironolactone 100 mg/day</p> <p>vs</p> <p>amiloride 20 mg/day</p> <p>vs</p> <p>amiloride 40 mg/day</p> <p>vs</p> <p>bendroflumethiazide* 2.5 mg/day</p> <p>vs</p> <p>bendroflumethiazide* 5 mg/day</p> <p>vs</p>	<p>pressure of 140/90 to 170/110 mm Hg, plasma renin of <math>\leq 12</math> mU/L, plasma aldosterone-renin ratio <math>&gt;750</math>, previous fall in SBP <math>\geq 20</math> mm Hg after 1 month of OL treatment with spironolactone 50 mg/day</p>		<p>between spironolactone 100 mg/day and bendroflumethiazide 5 mg/day</p> <p>Secondary: Change in blood pressure and plasma renin from baseline between amiloride and other diuretics and between lower and higher doses of each diuretic</p>	<p>Secondary: Spironolactone 50 mg/day-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (<math>P&lt;0.01</math>).</p> <p>Losartan 100 mg-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (<math>P&lt;0.05</math>).</p> <p>High-dose bendroflumethiazide- and amiloride-treated patients exhibited significantly greater reductions in blood pressure compared to the lower doses (<math>P&lt;0.05</math>).</p> <p>Spironolactone-treated patients exhibited a four-fold increase in baseline renin level compared to a two-fold increase observed in bendroflumethiazide-treated patients (<math>P=0.003</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo				
Maeda et al. <sup>113</sup> (2012) ARCH  Losartan and HCTZ (fixed-dose combination product)	MC, OL, OS, PRO  Patients 20 to 80 years of age with HTN uncontrolled by either ARB monotherapy or combination with and ARB and a calcium channel blocker	N=614  52 weeks	Primary: Change in blood pressure at 3 months  Secondary: Not reported	Primary: Blood pressure decreased significantly to 138.0/78.2 mm Hg by month three (P<0.001), and 36.2% of patients were able to achieve target blood pressure (P<0.05).  The hypotensive effect lasted for one year (P<0.001) and was found equally in patients receiving losartan-HCTZ and losartan-HCTZ plus a calcium channel blocker.  Secondary: Not reported
Ueda et al. <sup>114</sup> (abstract) (2012) MAPPY  Losartan and HCTZ 50-12.5 mg QD (fixed-dose combination product)  vs  losartan 100 mg QD	MC, OL, PG, PRO, RCT  Patients with morning HTN	N=216  Duration not specified	Primary: Change in baseline SBP, blood pressure control rate  Secondary: Safety	Primary: Morning SBP was reduced from 150.3±10.1 to 131.5±11.5 mm Hg with combination therapy (P<0.001) and from 151.0±9.3 to 142.5±13.6 mm Hg with high dose losartan therapy (P<0.001). The morning SBP reduction was significantly greater with combination therapy group compared to high dose losartan therapy (P<0.001).  Combination therapy decreased evening SBP from 141.6±13.3 to 125.3±13.1 mm Hg (P<0.001), and high dose losartan therapy decreased evening SBP from 138.9±9.9 to 131.4±13.2 mm Hg (P<0.01).  Although both therapies improved target blood pressure achievement rates in the morning and evening (P<0.001 for both), combination therapy significantly increased the achievement rates compared to high dose losartan therapy (P<0.001 and P<0.05, respectively).  Secondary: Combination therapy decreased urine albumin excretion (P<0.05) whereas high-dose therapy reduced serum uric acid. Both therapies indicated strong adherence and few adverse effects (P<0.001).
Salerno et al. <sup>115</sup> (2004)  Losartan and HCTZ 50-12.5 to	DB, RCT  Patients with severe HTN	N=585  6 weeks	Primary: Proportion of patients achieving goal blood pressure	Primary: Almost twice as many patients achieved goal blood pressure at four weeks on losartan 50 mg and HCTZ 12.5 mg vs losartan 50 to 100 mg monotherapy (P=0.002).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>100-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>losartan 50 to 100 mg QD</p> <p>Doses were titrated as needed to reach blood pressure goal (&lt;90 mm Hg).</p>			<p>Secondary: Adverse events</p>	<p>Almost three times as many patients achieved goal blood pressure at six weeks with losartan and HCTZ vs losartan monotherapy (P&lt;0.001).</p> <p>Adverse experiences on losartan and HCTZ (43%) were significantly less than with losartan monotherapy (53%).</p>
<p>Minami et al.<sup>116</sup> (2007)</p> <p>Losartan 50 mg/day and HCTZ 12.5 mg/day</p> <p>Candesartan 8 mg QD (n=10) or amlodipine 5 mg QD (n=5) administered to all patients for 2 months prior to switch to losartan plus HCTZ.</p>	<p>OL</p> <p>Japanese outpatients with essential HTN treated for ≥2 months with either candesartan or amlodipine and 24-hour ambulatory blood pressure ≥135/80 mm Hg</p>	<p>N=15</p> <p>12 months</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: In patients who had previously received candesartan, 24-hour blood pressure decreased significantly from 137/89 mm Hg to 126/81 mm Hg after three months (P&lt;0.05/P&lt;0.001) and to 123/81 mm Hg after 12 months (P&lt;0.01/P&lt;0.001) of treatment with losartan and HCTZ.</p> <p>In patients who had previously received amlodipine, 24-hour blood pressure decreased significantly from 137/81 to 125/75 mm Hg after three months (P&lt;0.05/P&lt;0.05) and to 124/77 mm Hg after 12 months (P&lt;0.05/P value not significant) of treatment with losartan and HCTZ.</p> <p>There were significant decreases in SBP during the daytime, nighttime and early morning after 12 months in both groups.</p> <p>No adverse changes in the indices of glucose or lipid metabolism were observed in either group.</p> <p>Secondary: Not reported</p>
<p>Lacourcière et al.<sup>117</sup> (2003) PROBE</p>	<p>DB, MC, OL, RCT</p> <p>Patients ≥18 years of age with mild-to-</p>	<p>N=597</p> <p>6 weeks</p>	<p>Primary: Mean changes in ambulatory DBP</p>	<p>Primary: During the last six hours of the dosing interval, telmisartan 40 mg and HCTZ 12.5 mg and telmisartan 80 mg and HCTZ 12.5 mg reduced mean DBP to a greater extent vs losartan 50 mg and HCTZ 12.5 mg. Treatment differences</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Losartan and HCTZ 50-12.5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and HCTZ 40-12.5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and HCTZ 80-12.5 mg QD (fixed-dose combination product)</p>	<p>moderate essential HTN</p>		<p>Secondary: Mean changes in ambulatory SBP, 24-hour DBP, safety</p>	<p>between the groups were 1.8 mm Hg (P&lt;0.05) and 2.5 mm Hg (P&lt;0.001) lower, respectively, with the telmisartan and HCTZ arms.</p> <p>Secondary: Telmisartan 40 mg and HCTZ 12.5 mg and telmisartan 80 mg and HCTZ 12.5 mg produced greater reductions in ambulatory SBP vs losartan 50 mg and HCTZ 12.5 mg of 2.5 and 3.4 mm Hg, respectively, during the last six hours of the dosing interval (P&lt;0.05), and of 2.1 and 3.4 mm Hg, respectively, over the entire 24-hour dosing interval (P&lt;0.05).</p> <p>Telmisartan 80 mg and HCTZ 12.5 mg also lowered mean 24-hour DBP by 2.3 mm Hg more than losartan 50 mg and HCTZ 12.5 mg (P&lt;0.001).</p> <p>All treatments were well tolerated.</p>
<p>Brunner et al.<sup>118</sup> (2006)</p> <p>Olmesartan 20 mg QD</p> <p>vs</p> <p>candesartan 8 mg QD</p>	<p>DB, RCT</p> <p>Patients with mainly mild-to-moderate HTN</p>	<p>N=635</p> <p>8 weeks</p>	<p>Primary: 24-hour antihypertensive efficacy (with particular emphasis on blood pressure control during the early morning period), proportion of patients who achieved various ABPM goals (SBP/DBP &lt;125/80 mm Hg)</p> <p>Secondary:</p>	<p>Primary: After eight weeks, significantly greater proportions of patients treated with olmesartan achieved 24-hour and daytime ABPM goals 25.6 and 18.3%, respectively) compared to candesartan (14.9%; P&lt;0.001 and 9.6%; P=0.002, respectively).</p> <p>During the last four hours of 24-hour ABPM, the proportion of patients who achieved goals was significantly greater with olmesartan (33.3%) than candesartan (22.9%; P&lt;0.001).</p> <p>Similarly, during the last two hours of 24-hour ABPM, the proportion of patients who achieved these blood pressure goals was higher with olmesartan (26.9 and 19.9%) compared to candesartan (19.6%; P=0.028 and 14.3%; P=0.061).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Not reported	Not reported
<p>Punzi et al.<sup>119</sup> (2012)</p> <p>Olmesartan 20 mg QD, up titrated to 40 mg QD</p> <p>vs</p> <p>losartan 50 mg QD, up titrated to 100 mg QD</p>	<p>DB, PRO, RCT</p> <p>Patients with HTN no previously treated or previously treated with antihypertensive medications</p>	<p>N=941</p> <p>8 weeks</p>	<p>Primary: Change in baseline seated cuff DBP at week 8</p> <p>Secondary: Mean change in seated cuff SBP at weeks 4 and 8 and seated cuff DBP at week 4, blood pressure target rates, safety</p>	<p>Primary: Olmesartan produced significantly greater LSM reductions in seated cuff DBP compared to losartan in treatment-naïve (-9.7±1.0 vs -6.6±1.0 mm Hg; P=0.0232) and treatment-experienced patients (-9.6±0.5 vs -7.3±0.5 mm Hg; P=0.0013).</p> <p>Secondary: Both treatment-naïve (-12.1±1.2 vs -8.5±1.3 mm Hg; P=0.0379) and treatment-experienced patients (-12.0±0.7 vs -8.5±0.7 mm Hg; P=0.0006) receiving olmesartan had significantly greater reductions in baseline cuff seated SBP compared to losartan at week 4. Similar results were observed at week eight (P=0.0178 and P=0.0016).</p> <p>A similar trend in significantly greater baseline reductions with olmesartan compared to losartan was observed at week four for seated cuff DBP in treatment-naïve (LSM difference, -2.3±1.10; P=0.0337) and treatment-experienced patients (LSM difference, -2.7±0.67; P&lt;0.0001).</p> <p>A significantly greater proportion of treatment-naïve patients receiving olmesartan achieved a seated cuff blood pressure goal of &lt;140/90 mm Hg with olmesartan compared to losartan (34.1 vs 19.0%; P=0.0109). Similar results were observed in treatment-experienced patients (31.0 vs 19.6%; P=0.0008).</p> <p>Treatment-emergent adverse events were reported in 30.5 and 31.4% of treatment-naïve and treatment-experienced patients receiving olmesartan. Corresponding proportions for losartan were 33.0 and 31.2%. Most events were mild to moderate in severity.</p>
<p>Oparil et al.<sup>120</sup> (2001)</p> <p>Olmesartan 20 mg QD</p> <p>vs</p> <p>irbesartan 150 mg</p>	<p>DB, MC, PG, RCT</p> <p>Patients ≥18 years old (mean age 52 years) with essential HTN (cuff DBP ≥100 mm Hg and ≤115 mm Hg and mean daytime DBP</p>	<p>N=588</p> <p>8 weeks</p>	<p>Primary: Change in seated cuff DBP at week 8</p> <p>Secondary: Change in seated cuff SBP at week 8, 24-hour DBP and SBP, adverse events</p>	<p>Primary: The mean reductions in seated cuff DBP at week eight were significantly greater with olmesartan (11.5 mm Hg) than with irbesartan (9.9 mm Hg; P=0.0412), losartan (8.2 mm Hg; P=0.0002) and valsartan (7.9 mm Hg; P&lt;0.0001).</p> <p>The clinical significance of a few mm Hg DBP difference between the groups is unknown.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD, losartan 50 mg QD, or valsartan 80 mg QD</p>	<p>≥90 mm Hg and &lt;120 mm Hg)</p>			<p>Secondary: Reductions of cuff SBP were not significantly different among the four ARBs and ranged from 8.4 to 11.3 mm Hg.</p> <p>The reduction in mean 24-hour DBP with olmesartan (8.5 mm Hg) was significantly greater than reductions with losartan and valsartan (6.2 and 5.6 mm Hg, respectively) and showed a trend toward significance when compared to irbesartan (7.4 mm Hg; P=0.087).</p> <p>The reduction in mean 24-hour SBP with olmesartan (12.5 mm Hg) was significantly greater than the reductions with losartan and valsartan (9.0 and 8.1 mm Hg, respectively) and equivalent to the reduction with irbesartan (11.3 mm Hg).</p> <p>All drugs were well tolerated with the incidence of adverse events reported in 30.6% of patients in the olmesartan group, 35.6% for irbesartan, 32.0% for losartan, and 44.8% for valsartan.</p>
<p>Chrysant et al.<sup>121</sup> (2004)</p> <p>Olmesartan 10 to 40 mg QD and HCTZ 12.5 to 25 mg QD</p> <p>vs</p> <p>olmesartan 10 to 40 mg QD</p> <p>vs</p> <p>HCTZ 12.5 to 25 mg QD</p> <p>vs</p>	<p>DB, RCT, factorial design</p> <p>Patients with a baseline mean seated DBP of 110 to 115 mm Hg</p>	<p>N=502</p> <p>8 weeks</p>	<p>Primary: Change in DBP at week 8</p> <p>Secondary: Change in SBP at week 8</p>	<p>Primary: Olmesartan and HCTZ produced greater reductions in seated DBP at week eight than did monotherapy with either component. All olmesartan and HCTZ combinations significantly reduced DBP compared to placebo in a dose-dependent manner.</p> <p>Reductions in mean trough DBP were 8.2, 16.4, and 21.9 mm Hg with placebo, olmesartan 20 mg plus HCTZ 12.5 mg, and olmesartan 40 mg plus HCTZ 25 mg, respectively.</p> <p>Secondary: Olmesartan and HCTZ produced greater reductions in seated SBP at week eight than did monotherapy with either component. All olmesartan and HCTZ combinations significantly reduced DBP compared to placebo in a dose-dependent manner.</p> <p>Reductions in mean trough SBP were 3.3, 20.1, and 26.8 mm Hg with placebo, olmesartan 20 mg plus HCTZ 12.5 mg, and olmesartan 40 mg plus HCTZ 25 mg, respectively.</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo				All treatments were well tolerated.
<p>Kereiakes et al.<sup>122</sup> (2007)</p> <p>Benazepril 10 mg/day for 2 weeks, then 20 mg/day for 2 weeks, then benazepril 20 mg/day plus amlodipine 5 mg/day for 4 weeks, then benazepril 20 mg/day plus amlodipine 10 mg/day for 4 weeks</p> <p>vs</p> <p>olmesartan 20 mg/day for 2 weeks, then 40 mg/day for 2 weeks then olmesartan and HCTZ 40-12.5 mg/day for 4 weeks increased to 40-25 mg for 4 weeks</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients with stage 2 HTN</p>	<p>N=190</p> <p>12 weeks</p>	<p>Primary: Change in mean seated SBP at the end of week 12</p> <p>Secondary: DBP at the end of week 12, percent of patients attaining blood pressure goals of &lt;140/90, &lt;130/85, and &lt;130/80 mm Hg</p>	<p>Primary: Patients treated with olmesartan and HCTZ experienced significantly greater reductions in mean seated SBP at week 12 than patients treated with benazepril plus amlodipine (least square mean change, -32.5 vs -26.5 mm Hg; P=0.024; least square mean treatment difference, -6.0 mm Hg; 95% CI, -11.1 to -0.8).</p> <p>Secondary: The least square mean change for reduction in DBP approached statistical significance with olmesartan and HCTZ compared to benazepril plus amlodipine at week 12 (P=0.056).</p> <p>The percentage of patients achieving goal rates at the end of the study for olmesartan and HCTZ and benazepril plus amlodipine were 66.3 and 44.7% (P=0.006) for &lt;140/90 mm Hg, 44.9 vs 21.2% (P=0.001) for &lt;130/85 mm Hg, and 32.6 and 14.1% (P=0.006) for &lt;130/80 mm Hg.</p> <p>Both treatments were well tolerated.</p>
<p>Chrysant et al.<sup>123</sup> (2008) COACH</p>	<p>DB, MC, PC, RCT</p> <p>Patients, age 18 years and older,</p>	<p>N=1,940</p> <p>8 weeks</p>	<p>Primary: Change from baseline in seated DBP at week 8</p>	<p>Primary: All active treatments and placebo resulted in significant decreases in seated DBP at week eight (P&lt;0.001). Reductions in seated DBP with monotherapy treatment ranged from -8.3 to -12.7 mm Hg; reductions with combination</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Olmesartan 10 to 40 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>olmesartan 10 to 40 mg and amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>placebo</p>	<p>with seated DBP of 95 to 120 mm Hg</p>		<p>Secondary: Change from baseline in seated SBP at week 8, mean change from baseline in seated DBP and SBP at weeks 2, 4, 6 and 8 without last observation carried forward, proportion of patients achieving BP goal (&lt;140/90 mm Hg or &lt;130/80 mm Hg), safety</p>	<p>therapy ranged from -13.8 to -19.0 mm Hg. All combinations reduced seated DBP significantly greater than either component as monotherapy at the same dosage (P&lt;0.001).</p> <p>Secondary: All active treatments and placebo resulted in significant decreases in seated SBP at week eight (P&lt;0.001 for treatment, P=0.024 for placebo). All combinations reduced seated SBP significantly greater either component as monotherapy at the same dosage (P&lt;0.001).</p> <p>The proportion of patients achieving goal blood pressures were: 20.0 to 36.3% of patients receiving olmesartan monotherapy, 21.1 to 32.5% of patients receiving amlodipine monotherapy, 35.0 to 53.2% of patients receiving combination therapy, and 8.8% of patients receiving placebo.</p> <p>Combination therapy resulted in significantly greater achievement of goal blood pressure than monotherapy (P&lt;0.005).</p> <p>No difference in overall rates of adverse events across the different treatment groups was seen. Nearly 27% of patients experienced a drug-related adverse event.</p> <p>Changes in laboratory values were not considered clinically significant nor followed a consistent pattern with treatment: none of the changes were considered clinically significant. Platelet counts increased significantly from baseline (statistically) for patients receiving amlodipine, however the increase was &lt;10% and not deemed clinically relevant.</p>
<p>Chrysant et al.<sup>124</sup> (2009) COACH</p> <p>Olmesartan 10 to 40 mg QD and amlodipine 5 to 10 mg QD</p> <p>HCTZ 12.5 to 25</p>	<p>OL, ES</p> <p>Patients ≥ 18 years of age with essential HTN (seated DBP ≥95 and &lt;120 mm Hg)</p>	<p>N=1,684</p> <p>44 weeks OL therapy (52 weeks total study duration including 8 week DB phase)</p>	<p>Primary: Reduction in mean sitting SBP DBP, change in mean sitting SBP and DBP, percentage of patients achieving blood pressure goal (&lt;140/90 mm Hg or</p>	<p>Primary: Mean sitting DBP decreased from 101.5 mm Hg at baseline to 81.9 mm Hg and mean sitting SBP decreased from 163.6 mm Hg at baseline to 131.2 mm Hg at week 52.</p> <p>Approximately 31% of patients remained on amlodipine 5 mg and olmesartan 40 mg. Increasing the dose of amlodipine to 10 mg in combination with olmesartan 40 mg produced further decreases in mean sitting DBP of 4.8 mm Hg and mean sitting SBP of 7.3 mm Hg. Addition of HCTZ 12.5 mg to amlodipine 10 mg and olmesartan 40 mg decreased mean sitting DBP by 4.5</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg could be added if blood pressure was not controlled (&lt;140/90 mm Hg or &lt;130/80 mm Hg in patients with diabetes).</p>			<p>&lt;130/80 mm Hg for patients with diabetes)</p>	<p>mm Hg and mean sitting SBP by 7.7 mm Hg. Doubling the HCTZ dose from 12.5 to 25 mg decreased mean sitting DBP and mean sitting SBP by an additional 6.0 mm Hg and 9.9 mm Hg, respectively. Patients who received the triple therapy had the greatest mean sitting SBP reduction (36.1 mm Hg).</p> <p>Approximately 67% of patients achieved blood pressure goal by week 52. The blood pressure goal achievement was 80% for amlodipine and olmesartan 5/40 mg, 70.6% for amlodipine and olmesartan 10/40 mg, 66.6% for amlodipine and olmesartan and HCTZ 10/40/12.5 mg, and 46.3% for amlodipine and olmesartan and HCTZ 10/ 40/25 mg.</p> <p>The addition of HCTZ 25 mg enabled more patients to achieve blood pressure targets of &lt;140/90 mm Hg (77.7%), &lt;130/85 mm Hg (47.5%), and &lt;130/80 mm Hg (36.4%) compared to the other treatment regimens.</p> <p>No major safety issues emerged with long-term therapy. The frequency of edema ranged from 8.9% in patients treated with amlodipine 5 mg and olmesartan 40 mg to 14.5% in patients treated with amlodipine 10 mg and olmesartan 40 mg plus HCTZ 25 mg. Other treatment-emergent adverse events experienced by ≥3% of patients included upper respiratory tract infection (6.5%), nasopharyngitis (5.2%), extremity pain (4.1%), sinusitis (3.6%), arthralgia (3.3%), and back pain (3.1%). headache (2.0%), hypotension (1.8%), and fatigue (1.6%). The incidence of cough was 0.4%.</p>
<p>Oparil et al.<sup>125</sup> (2009) COACH</p> <p>Amlodipine 5 to 10 mg QD and olmesartan 10 to 40 mg</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>DB, factorial, MC, PC, RCT</p> <p>Patients ≥18 years of age with seated DBP 95 to 120 mm Hg, with a subgroup analysis based on HTN (stage 1: SBP 140 to 159 mm Hg or DBP 90 to 99 mm Hg; stage 2: SBP ≥160 mm Hg or DBP ≥100 mm Hg)</p>	<p>N=1,940</p> <p>8 weeks</p>	<p>Primary: Mean change in DBP and SBP at week 8 for each subgroup</p> <p>Secondary: Proportion of patients achieving blood pressure goal (&lt;140/90 mm Hg or &lt;130/80 mm Hg)</p>	<p>Primary: Reductions in mean DBP as a result of combination treatment were similar between subgroups. Patients with stage 1 HTN achieved reductions of 14.8 to 15.8 mm Hg and patients with stage 2 HTN achieved reductions of 13.6 to 19.8 mm Hg. Reductions in mean SBP as a result of combination treatment resulted in greater reductions in patients with stage 2 HTN (25.1 to 32.7 mm Hg) compared to stage 1 HTN (17.7 to 23.7 mm Hg) (P value not reported).</p> <p>Reductions in mean DBP and SBP were similar between those with no prior antihypertensive treatment and those with prior hypertensive treatment.</p> <p>Secondary: The proportion of patients with stage 1 HTN who received combination treatment and achieved blood pressure goal was 65.6 to 80.0%, compared to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs olmesartan 10 to 40 mg QD vs placebo	and no prior antihypertensive medication			40.5 to 66.7% of those who received monotherapy (P<0.0001 across treatments).  The proportion of patients with stage 2 HTN who received combination treatment and achieved BP goal was 40.5 to 49.2%, compared to 13.1 to 29.2% of those who received monotherapy (P<0.0001).  Results of patients with baseline SBP ≥180 mm Hg were similar to other subgroups.
Braun et al. <sup>126</sup> (abstract) (2009)  Amlodipine 10 mg plus olmesartan 20 mg QD  If patients were uncontrolled after 4 weeks, they were changed to amlodipine and valsartan 10-160 mg QD.	OL, PRO  Patients with DBP 100 to 109 mm Hg	N=257  8 weeks	Primary: Reduction in SBP and DBP  Secondary: Adverse events	Primary: Following treatment with amlodipine and olmesartan, SBP/DBP decreased by 19.2±12.4/14.4±7.4 mm Hg.  The number of patients who progressed to treatment with amlodipine and valsartan was 175. Additional reductions in SBP of 7.9 mm Hg and DBP of 3.9 mm Hg were seen (P<0.0001 for both).  Secondary: Both treatments were well tolerated and reported adverse events were consistent with drug profiles.
Chrysant et al. <sup>127</sup> (2012) TRINITY  Olmesartan and amlodipine and HCTZ 40-10-25 mg/day (fixed-dose combination product)  vs	DB, MC, PG, RCT  Patients ≥18 years of age with mean sitting blood pressure ≥140/100 mm Hg or ≥160/90 mm Hg (off antihypertensive medication)	N=2,492  12 weeks	Primary: Change in baseline mean sitting DBP  Secondary: Change in baseline mean sitting SBP, blood pressure goal rate, safety	Primary: In both Black and non-Black patients, triple combination treatment resulted in significantly greater reductions in mean sitting DBP compared to combination therapies (P≤0.0001). Overall, triple combination treatment reduced LSM mean sitting blood pressure by -37.1/20.8 and -38.9/21.8 mm Hg in Black and non-Black patients at week 12 (P<0.0001 vs combination therapies).  Secondary: In both Black and non-Black patients, triple combination treatment resulted in significantly greater reductions in mean sitting SBP compared to combination therapies (P<0.0001).  A significantly greater proportion of patients receiving triple combination

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
component dual-combination treatments				<p>treatment achieved blood pressure goal compared to combination therapies, regardless of race.</p> <p>No new safety concerns were identified with any treatment. The majority of treatment emergent adverse events were mild to moderate in severity. Treatment emergent adverse events occurred in 366 (52.0%) and 921 (57.6%) Black and non-Black patients.</p>
<p>Chrysant et al.<sup>128</sup> (abstract) (2012) TRINITY</p> <p>Olmesartan and amlodipine and HCTZ 40-10-25 mg/day (fixed-dose combination product)</p> <p>vs</p> <p>component dual-combination treatments</p>	<p>Subgroup analysis</p> <p>Patients <math>\geq 18</math> years of age with HTN and diabetes</p>	<p>N=not reported</p> <p>12 weeks</p>	<p>Primary: Change in baseline blood pressure, blood pressure control rate</p> <p>Secondary: Safety</p>	<p>Primary: The prespecified changes in blood pressure from baseline for the diabetes subgroup receiving triple combination treatment were significantly greater compared to the dual-combination treatments (<math>P \leq 0.0013</math>).</p> <p>Significantly more patients with diabetes receiving triple combination treatment achieved goal blood pressure (<math>&lt; 130/80</math> mm Hg) compared to patients receiving dual combination treatments (<math>P \leq 0.0092</math>).</p> <p>Secondary: Most treatment-emergent adverse events were mild to moderate in severity.</p>
<p>Kereiakes et al.<sup>129</sup> (2011) TRINITY</p> <p>Olmesartan and amlodipine and HCTZ 40-10-25 mg/day (fixed-dose combination product)</p> <p>vs</p>	<p>ES, OL</p> <p>Patients <math>\geq 18</math> years of age with mean sitting blood pressure <math>\geq 140/100</math> mm Hg or <math>\geq 160/90</math> mm Hg (off antihypertensive medication)</p>	<p>N=2,112</p> <p>40 weeks</p>	<p>Primary: Efficacy, safety</p> <p>Secondary: Not reported</p>	<p>Primary: Mean changes in blood pressure from baseline to week 52 were comparable for all treatments. The proportion of patients receiving triple combination treatment who achieved blood pressure goals at week 52 ranged between 44.5 to 79.8% depending on the dose; lower doses were associated with a smaller proportion of patients achieving blood pressure goals.</p> <p>No new safety concerns were identified. Most adverse events and drug-related adverse events were considered to be of mild to moderate severity. One hundred and six patients reported a serious adverse event and five drug-related adverse events. Serious drug-related adverse events included acute renal insufficiency, presyncope, and hypotension in three patients; acute renal insufficiency with hyperkalemia in one patients; and syncope in one patient.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
component dual-combination treatments				Secondary: Not reported
Punzi, HA <sup>130</sup> (2014)  Once daily olmesartan medoxomil (OM)/amlodipine besylate (AM)/HCTZ 40/10/25 mg	OL, PRO, blinded-endpoint  Adults on 1, 2, or 3 antihypertensive medications and not at goal BP, defined as less than 140/90 mmHg or less than 130/80 mmHg if they had diabetes or renal disease	N=40  2 to 9 day screening period, followed by 4 to 6 weeks of open-label treatment	Primary: Mean change from baseline in 24 hour SBP ABPM at day 1  Secondary: Mean change from baseline in 24 hour DBP ABPM at day 1, the change from baseline in mean trough seated BP at weeks 1, 2, 3, and 4	Primary: At day 1, treatment with OM/AM/HCTZ resulted in a significant mean reduction from baseline in ambulatory SBP reduction of $5.55 \pm 1.3$ mmHg ( $P < 0.0001$ ).  Secondary: Significant proportion of patients (90%) receiving OM/AM/HCTZ achieved the seated BP goal of $< 140/90$ mmHg at week 4, with 97% achieving $< 140$ mmHg.  The proportion of patients achieving the 24 hour ambulatory BP target of $< 130/80$ mmHg was 84% at week 4. At day 1, for the secondary endpoints, treatment with OM/AM/HCTZ resulted in a significant mean reduction from baseline in ambulatory DBP of $2.55 \pm 1.0$ ( $P < 0.0052$ ), seated cuff SBP reduction of $9.78 \pm 1.5$ ( $P < 0.0001$ ), and seated cuff DBP reduction of $4.13 \pm 1.4$ ( $P < 0.0052$ ).
Sharma et al. <sup>131</sup> (2012)  Telmisartan  vs  placebo  All patients are receiving amlodipine	DB, PG, RCT  Patients with type 2 diabetes and stage 1 or 2 HTN	N=981  8 weeks	Primary: Change in mean seated trough cuff SBP at weeks 8  Secondary: Blood pressure goal rates; change in mean seated trough cuff SBP at weeks 1, 2, and 4; safety	Primary: After eight weeks, significantly greater reductions in mean seated trough cuff SBP was achieved with telmisartan compared to placebo ( $-29.0$ vs $-22.9$ mm Hg; $P < 0.0001$ ).  Secondary: After eight weeks, 71.4 and 53.8% of patients achieved blood pressure goal ( $< 140/90$ mm Hg) with telmisartan compared to placebo. A blood pressure goal of $< 130/80$ mm Hg was achieved by 36.4 and 17.9% of patients receiving telmisartan and placebo.  Significant reductions in mean seated trough cuff SBP with telmisartan were evidence from week one ( $P < 0.0001$ ) and continued throughout the trial.  The most common adverse events were peripheral edema, headache, and dizziness.
Williams et al. <sup>132</sup> (2009)	Pooled analysis: blinded endpoint,	N=1,613	Primary: Change from	Primary: A significantly greater reduction in mean ambulatory blood pressure during

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>PRISMA I and PRISMA II</p> <p>Ramipril 2.5 mg QD for 2 weeks then force titration to 5 mg QD for 6 weeks then 10 mg QD for 6 weeks</p> <p>vs</p> <p>telmisartan 40 mg QD for 2 weeks then force titration to 80 mg QD for 12 weeks</p>	<p>OL, PRO, RCT</p> <p>Patients <math>\geq 18</math> years of age with mild- to moderate HTN</p>	<p>14 weeks</p>	<p>baseline in mean ambulatory BP during the final 6 hours of the 24-hour dosing interval</p> <p>Secondary: Change from baseline in mean ambulatory blood pressure during the 24-hour dosing interval, morning, daytime and nighttime ambulatory blood pressure, 24-hour blood pressure load, treatment response, blood pressure control</p>	<p>the last six hours of the 24-hour dosing interval was observed with telmisartan 80 mg group compared to ramipril 5 and 10 mg (P&lt;0.0001).</p> <p>Secondary: Significantly greater reductions in mean 24-hour, morning, daytime, nighttime and 24-hour blood pressure load were observed with telmisartan 80 mg compared to ramipril 5 and 10 mg (P&lt;0.0001).</p> <p>Significantly greater reductions in treatment response and blood pressure control rates were observed with telmisartan 80 mg compared to ramipril 5 and 10 mg (P&lt;0.0001).</p>
<p>Karlberg et al.<sup>133</sup> (1999) TEES</p> <p>Enalapril 5 to 20 mg QD</p> <p>vs</p> <p>telmisartan 20 to 80 mg QD</p> <p>HCTZ 12.5 or 25 mg QD could be added to either group as needed to</p>	<p>DB, DD, MC, PG, RCT</p> <p>Patients <math>\geq 65</math> years of age with mild- to moderate HTN</p>	<p>N=278</p> <p>26 weeks</p>	<p>Primary: Change from baseline in supine SBP and DBP</p> <p>Secondary: Proportion of responders, safety</p>	<p>Primary: Both treatments had similar rates of HCTZ use.</p> <p>Both treatments showed comparable decreases in blood pressure. Mean changes in DBP were -12.8 mm Hg for telmisartan and -11.4 mm Hg for enalapril (P=0.074). Mean changes in SBP were -22.1 mm Hg for telmisartan and -20.1 mm Hg for enalapril (P=0.350).</p> <p>Secondary: Overall, 63 and 62% of patients responded to telmisartan and enalapril, respectively, with a DBP of &lt;90 mm Hg. Both regimens provided effective blood pressure lowering over the 24-hour dosing interval, as determined by ambulatory blood pressure monitoring.</p> <p>Both regimens were well tolerated; however, the enalapril group had a higher incidence of cough than the telmisartan group (15.8 vs 6.5%; P value</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
reach DBP goal ( $\leq 90$ mm Hg)				reported).
Xi et al. <sup>134</sup> (2008)  Telmisartan  vs  losartan	MA  Patients with HTN	N=1,832 (11 trials)  Variable duration	Primary: Reduction in DBP and SBP  Secondary: Therapeutic response of DBP and SBP, tolerability	Primary: Use of telmisartan resulted in a significant reduction in clinic DBP (WMD, 1.52; 95% CI, 0.85 to 2.19) and SBP (WMD, 2.77; 95% CI, 1.90 to 3.63) when compared to losartan.  Secondary: There was also a significant reduction in 24-hour mean ambulatory DBP (WMD, 2.49; 95% CI, 0.56 to 4.42) and SBP (WMD, 2.47; 95% CI, 0.40 to 4.55) with telmisartan as compared to losartan.  There was a significant increase in therapeutic response of DBP (RR, 1.14; 95% CI, 1.04 to 1.23) and SBP response (RR, 1.10; 95% CI, 1.01 to 1.20) with telmisartan as compared to losartan.  Both telmisartan and losartan were well tolerated.
Sharma et al. <sup>135</sup> (2007)  Telmisartan and amlodipine 40-5 mg QD (fixed-dose combination)  vs  amlodipine 5 mg QD	DB, MC, RCT  Patients 18 to 65 years of age with established stage II uncomplicated essential HTN	N=210  12 weeks	Primary: SBP/DBP reductions and responder rates (SBP/DBP <130/<80 mm Hg)  Secondary: Not reported	Primary: There was a significant reduction from baseline in mean SBP in both groups (telmisartan and amlodipine, from 176.3 to 128.0 mm Hg; amlodipine, from 171.8 to 143.4 mm Hg; both, P<0.05 vs baseline). There was a significant reduction in SBP from baseline in the telmisartan and amlodipine and amlodipine groups (-27.4% and -16.6%, respectively; P<0.05 within group and between groups).  There was a significant reduction from baseline in mean DBP in both treatment groups (telmisartan and amlodipine, from 100.9 to 93.8 mm Hg; amlodipine, from 99.7 to 94.3 mm Hg; both, P<0.05). There was a 20.2% reduction in mean DBP in the telmisartan and amlodipine group, which was significantly greater compared to the reduction of 12.7% observed in the amlodipine group (P<0.05 between groups and within both groups).  A total of 87.3% of patients receiving telmisartan and amlodipine reached the target SBP/DBP goal, compared to 69.3% of patients receiving amlodipine (P<0.05).  A total of 16.0% of patients in the telmisartan and amlodipine group



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>experienced adverse events compared to 15.4% of patients in the amlodipine group (P value not significant). The most common adverse events in the telmisartan and amlodipine group were peripheral edema (8.5%), headache (5.7%), dizziness and cough (3.8%), and diarrhea (1.9%).</p> <p>Secondary: Not reported</p>
<p>Littlejohn et al.<sup>136</sup> (2009)</p> <p>Telmisartan 20 to 80 mg and amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>telmisartan 20 to 80 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients ≥18 years of age with Stage 1 or 2 HTN (DBP ≥95 and ≤119 mm Hg)</p>	<p>N=2,607</p> <p>8 weeks</p>	<p>Primary: Change in the in-clinic seated diastolic BP</p> <p>Secondary: Change in the in-clinic seated SBP, DBP and SBP response (DBP &lt;90 mm Hg, decrease in DBP ≥10 mm Hg, SBP &lt;140 mm Hg, decrease in SBP ≥15 mm Hg), and BP control (DBP &lt;90 mm Hg and SBP &lt;140 mm Hg)</p>	<p>Primary: Both telmisartan (irrespective of amlodipine dosage; P&lt;0.0001) and amlodipine (irrespective of telmisartan dosage; P&lt;0.0001) significantly lowered the in-clinic DBP.</p> <p>The greatest reduction in blood pressure was with telmisartan 80 mg plus amlodipine 10 mg (SBP/DBP -26.4/-20.1 mm Hg; P&lt;0.05 vs both monotherapies).</p> <p>DBP and SBP response was achieved by 91.2 and 90.4% of patients in the telmisartan 80 mg plus amlodipine 10 mg group, respectively.</p> <p>More than 50% of patients treated with combination therapy achieved blood pressure control, with the highest percentages (76.5% [overall control] and 85.3% [DBP control]) being achieved by patients treated with telmisartan 80 mg plus amlodipine 10 mg.</p> <p>A total of 37.3% of patients reported at least one adverse event. The most commonly reported adverse events were headache (5.4%) and peripheral edema (4.4%). Headache was more frequent in the placebo group (10.9%) compared to the telmisartan monotherapy (5.9%), amlodipine monotherapy (6.0%), and combination therapy (4.7%). The incidence of peripheral edema was highest in the amlodipine 10-mg group (17.8%); however, this rate was lower when amlodipine was used in combination with telmisartan: 11.4% (telmisartan 20 mg and amlodipine 10 mg), 6.2% (telmisartan 40 mg and amlodipine 10 mg), and 11.3% (telmisartan 80 mg and amlodipine 10 mg).</p>
<p>Littlejohn et al.<sup>137</sup> (2009)</p> <p>Telmisartan and</p>	<p>DB, DD, MC, PC, PG, RCT</p> <p>Patients ≥18 years</p>	<p>N=1,078</p> <p>8 weeks</p>	<p>Primary: Change in DBP from baseline to study end point</p>	<p>Primary: Significant reductions in DBP were seen from baseline to study end for both dual therapy and monotherapy (P values not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amlodipine 40-5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 40-10 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 80-5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan and amlodipine 80-10 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>respective monotherapies, dosing frequency not specified</p>	<p>of age with stage 1 or 2 HTN (DBP <math>\geq</math>95 and <math>\leq</math>119 mm Hg), with a subgroup analysis including patients with DBP <math>\geq</math>100 mm Hg at baseline</p>	<p>N=858</p>	<p>Secondary: Change from baseline to study end in SBP; percent of patients achieving a DBP response (DBP &lt;90 mm Hg) and SBP response (SBP &lt;140 mm Hg or reduction from baseline <math>\geq</math>15 mm Hg); percent of patients achieving BP control (SBP/DBP &lt;140/&lt;90 mm Hg) and DBP control (&lt;90 mm Hg) and safety</p>	<p>Amlodipine 5 and 10 mg with telmisartan 40 and 80 mg significantly reduced DBP compared to respective monotherapies (P values not reported).</p> <p>Secondary: Amlodipine 5 and 10 mg with telmisartan 40 and 80 mg significantly reduced SBP compared to respective monotherapies (P values not reported).</p> <p>Combination therapy resulted in a greater DBP and SBP response than monotherapy (P values not reported).</p> <p>The highest rate of BP control was achieved with amlodipine 10 mg with telmisartan 80 mg.</p> <p>Rates of adverse events were similar between dual therapy and monotherapy. Incidences of adverse events were 4.40% with telmisartan monotherapy, 11.00% with amlodipine monotherapy and 11.75% with combination therapy. The most commonly reported events were headache and peripheral edema. Patients receiving amlodipine 10 mg had the highest incidence of peripheral edema; however rates were lower when amlodipine was used in combination with telmisartan.</p>
Neutel et al. <sup>138</sup>	DB, MC, PG, RCT	N=858	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2012) TEAMSTA</p> <p>Telmisartan and amlodipine 80-10 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>telmisartan 80 mg QD</p> <p>vs</p> <p>amlodipine 10 mg QD</p>	<p>Patients ≥18 years of age with severe HTN</p>	<p>8 weeks</p>	<p>Change in baseline blood pressure, blood pressure goal and response rates</p> <p>Secondary: Safety</p>	<p>Reductions in seated trough cuff blood pressure (-47.5/-18.7 mm Hg) were significantly greater with combination therapy compared to telmisartan (P&lt;0.001) or amlodipine (P=0.002). Significant reductions with combination therapy were observed at one, two, four, and six weeks.</p> <p>Blood pressure goal and response rates were consistently higher with combination therapy (50.4 and 91.4 to 99.7%) compared to monotherapy with either agent (24.1 and 69.3 to 91.5% and 35.6 and 83.9 to 98.5%).</p> <p>Secondary: Combination therapy was well tolerated and fewer adverse events were reported with combination therapy compared to amlodipine (12.6 vs 16.4%). Peripheral edema was reported more frequently with amlodipine compared to combination therapy (13.2 vs 9.3%).</p>
<p>Oparil et al.<sup>139</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>valsartan 160 to 320 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and valsartan 160 to 320 mg QD</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Men and women aged 18 years or over with stage 1-2 essential HTN (mean sitting DBP 95 to 109 mm Hg and 8-hr ambulatory DBP ≥90 mm Hg)</p>	<p>N=1,797</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, proportion of patients achieving a successful response to treatment (mean sitting DBP &lt;90 mm Hg and/or ≥10 mm Hg reduction from baseline) or achieving blood pressure control (mean sitting SBP/DBP &lt;140/90)</p>	<p>Primary: The combination of aliskiren 300 mg and valsartan 320 mg lowered mean sitting DBP from baseline by 12.2 mm Hg, significantly more than either monotherapy with aliskiren 300 mg (-9.0 mm Hg; P&lt;0.0001), valsartan 320 mg (-9.7 mm Hg; P&lt;0.0001) or with placebo (-4.1 mm Hg; P&lt;0.0001). Monotherapy with aliskiren or valsartan provided significantly greater reductions in mean sitting DBP than did placebo at week 8 (P&lt;0.0001 for all).</p> <p>Secondary: The combination of aliskiren 300 mg and valsartan 320 mg lowered mean sitting SBP from baseline by 17.2 mm Hg, significantly more than either monotherapy with aliskiren 300 mg (-13.0 mm Hg; P&lt;0.0001), valsartan 320 mg (-12.8 mm Hg; P&lt;0.0001), or with placebo (-4.6 mm Hg; P&lt;0.0001). Monotherapy with aliskiren or valsartan provided significantly greater reductions in mean sitting SBP than did placebo at week eight end point (all P&lt;0.0001).</p> <p>The proportion of patients achieving a successful response to treatment at week eight was significantly higher with the combination of aliskiren and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo			mm Hg), change in 24-hr ABPM, change in biomarkers, safety	<p>valsartan (66%) than with aliskiren alone (53%; P=0.0003) or valsartan alone (55%; P=0.0010). All active treatments were associated with significantly greater responder rates than placebo (30%; P&lt;0.0001 for all).</p> <p>The proportion of patients achieving blood pressure control was significantly greater in the combination group (49%) than in the aliskiren (37%; P=0.0005) or valsartan (34%; P&lt;0.0001) monotherapy groups. All active treatments were associated with significantly greater control rates than placebo (16%; P&lt;0.0001 for all).</p> <p>The combination of aliskiren and valsartan was significantly more effective in lowering mean 24-hr ambulatory SBP and DBP than was either agent alone (P&lt;0.0001 for all). The greater reductions in ambulatory blood pressure with aliskiren plus valsartan were maintained throughout the entire 24-hour dosing interval.</p> <p>Aliskiren and valsartan (P&lt;0.0001) and monotherapy with aliskiren (P&lt;0.0001) or valsartan (P=0.0002) provided significant increases in plasma renin concentrations versus placebo. Increases in plasma renin concentrations were significantly greater for the combination than aliskiren (P=0.0014) or valsartan (P&lt;0.0001) monotherapy.</p> <p>Valsartan monotherapy produced significantly greater increases in plasma renin activity than placebo (160 vs 18%; P=0.0003). By contrast, aliskiren alone significantly reduced plasma renin activity by 73% (P&lt;0.0001 vs placebo), while the combination of aliskiren plus valsartan led to a reduction in plasma renin activity of 44% (P&lt;0.0001 vs placebo).</p> <p>The combination of aliskiren and valsartan (-31%; P&lt;0.0001) and valsartan monotherapy (-25%; P=0.0007) provided significantly greater reductions in plasma aldosterone concentration than did placebo (7%), while aliskiren monotherapy had no significant effect (-5.9%; P=0.1059).</p> <p>Rates of adverse events and laboratory abnormalities were similar in all groups.</p>
Yarows et al. <sup>140</sup> (2008)	Post-hoc analysis of patients with stage 2	N=1,797	Primary: Change in mean	Primary: In patients with stage 2 HTN, significantly greater reductions in DBP were

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Aliskiren 150 mg QD for 4 weeks, followed by 300 mg QD for 4 weeks</p> <p>vs</p> <p>valsartan 160 mg QD for 4 weeks, followed by 320 mg QD for 4 weeks</p> <p>vs</p> <p>aliskiren and valsartan 150-160 mg QD for 4 weeks, followed by 300-320 mg QD for 4 weeks (fixed-dose combination products)</p> <p>vs</p> <p>placebo</p>	<p>HTN from Oparil et al.</p> <p>Men and women <math>\geq 18</math> years of age with stage 1 to 2 essential HTN (mean sitting DBP 95 to 109 mm Hg and 8-hour ambulatory DBP <math>\geq 90</math> mm Hg)</p>	<p>8 weeks</p>	<p>sitting DBP</p> <p>Secondary: Change in mean sitting SBP, proportion of patients achieving a successful response to treatment (mean sitting DBP <math>&lt; 90</math> mm Hg and/or <math>\geq 10</math> mm Hg reduction from baseline) or achieving blood pressure control (mean sitting SBP/DBP <math>&lt; 140/90</math> mm Hg)</p>	<p>demonstrated in the aliskiren and valsartan 300-320 mg group compared to either higher-dose monotherapy group (<math>P &lt; 0.05</math>) and placebo (<math>P &lt; 0.0001</math>).</p> <p>Secondary: In patients with stage 2 HTN, significantly greater reductions in SBP were demonstrated in the aliskiren and valsartan 300-320 mg group compared to either higher-dose monotherapy group (<math>P &lt; 0.05</math>) and placebo (<math>P &lt; 0.0001</math>).</p> <p>DBP and SBP reductions in both monotherapy groups were significantly greater compared to placebo (<math>P &lt; 0.0001</math>).</p> <p>The proportion of patients with stage 2 HTN achieving blood pressure control at week eight was significantly greater in the aliskiren and valsartan 300-320 mg group compared to both monotherapy groups and placebo (<math>P \leq 0.044</math>).</p> <p>Blood pressure control rates in the aliskiren group were significantly greater than placebo (<math>P &lt; 0.001</math>). No significant difference was observed between the valsartan monotherapy and placebo groups.</p>
<p>Pool et al.<sup>141</sup> (2007)</p> <p>Aliskiren 75 to 300 mg QD</p> <p>vs</p>	<p>DB, MC, PC, PG, RCT</p> <p>Men and women <math>\geq 18</math> years with mild-to-moderate essential HTN</p>	<p>N=1,123</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, safety</p>	<p>Primary: Aliskiren 300 mg significantly (<math>P &lt; 0.0001</math>) lowered mean sitting DBP compared with placebo. Reductions in mean sitting DBP for aliskiren 75 and 150 mg compared to placebo failed to reach statistical significance (<math>P = 0.052</math> and <math>P = 0.051</math>, respectively).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>valsartan 80 to 320 mg</p> <p>vs</p> <p>aliskiren 75 to 300 mg and valsartan 80 to 320 mg</p> <p>vs</p> <p>valsartan and HCTZ 160-12.5 mg QD (fixed-dose combination)</p> <p>vs</p> <p>placebo</p>	<p>(mean sitting DBP <math>\geq</math>95 mm Hg after a 3- to 4-week single-blind placebo run-in period)</p>			<p>Aliskiren 300 mg significantly (<math>P&lt;0.0001</math>) lowered mean sitting SBP compared with placebo.</p> <p>A statistically significant linear dose relationship was observed for the effect of aliskiren (75 to 300 mg) on mean sitting DBP (<math>P=0.0002</math>) and mean sitting SBP (<math>P=0.0005</math>). The effects of aliskiren monotherapy on mean sitting DBP and SBP across the 75 to 300 mg dose range were similar to the effects of valsartan 80 to 320 mg.</p> <p>Coadministration of aliskiren and valsartan produced a greater antihypertensive effect than either drug alone. Reductions in mean sitting DBP and SBP obtained with aliskiren 150 mg plus valsartan 160 mg and aliskiren 300 mg plus valsartan 320 mg were not significantly different from those observed with valsartan 160 mg plus HCTZ 12.5 mg.</p> <p>Responder rates were significantly greater than placebo for all 3 aliskiren monotherapy groups and for all aliskiren plus valsartan combinations. The proportion of responders with aliskiren 75 mg plus valsartan 80 mg was significantly greater than either component monotherapy (<math>P&lt;0.05</math>). There was no significant difference between the proportion of responders to aliskiren 150 mg plus valsartan 160 mg or aliskiren 300 mg plus valsartan 320 mg compared with valsartan 160 mg plus HCTZ 12.5 mg.</p> <p>Control rates were higher with aliskiren 300 mg compared with placebo and with valsartan 160 mg plus HCTZ 12.5 mg compared with aliskiren 150 mg plus valsartan 160 mg, but there were no significant differences between aliskiren plus valsartan combinations and the respective monotherapies.</p> <p>Aliskiren and valsartan were generally well tolerated either as monotherapy or in combination. The overall incidence of adverse events and rate of discontinuations because of adverse events were similar to placebo in all active treatment groups.</p>
<p>Geiger et al.<sup>142</sup> (2009)</p> <p>Aliskiren 150 to 300 mg QD, added to existing HCTZ</p>	<p>AC, DB, RCT</p> <p>Patients <math>\geq</math>18 years of age with mild to moderate essential HTN who were</p>	<p>N=641</p> <p>8 weeks</p>	<p>Primary: Change in DBP at week 8</p> <p>Secondary: Change SBP at</p>	<p>Primary: After eight weeks of therapy, the triple therapy showed significantly greater reductions in SBP and DBP compared with the other groups. The additional SBP and DBP reductions were 7 and 5 mm Hg, respectively compared to aliskiren and HCTZ (<math>P&lt;0.0001</math>), 3 and 2 mm Hg compared to valsartan and HCTZ (<math>P&lt;0.01</math>), and 15 and 10 mm Hg compared to HCTZ monotherapy</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>therapy</p> <p>vs</p> <p>valsartan 160 to 320 mg QD, added to existing HCTZ therapy</p> <p>vs</p> <p>aliskiren 150 to 300 mg and valsartan 160 to 320 mg QD, added to existing HCTZ therapy</p> <p>vs</p> <p>HCTZ 25 mg QD</p>	<p>taking HCTZ for 4 weeks with a DBP <math>\geq</math>95 mm Hg</p>		<p>week 8, change in DBP and SBP at week 4, proportion of patients achieving blood pressure control (SBP/DBP &lt;140/90 mm Hg), change in plasma renin activity, plasma renin concentration</p>	<p>(<math>P&lt;0.001</math>).</p> <p>Aliskiren and HCTZ and valsartan and HCTZ combination therapies were more effective compared to HCTZ monotherapy. Valsartan and HCTZ were more effective than aliskiren and HCTZ. SBP and DBP were reduced by 15 and 11 mm Hg, respectively in the aliskiren and HCTZ group. SBP and DBP were reduced by 18 and 14 mm Hg, respectively, in the valsartan and HCTZ group.</p> <p>Secondary: Blood pressure control rate was significantly higher with triple therapy compared to aliskiren and HCTZ (40.9%, <math>P&lt;0.001</math>), valsartan and HCTZ (48.7%, <math>P&lt;0.001</math>), and HCTZ monotherapy (20.5%, <math>P&lt;0.001</math>).</p> <p>At week four, a significantly greater blood pressure control rate was observed for the triple therapy group at lower doses (150-160-25 mg) compared to the respective doses of the other groups: aliskiren and valsartan and HCTZ (300-320-25 mg) group (56%) compared to aliskiren and HCTZ (36.6%, <math>P&lt;0.05</math>), valsartan and HCTZ (42.2%, <math>P&lt;0.05</math>), and HCTZ monotherapy (19.9%, <math>P&lt;0.01</math>).</p> <p>At week eight, plasma renin concentration was unchanged in the HCTZ group, but was significantly increased in other groups. A significant decrease in plasma renin activity from baseline was observed in the aliskiren and HCTZ group (<math>P&lt;0.001</math>) and a significant increase was observed in the valsartan and HCTZ (<math>P&lt;0.001</math>). In the HCTZ and triple therapy groups, there was no change in plasma renin activity (both <math>P&gt;0.75</math>).</p>
<p>Maciejewski et al.<sup>143</sup> (2006)</p> <p>Valsartan 80 to 160 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10</p>	<p>DB, PRO, RCT, XO</p> <p>African-Americans, older than 35 years, with baseline blood pressure &gt;140/90 mm Hg and not on antihypertensive treatment</p>	<p>N=20</p> <p>8 to 10 weeks for each arm with 2 week washout period before crossover</p>	<p>Primary: Comparison of 24 hour ABPM recordings</p> <p>Secondary: Magnitude of change from baseline in SBP and DBP with each</p>	<p>Primary: There was no difference between the groups based on 24 hour ABPM: SBP amlodipine <math>130\pm 8</math> vs valsartan <math>127\pm 17</math> (<math>P=0.350</math>) and DBP amlodipine <math>82\pm 5</math> vs valsartan <math>84\pm 16</math> (<math>P=0.430</math>).</p> <p>Secondary: There was no difference between groups in magnitude of change from baseline in blood pressure (amlodipine <math>-25\pm 8/-18\pm 7</math> vs valsartan <math>-25\pm 9/-16\pm 7</math>; <math>P=0.61</math>), and in percent of patients achieving goal blood pressure, 70% in the valsartan group and 75% in the amlodipine group (<math>P=0.62</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg QD  If blood pressure exceeded 140/90 while on highest treatment dose, HCTZ 12.5mg/day was added to the regimen.			treatment, percent of patients who achieved goal <140/<90 with each treatment based on clinic blood pressure measurements	
Ichihara et al. <sup>144</sup> (2006)  Valsartan 40 to 160 mg QD  vs  amlodipine 2.5 to 10 mg QD	RCT  Patients with untreated HTN (clinic SBP >140 mm Hg and/or DBP >90 mm Hg; or ABPM SBP >135 mm Hg and/or DBP >98 mm Hg)	N=100  12 months	Primary: ABPM and clinic blood pressure  Secondary: Pulse wave velocity, carotid intima-media thickness, urinary albumin excretion	Primary: Both treatments resulted in significant decreases in blood pressure, both ambulatory and clinic, over 12 months from baseline; blood pressure decreases were similar between treatment groups (between treatments: clinic SBP P=0.34; clinic DBP P=0.85; 24 hour ABPM P=0.14).  Blood pressure variability decreased significantly in the amlodipine group compared to the valsartan group, where there was no change in blood pressure variability (P<0.01).  Secondary: The decrease in pulse wave velocity was significant from baseline for both groups, but not significantly different from each other (P<0.05 from baseline).  Intima-media thickness was not changed significantly from baseline for either treatment (P>0.05 for both from baseline).  Urinary albumin excretion in the valsartan group decreased significantly both from baseline and compared to amlodipine treatment (P<0.05 from baseline, P value for comparison not reported).
Flynn et al. <sup>145</sup> (2008)  Phase 1 Valsartan low, medium or high dose (5 mg, 20 mg and 40 mg for <18	DB, MC, RCT  Children one to five years of age with SBP ≥95 <sup>th</sup> percentile for age, gender and height with minimum weight of	N=90  4 weeks (2 weeks of phase 1 and 2 weeks of phase 2)	Primary: Change in mean seated SBP from baseline to end of phase 1 and from end of phase 1 to the end of phase 2	Primary: In phase 1, valsartan significantly lowered SBP in all of the dose groups (SBP: low dose: -8.4 mm Hg; medium dose: -8.3 mm Hg; high dose: -8.6 mm Hg). From the end of phase 1 to the end of phase 2, subjects who remained on valsartan exhibited a mean reduction in seated SBP of -1.5 mm Hg, whereas placebo recipients had a mean increase in SBP of 1.5 mm Hg. The least-squares mean difference in the change in SBP between the pooled valsartan and placebo groups (-3.9 mm Hg) was significant (P=0.02).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
kg and 10 mg, 40 mg and 80 mg for $\geq 18$ kg, respectively) QD  Phase 2 Continue phase 1 valsartan dose  vs  placebo	8 kg; could have either untreated hypertension or inadequately controlled hypertension on current treatment		Secondary: Change in mean seated DBP from baseline to the end of phase 1 and change in mean seated DBP from the end of phase 1 to the end of phase 2	Secondary: In phase 1, valsartan significantly lowered DBP in all of the dose groups (low dose: -5.5 mm Hg; medium dose: -6.4 mm Hg; high dose: -5.5 mm Hg). From the end of phase 1 to the end of phase 2, a mean decrease in seated DBP was observed in the valsartan group (-2.5 mm Hg), whereas an increase of 2.0 mm Hg was observed in the placebo group. The least-squares mean difference in the change in DBP between the 2 pooled groups (-3.7 mm Hg) was significant (P=0.009).
Schaefer et al. <sup>146</sup> (2013)  Phase 1 Valsartan low (0.25 mg/kg), medium (1 mg/kg) and high (4 mg/kg) dose QD  Phase 2 Continue phase 1 valsartan dose  vs  placebo	DB, MC, PG, RCT  Children six months to five years of age (weight 6 to 40 kg) with a documented history of hypertension (mean sitting SBP of $\geq 95$ th percentile for age, sex, and height)	N=75  8 weeks (6 weeks of phase 1 and 2 weeks of phase 2)	Primary: Mean sitting SBP reduction over first 6 weeks  Secondary: Mean sitting DBP reduction over first 6 weeks and change from end of phase 1 to end of phase 2 for mean sitting SBP and DBP	Primary: At the end of phase 1 (six weeks), statistically significant reductions (P<0.05) from baseline in mean sitting SBP were observed for all the three doses (-8.3 mmHg for low dose, -10.3 mmHg for medium dose and -14.4 mmHg for high dose) of valsartan.  Secondary: Statistically significant reductions (P <0.05) from baseline in mean sitting DBP were observed for all the three doses of valsartan after the first six weeks as well.  During phase 2, an increase in mean sitting SBP was observed in both treatment groups. Least squares mean changes in SBP did not show any significant difference between pooled valsartan and placebo groups (LSM difference, 0.6; P=0.76). Small increases in mean sitting DBP from the end of phase 1 were observed in both treatment groups; the change being greater in the placebo group, but the difference between treatments was not statistically significant (LSM difference, -0.4; P=0.85).
Jankauskiene et al. <sup>147</sup> (2021)  Valsartan 0.25 mg/kg/day	DB, MC, RCT  Children one to five years of age with hypertension (mean SBP $\geq 95$ th percentile) with or	N=120  6 weeks	Primary: Change in mean SBP from baseline to 6 weeks  Secondary: Change in mean	Primary: A clinically and statistically significant reduction in mean SBP from baseline to Week 6 was observed with the valsartan 4 mg/kg group compared with the valsartan 0.25 mg/kg group (8.5 vs 4.1 mmHg; P=0.0157). A positive dose-response relationship for mean SBP reduction was observed between the 0.25 mg/kg and 4 mg/kg groups (P=0.0012).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>valsartan 4 mg/kg/day</p>	<p>without CKD</p>		<p>SBP from baseline to 6 weeks in patients with or without CKD groupsops, safety</p>	<p>Secondary:</p> <p>In the CKD subgroup, a significant reduction in mean SBP was observed with 4 mg/kg (9.2 mmHg) versus 0.25 mg/kg (1.2 mmHg; P=0.0096). In the non-CKD subgroup, a numerically greater decrease in mean SBP was observed with 4 mg/kg (7.8 mmHg) versus 0.25 mg/kg (6.9 mmHg; P=0.6531). Incidence of adverse events was lower with valsartan 4 mg/kg than 0.25 mg/kg (41.9% vs 51.6%) and similar between CKD and non-CKD subgroups (48.4% vs 45.3%) irrespective of dose.</p> <p>Increase in serum potassium (&gt;20% compared to baseline) was observed more frequently in patients with CKD compared to non-CKD patients.</p>
<p>Philipp et al.<sup>148</sup> (2007)</p> <p><u>Study 1</u></p> <p>Valsartan 40 to 320 mg QD and amlodipine 2.5 to 5 mg QD</p> <p>vs</p> <p>amlodipine 2.5 to 5 mg QD</p> <p>vs</p> <p>valsartan 40 to 320 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Males and females, ages 18 years and older with HTN (mean sitting DBP <math>\geq</math>95 mm Hg and &lt;110 mm Hg)</p>	<p>N=1,911</p> <p>8 weeks</p>	<p>Primary:</p> <p>Mean sitting DBP</p> <p>Secondary:</p> <p>Change in mean sitting SBP, response rate (proportion of patients with mean sitting DBP &lt;90 mm Hg or a <math>\geq</math>10 mm Hg reduction from baseline), control rate (proportion of patients with mean sitting DBP &lt;90 mm Hg), adverse events (combined with study 2)</p>	<p>Primary:</p> <p>All treatments significantly decreased mean sitting DBP from baseline (P&lt;0.05).</p> <p>Combination treatment resulted in significantly greater blood pressure reduction than either monotherapy (P&lt;0.05 for all combinations compared to respective doses of monotherapy except amlodipine 2.5 mg and valsartan 40 mg QD).</p> <p>Secondary:</p> <p>All treatments significantly decreased mean sitting SBP from baseline (P&lt;0.05).</p> <p>Combination treatment resulted in significantly greater blood pressure reduction than either monotherapy (P&lt;0.05 for all combinations compared to respective doses of monotherapy).</p> <p>Response rates were significantly different from placebo for all treatment groups (P&lt;0.05).</p> <p>Response rates for combination products were significantly different than each monotherapy for the following combinations: amlodipine 5 mg plus valsartan 80 mg, amlodipine 5 mg plus valsartan 40 mg and amlodipine 2.5 mg plus valsartan 80 mg (P&lt;0.05 for each combination compared to both monotherapy).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Response rates for all combinations produced significantly improved compared to either one of the monotherapies except amlodipine 2.5 mg plus valsartan 40 mg (P&lt;0.05 for each combination compared to one of the respective monotherapy).</p> <p>Control rates with therapy were significantly better than placebo, with the highest control rate achieved with amlodipine 5 mg plus valsartan 320 mg (P&lt;0.05 compared to placebo, P value not reported for others).</p> <p>Adverse event rates were not significantly different among combination treatment, amlodipine treatment, and placebo.</p> <p>Adverse event rates were significantly different between amlodipine plus valsartan and valsartan monotherapy (P&lt;0.05).</p> <p>The most commonly reported adverse events for combination treatment were: peripheral edema, headache, nasopharyngitis, upper respiratory tract infection and dizziness. Peripheral edema occurred significantly less frequently in the combination treatment group than the amlodipine monotherapy group (5.4 vs 8.7%; P=0.014) and significantly more frequently than in the valsartan monotherapy group (5.4 vs 2.1%; P&lt;0.001). Peripheral edema occurrence in the valsartan group was similar to the rate in the placebo group.</p>
<p>Philipp et al.<sup>148</sup> (2007)</p> <p><u>Study 2</u></p> <p>Valsartan 160 or 320 mg QD and amlodipine 10 mg QD</p> <p>vs</p> <p>amlodipine 10 mg QD</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Male and females, ages 18 years and older with hypertension (mean sitting DBP ≥95 mm Hg and &lt;110 mm Hg)</p>	<p>N=1,250</p> <p>8 weeks</p>	<p>Primary: Mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, response rate (proportion of patients with mean sitting DBP &lt;90 mm Hg or a ≥10 mm Hg reduction from baseline), control rate (proportion of</p>	<p>Primary: Mean sitting DBP was significantly reduced for both combination as compared to the individual components and to placebo (P&lt;0.05).</p> <p>Secondary: Response rates and control rates for combination treatments were significantly greater than valsartan monotherapy therapy and placebo therapy, but not different from amlodipine monotherapy (P&lt;0.05).</p> <p>Adverse event rates were not significantly different between combination treatment, amlodipine treatment and placebo.</p> <p>Adverse event rates were significantly different between amlodipine plus valsartan and valsartan monotherapy (P&lt;0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
valsartan 160 to 320 mg QD  vs  placebo			patients with mean sitting DBP <90 mm Hg), adverse events (combined with study 1)	
Sinkiewicz et al. <sup>149</sup> (2009)  Amlodipine and valsartan 10-160 mg or 5-160 mg QD (fixed-dose combination product)  vs  valsartan 160 mg QD	AC, DB, MC, RCT  Patients ≥18 years of age with essential HTN (mean sitting DBP ≥90 mm Hg and <110 mm Hg) who were inadequately controlled on valsartan 160 mg	N=947  8 weeks	Primary: Change from baseline in mean DBP  Secondary: Change from baseline in mean sitting SBP, responder rate (mean DBP <90 mm Hg or ≥10 mm Hg reduction from baseline), and DBP control rate (mean DBP < 90 mm Hg)	Primary: At week eight, a significantly greater reduction in mean DBP was observed with both amlodipine and valsartan combinations (10-160 mg: -11.5 mm Hg, 5-160 mg: -9.6 mm Hg; P<0.0001 for both) compared to valsartan monotherapy (-6.7 mm Hg).  Secondary: At week eight, a significantly greater reduction in mean SBP was observed in both amlodipine and valsartan combinations (10-160 mg: -14.3 mm Hg, 5-160 mg: -12.2 mm Hg; P<0.0001 for both) compared to valsartan monotherapy (-8.3 mm Hg).  Overall mean SBP/DBP reductions of 22.5/15.5 and 21.3/13.7 mm Hg were observed in the amlodipine and valsartan 10-160 and 5-160 mg treatment groups, respectively compared to 16.7/11.4 mm Hg in the valsartan 160 mg group. The amlodipine and valsartan 10-160 mg combination showed a significantly greater reduction in mean SBP/DBP compared to amlodipine and valsartan 5-160 mg (P<0.001).  Responder rates were higher in both amlodipine and valsartan groups (10-160 mg: 81% [P<0.0001]; 5-160 mg: 68% [P=0.0018], respectively) compared to valsartan monotherapy (57%).  Peripheral edema was the most frequent adverse event, which was reported in 9.1% of patients receiving amlodipine and valsartan (10-160 mg), 0.9% of patients receiving amlodipine and valsartan (5-160 mg), and 1.3% of patients receiving valsartan monotherapy.
Philipp et al (abstract). <sup>150</sup> (2011)	Post-hoc analysis  Patients with HTN	N=834  Not reported	Primary: Rate of blood pressure control	Primary: Two weeks after starting therapy, blood pressure control rates were greater with amlodipine and valsartan 10-320 mg/day (49%) vs monotherapies (32 to

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Amlodipine and valsartan 10-160 or 10-320 mg/day (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 10 mg/day</p> <p>vs</p> <p>valsartan 160 or 320 mg/day</p> <p>vs</p> <p>placebo</p>			<p>(&lt;140/90 mm Hg), change in baseline blood pressure</p> <p>Secondary: Safety</p>	<p>38%) and placebo (16%). Consistent results were observed in patients with stage 1 and 2 HTN. Among patients receiving combination therapy, statistically significant differences were observed at endpoint vs comparators. At all baseline blood pressure levels, the probability of achieving a blood pressure &lt;140/90 or &lt;130/80 mm Hg was greater with combination therapy compared to monotherapies and placebo.</p> <p>Secondary: Overall adverse events incidence was similar with combination therapy vs monotherapies and placebo.</p>
<p>Fogari et al.<sup>151</sup> (2009)</p> <p>Valsartan and amlodipine 160-5 to 10 mg/day (fixed-dose combination)</p> <p>vs</p> <p>irbesartan and HCTZ 300-12.5 to 25 mg/day (fixed-dose combination product)</p>	<p>Blind end endpoint, OL, PG, PRO, RCT</p> <p>Patients 75 to 89 years of age with moderate essential HTN (SBP <math>\geq</math>160, DBP &gt;95 to &lt;110 mm Hg)</p>	<p>N=94</p> <p>24 weeks</p>	<p>Primary: Proportion of patients achieving DBP &lt;90 mm Hg</p> <p>Secondary: Changes in ambulatory blood pressure, lying and standing changes in blood pressure, safety</p>	<p>Primary: The proportion of patients receiving valsartan and amlodipine and irbesartan and HCTZ who achieved blood pressure &lt;140/&lt;90 mm Hg was 82.9 and 85.1% (P value not significant between groups).</p> <p>Secondary: Both treatment combinations resulted in a significant decrease in ambulatory blood pressure without any differences between treatment groups (P&lt;0.001 from baseline, P&gt;0.05 between groups).</p> <p>Results were similar between groups for lying SBP/DBP but patients receiving irbesartan and HCTZ experienced greater changes in ambulatory blood pressure than those receiving valsartan and amlodipine (17.2/9.0 vs 10.1/1.9 mm Hg; P&lt;0.05 for SBP and P&lt;0.01 for DBP).</p> <p>Changes from baseline in serum potassium (decrease) and uric acid (increase)</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				were significant for those receiving irbesartan and HCTZ, but not valsartan and amlodipine (P<0.05 for irbesartan and HCTZ).
<p>Poldermans et al.<sup>152</sup> (2007)</p> <p>Valsartan 160 mg QD and amlodipine 5 to 10 mg QD</p> <p>vs</p> <p>lisinopril 10 to 20 mg and HCTZ 12.5 mg QD</p>	<p>AC, DB, MC, PG, RCT</p> <p>Males and females, ages 18 years and older with HTN (mean DBP <math>\geq</math>110 mm Hg and &lt;120 mm Hg)</p>	<p>N=130</p> <p>6 weeks</p>	<p>Primary: Safety/adverse events, vital signs, hematology, biochemistry variables</p> <p>Secondary: Efficacy (mean DBP, response rate, proportion of patients with mean DBP &lt;90 mm Hg or a <math>\geq</math>10 mm Hg reduction from baseline)</p>	<p>Primary: Both treatments were well tolerated, 26 (40.6%) of patients receiving amlodipine and valsartan and 21 (31.8%) of patients receiving lisinopril and HCTZ reported an adverse events and most were not considered drug related.</p> <p>Peripheral edema was reported more often in the amlodipine and valsartan group than the lisinopril and HCTZ group (7.7 vs 1.5%) and cough was reported less often in the amlodipine and valsartan group than the receiving lisinopril and hydrochlorothiazide group (1.6 vs 3.0%).</p> <p>No difference was found between the treatments in changes in laboratory values or biochemistry variables.</p> <p>Secondary: Both treatments led to a reduction in mean SBP and DBP (P&lt;0.0001 for both from baseline) but were not significantly different from each other. Mean blood pressure for each group at study end: amlodipine and valsartan 135.0/83.6 mm Hg and lisinopril and HCTZ 138.7/85.2 mm Hg.</p> <p>The response rate was similar among the groups (100 vs 95.5%; P value not significant).</p>
<p>White et al.<sup>153</sup> (2008)</p> <p>Val-DICTATE</p> <p>Valsartan and HCTZ 160-12.5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>HCTZ 25 mg QD</p>	<p>AC, MC, PG, RCT</p> <p>Patients with stage 1 to 2 HTN whose BP remained uncontrolled on HCTZ 12.5 mg</p>	<p>4 weeks</p> <p>Duration not reported</p>	<p>Primary: Percentage of patients whose clinic blood pressure values were &lt;140/90 mm Hg and blood pressure values</p> <p>Secondary: Not reported</p>	<p>Primary: A significantly higher proportion of hypertensive patients met blood pressure control levels in the valsartan and HCTZ group (37%) compared to the HCTZ group (16%; P&lt;0.001).</p> <p>Changes in SBP and DBP were significantly greater with valsartan and HCTZ (-12.4/-7.5 mm Hg) compared to HCTZ (-5.6/-2.1 mm Hg; P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Waeber et al.<sup>154</sup></p>	<p>OL, RCT</p>	<p>N=327</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2001)</p> <p>Valsartan 80 mg QD, which was switched to valsartan 80 mg and HCTZ 12.5 mg QD or valsartan 80 mg and benazepril 10 mg QD</p>	<p>Patients with mild-to-moderate uncontrolled HTN (DBP <math>\geq</math>90) while on valsartan monotherapy</p>	<p>4 weeks</p>	<p>Efficacy and safety</p> <p>Secondary: Not reported</p>	<p>The two combinations produced an additional blood pressure reduction compared to monotherapy (P&lt;0.001 for both), with similar DBP reductions reported for the two combination groups (-4.5 mm Hg with valsartan plus HCTZ and -3.3 mm Hg with valsartan plus benazepril).</p> <p>SBP reductions of -6.7 and -3.2 mm Hg with valsartan plus HCTZ and valsartan plus benazepril, respectively, were reported (P=0.1).</p> <p>At the end of the trial, the blood pressure of the responders to valsartan monotherapy was lower than that of patients requiring combination therapy.</p> <p>Valsartan given alone or in association with HCTZ or benazepril was well tolerated.</p> <p>Secondary: Not reported</p>
<p>Schweizer et al.<sup>155</sup> (2007)</p> <p>Valsartan and HCTZ 160-25 mg QD (fixed-dose combination)</p>	<p>OL</p> <p>Hypertensive patients not adequately controlled by free combination of candesartan and HCTZ for 4 weeks</p>	<p>N=197</p> <p>8 weeks</p>	<p>Primary: Reduction in mean sitting DBP between week 4 and 8</p> <p>Secondary: Reduction in mean sitting SBP from week 4 to 8</p>	<p>Primary: At baseline, DBP was 103.0 mm Hg. After four weeks of candesartan and HCTZ, DBP decreased to 93.8 mm Hg. Subsequent treatment with valsartan and HCTZ for four additional weeks reduced DBP to 88.7 mm Hg. This represented an additional decrease in DBP of 5.1 mm Hg (P&lt;0.0001).</p> <p>Secondary: The valsartan and HCTZ fixed-dose combination reduced SBP by 3.4 mm Hg (P=0.0029).</p>
<p>Lai et al.<sup>156</sup> (2011)</p> <p>Valsartan and HCTZ 80-12.5 mg QD (fixed-dose combination product)</p>	<p>MC, OS</p> <p>Asian patients with stage 1 or 2 essential HTN</p>	<p>N=7,567</p> <p>24 week (follow-up)</p>	<p>Primary: Safety, efficacy</p> <p>Secondary: Not reported</p>	<p>Primary: After 24 weeks, basal blood pressure was 155.9<math>\pm</math>13.3/96.3<math>\pm</math>10.1 mm Hg. SBP and DBP reductions were -25.4<math>\pm</math>15.2 and -14.9<math>\pm</math>13.5 mm Hg (P&lt;0.001).</p> <p>Response and control rates increased continuously from baseline to trial end (trial end: 94.3 and 73.6%, respectively).</p> <p>Based on a four point global assessment scale, 96.8% of patients and physicians reported good, very good, or excellent for subjective efficacy and tolerability assessments.</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
<p>Izzo Jr et al.<sup>157</sup> (2011) ValVET</p> <p>Valsartan and HCTZ 160-12.5 mg QD (fixed-does combination product)</p> <p>vs</p> <p>valsartan 160 mg QD</p> <p>vs</p> <p>HCTZ 12.5 mg QD</p> <p>All patients were allowed to up titrate study medication if blood pressure did not improve.</p>	<p>DB, RCT</p> <p>Patients <math>\geq</math>70 years of age with systolic HTN</p>	<p>N=384</p> <p>16 weeks</p>	<p>Primary: Change in baseline SBP at week 4</p> <p>Secondary: Time to blood pressure control</p>	<p>Primary: At week four, reductions in baseline SBP were significantly greater with combination therapy (-17.3 mm Hg) compared to valsartan (-8.6 mm Hg; P&lt;0.001). At this time, reductions with combination therapy and HCTZ were similar (-17.3 vs -13.6 mm Hg; P=0.096).</p> <p>Secondary: Median time to blood pressure control was significantly shorter with combination therapy compared to HCTZ (four vs eight weeks; P&lt;0.05) and valsartan (four vs 12 weeks; P&lt;0.0001).</p>
<p>Duprez et al.<sup>158</sup> (abstract) (2011) ValVET</p> <p>Valsartan and HCTZ 160-12.5 mg QD (fixed-does combination product)</p>	<p>Subgroup analysis</p> <p>Patients <math>\geq</math>70 years of age with systolic HTN</p>	<p>N=108</p> <p>Duration not specified</p>	<p>Primary: Change in ambulatory SBP</p> <p>Secondary: Safety</p>	<p>Primary: Initiation of treatment with combination valsartan and HCTZ reduced ambulatory blood pressure more effectively compared to monotherapy with either valsartan or HCTZ throughout daytime, night-time, and 24 hr monitoring periods, as well as during the last four to six hour dosing periods.</p> <p>Twenty-four hour ambulatory blood pressure was reduced from 141.1/76.5 to 125.8/69.2 mm Hg by week four with combination valsartan and HCTZ compared to reductions from 142.2/78.7 to 139.1/77.5 mm Hg with HCTZ and 142.2/78.3 to 136.4/75.1 mm Hg with valsartan (P&lt;0.01 for all).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>valsartan 160 mg QD</p> <p>vs</p> <p>HCTZ 12.5 mg QD</p> <p>All patients were allowed to up titrate study medication if blood pressure did not improve.</p>				<p>Secondary:</p> <p>In the overall study, tolerability was similar among the three treatment groups.</p>
<p>Fogari et al.<sup>159</sup> (2006)</p> <p>Valsartan 160 mg vs olmesartan 20 mg</p> <p>All patients were also receiving HCTZ 12.5 mg QD.</p>	<p>PG, PRO, RCT</p> <p>Hypertensive patients aged 35 to 75 years with DBP 90 to 110 mm Hg after 4 weeks of monotherapy on either valsartan or olmesartan</p>	<p>N=130</p> <p>8 weeks (4 weeks of combination therapy)</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>Both combinations induced a greater ambulatory blood pressure reduction than monotherapy. However, mean reduction from baseline in the valsartan and HCTZ-treated patients (-21.5/-14.6 mm Hg for 24 hours, -21.8/-14.9 mm Hg for daytime, and -20.4/-13.7 mm Hg for nighttime SBP/DBP) was greater than in the olmesartan and HCTZ-treated patients (-18.8/-12.3 mm Hg for 24 hours, -19.3/-12.8 mm Hg for daytime, and -17.4/-10.6 mm Hg for nighttime SBP/DBP). The difference between the effects of the two treatments was significant (P&lt;0.01).</p> <p>Plasma concentrations of HCTZ were significantly greater with valsartan than with olmesartan at each determination time (P&lt;0.05).</p> <p>Secondary: Not reported</p>
<p>White et al.<sup>160</sup> (2008)</p> <p>Valsartan 160 mg and HCTZ 25 mg</p>	<p>DB, PC, RCT</p> <p>Hypertensive patients</p>	<p>N=1,181</p> <p>8 weeks</p>	<p>Primary: Changes in DBP and SBP at 8 weeks</p> <p>Secondary:</p>	<p>Primary:</p> <p>Changes from baseline in blood pressure following telmisartan and HCTZ (-24.6/-18.2 mm Hg) were significantly greater than both valsartan and HCTZ (-22.5/-17.0 mm Hg; P=0.017 for SBP and P=0.025 for DBP), and placebo (-4.1/-6.1 mm Hg; P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD</p> <p>vs</p> <p>telmisartan 80 mg and HCTZ 25 mg QD</p> <p>vs</p> <p>placebo</p>			Safety	<p>Secondary:</p> <p>The total number of patients with at least one adverse event reported was similar among the 3 treatment groups and was 37% for valsartan and HCTZ, 36% for telmisartan and HCTZ, and 42% for placebo.</p>
<p>Sharma et al.<sup>161</sup> (2007) SMOOTH</p> <p>Valsartan 160 mg for 4 weeks</p> <p>vs</p> <p>telmisartan 80 mg for 4 weeks</p> <p>After 4 weeks, all patients received add-on HCTZ 12.5 mg QD for 6 six weeks.</p>	<p>MC, OL, PRO, RCT, blinded-end point</p> <p>Men and women aged <math>\geq 30</math> years with mild-to-moderate HTN (mean seated SBP 140 to 179 mm Hg and/or DBP 95 to 109 mm Hg), with type 2 diabetes and BMI <math>&gt;27</math> kg/m<sup>2</sup></p>	<p>N=840</p> <p>10 weeks</p>	<p>Primary: Change in mean ambulatory SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: At 10 weeks, telmisartan and HCTZ provided significantly greater reductions in the last six hours of mean ambulatory blood pressure (differences in SBP were 3.9 mm Hg; <math>P&lt;0.0001</math> and differences in DBP were 2.0 mm Hg; <math>P=0.0007</math>).</p> <p>Telmisartan and HCTZ also produced significantly greater reductions than valsartan and HCTZ in 24-hour mean ambulatory blood pressure (differences in SBP were 3.0 mm Hg; <math>P=0.0002</math> and differences in DBP were 1.6 mm Hg; <math>P=0.0006</math>) and during morning, daytime and nighttime periods (<math>P&lt;0.003</math>).</p> <p>Both treatments were well tolerated.</p> <p>Secondary: Not reported</p>
<p>Calhoun et al.<sup>162</sup> (2009)</p> <p>Valsartan and HCTZ 320-25 mg QD (fixed-dose combination product)</p>	<p>DB, MC, RCT</p> <p>Patients 18 to 85 years of age with moderate to severe essential HTN</p>	<p>N=2,271</p> <p>8 weeks</p>	<p>Primary: Difference in mean sitting diastolic blood pressure and mean sitting systolic blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: At each assessment after week three, a significantly greater proportion of patients receiving triple therapy achieved overall blood pressure control (<math>&lt;140/90</math> mm Hg) compared to those receiving any of the dual therapies (all <math>P&lt;0.0001</math>).</p> <p>At end point, 70.8% of patients in the triple-therapy group achieved control, compared to 48.3% for valsartan and HCTZ, 54.1% for amlodipine and valsartan, and 44.8% for amlodipine and HCTZ (all <math>P&lt;0.0001</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>amlodipine and valsartan 10-320 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and HCTZ 10-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and valsartan and HCTZ 10-320-25 mg QD (fixed-dose combination product)</p>				<p>Triple therapy with amlodipine and valsartan and HCTZ improved blood pressure control significantly better than any of the dual therapies.</p> <p>Secondary: Not reported</p>
<p>Calhoun et al.<sup>163</sup> (2009)</p> <p>Valsartan and HCTZ 320-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and valsartan 10-320</p>	<p>Secondary analysis</p> <p>Patients 18 to 85 years of age with moderate to severe HTN (mean SBP/DBP <math>\geq 145/\geq 100</math> mm Hg)</p>	<p>N=2,271</p> <p>8 weeks</p>	<p>Primary: Proportion and mean SBP of patients with mean SBP reductions <math>\geq 60</math>, <math>\geq 50</math>, <math>\geq 40</math>, <math>\geq 30</math> and <math>\geq 20</math> mm Hg at week three and at the end of the study</p> <p>Secondary: Changes from baseline in mean</p>	<p>Primary: The proportion of patients with mean SBP reductions <math>\geq 20</math> mm Hg was greater with triple therapy than dual therapy at week three (74.5 vs 58.8 to 65.5%) and at study endpoint (87.6 vs 75.8 to 81.5%).</p> <p>More patients who received triple therapy, as compared to dual therapy, achieved mean SBP reductions of <math>\geq 30</math>, <math>\geq 40</math>, <math>\geq 50</math> and <math>\geq 60</math> mm Hg at week three and at study endpoint (P value not reported).</p> <p>In patients with severe SBP (<math>\geq 180</math> mm Hg), triple therapy resulted in significantly greater reductions than those for each dual therapy at week three (P&lt;0.01), except for amlodipine/valsartan (P=0.11).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and HCTZ 10-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine and valsartan and HCTZ 10-320-25 mg QD (fixed-dose combination product)</p>			<p>SBP based upon baseline severity, SBP control rates, safety</p>	<p>Secondary:</p> <p>Patients with higher baseline mean SBP had greater reductions in mean SBP than those with lower baseline mean SBP. Changes in mean SBP were significantly greater for triple therapy than dual therapy for all baseline SBP (P&lt;0.05), except for valsartan and HCTZ and amlodipine and HCTZ in patients with baseline mean SBP 150 to &lt;160 mm Hg (P value not reported).</p> <p>Significantly more patients (91.8%) receiving triple therapy achieved SBP control (<math>\geq 20</math> mm Hg reduction or mean SBP &lt;140 mm Hg) compared to those receiving amlodipine and HCTZ (80.1%), valsartan and HCTZ (80.8%) or valsartan and amlodipine (85.7%) (P&lt;0.01 for all).</p> <p>The overall incidence of adverse events was comparable across treatments, regardless of baseline blood pressure severity.</p>
<p>Karotsis et al.<sup>164</sup> (2006)</p> <p>Valsartan 80 mg QD</p> <p>vs</p> <p>lisinopril 10 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 mg QD</p> <p>vs</p>	<p>RCT</p> <p>Patients 25 to 79 years of age with uncontrolled HTN (average office blood pressure &gt;140/90 mm Hg for all or &gt;153/85 mm Hg for diabetics or patients &lt;65 years of age, confirmed on 2 office visits <math>\geq 1</math> week apart) after <math>\geq 4</math> weeks of OL monotherapy with diltiazem at 240 mg</p>	<p>N=211</p> <p>8 weeks</p>	<p>Primary: Blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>There was a significant decline in both office and home SBP and DBP during the trial with all treatments. The antihypertensive effect was more pronounced and reached significance when home blood pressure monitoring was used in comparison to office blood pressure without the white-coat effect (P&lt;0.001 for all blood pressure changes). With or without the white-coat effect, blood pressure still declined and the differences were significant (P&lt;0.0001 for all blood pressure changes).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
felodipine 5 mg QD  All patients also received diltiazem 240 mg QD.	QD			
Conlin et al. <sup>165</sup> (2000) PREVAIL  Candesartan 8 to 16 mg QD, irbesartan 150 to 300 mg QD, losartan 50 to 100 mg QD, and valsartan 80 to 160 mg QD  vs  another ARB  vs  ARB plus low-dose HCTZ	MA  Patients with HTN	N=11,281 (43 trials)  Duration varied	Primary: Weighted average for SBP and DBP reduction with ARB monotherapy, dose titration, and with the addition of low-dose HCTZ were calculated; responder rates  Secondary: Not reported	Primary: The absolute weighted-average reductions in DBP (8.2 to 8.9 mm Hg) and SBP (10.4 to 11.8 mm Hg) for ARB monotherapy were comparable for all ARBs. Responder rates for ARB monotherapy were 48 to 55%.  Dose titration resulted in slightly greater blood pressure reductions and an increase in responder rates of 53 to 63%.  ARB and HCTZ combinations produced substantially greater reductions in SBP (16.1 to 20.6 mm Hg) and DBP (9.9 to 13.6 mm Hg) than ARB monotherapy. Responder rates for ARB and HCTZ combinations were 56 to 70%.  The authors concluded that candesartan, irbesartan, losartan, and valsartan produced comparable antihypertensive efficacy when administered at their recommended doses, a near flat dose response when titrating from starting to maximum recommended dose, and substantial potentiation of the antihypertensive effect with addition of HCTZ.  Secondary: Not reported
Stanton et al. <sup>166</sup> (2010)  Aliskiren 300 mg QD  vs  irbesartan,	MA  Adults with mild to moderate essential HTN	N=4,877 (8 trials)  4 to 12 weeks	Primary: Paradoxical blood pressure rises, as well as the percentage of patients with SBP increases (>10 or >20 mm Hg) or DBP increases (>5	Primary: There were no significant differences among the pooled aliskiren, irbesartan, losartan, valsartan, ramipril, and HCTZ groups in the incidence of SBP increases >10 mm Hg (P=0.30) and >20 mm Hg (P=0.28) or DBP increases >5 mm Hg (P=0.65) and >10 mm Hg (P=0.5).  Increases in SBP and DBP occurred significantly more frequently in the pooled placebo group than the aliskiren group (P<0.001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
losartan, valsartan, ramipril, HCTZ, placebo			or >10 mm Hg) from baseline  Secondary: Not reported	Secondary: Not reported
Lindholm et al. <sup>167</sup> (2005)  Other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)  or  placebo  vs  β-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol, or propranolol)	MA  13 RCTs evaluating the treatment of primary HTN with a β-blocker as first-line treatment (in ≥50% of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both	N=105,951  2.1 to 10.0 years	Primary: Stroke, MI, all-cause mortality  Secondary: Not reported	Primary: The RR of stroke was 16% higher with β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non β-blockers (RR, 1.26%; 95% CI, 1.15 to 1.38; P<0.0001).  The relative risk of MI was 2% higher for β-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).  The RR of all-cause mortality was 3% higher for β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).  Secondary: Not reported
Van Bortel et al. <sup>168</sup>	MA	N=2,653	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2008)</p> <p>ACE inhibitor, ARB, <math>\beta</math>-blocker, calcium channel blocker, or placebo</p> <p>vs</p> <p>nebivolol</p>	<p>12 RCTs involving &gt;25 patients with essential HTN where nebivolol 5 mg QD was compared to placebo or other active drugs for &gt;1 month</p>	<p>Duration varied</p>	<p>Antihypertensive effect and tolerability</p> <p>Secondary: Not reported</p>	<p>Overall, higher response rates were observed with nebivolol than all other antihypertensive agents combined (OR, 1.41; 95% CI, 1.15 to 1.73; P=0.001) and compared to the ACE inhibitors (OR, 1.92; 1.30 to 2.85; P=0.001), but response rates to nebivolol were similar to <math>\beta</math>-blockers (OR, 1.29; 95% CI, 0.81 to 2.04; P=0.283), calcium channel blockers (OR, 1.19; 95% CI, 0.83 to 1.70; P=0.350) and losartan (OR, 1.35; 95% CI, 0.84 to 2.15; P=0.212).</p> <p>Overall, a higher percentage of patients obtained normalized blood pressure with nebivolol compared to the other antihypertensive agents combined (OR, 1.35; 95% CI, 1.07 to 1.72; P=0.012). A higher percentage of patient receiving nebivolol obtained normalized blood pressure compared to losartan (OR, 1.98; 95% CI, 1.24 to 3.15; P=0.004) and calcium channel blockers (OR, 1.96; 95% CI, 1.05 to 1.96; P=0.024), but not when compared to other <math>\beta</math>-blockers (OR, 1.29; 95% CI, 0.81 to 1.65; P=0.473).</p> <p>Overall, the percentage of adverse events was significantly lower with nebivolol compared to the other antihypertensive agents combined (OR, 0.59; 95% CI, 0.48 to 0.72; P&lt;0.001) and similar to placebo (OR, 1.16; 95% CI, 0.76 to 1.67; P=0.482). In comparing nebivolol to the individual treatments, nebivolol had a lower percentage of adverse events compared to losartan (OR, 0.52; 95% CI, 0.30 to 0.89; P=0.016), the other <math>\beta</math>-blockers (OR, 0.56; 95% CI, 0.36 to 0.85; P=0.007) and calcium channel blockers (OR, 0.49; 95% CI 0.33 to 0.72; P&lt;0.001), but was similar to ACE inhibitors (OR, 0.75; 95% CI 0.52 to 1.08).</p> <p>Secondary: Not reported</p>
<p>Wiysonge et al.<sup>169</sup> (2007)</p> <p>Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system</p>	<p>MA</p> <p>13 RCTs evaluating patients <math>\geq</math>18 years of age with HTN</p>	<p>N=91,561</p> <p>Duration varied</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions</p>	<p>Primary: There was not a significant difference observed in all-cause mortality between <math>\beta</math>-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with <math>\beta</math>-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>inhibitors)</p> <p>vs</p> <p>β-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)</p>				<p>There was a significant decrease in stroke observed with β-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between β-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07], diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with β-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of β-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with β-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>
<p>Baguet et al.<sup>170</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ,</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the β-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)</p> <p>Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>The weighted average reduction of SBP and DBP for each drug class were as follows:            Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively.            β-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively.            Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively.            ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.            ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively.            Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary: Not reported</p>
<b>Miscellaneous</b>				
<p>Papademetriou et al.<sup>171</sup> (2004) SCOPE</p> <p>Candesartan 16 mg/day vs</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients 7 to 89 years old with isolated systolic HTN (SBP &gt;160 mm Hg and DBP &lt;90 mm Hg) and MMSE scores ≥24</p>	<p>N=1,518 3.7 years</p>	<p>Primary: First major coronary event including cardiovascular death, nonfatal MI, or nonfatal stroke</p> <p>Secondary: cardiovascular death, nonfatal and</p>	<p>Primary: There was no difference in the first major cardiovascular event between patients (with isolated systolic hypertension) who were treated with candesartan vs placebo (RR, 0.89; 95% CI, 0.65 to 1.21; P&gt;0.20).</p> <p>Secondary: A total of 20 fatal/nonfatal strokes occurred in the candesartan group and 35 in the control group (RR, 0.58; 95% CI, 0.33 to 1.0) for a RR reduction of 42% (P=0.050 unadjusted and P=0.049 adjusted for baseline risk).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo in addition to conventional therapy (diuretics, ACE inhibitors, $\beta$ -blockers, calcium channel blockers)			fatal stroke and MI	There were no marked or statistically significant differences between the treatment groups in other cardiovascular end points or all-cause mortality.
Ogihara et al. <sup>172</sup> (2008) CASE-J Candesartan 4 to 12 mg QD vs amlodipine 2.5 to 10 mg QD	AC, MC, OL, RCT  Patients with high risk HTN (SBP $\geq 140$ mm Hg or DBP $\geq 90$ mm Hg in patients $< 70$ years old or SBP $\geq 160$ mm Hg or DBP $\geq 90$ mm Hg in patients $\geq 70$ years old), with either type 2 diabetes, history of stroke or ischemic attack, left ventricular hypertrophy, proteinuria or serum creatinine $\geq 1.3$ mg/dL	N=4,703  Up to 4 years	Primary: First fatal or nonfatal cardiovascular event  Secondary: All-cause death, new-onset diabetes, discontinuation due to adverse events	Primary: A total of 134 patients experienced a cardiovascular event in each treatment regimen (HR, 1.0; 95% CI, 0.78 to 1.27; P=0.969).  Secondary: All-cause death rates did not differ between treatments, 73 deaths in the candesartan group and 86 in the amlodipine group.  New-onset diabetes occurred in significantly fewer patients in the candesartan group than the amlodipine group (HR, 0.64; 95% CI, 0.43 to 0.97; P=0.033).  A total of 125 (5.4%) patients in the candesartan group and 134 (5.8%) of patients in the amlodipine group discontinued due to adverse events.
Taniguchi et al. <sup>173</sup> (2006)  Candesartan 8 mg in addition to spironolactone 25 mg QD for 6 months, after 6 months of candesartan monotherapy	DB, RCT, XO  Patients, 67 years of age on average, with essential HTN and left ventricular hypertrophy	N=97  1 year	Primary: Change in blood pressure and relative wall thickness  Secondary: Not reported	Primary: Both study groups experienced a statistically significant reduction in blood pressure from baseline (P<0.05).  While candesartan was associated with a significant reduction in relative wall thickness among patients with concentric left ventricular remodeling or hypertrophy (P<0.05), the addition of spironolactone did not provide additional benefit.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(combination group)  vs  candesartan 8 mg daily for 12 months				
Montalescot et al. <sup>174</sup> (2009) ARCHIPELAGO  Enalapril 10 mg QD, followed by 20 mg QD on day 15  vs  irbesartan 150 mg QD, followed by 300 mg QD on day 15	AC, DB, MC, RCT  Adults with non-ST elevation ACS	N=429  60 days	Primary: Change from baseline in high-sensitivity C-reactive protein at day 60  Secondary: Changes in other inflammatory markers such as troponin I	Primary: High-sensitivity C-reactive protein levels were comparable in both treatment groups (irbesartan: 15.2 mg/L at baseline, 6.5 mg/L at day 60; absolute change of -8.7 mg/L; enalapril: 12.6 mg/L at baseline, 5.5 mg/L at day 60; absolute change of -7.1 mg/L, P value not significant).  Secondary: Similarly, mean levels of markers of myocardial injury (troponin I) and endothelial dysfunction (microalbuminuria) also decreased from baseline to day 60, with no significant differences between treatment groups.
Solomon et al. <sup>175</sup> (2009) ALLAY  Losartan 100 mg QD  vs  aliskiren 300 mg QD  vs	AC, RCT  Adults with HTN and increased left ventricular wall thickness	N=465  9 months	Primary: Change in left ventricular mass  Secondary: Not reported	Primary: There were reductions in left ventricular mass from baseline in all treatment groups, with 4.9-g/m <sup>2</sup> (5.4%), 4.8-g/m <sup>2</sup> (4.7%), and 5.8-g/m <sup>2</sup> (6.4%) reductions in the aliskiren, losartan, and combination arms, respectively (P<0.0001 for all treatment groups).  The reduction in left ventricular mass in the combination group was not significantly different from that with losartan alone (P=0.52).  The difference in left ventricular mass regression between the aliskiren and losartan arms was within the prespecified non-inferiority margin, suggesting that aliskiren was as effective as losartan in reducing left ventricular hypertrophy (P<0.0001 for non-inferiority).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
aliskiren 300 mg and losartan 100 mg QD				Secondary: Not reported
Fliser et al. <sup>176</sup> (2004) EUTOPIA  Olmesartan 20 mg/day  vs  placebo  All patients received pravastatin 20 mg/day after six weeks of therapy.	DB, PC, PG, RCT  Patients ≥18 years old with HTN, atherosclerotic disease, type 2 diabetes mellitus, and/or LDL-C between 3.89 to 6.48 mmol/L	N=199  12 weeks	Primary: Evaluate anti-inflammatory effects of olmesartan using a panel of inflammation markers: high-sensitivity C-reactive protein, high-sensitivity tumor necrosis factor- $\alpha$ , interleukin-6  Secondary: Not reported	Primary: After six weeks of therapy, olmesartan treatment significantly reduced serum levels of C-reactive protein (-15.1%; P<0.05), tumor necrosis factor- $\alpha$ (-8.9%; P<0.02), interleukin-6 (-14.0%; P<0.05) and monocyte chemotactic protein-1 (-6.5%; P<0.01), whereas placebo treatment had no major effect on inflammation markers.  After 12 weeks of therapy, C-reactive protein (-21.1%; P<0.02), tumor necrosis factor- $\alpha$ (-13.6%; P<0.01), and interleukin-6 (-8.0%; P<0.01) decreased further with olmesartan and pravastatin cotherapy, but treatment with pravastatin alone did not significantly alter inflammation markers.  In contrast, addition of pravastatin led to a significant (P<0.001) reduction in LDL-C in the olmesartan and placebo groups (-15.1 and -12.1%, respectively).  Secondary: Not reported
Rosendorff et al. <sup>177</sup> (2009)  Olmesartan 20 to 40 mg QD  vs  amlodipine 5 to 10 mg QD	DB, AC, RCT  Adults with HTN and left ventricular hypertrophy	N=102  52 weeks	Primary: Change in left ventricular mass from baseline to 52 weeks  Secondary: Change in left ventricular mass after 26 weeks of treatment	Primary: Mean±SD left ventricular masses of 252.9±73.06 g in the olmesartan group and 236.9±59.94 g in the amlodipine group at baseline were decreased to 248.2±69.31 and 223.9±53.18 g, respectively, after 52 weeks of therapy. Neither of these changes was significantly different from baseline, and the difference between the two treatment groups was not significant.  Secondary: At 26 weeks, adjusted percent changes in left ventricular mass were 8.0% with olmesartan and 6.0% with amlodipine. Changes occurring at the 26-week assessment were not significantly different from baseline or from each other.
ONTARGET Investigators <sup>178</sup> (2008)  Ramipril 10 mg/day	DB, MC, PC, RCT  Patients with coronary, peripheral, or cerebrovascular disease or diabetes	N=25,620  56 months (median follow-up)	Primary: Death from cardiovascular causes, MI, stroke or hospitalization for heart failure	Primary: The primary outcome occurred in 16.5, 16.7, and 16.3% of patients receiving ramipril, telmisartan and combination therapy, respectively.  Secondary: The composite of death from cardiovascular causes, MI or stroke occurred in

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  telmisartan 80 mg/day  vs  ramipril 10 mg/day and telmisartan 80 mg/day	with end-organ damage		Secondary: Composite of death from cardiovascular causes, MI or stroke; heart failure, worsening or new angina, new diagnosis diabetes mellitus, new atrial fibrillation, renal impairment, revascularization procedures	<p>14.1% of patients in the ramipril group and 13.9% of patients in the telmisartan group (RR, 0.99; 95% CI, 0.91 to 1.17; P=0.001 for non-inferiority). Combination therapy was not significantly better than ramipril alone (RR, 0.99; 95% CI, 0.92 to 1.07).</p> <p>There were no significant differences in the rates of secondary outcomes, except for renal dysfunction, which occurred in 10.2% of patients receiving ramipril, 10.6% of patients receiving telmisartan and 13.5% of patients receiving combination therapy (P&lt;0.001 vs ramipril; P value not reported vs telmisartan).</p> <p>As compared to the ramipril group, the telmisartan group had lower rates of cough (1.1 vs 4.2%; P&lt;0.001) and angioedema (0.1 vs 0.3%; P=0.01) and a higher rate of hypotensive symptoms (2.6 vs 1.7%; P&lt;0.001); the rate of syncope was the same in the two groups (0.2%).</p> <p>As compared to the ramipril group, combination therapy had an increased risk of hypotensive symptoms (4.8 vs 1.7%; P&lt;0.001), syncope (0.3 vs 0.2%; P=0.03) and renal dysfunction (13.5 vs 10.2%; P&lt;0.001).</p>
Mann et al. <sup>179</sup> (2013) ONTARGET  Ramipril with telmisartan	Subanalysis  Patients in the ONTARGET trial with diabetes mellitus	N=3163 with CKD N=6465 no CKD  56 months	Primary: Composite of death from cardiovascular cause, nonfatal MI, nonfatal stroke or hospitalization for CHF  Secondary: composite renal outcome for this analysis was defined posthoc as chronic dialysis (>2 months) or a doubling of baseline serum creatinine	Primary: The stroke rate in all participants with diabetes was not different between the treatment groups, 1.19 and 1.22 per 100 patient-years in those on dual and monotherapy, respectively (HR, 0.99; 95% CI, 0.82 to 1.20). The results were consistent in those with or without renal disease (P value for interaction =0.60; 1.59 vs 1.55 and 1.01 vs 1.08 strokes per 100 patient-years, respectively). Results for other major outcomes indicated no differences and no interaction of renal subgroups with treatment effects.  Secondary: Dialysis-dependent acute kidney injury tended to occur more frequently in those allocated to dual than with monotherapy, 0.14 vs 0.08 cases per 100 patient-years, (HR, 1.55; 95% CI, 0.84 to 2.85), and hyperkalemia was more frequent, 1.82 vs 1.07 cases per 100 patient-years (HR, 1.71; 95% CI, 1.44 to 2.02). Both adverse outcomes were more frequent in those with renal disease; however, the excess due to dual therapy was similar in those with and without renal disease.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Julius et al.<sup>180</sup> (2004) VALUE</p> <p>Valsartan 80 to 160 mg QD</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD</p>	<p>DB, PG, RCT</p> <p>Patients ≥50 years old with treated or untreated HTN and history of cardiovascular disease, stroke, or diabetes, previous medications were discontinued at trial onset</p>	<p>N=15,245</p> <p>4.2 years (mean)</p>	<p>Primary: Time to first cardiac event (cardiac morbidity and mortality)</p> <p>Secondary: Fatal and nonfatal MI, fatal and nonfatal heart failure and fatal and nonfatal stroke, all-cause mortality, new onset diabetes</p>	<p>Primary: There were no differences in the primary composite end point between the valsartan and amlodipine groups (10.6 vs 10.4%; P=0.49).</p> <p>Secondary: There was a higher incidence of myocardial infarction (4.8 vs 4.1%; P=0.02) in patients receiving valsartan than amlodipine.</p> <p>There was no difference in the incidence of heart failure (4.6 vs 5.3%; P=0.12), stroke (4.2 vs 3.7%; P=0.08), and all-cause mortality (11 vs 10.8%; P=0.45) between valsartan- and amlodipine-treated patients.</p> <p>New onset diabetes occurred less with valsartan (13.1%) vs amlodipine (16.4%; P&lt;0.001).</p> <p>Combined target blood pressure (&lt;140/90 mm Hg) was achieved in 58% and 62% of patients receiving valsartan and amlodipine, respectively.</p>
<p>Zanchetti et al.<sup>181</sup> (2006) VALUE</p> <p>Amlodipine 5 mg QD</p> <p>vs</p> <p>valsartan 80 mg QD</p>	<p>Subgroup analysis of VALUE</p> <p>Patients with HTN</p>	<p>N=15,245</p> <p>4.2 years</p>	<p>Primary: Time to first cardiac event, analyzed by subgroup</p> <p>Secondary: MI, heart failure and stroke</p>	<p>Primary: The only significant result of the analyses by subgroup for time to first cardiac event was sex; women in the valsartan group experienced more cardiac events as compared to men in the valsartan group (HR for women, 1.21; 95% CI, 1.3 to 1.42; HR for men, 0.94; 95% CI, 0.82 to 1.17; P=0.016).</p> <p>The VALUE trial showed no difference in the primary outcome as well as in cardiac morbidity and mortality between amlodipine treatment and valsartan treatment. SBP and DBP were lower, as was incidence of MI, in the amlodipine treatment group as compared to the valsartan group.</p> <p>Secondary: Male patients treated with valsartan had a significantly lower incidence of heart failure than males treated with amlodipine (P&lt;0.001 for male vs female difference; for men, HF rates with valsartan were 4.1% vs amlodipine 5.8% [HR, 0.73; 95% CI, 0.60 to 0.88]; for women, rates were valsartan 5.3% vs amlodipine 4.6%, [HR, 1.18; 95% CI, 0.95 to 1.47]).</p> <p>Patients without a history of stroke had a greater reduction in stroke risk if treated with amlodipine (valsartan 3.4% vs amlodipine 2.6%; HR, 1.34; 95%</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				CI, 19 to 1.65).
<p>Sawada et al.<sup>182</sup> (2009) KYOTO HEART</p> <p>Valsartan up to 160 mg QD plus an additional antihypertensive agent (other than an ACE inhibitor) if necessary to reach target blood pressure &lt;140/90 or &lt;130/80 mm Hg</p> <p>vs</p> <p>antihypertensive agents (other than ACE inhibitors and ARBs) to reach target blood pressure &lt;140/90 or &lt;130/80 mm Hg</p>	<p>MC, OL, BE, RCT</p> <p>Japanese adults with uncontrolled HTN and coronary artery disease, cerebral vascular disease, or peripheral vascular disease.</p>	<p>N=3,031</p> <p>Median 3.27 years</p>	<p>Primary: New onset cardiovascular or cerebrovascular events (stroke, TIA, acute MI, unstable angina, aortic aneurysm, emergency thrombosis, lower limb arterial obstruction, transition to dialysis)</p> <p>Secondary: Not reported</p>	<p>Primary: In both groups, blood pressure was identical at baseline and at the end of study (157/88 and 133/76, respectively).</p> <p>The primary endpoint was recorded in fewer patients given valsartan add-on (5.5%) than in those given additional non-ARB treatment (10.2%; HR, 0.55; 95% CI, 0.42-0.72; P=0.00001).</p> <p>The difference in the number of primary endpoints was mainly attributable to reduced frequency of stroke and TIA, and unstable angina. These benefits cannot be explained by a difference in blood pressure control.</p> <p>Secondary: Not reported</p>
<p>The GISSI-AF Investigators<sup>183</sup> (2009) GISSI-AF</p> <p>Valsartan up to 320 mg QD</p> <p>vs</p> <p>placebo</p>	<p>MC, DB, PC, RCT</p> <p>Adults in sinus rhythm who had a recent history of documented atrial fibrillation</p>	<p>N=1,442</p> <p>1 year</p>	<p>Primary: Time to a first occurrence of atrial fibrillation and proportion of patients who had more than one recurrence of atrial fibrillation over the course of 1 year</p> <p>Secondary:</p>	<p>Primary: Atrial fibrillation recurred in 371 of the 722 patients (51.4%) in the valsartan group, as compared to 375 of 720 (52.1%) in the placebo group (adjusted HR, 0.97; 96% CI, 0.83 to 1.14; P=0.73).</p> <p>More than one episode of atrial fibrillation occurred in 194 of 722 patients (26.9%) in the valsartan group and in 201 of 720 (27.9%) in the placebo group (adjusted OR, 0.89; 99% CI, 0.64 to 1.23; P=0.34).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Not reported	
The Navigator Study Group <sup>184</sup> (2010) NAVIGATOR  Valsartan up to 160 mg QD or matching placebo  and  nateglinide or matching placebo	DB, MC, RCT  Adults with impaired glucose tolerance and established cardiovascular disease or cardiovascular risk factors.	N=9,306  5 years	Primary: Incidence of diabetes and a composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for heart failure, arterial revascularization, or hospitalization for unstable angina  Secondary: Not reported	Primary: The cumulative incidence of diabetes was 33.1% in the valsartan group, as compared to 36.8% in the placebo group (HR in the valsartan group, 0.86; 95% CI, 0.80 to 0.92; P<0.001).  Valsartan, as compared to placebo, did not significantly reduce the incidence of the composite cardiovascular outcome (14.5% vs 14.8%; HR, 0.96; 95% CI, 0.86 to 1.07; P=0.43).  Secondary: Not reported
Blood Pressure Lowering Treatment Trialists' Collaboration <sup>185</sup> (2007)  ACE inhibitors (17 trials)  vs  ARBs (9 trials)	MA  Patients with high blood pressure, diabetes, history or CHD or cerebrovascular disease	N=146,838 (26 trials)  Variable duration	Primary: Nonfatal MI or death from CHD, including sudden death; heart failure causing death or requiring hospitalization; nonfatal stroke or death from cerebrovascular disease  Secondary: Not reported	Primary: From a total of 146,838 individuals with high blood pressure or an elevated risk of cardiovascular disease, major cardiovascular events were documented in 22,666 patients during follow-up. The analyses showed comparable blood pressure-dependent reductions in risk with ACE inhibitors and ARBs (P≥0.3 for all three outcomes).  ACE inhibitors produced a blood pressure-independent reduction in the relative risk of CHD of approximately 9% (95% CI, 3 to 14%). No similar effect was detected for ARBs, and there was some evidence of a difference between ACE inhibitors and ARBs in this regard (P=0.002).  For both stroke and heart failure, there was no evidence of any blood pressure-independent effects of either ACE inhibitors or ARBs.  Secondary: Not reported

\*Agent not available in the United States.

Study regimen abbreviations: QD=once daily, SR=sustained-release, TID=three times daily

Study design abbreviations: AC=active comparator, DB=double blind, DD=double dummy, ES=extension study, MA=meta-analysis, MC=multicenter, OL=open-label, OS=observational, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, XO=cross-over



Miscellaneous abbreviations: ABPM=ambulatory blood pressure monitoring, ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor blocker, ACS=acute coronary syndrome, BMI=body mass index, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, ESRD=end stage renal disease, eGFR=estimated glomerular filtration rate, GFR=glomerular filtration rate, HbA<sub>1c</sub>=glycosylated hemoglobin, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HR=hazard ratio, HTN=hypertension, LDL-C=low-density lipoprotein cholesterol, LSM=least squares mean, LVEF=left ventricular ejection fraction, MAP=mean arterial pressure, MI=myocardial infarction, MMSE=Mini Mental State Examination, NYHA=New York Heart Association, OR=odds ratio, RR=relative risk, QOL=quality of life, Sc=serum creatinine, SD=standard deviation, SBP=systolic blood pressure, TC=total cholesterol, TG=triglycerides, TIA=transient ischemic attack, UAER=urinary albumin excretion rate, WHO=World Health Organization, WMD=weighted mean difference

**Additional Evidence**

Dose Simplification

Sung et al. evaluated adherence rates with a triple component single pill regimen of olmesartan 20 mg/amlodipine 5 mg/hydrochlorothiazide 12.5 mg compared to the equivalent two-pill regimen of olmesartan 20 mg/hydrochlorothiazide 12.5 mg plus amlodipine 5 mg for 12 weeks. The single-pill group had significantly higher percentage of doses taken than the two-pill group with median (25 to 75 percentile) of 95.1 (95% confidence interval [CI], 86.7 to 100.0) versus 92.1 (95% CI, 73.0 to 97.3) as well as a significantly higher prescribed dose taken correctly with median (25 to 75 percentile) of 91.0 (95% CI, 79.4 to 96.5) versus 88.6 (95% CI, 69.2 to 96.3), P = 0.04 for both comparisons.<sup>186</sup>

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 12. Relative Cost of the Angiotensin II Receptor Antagonists**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Azilsartan	tablet	Edarbi®	\$\$\$\$\$	N/A
Candesartan	tablet	Atacand®*	\$\$\$\$\$	\$\$
Irbesartan	tablet	Avapro®*	\$\$\$\$\$	\$
Losartan	tablet	Cozaar®*	\$\$\$\$	\$
Olmesartan	tablet	Benicar®*	\$\$\$\$\$	\$\$
Telmisartan	tablet	Micardis®*	\$\$\$\$	\$
Valsartan	tablet	Diovan®*	\$\$\$\$\$	\$
<b>Combination Products</b>				
Azilsartan and chlorthalidone	Tablet	Edarbyclor®	\$\$\$\$	N/A
Candesartan and HCTZ	tablet	Atacand HCT®*	\$\$\$\$\$	\$\$\$\$
Irbesartan and HCTZ	tablet	Avalide®*	\$\$\$\$\$	\$

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Losartan and HCTZ	tablet	Hyzaar <sup>®*</sup>	\$\$\$\$	\$
Olmesartan and amlodipine and hydrochlorothiazide	tablet	Tribenzor <sup>®*</sup>	\$\$\$\$	\$\$\$\$
Olmesartan and HCTZ	tablet	Benicar HCT <sup>®*</sup>	\$\$\$\$	\$
Telmisartan and amlodipine	tablet	Twynsta <sup>®*</sup>	\$\$\$\$	\$\$\$\$
Telmisartan and HCTZ	tablet	Micardis HCT <sup>®*</sup>	\$\$\$\$	\$\$\$
Valsartan and HCTZ	tablet	Diovan HCT <sup>®*</sup>	\$\$\$\$	\$

\*Generic is available in at least one dosage form or strength.  
HCTZ=hydrochlorothiazide, N/A=not available

## X. Conclusions

All of the angiotensin II receptor blockers (ARBs) are approved for the treatment of hypertension. Some of the products are also approved for the treatment of diabetic nephropathy (irbesartan and losartan), heart failure (candesartan and valsartan), post-myocardial infarction (valsartan), as well as cardiovascular and cerebrovascular risk reduction (telmisartan and losartan, respectively).<sup>3-20</sup> The ARBs are available as single entity products, and most are also available in combination with hydrochlorothiazide. Azilsartan is available in combination with chlorthalidone, telmisartan is available in combination with amlodipine, and olmesartan is available in combination with amlodipine and hydrochlorothiazide (triple therapy). There are other ARBs that are available in combination with amlodipine (olmesartan and valsartan); however, these products are included in the dihydropyridines class review (AHFS Class 242808). All single entity products with the exception of azilsartan are available generically. Fixed-dose combination products are available in a generic formulation with the exception of azilsartan-chlorthalidone.

National and international guidelines recommend the use of ACE inhibitors or ARBs in patients with cerebrovascular disease, coronary artery disease, heart failure, hypertension, left ventricular dysfunction, left ventricular hypertrophy, diabetes, diabetic nephropathy, previous myocardial infarction, and renal disease.<sup>22-40</sup> In general, guidelines do not give preference to one ARB over another.<sup>22-40</sup> Some of the guidelines specifically recommend the use of ACE inhibitors as initial therapy, with the subsequent use of ARBs in patients who do not tolerate ACE inhibitors.<sup>22,24-31,36</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>32-37</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>32</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>32-39</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>32-36</sup>

Numerous clinical trials have shown that the ARBs can effectively lower systolic and diastolic blood pressure, administered alone or in combination with other antihypertensive agents. Some comparative trials have demonstrated slight differences in blood pressure effects among the various ARBs; however, the clinical significance of these differences remains to be established.<sup>75-170</sup> Guidelines do not give preference to one ARB over another for the treatment of hypertension. Most patients will require more than one antihypertensive agent to achieve blood pressure goals.<sup>32-36</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>34-35,38,186</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations.

ARBs have been shown to reduce cardiovascular morbidity and mortality, as well as preserve renal function.<sup>41-64</sup> The use of losartan also decreases the risk of stroke in patients with hypertension and left ventricular hypertrophy.<sup>6,13</sup> It should be noted that the ACE inhibitors have also been shown to positively impact these endpoints as well (please refer to ACE inhibitor class review for additional information). Several studies comparing ARBs and ACE inhibitors have demonstrated similar efficacy with regards to cardiovascular events, heart failure and the rate of progression of nephropathy.<sup>41,42,44,46,51-55,58,64,66,71-74,178</sup> ACE inhibitors inhibit the

breakdown of bradykinin, which may lead to the development of a persistent non-productive cough. The ARBs do not increase bradykinin and may be better tolerated in some patients.<sup>20,21</sup>

The FDA has evaluated data from two clinical trials (ROADMAP and ORIENT) in which patients with type 2 diabetes who were taking olmesartan had a higher rate of death from cardiovascular causes compared to those who were taking placebo. After the review was completed in April 2011, the FDA has determined that the benefits of olmesartan continue to outweigh its potential risks when used for the treatment of patients with high blood pressure according to the approved drug label. Of note, olmesartan is not recommended as a treatment to delay or prevent protein in the urine in diabetic patients.<sup>187</sup> In June of 2011, the FDA also concluded that a review of a meta-analysis of 31 randomized-controlled trials comparing ARBs to other treatments found no evidence of an increased risk of incident (new) cancer, cancer-related death, breast cancer, lung cancer, or prostate cancer in patients receiving ARBs.<sup>188</sup>

At this time, there is insufficient evidence to conclude that the angiotensin II receptor antagonists offer a significant clinical advantage over other alternatives in general use. Therefore, all brand angiotensin II receptor antagonists within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand angiotensin II receptor antagonist is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
2. Givertz MM, Colucci WS. Pharmacologic therapy of heart failure with reduced ejection fraction: Mechanisms of action. In: Post TW (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2022 [cited 2022 Jan]. Available from: <http://www.uptodate.com/utd/index.do>.
3. Edarbi® [package insert]. Atlanta (GA): Arbor Pharmaceuticals, LLC; 2020 Mar.
4. Atacand® [package insert]. Wilmington (DE): AstraZeneca LP; 2020 Jun.
5. Avapro® [package insert]. Bridgewater (NJ): Sanofi-Aventis US LLC; 2021 Sep.
6. Cozaar® [package insert]. Whitehouse Station (NJ): Merck & Co, Inc; 2019 Nov.
7. Benicar® [package insert]. Parsippany (NJ): Daiichi Sankyo, Inc; 2019 Oct.
8. Micardis® [package insert]. Ridgefield (CT): Boehringer Ingelheim Pharmaceuticals, Inc; 2020 Jul.
9. Diovan® [package insert]. East Hanover (NJ): Novartis Pharmaceutical Corp.; 2021 Jan.
10. Edarbyclor® [package insert]. Atlanta (GA): Arbor Pharmaceuticals, LLC; 2020 Mar.
11. Atacand HCT® [package insert]. Wilmington (DE): AstraZeneca LP; 2020 May.
12. Avalide® [package insert]. Bridgewater (NJ): Sanofi-Aventis US LLC; 2021 Sep.
13. Hyzaar® [package insert]. Whitehouse Station (NJ): Merck & Co, Inc; 2020 Aug.
14. Tribenzor® [package insert]. Parsippany (NJ): Daiichi Sankyo, Inc.; 2020 Oct.
15. Benicar HCT® [package insert]. Parsippany (NJ): Daiichi Sankyo, Inc.; 2020 May.
16. Twynsta® [package insert]. Ridgefield (CT): Boehringer Ingelheim Pharmaceuticals, Inc; 2018 Feb.
17. Micardis HCT® [package insert]. Ridgefield (CT): Boehringer Ingelheim Pharmaceuticals; 2020 Aug.
18. Diovan HCT® [package insert]. East Hanover (NJ): Novartis Pharmaceutical Corp.; 2020 Aug.
19. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
20. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
21. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
22. Fraker T, Fihn S, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of chronic stable angina: a report of the American College of Cardiology/American Heart Association task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *Circulation*. 2007 Dec 4;116(23):2762-72.
23. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425.
24. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline From the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
25. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
26. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42(14);7 April 2021:1289–1367. <https://doi.org/10.1093/eurheartj/ehaa575>.
27. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.

28. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
29. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].
30. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
31. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
32. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
33. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, Ramirez A, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
34. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, Butalia S, Leung AA, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
35. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
36. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
37. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
38. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
39. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
40. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl.1).
41. Pfeffer MA, McMurray JJV, Velazquez EJ, et al; for the Valsartan in Acute Myocardial Infarction Trial Investigators. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med*. 2003 Nov 13;349:1893-906.
42. Dickstein K, Kjekshus J, and the OPTIMAAL Steering Committee, for the OPTIMAAL Study Group. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomized trial. *Lancet*. 2002 Sep 7;360:752-60.
43. Graham DJ, Zhou EH, McKean S, et al. Cardiovascular and mortality risk in elderly Medicare beneficiaries treated with olmesartan versus other angiotensin receptor blockers. *Pharmacoepidemiol Drug Saf*. 2014 Apr;23(4):331-9.
44. White WB, Cuadra RH, Lloyd E, Bakris GL, Kupfer S. Effects of azilsartan medoxomil compared with olmesartan and valsartan on ambulatory and clinic blood pressure in patients with type 2 diabetes and prediabetes. *J Hypertens*. 2016 Apr;34(4):788-97.

45. Morgensen CE, Neldam S, Tikkanen I, et al. Randomized controlled trial of dual blockade of renin-angiotensin system in patients with hypertension, microalbuminuria, and non-insulin dependent diabetes: the candesartan and lisinopril microalbuminuria (CALM) study. *BMJ*. 2000 Dec 9;321:1440-4.
46. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, et al; Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med*. 2001 Sep 20;345(12):851-60.
47. Parving HH, Lehnert H, Bröchner-Mortensen J, Gomis R, Andersen S, Arner P; Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria Study Group. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med*. 2001 Sep 20;345(12):870-8.
48. Persson F, Rossing P, Reinhard H, et al. Renal effects of aliskiren compared with and in combination with irbesartan in patients with type 2 diabetes, hypertension, and albuminuria. *Diabetes Care* 2009;32:1873-9.
49. Chrysostomou A, Pedagogos E, MacGregor L, Becker GJ. Double-blind, placebo-controlled study on the effect of the aldosterone receptor antagonist spironolactone in patients who have persistent proteinuria and are on long-term angiotensin-converting enzyme inhibitor therapy, with or without an angiotensin II receptor blocker. *Clin J Am Soc Nephrol*. 2006 Mar;1(2):256-62.
50. Bianchi S, Bigazzi R, Campese VM. Intensive versus conventional therapy to slow the progression of idiopathic glomerular disease. *Am J Kidney Dis* 2010;55:671-81.
51. Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, et al; RENAAL Study Investigators. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med*. 2001 Sep 20;345(12):861-9.
52. Kiernan MS, Gregory D, Sarnak MJ, Rossignol P, Massaro J, Kociol R, et al. Early and late effects of high-versus low-dose angiotensin receptor blockade on renal function and outcomes in patients with chronic heart failure. *JACC Heart Fail*. 2015 Mar;3(3):214-23.
53. Hou FF, Xie D, Zhang X, et al. Renoprotection of optimal antiproteinuric doses (ROAD) study: a randomized controlled study of benazepril and losartan in chronic renal insufficiency. *J Am Soc Nephrol*. 2007;18:1889-98.
54. Fried LF, Emanuele N, Zhang JH, et al. Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med*. 2013 Nov 14;369(20):1892-903.
55. Nakao N, Yoshimura A, Morita H, et al. Combination treatment of angiotensin-II receptor blocker and angiotensin-converting enzyme inhibitor in non-diabetic renal disease (COOPERATE): a randomized controlled trial. *Lancet*. 2003 Jan 11;361:117-24.
56. Mann JF, Schmeider RE, Dyal L. Effect of telmisartan on renal outcomes: a randomized trial. *Ann Intern Med* 2009;151:1-10.
57. Foulquier S, Böhm M, Schmieder R, et al. Impact of telmisartan on cardiovascular outcome in hypertensive patients at high risk: a Telmisartan Randomised Assessment Study in ACE iNtolerant subjects with cardiovascular Disease subanalysis. *J Hypertens*. 2014 Jun;32(6):1334-41.
58. Barnett AH, Bain SC, Bouter P, et al. Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N Engl J Med*. 2004 Nov 4;351:1952-61.
59. Galle J, Schwedhelm E, Pinnetti S, Boger Rh, Wanner C; on behalf of the VIVALDI Investigators. Antiproteinuric effects of angiotensin receptor blockers: telmisartan versus valsartan in hypertensive patients with type 2 diabetes mellitus and overt nephropathy. *Nephrol Dial Transplant* 2008;23:3174-83.
60. Fogari R, Derosa G, Zoppi A, et al. Effect of telmisartan-amlodipine combination at different doses on urinary albumin excretion in hypertensive diabetic patients with microalbuminuria. *Am J Hypertens* 2007;20:417-22.
61. Viberti G, Wheeldon NM; MicroAlbuminuria Reduction With VALsartan (MARVAL) Study Investigators. Microalbuminuria reduction with valsartan in patients with type 2 diabetes mellitus: a blood pressure-independent effect. *Circulation*. 2002 Aug 6;106(6):672-8.
62. Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet*. 2005 Dec 10;366:2026-33.
63. Strippoli GFM, Craig M, Deeks JJ, et al. Effects of angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists on mortality and renal outcomes in diabetic nephropathy: systematic review. *BMJ*. 2004 Sep 30; 329:828-38.
64. Strippoli GFM, Bonifati C, Craig M, Navaneethan DS, Craig JC. Angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists for preventing the progression of diabetic kidney disease (review). *Cochrane Database Syst Rev*. 2006 Oct 18;(4):CD006257.
65. Cohn JN, Tognoni G; Valsartan Heart Failure Trial Investigators. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med*. 2001 Dec 6;345(23):1667-75.

66. Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, Michelson EL, et al; CHARM Investigators and Committees. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet*. 2003 Sep 6;362(9386):759-66.
67. McMurray JJ, Ostergren J, Swedberg K, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet*. 2003 Sep 6;362(9386):767-71.
68. Granger CB, McMurray JJ, Yusuf S, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet*. 2003 Sep 6;362(9386):772-6.
69. Yusuf S, Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *Lancet*. 2003 Sep 6;362(9386):777-81.
70. Castagno D, Skali H, Takeuchi M, Swedberg K, Yusuf S, Granger CB, et al. Association of heart rate and outcomes in a broad spectrum of patients with chronic heart failure. *J Am Coll Cardiol*. 2012;59:1785-95.
71. Pitt B, Segal R, Martinez FA, et al; on behalf of ELITE Study Investigators. Randomized trial of losartan versus captopril in patients over 65 with heart failure (Evaluation of Losartan in the Elderly Study, ELITE). *Lancet*. 1997 Mar 15;349:747-52.
72. Pitt B, Poole-Wilson PA, Segal R, et al; on behalf of the ELITE II Investigators. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomized trial—the Losartan Heart Failure Survival Study, ELITE II. *Lancet*. 2000 May 6;355:1582-7.
73. McKelvie RS, Yusuf S, Pericak D, et al. Comparison of candesartan, enalapril, and their combination in congestive heart failure: randomized evaluation of strategies for left ventricular dysfunction (RESOLVD) pilot study: the RESOLVD Pilot Study Investigators. *Circulation*. 1999 Sep 7;100:1056-64.
74. Lee VC, Rhew DC, Dylan M, Badamgarav E, Braunstein GD, Weingarten SR. Meta-analysis: angiotensin-receptor blockers in chronic heart failure and high-risk acute myocardial infarction. *Ann Intern Med*. 2004 Nov 2;141(9):693-704.
75. Rakugi H, Enya K, Sugiura K, Ikeda Y. Comparison of the efficacy and safety of azilsartan with that of candesartan cilexetil in Japanese patients with grade I-II essential hypertension: a randomized, double-blind clinical study. *Hypertension Research*. 2012;35:552-558.
76. Sica D, White WB, Weber MA, Bakris GL, Perez A, Cao C, et al. Comparison of the novel angiotensin II receptor blocker azilsartan medoxomil vs valsartan by ambulatory blood pressure monitoring. *J Clin Hypertens (Greenwich)*. 2011;13:467-472.
77. Cushman WC, Bakris GL, White WB, Weber MA, Sica D, Roberts A, et al. Azilsartan medoxomil plus chlorthalidone reduces blood pressure more effectively than olmesartan plus hydrochlorothiazide in stage 2 systolic hypertension. *Hypertension*. 2012;60:310-318.
78. Bönner G, Bakris GL, Sica D, et al. Antihypertensive efficacy of the angiotensin receptor blocker azilsartan medoxomil compared with the angiotensin-converting enzyme inhibitor ramipril. *J Hum Hypertens*. 2013 Aug;27(8):479-86.
79. Handley A, Lloyd E, Roberts A, Barger B. Safety and tolerability of azilsartan medoxomil in subjects with essential hypertension: a one-year, phase 3, open-label study. *Clin Exp Hypertens*. 2016;38(2):180-8.
80. Kipnes MS, Handley A, Lloyd E, Barger B, Roberts A. Safety, tolerability, and efficacy of azilsartan medoxomil with or without chlorthalidone during and after 8 months of treatment for hypertension. *J Clin Hypertens (Greenwich)*. 2015 Mar;17(3):183-92.
81. Neutel JM, Cushman WC, Lloyd E, Barger B, Handley A. Comparison of long-term safety of fixed-dose combinations azilsartan medoxomil/chlorthalidone vs olmesartan medoxomil/hydrochlorothiazide. *J Clin Hypertens (Greenwich)*. 2017 Sep;19(9):874-883.
82. Cushman WC, Bakris GL, White WB, Weber MA, Sica D, Roberts A, et al. A randomized titrate-to-target study comparing fixed-dose combinations of azilsartan medoxomil and chlorthalidone with olmesartan and hydrochlorothiazide in stage-2 systolic hypertension. *J Hypertens*. 2018 Apr;36(4):947-956.
83. Lithell H, Hansson L, Skoog I, Elmfeldt D, Hofman A, Olofsson B, et al; SCOPE Study Group. The Study on Cognition and Prognosis in the Elderly (SCOPE): principal results of a randomized double-blind intervention trial. *J Hypertens*. 2003 May;21(5):875-86.
84. Baguet JP, Nisse-Durgeat S, Mouret S, Asmar R, Mallion JM. A placebo-controlled comparison of the efficacy and tolerability of candesartan cilexetil, 8 mg, and losartan, 50 mg, as monotherapy in patients with essential hypertension, using 36-h ambulatory blood pressure monitoring. *Int J Clin Pract* 2006;60:391-8.



85. Ohma KP, Milon H, Valnes K. Efficacy and tolerability of a combination tablet of candesartan cilexetil and hydrochlorothiazide in insufficiently controlled primary hypertension--comparison with a combination of losartan and hydrochlorothiazide. *Blood Press.* 2000;9(4):214-20.
86. Mengden T, Hubner R, Bramlage P. Office and ambulatory blood pressure control with a fixed-dose combination of candesartan and hydrochlorothiazide in previously uncontrolled hypertensive patients: results of the CHILI CU Soon. *Vascular Health and Risk Management.* 2011;7:761-769.
87. McInnes GT, O'Kane KP, Istad H, Keinänen-Kiukaanniemi S, Van Mierlo HF. Comparison of the AT1-receptor blocker, candesartan cilexetil, and the ACE inhibitor, lisinopril, in fixed combination with low dose hydrochlorothiazide in hypertensive patients. *J Hum Hypertens.* 2000 Apr;14(4):263-9.
88. Hosaka M, Metoki H, Satoh M, Ohkubo T, Asayama K, Kikuya M, et al. Randomized trial comparing the velocities of the antihypertensive effects on home blood pressure of candesartan and candesartan with hydrochlorothiazide. *Hypertens Res.* 2015 Oct;38(10):701-7.
89. Fogari R, Mugellini A, Derosa G; CANDIA Study Group. Efficacy and tolerability of candesartan cilexetil/hydrochlorothiazide and amlodipine in patients with poorly controlled mild-to-moderate essential hypertension. *J Renin Angiotensin Aldosterone Syst.* 2007 Sep;8(3):139-44.
90. Gradman AH, Schmieder RE, Lins RL, et al. Aliskiren, a novel orally effective renin inhibitor, provides dose-dependent antihypertensive efficacy and placebo-like tolerability in hypertensive patients. *Circulation* 2005;111:1012-8.
91. Jordan J, Engeli S, Boye S, et al. Direct renin inhibition with aliskiren in obese patients with arterial hypertension. *Hypertension.* 2007 May;49:1047-55.
92. O'Brien E, Barton J, Nussberger J, et al. Aliskiren reduces blood pressure and suppresses plasma renin activity in combination with a thiazide diuretic, an angiotensin-converting enzyme inhibitor, or an angiotensin receptor blocker. *Hypertension.* 2007 Feb;49(2):276-84.
93. Derosa G, Cicero AFG, Gaddi A, et al. Effects of doxazosin and irbesartan on blood pressure and metabolic control in patients with type 2 diabetes and hypertension. *J Cardiovasc Pharmacol.* 2005;45(6):599-604.
94. Neutel JM, Franklin SS, Oparil S, Bhaumik A, Ptaszynska A, Lapuerta P. Efficacy and safety of irbesartan/HCTZ combination therapy as initial treatment for rapid control of severe hypertension. *J Clin Hypertens (Greenwich).* 2006 Dec;8(12):850-7.
95. Neutel JM. A comparison of the efficacy and safety of irbesartan/hydrochlorothiazide combination therapy with irbesartan monotherapy in the treatment of moderate or severe hypertension in diabetic and obese hypertensive patients: a post-hoc analysis review (abstract). *Postgrad Med.* 2011 Jul;123(4):126-34.
96. Neutel JM, Franklin SS, Lapuerta P, Bhaumik A, Ptaszynska A. A comparison of the efficacy and safety of irbesartan/HCTZ combination therapy with irbesartan and HCTZ monotherapy in the treatment of moderate hypertension. *J Hum Hypertens* 2008;22:266-74.
97. Weir MR, Neutel JM, Bhaumik A, De Obaldia ME, Lapuerta P. The efficacy and safety of initial use of irbesartan/hydrochlorothiazide fixed-dose combination in hypertensive patients with and without high CV risk. *J Clin Hypertens (Greenwich).* 2007 Dec;9(12 Suppl 5):23-30.
98. Bobrie G, Delonca J, Moulin C, et al; for the comparative study of efficacy of irbesartan/HCTZ with valsartan/HCTZ using home blood pressure monitoring in the treatment of mild-to-moderate hypertension (COSIMA) investigators. *Am J Hypertens.* 2005;18:1482-8.
99. Stanton A, Jensen C, Nussberger J, et al. Blood pressure lowering in essential hypertension with an oral renin inhibitor, aliskiren. *Hypertension.* 2003 Dec;42:1137-43.
100. Ribeiro AB, Mion D Jr, Marin MJ, et al; Latin American Hypertension Study (LAMHYST) Group. Antihypertensive efficacy of amlodipine and losartan after two 'missed' doses in patients with mild to moderate essential hypertension. *J Int Med Res.* 2007 Nov-Dec;35(6):762-72.
101. Oparil S, Barr E, Elkins M, Liss C, Vrecenak A, Edelman J. Efficacy, tolerability, and effects on quality of life of losartan, alone or with hydrochlorothiazide, versus amlodipine, alone or with hydrochlorothiazide, in patients with essential hypertension. *Clin Ther.* 1996 Jul-Aug;18(4):608-25.
102. Dahlöf B, Devereux RB, Kjeldsen SE, Julius S, Beevers G, de Faire U, et al; LIFE Study Group. Cardiovascular morbidity and mortality in the Losartan Intervention For End point reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet.* 2002 Mar 23;359(9311):995-1003.
103. Julius S, Alderman MH, Beevers G, et al. Cardiovascular risk reduction in hypertensive black patients with left ventricular hypertrophy: the LIFE study. *J Am Coll Cardiol* 2004;43:1047-55.
104. Lindholm LH, Ibsen H, Dahlöf B, Devereux RB, Beevers G, de Faire U, et al; LIFE Study Group. Cardiovascular morbidity and mortality in patients with diabetes in the Losartan Intervention For End point reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002;359:1004-10.

105. Kjeldsen SE, Dahlöf B, Devereux RB, Julius S, Aurup P, Edelman J, et al, Snapinn S, Wedel H; LIFE (Losartan Intervention for End point Reduction) Study Group. Effects of losartan on cardiovascular morbidity and mortality in patients with isolated systolic hypertension and left ventricular hypertrophy: a Losartan Intervention for End point Reduction (LIFE) substudy. *JAMA*. 2002 Sep 25;288(12):1491-8.
106. Fossum E, Olsen M, Hoiegggen A, Wachtell K, Reims H, Kjeldsen S et al. Long-term effects of a losartan-compared with and atenolol-based treatment regimen on carotid artery plaque development in hypertensive patients with left ventricular hypertrophy: ICARUS, a LIFE substudy. *J Clin Hypertens*. 2006;8:169-73.
107. Kizer J, Dahlof B, Kjeldsen S, Julius S, Beevers G, de Faire U et al. Stroke reduction in hypertensive adults with cardiac hypertrophy randomized to losartan versus atenolol: the losartan intervention for endpoint reduction in hypertension study. *Hypertension*. 2005;45:46-52.
108. Wachtell K, Lehto M, Gerds E, Olsen M, Hornestam B, Dahlof B et al. Angiotensin II receptor blockade reduces new-onset atrial fibrillation and subsequent stroke compared to atenolol: the losartan intervention for endpoint reduction in hypertension (LIFE) study. *J Am Coll Cardiol*. 2005;45:712-9.
109. Wachtell K, Hornestam B, Lehto M, Slotwiner D, Gerds E, Olsen M et al. Cardiovascular morbidity and mortality in hypertensive patients with a history of atrial fibrillation: the losartan intervention for endpoint reduction in hypertension (LIFE) study. *J Am Coll Cardiol*. 2005;45:705-11.
110. Van Bortel LM, Bulpitt CJ, Fici F. Quality of life and antihypertensive effect with nebivolol and losartan. *Am J Hypertens*. 2005 Aug;18(8):1060-6.
111. Flack JM, Oparil S, Pratt JH et al. Efficacy and tolerability of eplerenone and losartan in hypertensive black and white patients. *Journal of the American College of Cardiology* 2003;41(7):1148-55.
112. Hood SJ, Taylor KP, Ashby MJ, Brown MJ. The spironolactone, amiloride, losartan, and thiazide (SALT) double-blind crossover trial in patients with low-renin hypertension and elevated aldosterone-renin ratio. *Circulation*. 2007 Jul 17;116(3):268-75.
113. Maeda K, Adachi M, Kinoshita A, Koh N, Miura Y, Murohara T. Efficacy and safety of the losartan-hydrochlorothiazide combination tablet in patients with hypertension uncontrolled by angiotensin II receptor antagonist therapy: the Aichi Research on Combination Therapy for Hypertension (ARCH) study. *Intern Med*. 2012;51:1167-1175.
114. Ueda T, Kai H, Imaizumi T, MAPPY Study Investigators. Losartan/hydrochlorothiazide combination vs. high-dose losartan in patients with morning hypertension—a prospective, randomized, open-labeled, parallel-group, multicenter trial (abstract). *Hypertens Res*. 2012 Jul;35(7):708-14.
115. Salerno CM, Demopoulos L, Mukherjee R, Gradman AH. Combination angiotensin receptor blocker/hydrochlorothiazide as initial therapy in the treatment of patients with severe hypertension. *J Clin Hypertens (Greenwich)*. 2004 Nov;6(11):614-20.
116. Minami J, Abe C, Akashiba A, Takahashi T, Kameda T, Ishimitsu T, Matsuoka H. Long-term efficacy of combination therapy with losartan and low-dose hydrochlorothiazide in patients with uncontrolled hypertension. *Int Heart J*. 2007 Mar;48(2):177-86.
117. Lacourcière Y, Gil-Extremera B, Mueller O, Byrne M, Williams L. Efficacy and tolerability of fixed-dose combinations of telmisartan plus HCTZ compared with losartan plus HCTZ in patients with essential hypertension. *Int J Clin Pract*. 2003 May;57(4):273-9.
118. Brunner HR, Arakawa K. Antihypertensive efficacy of olmesartan medoxomil and candesartan cilexetil in achieving 24-hour blood pressure reductions and ambulatory blood pressure goals. *Clin Drug Investig*. 2006;26(4):185-93.
119. Punzi HA, Lewin A, Li W, Chavanu KJ. Efficacy/safety of olmesartan medoxomil versus losartan potassium in naïve versus previously treated subjects with hypertension. *Adv Ther*. 2012;29(6):524-537.
120. Brunner HR, Arakawa K. Antihypertensive efficacy of olmesartan medoxomil and candesartan cilexetil in achieving 24-hour blood pressure reductions and ambulatory blood pressure goals. *Clin Drug Investig*. 2006;26(4):185-93.
121. Chrysant SG, Weber MA, Wang AC, Hinman DJ. Evaluation of antihypertensive therapy with the combination of olmesartan medoxomil and hydrochlorothiazide. *Am J Hypertens*. 2004 Mar;17(3):252-9.
122. Kereiakes DJ, Neutel JM, Punzi HA, et al. Efficacy and safety of olmesartan medoxomil and hydrochlorothiazide compared with benazepril and amlodipine besylate. *Am J Cardiovasc Drugs*. 2007;7(5):36-72.
123. Chrysant SG, Melino M, Karki S, Lee J, Heyrman R. The combination of olmesartan medoxomil and amlodipine besylate in controlling high blood pressure: COACH, a randomized, double-blind, placebo-controlled, 8-week factorial safety and efficacy study. *Clin Ther*. 2008;30:587-604.

124. Chrysant SG, Oparil S, Melino M, et al. Efficacy and safety of long-term treatment with the combination of amlodipine besylate and olmesartan medoxomil in patients with hypertension. *J Clin Hypertens*. (Greenwich) 2009;11:475-82.
125. Oparil S, Lee J, Karki S, Melino M. Subgroup analyses of an efficacy and safety study of concomitant administration of amlodipine besylate and olmesartan medoxomil: evaluation by baseline hypertension stage and prior antihypertensive medication use. *J Cardiovasc Pharmacol*. 2009;54(5):427-36.
126. Braun N, Ulmer HJ, Handrock R, Klebs S. Efficacy and safety of the single pill combination of amlodipine 10 mg plus valsartan 160 mg in hypertensive patients not controlled by amlodipine 10 mg plus olmesartan 20 mg in free combination (abstract). *Current Medical Research & Opinion*. 2009;25(2):421-30.
127. Chrysant SG, Littlejohn T III, Izzo JL Jr, Kereiakes DJ, Oparil S, Melino M, et al. Triple-combination therapy with olmesartan, amlodipine, and hydrochlorothiazide in black and non-black study participants with hypertension. *Am J Cardiovasc Drugs*. 2012;12(4):233-243.
128. Chrysant SG, Izzo JL Jr, Kereiakes DJ, Littlejohn T 3<sup>rd</sup>, Oparil S, Melino M, et al. Efficacy and safety of triple-combination therapy with olmesartan, amlodipine, and hydrochlorothiazide in study participants with hypertension and diabetes: a subpopulation analysis of the TRINITY study (abstract). *J Am Soc Hypertens*. 2011 Mar-Apr;6(2):132-41.
129. Kereiakes DJ, Chrysant SG, Izzo JL Jr, Littlejohn T III, Oparil S, Melino M, et al. Long-term efficacy and safety of triple-combination therapy with olmesartan medoxomil and amlodipine besylate and hydrochlorothiazide for hypertension. *J Clin Hypertens* (Greenwich). 2012;14:149-157.
130. Punzi HA. Efficacy and safety of olmesartan/amlodipine/hydrochlorothiazide in patients with hypertension not at goal with mono, dual or triple drug therapy: results of the CHAMPiOn study. *Ther Adv Cardiovasc Dis*. 2014 Feb;8(1):12-21.
131. Sharma AM, Bakris G, Neutel JM, Littlejohn TW, Kobe M, Ting N, et al. Single-pill combination of telmisartan/amlodipine versus amlodipine monotherapy in diabetic hypertensive patients: an 8-week randomized, parallel-group, double-blind trial. *Clin Ther*. 2012;34(3):537-551.
132. Williams B, Lacourcière Y, Schumacher H, Gosse P, Neutel JM. Antihypertensive efficacy of telmisartan vs ramipril over the 24-h dosing period, including the critical early morning hours: a pooled analysis of the PRISMA I and II randomized trials. *J Hum Hypertens*. 2009 Sep;23(9):610-9.
133. Karlberg BE, Lins LE, Hermansson K. Efficacy and safety of telmisartan, a selective AT1 receptor antagonist, compared with enalapril in elderly patients with primary hypertension. TEES Study Group. *J Hypertens*. 1999 Feb;17(2):293-302.
134. Xi GL, Cheng JW, Lu GC. Meta-analysis of randomized controlled trials comparing telmisartan with losartan in the treatment of patients with hypertension. *Am J Hypertens*. 2008 May;21(5):546-52.
135. Sharma A, Bagchi A, Kinagi SB, et al. Results of a comparative, phase III, 12-week, multicenter, prospective, randomized, double-blind assessment of the efficacy and tolerability of a fixed-dose combination of telmisartan and amlodipine versus amlodipine monotherapy in Indian adults with stage II hypertension. *Clin Ther* 2007;29:2667-76.
136. Littlejohn TW 3<sup>rd</sup>, Majul CR, Olvera R, et al. Results of treatment with telmisartan-amlodipine in hypertensive patients. *J Clin Hypertens* (Greenwich) 2009;11:207-13.
137. Littlejohn T, Majul C, Olver R, Seeber M, Kobe M, Guthrie R et al. Telmisartan plus amlodipine in patients with moderate or severe hypertension: results from a subgroup analysis of a randomized, placebo-controlled, parallel-group, 4x4 factorial study. *Postgrad Med*. 2009;121(2):5-14.
138. Neutel JM, Mancina G, Black HR, Dahlof B, Defeo H, Ley L, et al. Single-pill combination of telmisartan/amlodipine in patients with severe hypertension: results from the TEAMSTA severe HTN study. *J Clin Hypertens* (Greenwich). 2012 Apr;14(4):206-215.
139. Oparil S, Yarows S, Patel S, et al. Efficacy and safety of combined use of aliskiren and valsartan in patients with hypertension: a randomized, double-blind trial. *Lancet*. 2007;370:221-9.
140. Yarows S, Oparil S, Patel S, Fang H, Zhang J. Aliskiren and valsartan in stage 2 hypertension; subgroup analysis of a randomized, double-blind study. *Adv Ther*. 2008;25(12):1288-302.
141. Pool JL, Schmieder RE, Azizi M, et al. Aliskiren, an orally effective renin inhibitor, provides antihypertensive efficacy alone and in combination with valsartan. *Am J Hypertens* 2007;20(1):11-20.
142. Geiger H, Barranco E, Gorostidi M, et al. Combination therapy with various combinations of aliskiren, valsartan, and hydrochlorothiazide in hypertensive patients not adequately responsive to hydrochlorothiazide alone. *J Clin Hypertens* (Greenwich). 2009;11:324-32.
143. Maciejewski S, Mohiuddin SM, Packard KA, et al. Randomized, double-blind, crossover comparison of amlodipine and valsartan in African-Americans with hypertension using 24-hour ambulatory blood pressure monitoring. *Pharmacotherapy*. 2006 Jul;26(7):889-95.

144. Ichihara A, Kaneshiro Y, Takemitsu T, Sakoda M. Effects of amlodipine and valsartan on vascular damage and ambulatory blood pressure in untreated hypertensive patients. *J Hum Hypertens*. 2006 Oct;20(10):787-94.
145. Flynn JT, Meyers KE, Neto JP, Meneses RP, Zurowska A, Bagga A et al. Efficacy and safety of the Angiotensin receptor blocker valsartan in children with hypertension aged 1 to 5 years. 2008 Aug;52(2):222-8. doi: 10.1161/HYPERTENSIONAHA.108.111054. Epub 2008 Jun 30.
146. Schaefer F, Coppo R, Bagga A, Senguttuvan P, Schlosshauer F, Zhang Y et al. Efficacy and safety of valsartan in hypertensive children 6 months to 5 years of age. *J Hypertens*. 2013 May;31(5):993-1000. doi: 10.1097/HJH.0b013e32835f5721.
147. Jankauskiene A, Drozd D, Wasilewska A, Paula-Bernardes R, Glazer R, Valentin M et al. Efficacy and safety of valsartan in children aged 1-5 years with hypertension, with or without chronic kidney disease: a randomized, double-blind study followed by open-label phase. *Curr Med Res Opin*. 2021 Dec;37(12):2113-2122. doi:10.1080/03007995.2021.1982681. Epub 2021 Oct 9.
148. Philipp T, Smith TR, Glazer R, et al. Two multicenter, 8-week, randomized, double-blind, placebo-controlled, parallel-group studies evaluating the efficacy and tolerability of amlodipine and valsartan in combination and as monotherapy in adult patients with mild to moderate essential hypertension. *Clin Ther*. 2007 Apr;29(4):563-80.
149. Sinkiewicz W, Glazer RD, Kavoliuniene A, et al. Efficacy and tolerability of amlodipine/valsartan combination therapy in hypertensive patients not adequately controlled on valsartan monotherapy. *Curr Med Res Opin* 2009;25:315-24.
150. Philipp T, Glazer RD, Wernsing M, Yen J. Initial combination therapy with amlodipine/valsartan compared with monotherapy in the treatment of hypertension (abstract). *J Amer Soc Hyperten*. 2011;5(5):417-424.
151. Fogari R, Zoppi A, Mugellini A, Corradi L, Lazzari P, Preti P et al. Efficacy and safety of two treatment combinations of hypertension in very elderly patients. *Archives of Gerontology and Geriatrics*. 2009;48:401-5.
152. Poldermans D, Glazer R, Karagiannis S, et al. Tolerability and blood pressure-lowering efficacy of the combination of amlodipine plus valsartan compared with lisinopril plus hydrochlorothiazide in adult patients with stage 2 hypertension. *Clin Ther*. 2007 Feb;29(2):279-89.
153. White WB, Calhoun DA, Samuel R, et al. Improving blood pressure control: increase the dose of diuretic or switch to a fixed-dose angiotensin receptor blocker/diuretic? The valsartan hydrochlorothiazide diuretic for initial control and titration to achieve optimal therapeutic effect (Val-DICTATE) trial. *J Clin Hypertens (Greenwich)* 2008;10:450-8.
154. Waeber B, Aschwanden R, Sadecky L, Ferber P. Combination of hydrochlorothiazide or benazepril with valsartan in hypertensive patients unresponsive to valsartan alone. *J Hypertens*. 2001 Nov;19(11):2097-104.
155. Schweizer J, Hilsmann U, Neumann G, Handrock R, Klebs S. Efficacy and safety of valsartan 160/HCTZ 25 mg in fixed combination in hypertensive patients not controlled by candesartan 32 mg plus HCTZ 25 mg in free combination. *Curr Med Res Opin*. 2007 Nov;23(11):2877-85.
156. Lai WT, Park JE, Dongre N, Wang J. Efficacy, safety, and tolerability of valsartan/hydrochlorothiazide in Asian patients with essential hypertension. *Adv Ther*. 2011;28(5):427-438.
157. Izzo JL Jr, Weintraub HS, Duprez DA, Purkayastha D, Zappe D, Smauel R, et al. Treating systolic hypertension in very elderly with valsartan-hydrochlorothiazide vs either monotherapy: ValVET primary results. 2011;13:722-730.
158. Duprenz DA, Weintraub HS, Cushman WC, Purkayastha D, Zappe D, Samuel R, et al. Effect of valsartan, hydrochlorothiazide, and their combination on 24-h ambulatory blood pressure response in elderly patients with systolic hypertension: a ValVET substudy. *Blood Press Monitor*. 2011 Aug;16(4):186-96.
159. Fogari R, Zoppi A, Mugellini A, et al. Hydrochlorothiazide added to valsartan is more effective than when added to olmesartan in reducing blood pressure in moderately hypertensive patients inadequately controlled by monotherapy. *Adv Ther*. 2006 Sep-Oct;23(5):680-95.
160. White WB, Murwin D, Chrysant SG, et al. Effects of the angiotensin II receptor blockers telmisartan versus valsartan in combination with hydrochlorothiazide: a large, confirmatory trial. *Blood Press Monit* 2008;13:21-7.
161. Sharma AM, Davidson J, Koval S, Lacourciere Y. Telmisartan/hydrochlorothiazide versus valsartan/hydrochlorothiazide in obese hypertensive patients with type 2 diabetes: the SMOOTH study. *Cardiovasc Diabetol*. 2007 Oct 2;6:28.
162. Calhoun DA, Lacourciere Y, Chiang YT, Glazer RD. Triple antihypertensive therapy with amlodipine, valsartan, and hydrochlorothiazide: A randomized controlled trial. *Hypertension* 2009;54:32-9.
163. Calhoun D, Crikelaire N, Yen J, Glazer R. Amlodipine/valsartan/hydrochlorothiazide triple combination therapy in moderate/severe hypertension: secondary analyses evaluating efficacy and safety. *Adv Ther*. 2009;26(11):1012-23.

164. Karotsis AK, Symeonidis A, Mastorantonakis SE, Stergiou GS. Additional antihypertensive effect of drugs in hypertensive subjects uncontrolled on diltiazem monotherapy: a randomized controlled trial using office and home blood pressure monitoring. *Clin Exp Hypertens*. 2006;28(7):655-62.
165. Conlin PR, Spence JD, Williams B, Ribeiro AB, Saito I, Benedict C, et al. Angiotensin II antagonists for hypertension: are there differences in efficacy? *Am J Hypertens*. 2000 Apr;13(4 Pt 1):418-26.
166. Stanton AV, Gradman AH, Schmieder RE, et al. Aliskiren monotherapy does not cause paradoxical blood pressure rises. Meta-analysis of data from 8 clinical trials. *Hypertension* 2010;55:54-60.
167. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*. 2005 Oct 29-Nov 4;366(9496):1545-53.
168. Van Bortel LM, Fici F, Mascagni F. Efficacy and tolerability of nebivolol compared with other antihypertensive drugs: a meta-analysis. *Am J Cardiovasc Drugs*. 2008;8(1):35-44.
169. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
170. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig*. 2007;27(11):735-53.
171. Papademetriou V, Farsang C, Elmfeldt D, et al. Stroke prevention with the angiotensin II type 1-receptor blocker candesartan in elderly patients with isolated systolic hypertension; the study on cognition and prognosis in the elderly (SCOPE). *J Am Coll Cardiol*. 2004;44:1175-80.
172. Ogihara T, Nakao K, Fukui T, et al; Candesartan Antihypertensive Survival Evaluation in Japan Trial Group. Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. *Hypertension*. 2008 Feb;51(2):393-8.
173. Taniguchi I, Kawai M, Date T, Yoshida S, Seki S, Taniguchi M, Shimizu M, Mochizuki S. Effects of spirinolactone during an angiotensin II receptor blocker treatment on the left ventricular mass reduction in hypertensive patients with concentric left ventricular hypertrophy. *Circ J*. 2006 Aug; 70(8):995-1000.
174. Montalescot G, Drexler H, Gallo R, et al. Effect of irbesartan and enalapril in non-ST elevation acute coronary syndrome: results of the randomized, double-blind ARCHPELAGO study. *Euro Heart J* 2009;30:2733-41.
175. Solomon SD, Appelbaum E, Manning WJ, et al. Effect of the direct renin inhibitor aliskiren, the angiotensin receptor blocker losartan, or both on left ventricular mass in patients with hypertension and left ventricular hypertrophy. *Circulation* 2009;119:530-7.
176. Fliser D, Buchholz K, Haller H; European Trial on Olmesartan and Pravastatin in Inflammation and Atherosclerosis (EUTOPIA) Investigators. Antiinflammatory effects of angiotensin II subtype 1 receptor blockade in hypertensive patients with microinflammation. *Circulation*. 2004 Aug 31;110(9):1103-7.
177. Rosendorff C, Dubiel R, Xu J, Chavanu KJ. Comparison of olmesartan medoxomil versus amlodipine besylate on regression of ventricular and vascular hypertrophy. *Am J Cardiol* 2009;104:359-65.
178. ONTARGET Investigators. Telmisartan, ramipril, or both in patients at high risk for vascular events. *N Engl J Med*. 2008 Apr 10;358(15):1547-59.
179. Mann JF, Anderson C, Gao P, et al. Dual inhibition of the renin-angiotensin system in high-risk diabetes and risk for stroke and other outcomes: results of the ONTARGET trial. *J Hypertens*. 2013 Feb;31(2):414-21.
180. Julius S, Kjeldsen SE, Weber M, Brunner HR, Ekman S, Hansson L, et al; VALUE trial group. Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomised trial. *Lancet*. 2004 Jun 19;363(9426):2022-31.
181. Zanchetti A, Julius S, Kjeldsen S, et al. Outcomes in subgroups of hypertensive patients treated with regimens based on valsartan and amlodipine: An analysis of findings from the VALUE trial. *J Hypertens*. 2006 Nov;24(11):2163-8.
182. Sawada T, Yamada H, Dahlöf B, et al. Effects of valsartan on morbidity and mortality in uncontrolled hypertensive patients with high cardiovascular risk: Kyoto Heart Study. *Euro Heart J* 2009;30:2461-9.
183. The GISSI-AF Investigators. Valsartan for prevention of recurrent atrial fibrillation. *N Engl J Med* 2009;360:1606-17.
184. The Navigator Study Group. Effect of valsartan on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010;362:1477-90.
185. Blood Pressure Lowering Treatment Trialists' Collaboration. Blood pressure-dependent and independent effects of agents that inhibit the renin-angiotensin system. *J Hypertens*. 2007 Jun;25(5):951-8.
186. Sung J, Ahn KT, Cho BR, Lee SY, Kim BJ, Kim DK et al. Adherence to triple-component antihypertensive regimens is higher with single-pill than equivalent two-pill regimens: A randomized controlled trial. *Clin Transl Sci*. 2021 May;14(3):1185-1192. doi: 10.1111/cts.12979. Epub 2021 Feb 13.

187. US FDA Medwatch [database on the internet]. Benicar (olmesartan) ongoing safety review. Rockville (MD): Food and Drug Administration (US); 2011 Apr 14. Available from:  
<http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm215249.htm>.
188. US FDA MedWatch [database on the internet]. Angiotensin Receptor Blockers (ARBs): Drug Safety Communication- Drug Safety Review Completed. Rockville (MD): Food and Drug Administration (US); 2011 Jun 2. Available from:  
<http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm219185.htm>.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Mineralocorticoid (Aldosterone) Receptor Antagonists  
AHFS Class 243220  
May 4, 2022**

**I. Overview**

Aldosterone is a component of the renin-angiotensin-aldosterone (RAAS) system, which is responsible for the regulation of extracellular volume and blood pressure. Upon binding to the mineralocorticoid receptor on the distal renal tubule, aldosterone activates the sodium-potassium exchange pump, leading to sodium and water retention, as well as potassium excretion. Increased levels of aldosterone are present in both primary and secondary hyperaldosteronism. Heart failure, hepatic cirrhosis, and the nephrotic syndrome are edematous conditions, which can lead to secondary aldosteronism. Volume depletion and sodium loss due to diuretic therapy may also cause secondary aldosteronism.<sup>1,2</sup>

The mineralocorticoid (aldosterone) receptor antagonists, eplerenone and spironolactone, are approved for the treatment of edema, heart failure, hypertension, hypokalemia, and primary hyperaldosteronism. Eplerenone and spironolactone bind to mineralocorticoid receptors, which blocks the binding of aldosterone.<sup>1-6</sup> They are available as single entity agents, and spironolactone is also available in combination with hydrochlorothiazide. Hydrochlorothiazide inhibits the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride.<sup>1,2</sup> Finerenone is a novel, non-steroidal, selective mineralocorticoid receptor antagonist that has shown to reduce albuminuria in type 2 diabetes (T2D) patients with chronic kidney disease (CKD), while revealing only a low risk of hyperkalemia. In contrast to steroidal mineralocorticoid receptor antagonists such as spironolactone, finerenone is more selective for the mineralocorticoid receptor.<sup>7-9</sup>

The mineralocorticoid (aldosterone) receptor antagonists that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the agents except finerenone are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Mineralocorticoid (Aldosterone) Receptor Antagonists Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Eplerenone	tablet	Inspra®*	eplerenone
Finerenone	tablet	Kerendia®	none
Spironolactone	suspension, tablet	Aldactone®*, Carospir®	spironolactone
<b>Combination Products</b>			
Spironolactone and hydrochlorothiazide	tablet	Aldactazide®*	spironolactone and hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the mineralocorticoid (aldosterone) receptor antagonists are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Mineralocorticoid (Aldosterone) Receptor Antagonists**

Clinical Guideline	Recommendations
American College of Cardiology/American Heart Association: <b>2007 Chronic Angina Focused Update of</b>	<ul style="list-style-type: none"> <li>Aspirin should be started at 75 to 162 mg/day and continued indefinitely in all patients, unless contraindicated.</li> <li>Use of warfarin in conjunction with aspirin and/or clopidogrel is associated with an increased risk of bleeding and should be monitored closely.</li> </ul>

Clinical Guideline	Recommendations
<p><b>the 2002 Guidelines for the Management of Patients With Chronic Stable Angina (2007)<sup>10</sup></b></p>	<ul style="list-style-type: none"> <li>• Patients with hypertension and established coronary artery disease (CAD) should be treated with blood pressure medication(s) as tolerated, including angiotensin-converting enzyme inhibitors (ACE inhibitors) and/or <math>\beta</math>-adrenergic antagonists (<math>\beta</math>-blockers) with the addition of other medications as needed to achieve blood pressure goals of &lt;140/90 or &lt;130/80 mm Hg for patients with chronic kidney disease or diabetes.</li> <li>• Long-acting calcium-channel blocking agents or long-acting nitrates may be used if <math>\beta</math>-blockers are contraindicated. Immediate-release and short-acting dihydropyridine calcium channel blockers can increase adverse cardiac events and should not be used.</li> <li>• Long-acting calcium channel blockers or long-acting nitrates may be used with <math>\beta</math>-blockers if initial treatment is not successful.</li> <li>• ACE inhibitors should be used indefinitely in patients with a left ventricular ejection fraction (LVEF) of <math>\leq 40\%</math> and in those with hypertension, diabetes or chronic kidney disease, unless contraindicated.</li> <li>• ACE inhibitors should also be used indefinitely in patients at lower risk (mildly reduced or normal LVEF in whom cardiovascular risk factors remain well controlled and revascularization has been performed), unless contraindicated.</li> <li>• Angiotensin II receptor blockers (ARBs) are recommended in patients with hypertension, those who have an indication for an ACE inhibitor and are intolerant to them, who have heart failure, or who have had a myocardial infarction (MI) and have a LVEF of <math>\leq 40\%</math>.</li> <li>• ARBs may be considered in combination with an ACE inhibitor for heart failure due to left ventricular systolic dysfunction.</li> <li>• Aldosterone blockade is recommended in patients post-MI without significant renal dysfunction or hyperkalemia who are already receiving therapeutic doses of an ACE inhibitor and a <math>\beta</math>-blocker, have a LVEF <math>\leq 40\%</math> and have either diabetes or heart failure.</li> <li>• It is beneficial to start and continue <math>\beta</math>-blocker therapy indefinitely in all patients who have had a MI, acute coronary syndrome or left ventricular dysfunction with or without heart failure symptoms, unless contraindicated.</li> <li>• Annual influenza vaccination is recommended in patients with cardiovascular disease.</li> </ul>
<p><b>European Society of Cardiology: Guidelines for the Diagnosis and Management of Chronic Coronary Syndromes (2019)<sup>11</sup></b></p>	<p><b>Pharmacological management of stable coronary artery disease (CAD) patients</b></p> <ul style="list-style-type: none"> <li>• The two aims of the pharmacological management of stable CAD patients are to obtain relief of symptoms and to prevent CV events.</li> <li>• Optimal medical treatment indicates at least one drug for angina/ischemia relief plus drugs for event prevention.</li> <li>• It is recommended to educate patients about the disease, risk factors and treatment strategy.</li> <li>• It is indicated to review the patient's response soon after starting therapy.</li> <li>• Concomitant use of a proton pump inhibitor is recommended in patients receiving aspirin monotherapy, DAPT, or DOAC monotherapy who are at high risk of gastrointestinal bleeding.</li> <li>• Lipid-lowering drugs: if goals are not met on maximum tolerated dose of a statin, consideration of combination therapy with ezetimibe or a PCSK9 inhibitor is recommended</li> <li>• ACE inhibitors should be considered in patients at a very high risk of cardiovascular adverse events</li> <li>• Angina/ischemia relief:             <ul style="list-style-type: none"> <li>○ Short-acting nitrates are recommended.</li> <li>○ First-line treatment is indicated with <math>\beta</math>-blockers and/or calcium channel blockers to control heart rate and symptoms.</li> <li>○ Long-acting nitrates should be considered as a second-line treatment option when initial therapy with a beta-blocker and/or a non-DHP-</li> </ul> </li> </ul>



Clinical Guideline	Recommendations
	<p>calcium channel blocker is contraindicated, poorly tolerated, or inadequate in controlling angina symptoms</p> <ul style="list-style-type: none"> <li>○ Nicorandil, ranolazine, ivabradine, or trimetazidine should be considered as a second-line treatment to reduce angina frequency and improve exercise tolerance in subjects who cannot tolerate, have contraindications to, or whose symptoms are not adequately controlled by beta-blockers, CCBs, and long-acting nitrates.</li> <li>○ According to comorbidities/tolerance, it is indicated to use second-line therapies as first-line treatment in selected patients.</li> <li>○ In asymptomatic patients with large areas of ischaemia (&gt;10%) <math>\beta</math>-blockers should be considered.</li> <li>○ In patients with vasospastic angina, calcium channel blockers and nitrates should be considered and beta-blockers avoided.</li> </ul> <ul style="list-style-type: none"> <li>● Event prevention:           <ul style="list-style-type: none"> <li>○ Low-dose aspirin daily is recommended in all stable CAD patients.</li> <li>○ Clopidogrel is indicated as an alternative in case of aspirin intolerance.</li> <li>○ Statins are recommended in all stable CAD patients.</li> <li>○ It is recommended to use ACE inhibitors (or ARBs) if presence of other conditions (e.g., heart failure, hypertension or diabetes).</li> </ul> </li> </ul> <p><u>Treatment in patients with microvascular angina</u></p> <ul style="list-style-type: none"> <li>● It is recommended that all patients receive secondary prevention medications including aspirin and statins.</li> <li>● <math>\beta</math>-blockers are recommended as a first-line treatment.</li> <li>● Calcium antagonists are recommended if <math>\beta</math>-blockers do not achieve sufficient symptomatic benefit or are not tolerated.</li> <li>● ACE inhibitors or nicorandil may be considered in patients with refractory symptoms.</li> <li>● Xanthine derivatives or nonpharmacological treatments such as neurostimulatory techniques may be considered in patients with symptoms refractory to the above listed drugs.</li> </ul> <p><u>Stenting and peri-procedural antiplatelet strategies in stable CAD patients</u></p> <ul style="list-style-type: none"> <li>● Drug-eluting stent (DES) is recommended in stable CAD patients undergoing stenting if there is no contraindication to prolonged dual antiplatelet therapy (DAPT).</li> <li>● Aspirin is recommended for elective stenting.</li> <li>● Clopidogrel is recommended for elective stenting.</li> <li>● Prasugrel or ticagrelor should be considered in patients with stent thrombosis on clopidogrel without treatment interruption.</li> <li>● GP IIb/IIIa antagonists should be considered for bailout situation only.</li> <li>● Platelet function testing or genetic testing may be considered in specific or high risk situations (e.g., prior history of stent thrombosis; compliance issue; suspicion of resistance; high bleeding risk) if results may change the treatment strategy.</li> <li>● Prasugrel or ticagrelor may be considered in specific high risk situations of elective stenting (e.g., left main stenting, high risk of stent thrombosis, diabetes).</li> <li>● Pretreatment with clopidogrel (when coronary anatomy is not known) is not recommended.</li> <li>● Routine platelet function testing (clopidogrel and aspirin) to adjust antiplatelet therapy before or after elective stenting is not recommended.</li> <li>● Prasugrel or ticagrelor is not recommended in low risk elective stenting.</li> <li>● After uncomplicated PCI, early cessation (<math>\leq 1</math> week) of aspirin, and continuation of dual therapy with oral anticoagulation therapy and clopidogrel should be considered if the risk of stent thrombosis is low</li> <li>● Triple therapy with aspirin, clopidogrel, and a DOAC for <math>\geq 1</math> month should be</li> </ul>

Clinical Guideline	Recommendations
	<p>considered when the risk of stent thrombosis outweighs the bleeding risk, with a total of no more than six months</p> <p><u>Follow-up of revascularized stable coronary artery disease patients</u></p> <ul style="list-style-type: none"> <li>• It is recommended that all revascularized patients receive a secondary prevention and be scheduled for follow-up visit.</li> <li>• It is recommended to instruct patients before discharge about return to work and resumption of full activities. Patients have to be advised to seek immediate medical contact if symptoms (re-) occur.</li> <li>• Single antiplatelet therapy, usually aspirin, is recommended indefinitely.</li> <li>• DAPT is indicated after bare metal stent (BMS) for at least one month.</li> <li>• DAPT is indicated for six to 12 months after 2nd generation DES.</li> <li>• DAPT may be used for more than one year in patients at high ischemic risk (e.g., stent thrombosis, recurrent acute coronary syndrome on DAPT, post MI/diffuse CAD) and low bleeding risk.</li> <li>• DAPT for one to three months may be used after DES implantation in patients at high bleeding risk or with undeferrable surgery or concomitant anticoagulant treatment.</li> </ul> <p><u>Antithrombotic therapy in patients with chronic coronary syndrome:</u></p> <ul style="list-style-type: none"> <li>• Addition of a second antithrombotic drug to aspirin for long-term secondary prevention should be considered in patients with at least a moderately increased risk of ischemic events and without high bleeding risk</li> <li>• When oral anticoagulation is initiated in patients with AF, a DOAC is recommended in preference to VKA therapy.</li> </ul>
<p>American College of Physicians/ American College of Cardiology Foundation/ American Heart Association/ American Association for Thoracic Surgery/ Preventive Cardiovascular Nurses Association/ Society of Thoracic Surgeons: <b>Management of Stable Ischemic Heart Disease (2012)</b><sup>12</sup></p>	<p><u>Medical therapy to prevent MI and death in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Aspirin 75 to 162 mg daily should be continued indefinitely in the absence of contraindications.</li> <li>• Treatment with clopidogrel is a reasonable option when aspirin is contraindicated.</li> <li>• Dipyridamole should not be used as antiplatelet therapy.</li> <li>• Beta-blocker therapy should be initiated and continued for three years in all patients with normal left ventricular (LV) function following MI or acute coronary syndromes.</li> <li>• Metoprolol succinate, carvedilol, or bisoprolol should be used for all patients with systolic LV dysfunction (ejection fraction <math>\leq 40\%</math>) with heart failure or prior MI, unless contraindicated.</li> <li>• ACE inhibitors should be prescribed in all patients with stable IHD who also have hypertension, diabetes, LV systolic dysfunction (ejection fraction <math>\leq 40\%</math>), and/or chronic kidney disease, unless contraindicated.</li> <li>• Angiotensin-receptor blockers (ARBs) are recommended for patients with stable IHD who have hypertension, diabetes, LV systolic dysfunction, or chronic kidney disease and have indications for, but are intolerant of, ACE inhibitors.</li> <li>• Patients should receive an annual influenza vaccine.</li> </ul> <p><u>Medical therapy for relief of symptoms in patients with stable IHD</u></p> <ul style="list-style-type: none"> <li>• Beta-blockers are recommended as initial therapy for relief of symptoms.</li> <li>• Calcium channel blockers or long-acting nitrates should be prescribed for relief of symptoms when <math>\beta</math>-blockers are contraindicated or cause unacceptable side effects.</li> <li>• Calcium channel blockers or long-acting nitrates, in combination with <math>\beta</math>-blockers, should be prescribed for relief of symptoms when initial treatment with <math>\beta</math>-blockers is unsuccessful.</li> <li>• Nitroglycerin or nitroglycerin spray should be used for immediate relief of angina.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>Ranolazine is a fourth-line agent reserved for patients who have contraindications to, do not respond to, or cannot tolerate <math>\beta</math>-blockers, calcium-channel blockers, or long-acting nitrates.</li> </ul>
<p>American College of Cardiology Foundation/American Heart Association: <b>2014 American Heart Association/ American College of Cardiology Foundation Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes (2014)</b><sup>13</sup></p>	<p><u>Early hospital care- standard medical therapies</u></p> <ul style="list-style-type: none"> <li>Supplemental oxygen should be administered to patients with non-ST-elevation acute coronary syndrome (NSTEMI-ACS) with arterial oxygen saturation &lt;90%, respiratory distress, or other high risk features of hypoxemia.</li> <li>Anti-ischemic and analgesic medications <ul style="list-style-type: none"> <li>Nitrates <ul style="list-style-type: none"> <li>Patients with NSTEMI-ACS with continuing ischemic pain should receive sublingual nitroglycerin (0.3 to 0.4 mg) every 5 minutes for up to three doses, after which an assessment should be made about the need for intravenous nitroglycerin.</li> <li>Intravenous nitroglycerin is indicated for patients with NSTEMI-ACS for the treatment of persistent ischemia, heart failure, or hypertension.</li> <li>Nitrates should not be administered to patients who recently received a phosphodiesterase inhibitor, especially within 24 hours of sildenafil or vardenafil, or within 48 hours of tadalafil.</li> </ul> </li> <li>Analgesic therapy <ul style="list-style-type: none"> <li>In the absence of contraindications, it may be reasonable to administer morphine sulphate intravenously to patients with NSTEMI-ACS if there is continued ischemic chest pain despite treatment with maximally tolerated anti-ischemic medications.</li> <li>Nonsteroidal anti-inflammatory drugs (NSAIDs) (except aspirin) should not be initiated and should be discontinued during hospitalization due to the increased risk of major adverse cardiac event associated with their use</li> </ul> </li> <li>Beta-adrenergic blockers <ul style="list-style-type: none"> <li>Oral <math>\beta</math>-blocker therapy should be initiated within the first 24 hours in patients who do not have any of the following: 1) signs of HF, 2) evidence of low-output state, 3) increased risk for cardiogenic shock, or 4) other contraindications to <math>\beta</math>-blockade (e.g., PR interval &gt;0.24 second, second- or third-degree heart block without a cardiac pacemaker, active asthma, or reactive airway disease)</li> <li>In patients with concomitant NSTEMI-ACS, stabilized heart failure, and reduced systolic function, it is recommended to continue <math>\beta</math>-blocker therapy with one of the three drugs proven to reduce mortality in patients with heart failure: sustained-release metoprolol succinate, carvedilol, or bisoprolol.</li> <li>Patients with documented contraindications to <math>\beta</math>-blockers in the first 24 hours should be re-evaluated to determine subsequent eligibility.</li> </ul> </li> <li>Calcium channel blockers (CCBs) <ul style="list-style-type: none"> <li>In patients with NSTEMI-ACS, continuing or frequently recurring ischemia, and a contraindication to <math>\beta</math>-blockers, a nondihydropyridine CCB (e.g., verapamil or diltiazem) should be given as initial therapy in the absence of clinically significant LV dysfunction, increased risk for cardiogenic shock, PR interval &gt;0.24 seconds, or second or third degree atrioventricular block without a cardiac pacemaker.</li> <li>Oral nondihydropyridine calcium antagonists are recommended in patients with NSTEMI-ACS who have recurrent ischemia in the absence of contraindications, after appropriate use of <math>\beta</math>-blockers and nitrates.</li> <li>CCBs are recommended for ischemic symptoms when <math>\beta</math>-blockers are not successful, are contraindicated, or cause unacceptable side effects.</li> <li>Long-acting CCBs and nitrates are recommended in patients with coronary artery spasm.</li> <li>Immediate-release nifedipine should not be administered to patients</li> </ul> </li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>with NSTEMI-ACS in the absence of <math>\beta</math>-blocker therapy.</p> <ul style="list-style-type: none"> <li>○ Other anti-ischemic interventions           <ul style="list-style-type: none"> <li>▪ Ranolazine is currently indicated for treatment of chronic angina; however, it may also improve outcomes in NSTEMI-ACS patients due to a reduction in recurrent ischemia.</li> </ul> </li> <li>○ Cholesterol management           <ul style="list-style-type: none"> <li>▪ High-intensity statin therapy should be initiated or continued in all patients with NSTEMI-ACS and no contraindications to its use. Treatment with statins reduces the rate of recurrent MI, coronary heart disease mortality, need for myocardial revascularization, and stroke.</li> <li>▪ It is reasonable to obtain a fasting lipid profile in patients with NSTEMI-ACS, preferably within 24 hours of presentation.</li> </ul> </li> <li>● Inhibitors of renin-angiotensin-aldosterone system           <ul style="list-style-type: none"> <li>○ ACE inhibitors should be started and continued indefinitely in all patients with LVEF &lt;0.40 and in those with hypertension, diabetes mellitus, or stable CKD, unless contraindicated.</li> <li>○ ARBs are recommended in patients with heart failure or myocardial infarction with LVEF &lt;0.40 who are ACE inhibitor intolerant.</li> <li>○ Aldosterone-blockade is recommended in patients post-MI without significant renal dysfunction (creatinine &gt;2.5 mg/dL in men or &gt;2.0 mg/dL in women) or hyperkalemia (K &gt;5.0 mEq/L) who are receiving therapeutic doses of ACE inhibitor and <math>\beta</math>-blocker and have a LVEF &lt;0.40, diabetes mellitus, or heart failure.</li> </ul> </li> <li>● Initial antiplatelet/anticoagulant therapy in patients with definite or likely NSTEMI-ACS treated with an initial invasive or ischemia-guided strategy           <ul style="list-style-type: none"> <li>○ Non-enteric coated, chewable aspirin (162 to 325 mg) should be given to all patients with NSTEMI-ACS without contraindications as soon as possible after presentation, and a maintenance dose of aspirin (81 to 162 mg/day) should be continued indefinitely.</li> <li>○ In patients who are unable to take aspirin because of hypersensitivity or major gastrointestinal intolerance, a loading dose of clopidogrel followed by a daily maintenance dose should be administered.</li> <li>○ A P2Y<sub>12</sub> receptor inhibitor (clopidogrel or ticagrelor) in addition to aspirin should be administered for up to 12 months to all patients with NSTEMI-ACS without contraindications who are treated with an early invasive or ischemia-guided strategy. Options include:               <ul style="list-style-type: none"> <li>▪ Clopidogrel: 300 or 600 mg loading dose, then 75 mg daily.</li> <li>▪ Ticagrelor: 180 mg loading dose, then 90 mg twice daily.</li> <li>▪ It is reasonable to use ticagrelor in preference to clopidogrel for P2Y<sub>12</sub> treatment in patients with NSTEMI-ACS who undergo an early invasive or ischemia-guided strategy.</li> <li>▪ In patients with NSTEMI-ACS treated with an early invasive strategy and dual antiplatelet therapy (DAPT) with intermediate/high-risk features (e.g., positive troponin), a GP IIb/IIIa inhibitor may be considered as part of initial antiplatelet therapy. Preferred options are eptifibatid or tirofiban.</li> </ul> </li> </ul> </li> </ul> <p><u>Percutaneous coronary intervention (PCI)- Antiplatelet and anticoagulant therapy</u></p> <ul style="list-style-type: none"> <li>● Antiplatelet agents           <ul style="list-style-type: none"> <li>○ Patients already taking daily aspirin before PCI should take 81 to 325 mg non-enteric coated aspirin before PCI</li> <li>○ Patients not on aspirin therapy should be given non-enteric coated aspirin 325 mg as soon as possible before PCI.</li> <li>○ After PCI, aspirin should be continued indefinitely.</li> <li>○ A loading dose of a P2Y<sub>12</sub> inhibitor should be given before the procedure in patients undergoing PCI with stenting. Options include clopidogrel 600 mg,</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>prasugrel 60 mg, or ticagrelor 180 mg.</p> <ul style="list-style-type: none"> <li>○ In patients with NSTEMI-ACS and high-risk features (e.g., elevated troponin) not adequately pretreated with clopidogrel or ticagrelor, it is useful to administer a GP IIb/IIIa inhibitor (abciximab, double-bolus eptifibatide, or high-dose bolus tirofiban) at the time of PCI.</li> <li>○ In patients receiving a stent (bare metal or drug eluting) during PCI, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months. Options include clopidogrel 75 mg daily, prasugrel 10 mg daily, or ticagrelor 90 mg twice daily.</li> </ul> <ul style="list-style-type: none"> <li>● Anticoagulant therapy           <ul style="list-style-type: none"> <li>○ An anticoagulant should be administered to patients with NSTEMI-ACS undergoing PCI to reduce the risk of intracoronary and catheter thrombus formation.</li> <li>○ Intravenous unfractionated heparin (UFH) is useful in patients with NSTEMI-ACS undergoing PCI.</li> <li>○ Bivalirudin is useful as an anticoagulant with or without prior treatment with UFH.</li> <li>○ An additional dose of 0.3 mg/kg intravenous enoxaparin should be administered at the time of PCI to patients with NSTEMI-ACS who have received fewer than two therapeutic subcutaneous doses or received the last subcutaneous enoxaparin dose eight to 12 hours before PCI.</li> <li>○ If PCI is performed while the patient is on fondaparinux, an additional 85 IU/kg of UFH should be given intravenously immediately before PCI because of the risk of catheter thrombosis (60 IU/kg IV if a GP IIb/IIIa inhibitor used with UFH dosing based on the target-activated clotting time).</li> <li>○ Anticoagulant therapy should be discontinued after PCI unless there is a compelling reason to continue.</li> </ul> </li> <li>● Timing of CABG in relation to use of antiplatelet agents           <ul style="list-style-type: none"> <li>○ Non-enteric coated aspirin (81 to 325 mg daily) should be administered preoperatively to patients undergoing CABG.</li> <li>○ In patients referred for elective CABG, clopidogrel and ticagrelor should be discontinued for at least five days before surgery and prasugrel for at least seven days before surgery.</li> <li>○ In patients referred for urgent CABG, clopidogrel and ticagrelor should be discontinued for at least 24 hours to reduce major bleeding.</li> <li>○ In patients referred for CABG, short-acting intravenous GP IIb/IIIa inhibitors (eptifibatide or tirofiban) should be discontinued for at least 2 to 4 hours before surgery and abciximab for at least 12 hours before to limit blood loss and transfusion.</li> </ul> </li> </ul> <p><u>Late hospital care, hospital discharge, and posthospital discharge care</u></p> <ul style="list-style-type: none"> <li>● Medications at discharge           <ul style="list-style-type: none"> <li>○ Medications required in the hospital to control ischemia should be continued after hospital discharge in patients with NSTEMI-ACS who do not undergo coronary revascularization, patients with incomplete or unsuccessful revascularization, and patients with recurrent symptoms after revascularization. Titration of the doses may be required.</li> <li>○ All patients who are post-NSTEMI-ACS should be given sublingual or spray nitroglycerin with verbal and written instructions for its use.</li> <li>○ Before hospital discharge, patients with NSTEMI-ACS should be informed about symptoms of worsening myocardial ischemia and MI and should be given verbal and written instructions about how and when to seek emergency care for such symptoms.</li> <li>○ Before hospital discharge, patients who are post-NSTEMI-ACS and/or designated responsible caregivers should be provided with easily understood and culturally sensitive verbal and written instructions about medication</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>type, purpose, dose, frequency, side effects, and duration of use.</p> <ul style="list-style-type: none"> <li>○ For patients who are post–NSTE-ACS and have initial angina lasting more than one minute, nitroglycerin (one dose sublingual or spray) is recommended if angina does not subside within three to five minutes; call 9-1-1 immediately to access emergency medical services.</li> <li>○ If the pattern or severity of angina changes, suggesting worsening myocardial ischemia (e.g., pain is more frequent or severe or is precipitated by less effort or occurs at rest), patients should contact their clinician without delay to assess the need for additional treatment or testing.</li> <li>○ Before discharge, patients should be educated about modification of cardiovascular risk factors.</li> </ul> <ul style="list-style-type: none"> <li>● Late hospital and post-hospital oral antiplatelet therapy           <ul style="list-style-type: none"> <li>○ Aspirin should be continued indefinitely. The dose should be 81 mg daily in patients treated with ticagrelor and 81 to 325 mg daily in all other patients.</li> <li>○ In addition to aspirin, a P2Y<sub>12</sub> inhibitor (either clopidogrel or ticagrelor) should be continued for up to 12 months in all patients with NSTE-ACS without contraindications who are treated with an ischemia-guided strategy.</li> <li>○ In patients receiving a stent (bare-metal stent or DES) during PCI for NSTE-ACS, P2Y<sub>12</sub> inhibitor therapy should be given for at least 12 months.</li> </ul> </li> <li>● Combined oral anticoagulant therapy and antiplatelet therapy in patients with NSTE-ACS           <ul style="list-style-type: none"> <li>○ The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor in patients with NSTE-ACS should be minimized to the extent possible to limit the risk of bleeding.</li> <li>○ Proton pump inhibitors should be prescribed in patients with NSTE-ACS with a history of gastrointestinal bleeding who require triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor.</li> </ul> </li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation (2020)</b><sup>14</sup></p>	<p><u>Pharmacological treatment of ischemia</u></p> <ul style="list-style-type: none"> <li>● Sublingual or intravenous nitrates and early initiation of β-blocker treatment is recommended in patients with ongoing ischemic symptoms and without contraindications.</li> <li>● Continuation of chronic β-blocker therapy is recommended unless the patient is in overt heart failure</li> <li>● Sublingual or intravenous nitrates are recommended to relieve angina; intravenous treatment is recommended in patients with recurrent angina, uncontrolled hypertension, or signs of heart failure.</li> <li>● In patients with suspected/confirmed vasospastic angina, calcium channel blockers, and nitrates should be considered and β-blockers avoided.</li> </ul> <p><u>Recommendations for platelet inhibition in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>● Aspirin is recommended for all patients without contraindications at an initial oral loading dose of 150 to 300 mg (in aspirin-naïve patients) and a maintenance dose of 75 to 100 mg/day long-term regardless of treatment strategy.</li> <li>● A P2Y<sub>12</sub> inhibitor is recommended, in addition to aspirin, for 12 months unless there are contraindications such as excessive risks of bleeds.           <ul style="list-style-type: none"> <li>○ Ticagrelor (180 mg loading dose, 90 mg twice daily) is recommended, in the absence of contraindication, for all patients at moderate-to-high risk of ischemic events (e.g., elevated cardiac troponins), regardless of initial treatment strategy and including those pretreated with clopidogrel (which should be discontinued when ticagrelor is started).</li> <li>○ Prasugrel (60 mg loading dose, 10 mg daily dose) is recommended in patients who are proceeding to PCI if no contraindication. Prasugrel should be considered in preference to ticagrelor in NSTE-ACS patients who proceed to PCI.</li> <li>○ Clopidogrel (300 to 600 mg loading dose, 75 mg daily dose) is</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<p>recommended for patients who cannot receive ticagrelor or prasugrel or who require oral anticoagulation.</p> <ul style="list-style-type: none"> <li>• P2Y<sub>12</sub> inhibitor administration for a shorter duration of three to six months after DES implantation may be considered in patients deemed at high bleeding risk.</li> <li>• Pre-treatment with a P2Y<sub>12</sub> inhibitor may be considered in patients with NSTEMI-ACS who are not planned to undergo an early invasive strategy.</li> <li>• It is not recommended to administer routine pre-treatment with a P2Y<sub>12</sub> inhibitor in patients in whom coronary anatomy is not known.</li> <li>• It is not recommended to administer prasugrel in patients whom coronary anatomy is not known.</li> <li>• GPIIb/IIIa inhibitors during PCI should be considered for bailout situations or thrombotic complications.</li> <li>• Cangrelor may be considered in P2Y<sub>12</sub> inhibitor-naïve patients undergoing PCI.</li> <li>• It is not recommended to administer GPIIb/IIIa inhibitors in patients whom coronary anatomy is not known.</li> <li>• P2Y<sub>12</sub> inhibitor administration in addition to aspirin beyond one year may be considered after careful assessment of the ischemic and bleeding risks of the patient.</li> </ul> <p><u>Recommendations for anticoagulation in non-ST-elevation acute coronary syndromes</u></p> <ul style="list-style-type: none"> <li>• Parenteral anticoagulation is recommended at the time of diagnosis according to both ischemic and bleeding risks.</li> <li>• Fondaparinux is recommended as having the most favorable efficacy-safety profile regardless of the management strategy.</li> <li>• Bivalirudin is recommended as an alternative to UFH plus GPIIb/IIIa inhibitors during PCI.</li> <li>• UFH is recommended in patients undergoing PCI who did not receive any anticoagulant.</li> <li>• In patients on fondaparinux undergoing PCI, a single intravenous bolus of UFH is recommended during the procedure.</li> <li>• Enoxaparin or UFH are recommended when fondaparinux is not available.</li> <li>• Enoxaparin should be considered as an anticoagulant for PCI in patients pretreated for PCI with subcutaneous enoxaparin.</li> <li>• Additional activated clotting time-guided intravenous boluses of UFH during PCI may be considered following initial UFH treatment.</li> <li>• Discontinuation of anticoagulation should be considered after PCI, unless otherwise indicated.</li> <li>• Crossover between UFH and LMWH is not recommended.</li> <li>• In NSTEMI patients with no prior stroke/TIA and at high ischemic risk as well as low bleeding risk receiving aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily for approximately one year) may be considered after discontinuation of parenteral anticoagulation.</li> </ul> <p><u>Recommendations for combining antiplatelet agents and anticoagulants in non-ST-elevation acute coronary syndrome patients requiring chronic oral anticoagulation</u></p> <ul style="list-style-type: none"> <li>• In patients with a firm indication for oral anticoagulation (e.g., atrial fibrillation with a CHADS<sub>2</sub>-VASc score ≥2, recent VTE, mechanical valve prosthesis), oral anticoagulation is recommended in addition to antiplatelet therapy.</li> <li>• An early invasive coronary angiography (within 24 hours) should be considered in moderate- to high-risk patients, irrespective of oral anticoagulant exposure, to expedite treatment allocation (medical vs PCI vs CABG) and to determine optimal antithrombotic regimen.</li> <li>• Initial dual antiplatelet therapy with aspirin plus a P2Y<sub>12</sub> inhibitor in addition to oral anticoagulation before coronary angiography is not recommended.</li> <li>• During PCI, additional parenteral anticoagulation is recommended, irrespective</li> </ul>

Clinical Guideline	Recommendations
	<p>of the timing of the last dose of all non-vitamin K antagonist oral anticoagulants (NOACs) and if INR is &lt;2.5 in VKA-treated patients.</p> <ul style="list-style-type: none"> <li>• Uninterrupted therapeutic anticoagulation with VKA or NOACs should be considered during the periprocedural phase.</li> <li>• Periprocedural DAPT administration consisting of aspirin and clopidogrel up to one week is recommended</li> <li>• Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months</li> <li>• Following coronary stenting, dual (oral) antiplatelet therapy (DAPT) including new P2Y<sub>12</sub> inhibitors should be considered as an alternative to triple therapy for patients with non-ST-elevation acute coronary syndromes and atrial fibrillation with a CHADS<sub>2</sub>-VASc score of 1 (in males) or 2 (in females).</li> <li>• If at low bleeding risk (HAS-BLED ≤2), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for six months, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months.</li> <li>• If at high bleeding risk (HAS-BLED ≥3), triple therapy with oral anticoagulant, aspirin, and clopidogrel should be considered for one month, followed by oral anticoagulant and aspirin or clopidogrel continued up to 12 months irrespective of the stent type.</li> <li>• Dual therapy with oral anticoagulant and clopidogrel may be considered as an alternative to triple antithrombotic therapy in selected patients (HAS-BLED ≥3 and low risk of stent thrombosis).</li> <li>• The use of ticagrelor or prasugrel as part of triple therapy is not recommended.</li> <li>• In medically managed patients, one antiplatelet agent in addition to oral anticoagulant should be considered for up to one year.</li> </ul> <p><u>Recommendations for post-interventional and maintenance treatment</u></p> <ul style="list-style-type: none"> <li>• In patients with NSTEMI-ACS with coronary stent implantation, DAPT with a P2Y<sub>12</sub> inhibitor on top of aspirin is recommended for 12 months unless there are contraindications such as excessive risk of bleeding.</li> <li>• Adding a second anti-thrombotic agent to aspirin for extended long-term secondary prevention should be considered in patients with a moderate to high risk of ischemic events and without increased risk of major bleeding.</li> <li>• After stent implantation with high risk of bleeding, discontinuation of P2Y<sub>12</sub> inhibitor therapy after three months should be considered</li> <li>• After stent implantation in patients undergoing DAPT, stopping aspirin after three to six months should be considered, depending on balance between ischemic and bleeding risk.</li> <li>• De-escalation of P2Y<sub>12</sub> inhibitor treatment may be considered as an alternative DAPT strategy, especially for ACS patients deemed unsuitable for potent platelet inhibition.</li> </ul>
<p>American College of Cardiology/American Heart Association:  <b>Guideline for the Management of ST-Elevation Myocardial Infarction (2013)</b><sup>15</sup></p>	<p><u>Routine medical therapies: β-blockers</u></p> <ul style="list-style-type: none"> <li>• Oral β-blockers should be initiated within the first 24 hours in patients with an ST-segment elevation myocardial infarction (STEMI) who do not have any of the following: 1) signs of heart failure, 2) evidence of a low-output state, 3) increased risk of cardiogenic shock, 4) other contraindications to use of oral β-blockers (e.g., PR interval &gt;24 seconds, second or third degree heart block, active asthma, reactive airway disease).</li> <li>• β-blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use.</li> <li>• Patients with initial contraindications to the use of β-blockers in the first 24 hours after STEMI should be re-evaluated to determine their subsequent eligibility.</li> <li>• It is reasonable to administer intravenous β-blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia.</li> </ul>



Clinical Guideline	Recommendations
	<p><u>Routine medical therapies: renin-angiotensin-aldosterone system inhibitors</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor should be administered within the first 24 hours to all patients with ST-segment elevation myocardial infarction with anterior location, heart failure, or ejection fraction <math>\leq 40\%</math>, unless contraindicated.</li> <li>• An ARB should be given to patients who have indications for but are intolerant of ACE inhibitors.</li> <li>• ACE inhibitors are reasonable for all patients with no contraindications to their use.</li> <li>• An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and <math>\beta</math>-blocker and who have an EF <math>\leq 40\%</math> and either symptomatic heart failure or diabetes.</li> </ul> <p><u>Routine medical therapies: Lipid management</u></p> <ul style="list-style-type: none"> <li>• High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use.</li> <li>• It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation.</li> </ul>
<p>European Society of Cardiology: <b>Management of Acute Myocardial Infarction in Patients Presenting with ST-segment Elevation (2017)</b><sup>16</sup></p>	<p><u>Routine therapies in the acute, subacute and long term phase of ST-elevation myocardial infarction (STEMI)</u></p> <ul style="list-style-type: none"> <li>• Antiplatelet therapy with low dose aspirin (75 to 100 mg) is indicated indefinitely after STEMI.</li> <li>• Dual antiplatelet therapy with a combination of aspirin and prasugrel or aspirin and ticagrelor is recommended for 12 months after percutaneous coronary intervention (PCI), unless there are contraindications such as excessive risk of bleeding.</li> <li>• A proton pump inhibitor (PPI) in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.</li> <li>• In patients with an indication for oral anticoagulation, oral anticoagulants are indicated in addition to antiplatelet therapy.</li> <li>• In patients who are at high risk of severe bleeding complications, discontinuation of P2Y<sub>12</sub> inhibitor therapy after six months should be considered.</li> <li>• In STEMI patients with stent implantation and an indication for oral anticoagulation, triple therapy (oral anticoagulant, aspirin, and clopidogrel) should be considered for one to six months (according a balance between the estimated risk of recurrent coronary events and bleeding).</li> <li>• In patients with left ventricular thrombus, anticoagulation should be instituted for a minimum of six months, guided by repeated imaging.</li> <li>• In selected patients who receive aspirin and clopidogrel, low-dose rivaroxaban (2.5 mg twice daily) may be considered if the patient is at low bleeding risk.</li> <li>• Dual antiplatelet therapy should be used up to one year in patients with STEMI who did not receive a stent unless there are contraindications such as excessive risk of bleeding.</li> <li>• In high ischemic-risk patients (age <math>\geq 50</math> years, and at least one of the following risk factors: age <math>\geq 65</math> years, diabetes mellitus on medication, prior spontaneous MAI, multivessel CAD, or chronic renal dysfunction with eGFR <math>&lt; 60</math> mL/min) who have tolerated dual antiplatelet therapy without a bleeding complication, treatment with dual antiplatelet therapy in the form of ticagrelor 60 mg twice a day on top of aspirin for longer than 12 months may be considered for up to three years.</li> <li>• The use of ticagrelor or prasugrel is not recommended as part of triple antithrombotic therapy with aspirin and oral anticoagulation.</li> <li>• Oral treatment with <math>\beta</math>-blockers should be considered during hospital stay and continued thereafter in all patients without contraindications.</li> <li>• Oral treatment with <math>\beta</math>-blockers is indicated in patients with heart failure or left</li> </ul>

Clinical Guideline	Recommendations
	<p>ventricular dysfunction, LVEF <math>\leq</math>40% unless contraindicated.</p> <ul style="list-style-type: none"> <li>• Intravenous <math>\beta</math>-blockers must be avoided in patients with hypotension or acute heart failure or AV block or severe bradycardia.</li> <li>• Intravenous <math>\beta</math>-blockers should be considered at the time of presentation in patients undergoing primary PCI without contraindications, with high blood pressure, tachycardia, and no signs of heart failure.</li> <li>• A fasting lipid profile must be obtained in all STEMI patients, as soon as possible after presentation.</li> <li>• It is recommended to initiate or continue high dose statins early after admission in all STEMI patients without contraindication or history of intolerance, regardless of initial cholesterol values and maintain it long-term.</li> <li>• An LDL-C goal of <math>&lt;1.8</math> mmol/L (70 mg/dL) or a reduction of at least 50% if the baseline LDL-C is between 1.8 to 3.5 mmol/L (70 to 135 mg/dL) is recommended.</li> <li>• In patients with LDL-C <math>&gt;1.8</math> mmol/L (<math>&gt;70</math> mg/dL) despite a maximally tolerated statin dose who remain at high risk, further therapy to reduce LDL-C should be considered.</li> <li>• ACE inhibitors are indicated starting within the first 24 hours of STEMI in patients with evidence of heart failure, LV systolic dysfunction, diabetes or an anterior infarct.</li> <li>• An ARB, preferably valsartan, is an alternative to ACE inhibitors in patients with heart failure or LV systolic dysfunction, particularly those who are intolerant to ACE inhibitors.</li> <li>• ACE inhibitors should be considered in all patients in the absence of contraindications.</li> <li>• Aldosterone antagonists, e.g. eplerenone, are indicated in patients with an ejection fraction <math>\leq</math>40% and heart failure or diabetes, provided no renal failure or hyperkalemia.</li> </ul>
<p>American College of Cardiology/ American Heart Association: <b>Guideline on the Primary Prevention of Cardiovascular Disease (2019)</b><sup>17</sup></p>	<p><u>Top 10 messages for the primary prevention of cardiovascular disease</u></p> <ul style="list-style-type: none"> <li>• The most important way to prevent atherosclerotic vascular disease, heart failure, and atrial fibrillation is to promote a healthy lifestyle throughout life.</li> <li>• A team-based care approach is an effective strategy for the prevention of cardiovascular disease. Clinicians should evaluate the social determinants of health that affect individuals to inform treatment decisions.</li> <li>• Adults who are 40 to 75 years of age and are being evaluated for cardiovascular disease prevention should undergo 10-year atherosclerotic cardiovascular disease (ASCVD) risk estimation and have a clinician–patient risk discussion before starting on pharmacological therapy, such as antihypertensive therapy, a statin, or aspirin. In addition, assessing for other risk-enhancing factors can help guide decisions about preventive interventions in select individuals, as can coronary artery calcium scanning.</li> <li>• All adults should consume a healthy diet that emphasizes the intake of vegetables, fruits, nuts, whole grains, lean vegetable or animal protein, and fish and minimizes the intake of trans fats, processed meats, refined carbohydrates, and sweetened beverages. For adults with overweight and obesity, counseling and caloric restriction are recommended for achieving and maintaining weight loss.</li> <li>• Adults should engage in at least 150 minutes per week of accumulated moderate-intensity physical activity or 75 minutes per week of vigorous-intensity physical activity.</li> <li>• For adults with type 2 diabetes mellitus, lifestyle changes, such as improving dietary habits and achieving exercise recommendations, are crucial. If medication is indicated, metformin is first-line therapy, followed by consideration of a sodium-glucose cotransporter 2 inhibitor or a glucagon-like peptide-1 receptor agonist.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• All adults should be assessed at every healthcare visit for tobacco use, and those who use tobacco should be assisted and strongly advised to quit.</li> <li>• Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.</li> <li>• Statin therapy is first-line treatment for primary prevention of ASCVD in patients with elevated low-density lipoprotein cholesterol levels (<math>\geq 190</math> mg/dL), those with diabetes mellitus, who are 40 to 75 years of age, and those determined to be at sufficient ASCVD risk after a clinician–patient risk discussion.</li> <li>• Nonpharmacological interventions are recommended for all adults with elevated blood pressure or hypertension. For those requiring pharmacological therapy, the target blood pressure should generally be <math>&lt;130/80</math> mm Hg.</li> </ul> <p><u>Adults with Type 2 Diabetes Mellitus</u></p> <ul style="list-style-type: none"> <li>• For all adults with T2DM, a tailored nutrition plan focusing on a heart-healthy dietary pattern is recommended to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• Adults with T2DM should perform at least 150 minutes per week of moderate-intensity physical activity or 75 minutes of vigorous-intensity physical activity to improve glycemic control, achieve weight loss if needed, and improve other ASCVD risk factors.</li> <li>• For adults with T2DM, it is reasonable to initiate metformin as first-line therapy along with lifestyle therapies at the time of diagnosis to improve glycemic control and reduce ASCVD risk.</li> <li>• For adults with T2DM and additional ASCVD risk factors who require glucose-lowering therapy despite initial lifestyle modifications and metformin, it may be reasonable to initiate a sodium-glucose cotransporter 2 (SGLT-2) inhibitor or a glucagon-like peptide-1 receptor (GLP-1R) agonist to improve glycemic control and reduce CVD risk.</li> </ul> <p><u>Adults with high blood cholesterol</u></p> <ul style="list-style-type: none"> <li>• In adults at intermediate risk (<math>\geq 7.5\%</math> to <math>&lt;20\%</math> 10-year ASCVD risk), statin therapy reduces risk of ASCVD, and in the context of a risk discussion, if a decision is made for statin therapy, a moderate-intensity statin should be recommended.</li> <li>• In intermediate risk (<math>\geq 7.5\%</math> to <math>&lt;20\%</math> 10-year ASCVD risk) patients, LDL-C levels should be reduced by 30% or more, and for optimal ASCVD risk reduction, especially in patients at high risk (<math>\geq 20\%</math> 10-year ASCVD risk), levels should be reduced by 50% or more.</li> <li>• In adults 40 to 75 years of age with diabetes, regardless of estimated 10-year ASCVD risk, moderate-intensity statin therapy is indicated.</li> <li>• In patients 20 to 75 years of age with an LDL-C level of 190 mg/dL (<math>\geq 4.9</math> mmol/L) or higher, maximally tolerated statin therapy is recommended.</li> <li>• In adults with diabetes mellitus who have multiple ASCVD risk factors, it is reasonable to prescribe high-intensity statin therapy with the aim to reduce LDL-C levels by 50% or more.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt;20\%</math> 10-year ASCVD risk) adults, risk-enhancing factors favor initiation or intensification of statin therapy.</li> <li>• In intermediate-risk (<math>\geq 7.5\%</math> to <math>&lt;20\%</math> 10-year ASCVD risk) adults or selected borderline-risk (5% to <math>&lt;7.5\%</math> 10-year ASCVD risk) adults in whom a coronary artery calcium score is measured for the purpose of making a treatment decision, AND <ul style="list-style-type: none"> <li>○ If the coronary artery calcium score is zero, it is reasonable to withhold statin therapy and reassess in five to 10 years, as long as higher-risk conditions are absent (e.g., diabetes, family history of premature CHD, cigarette smoking);</li> </ul> </li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>○ If coronary artery calcium score is one to 99, it is reasonable to initiate statin therapy for patients <math>\geq 55</math> years of age;</li> <li>○ If coronary artery calcium score is 100 or higher or in the 75th percentile or higher, it is reasonable to initiate statin therapy.</li> </ul> <ul style="list-style-type: none"> <li>● In patients at borderline risk (5% to <math>&lt;7.5\%</math> 10-year ASCVD risk), in risk discussion, the presence of risk-enhancing factors may justify initiation of moderate-intensity statin therapy.</li> </ul> <p><u>Adults with high blood pressure or hypertension</u></p> <ul style="list-style-type: none"> <li>● In adults with elevated blood pressure (BP) or hypertension, including those requiring antihypertensive medications nonpharmacological interventions are recommended to reduce BP. These include:           <ul style="list-style-type: none"> <li>○ weight loss;</li> <li>○ a heart-healthy dietary pattern;</li> <li>○ sodium reduction;</li> <li>○ dietary potassium supplementation;</li> <li>○ increased physical activity with a structured exercise program; and</li> <li>○ limited alcohol.</li> </ul> </li> <li>● In adults with an estimated 10-year ASCVD risk (ACC/AHA pooled cohort equations to estimate 10-year risk of ASCVD) of 10% or higher and an average systolic BP (SBP) of 130 mm Hg or higher or an average diastolic BP (DBP) of 80 mm Hg or higher, use of BP-lowering medications is recommended for primary prevention of CVD.</li> <li>● In adults with confirmed hypertension and a 10-year ASCVD event risk of 10% or higher, a BP target of less than 130/80 mm Hg is recommended.</li> <li>● In adults with hypertension and chronic kidney disease, treatment to a BP goal of less than 130/80 mm Hg is recommended.</li> <li>● In adults with T2DM and hypertension, antihypertensive drug treatment should be initiated at a BP of 130/80 mm Hg or higher, with a treatment goal of less than 130/80 mm Hg.</li> <li>● In adults with an estimated 10-year ASCVD risk <math>&lt;10\%</math> and an SBP of 140 mm Hg or higher or a DBP of 90 mm Hg or higher, initiation and use of BP-lowering medication are recommended.</li> <li>● In adults with confirmed hypertension without additional markers of increased ASCVD risk, a BP target of less than 130/80 mm Hg may be reasonable.</li> </ul> <p><u>Recommendations for treatment of tobacco use</u></p> <ul style="list-style-type: none"> <li>● All adults should be assessed at every healthcare visit for tobacco use and their tobacco use status recorded as a vital sign to facilitate tobacco cessation.</li> <li>● To achieve tobacco abstinence, all adults who use tobacco should be firmly advised to quit.</li> <li>● In adults who use tobacco, a combination of behavioral interventions plus pharmacotherapy is recommended to maximize quit rates.</li> <li>● In adults who use tobacco, tobacco abstinence is recommended to reduce ASCVD risk.</li> <li>● To facilitate tobacco cessation, it is reasonable to dedicate trained staff to tobacco treatment in every healthcare system.</li> <li>● All adults and adolescents should avoid secondhand smoke exposure to reduce ASCVD risk.</li> </ul> <p><u>Recommendations for aspirin use</u></p> <ul style="list-style-type: none"> <li>● Low-dose aspirin (75 to 100 mg orally daily) might be considered for the primary prevention of ASCVD among select adults 40 to 70 years of age who are at higher ASCVD risk but not at increased bleeding risk.</li> <li>● Low-dose aspirin (75 to 100 mg orally daily) should not be administered on a</li> </ul>

Clinical Guideline	Recommendations
	<p>routine basis for the primary prevention of ASCVD among adults &gt;70 years of age.</p> <ul style="list-style-type: none"> <li>• Low-dose aspirin (75 to 100 mg orally daily) should not be administered for the primary prevention of ASCVD among adults of any age who are at increased risk of bleeding.</li> </ul>
<p>American Heart Association/American College of Cardiology/Heart Failure Society of America: <b>2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)</b><sup>18</sup></p>	<p><b>Treatment of Stage A heart failure (HF)</b></p> <ul style="list-style-type: none"> <li>• Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>• In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>• In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>• Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><b>Treatment of Stage B heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>• In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>• In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an</li> </ul>

Clinical Guideline	Recommendations
	<p>ARNI is not feasible. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
European Society of Cardiology:	Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction

Clinical Guideline	Recommendations
<p><b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>19</sup></p>	<ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF</li> </ul>



Clinical Guideline	Recommendations
	<p>hospitalization and death.</p> <ul style="list-style-type: none"> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
Eighth Joint National Committee (JNC 8):	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure</li> </ul>

Clinical Guideline	Recommendations
<p><b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>20</sup></p>	<p>≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</p> <ul style="list-style-type: none"> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p><b>International Society of Hypertension: Global Hypertension Practice Guidelines (2020)</b><sup>21</sup></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation</li> </ul>

Clinical Guideline	Recommendations
	<p>programs are advised.</p> <ul style="list-style-type: none"> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week. Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment           <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment           <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy           <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for</p>	<p>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</p>

Clinical Guideline	Recommendations
<p><b>the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)</b><sup>22</sup></p>	<ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq 100</math> mmHg or average systolic blood pressure (SBP) measurements of <math>\geq 160</math> mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq 90</math> mmHg or for average SBP readings <math>\geq 140</math> mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC).             <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are:                 <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE</li> </ul>

Clinical Guideline	Recommendations
	<p>inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</p> <ul style="list-style-type: none"> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt; 40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening</li> </ul>

Clinical Guideline	Recommendations
	<p>renal function. Additional therapies may also include dihydropyridine CCBs.</p> <ul style="list-style-type: none"> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR ≤30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours)             <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke             <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours)             <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in</li> </ul>

Clinical Guideline	Recommendations
	<p>cases of complex lesions less amenable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</p> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during</li> </ul>



Clinical Guideline	Recommendations
	<p>pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</p> <ul style="list-style-type: none"> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>23</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be</li> </ul>



Clinical Guideline	Recommendations
	<p>accompanied by close follow-up.</p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of</li> </ul>

Clinical Guideline	Recommendations
	<p>gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</p> <ul style="list-style-type: none"> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul>

Clinical Guideline	Recommendations
	<p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>24</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African–Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>

Clinical Guideline	Recommendations
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>25</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>26</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and</li> </ul>

Clinical Guideline	Recommendations
	<p>direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</p> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>• The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>27</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>• Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>• Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>• Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>• For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>• For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>• Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event</li> </ul>

Clinical Guideline	Recommendations
	<p>to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</p> <ul style="list-style-type: none"> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal <math>&lt;130/80</math> mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal <math>&lt;130</math> mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP <math>&lt;140</math> mmHg and a DBP <math>&lt;90</math> mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal <math>&lt;130/80</math> mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target <math>&lt;130/80</math> mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul>



Clinical Guideline	Recommendations
	<p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP ≥180 mmHg or DBP ≥110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>28</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (≥140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure ≥180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk ≥15%), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it</li> </ul>

Clinical Guideline	Recommendations
	<p>can be safely attained.</p> <ul style="list-style-type: none"> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq</math>300 mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><u>Chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq</math>300 mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq</math>300 mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq</math>25</li> </ul>

Clinical Guideline	Recommendations
	<p>mL/min/1.73 m<sup>2</sup> or ≥300 mg/g, respectively.</p> <ul style="list-style-type: none"> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium–glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (≤30%) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin–to–creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin–to–creatinine ratio ≥300 mg/g creatinine and/or estimated glomerular filtration rate &lt;60 mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin–to–creatinine ratio (&lt;30 mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate &lt;30 mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Association for the Study of Liver Diseases: <b>Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance (2021)</b><sup>29</sup></p>	<p><b>Treatment of ascites</b></p> <ul style="list-style-type: none"> <li>• Moderate sodium restriction (2 g or 90 mmol/day) and diuretics (spironolactone with or without furosemide) are the first-line treatment in patients with cirrhosis and grade 2 ascites.</li> <li>• After ascites is adequately mobilized, attempts should be made to taper the diuretics to the lowest dose necessary to maintain minimal or no ascites to prevent the development of adverse effects.</li> <li>• Fluid restriction is not necessary unless serum sodium is &lt;125 mmol/L.</li> <li>• In patients receiving diuretics, body weight and serum creatinine and sodium should be regularly monitored to assess response and to detect the development of adverse effects.</li> <li>• Human albumin solution (20 to 40 g/week) or baclofen administration (10 mg/day, with a weekly increase of 10 mg/day, up to 30 mg/day) can be considered in cases of severe muscle cramps.</li> <li>• Large-volume paracentesis is the first-line treatment of grade 3 ascites. After paracentesis, sodium restriction and diuretics should be started.</li> <li>• Referral for liver transplant evaluation should be considered in patients with grade 2 or 3 ascites.</li> <li>• Nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers should be avoided in patients with cirrhosis and ascites.</li> <li>• Aminoglycosides should be avoided whenever possible in the treatment of bacterial infections.</li> <li>• For patients with cirrhosis and diuretic-responsive ascites, controversial data</li> </ul>

Clinical Guideline	Recommendations
	suggest potential benefits of long-term infusion of human albumin solution. At present, no recommendation can be made for its use in routine clinical practice.
Endocrine Society: <b>The Management of Primary Aldosteronism: Case Detection, Diagnosis, and Treatment (2016)</b> <sup>30</sup>	<ul style="list-style-type: none"> <li>• Unilateral laparoscopic adrenalectomy is recommended for patients with documented unilateral primary aldosteronism (i.e., aldosterone-producing adenoma [APA] or unilateral adrenal hyperplasia [UAH]). If a patient is unable or unwilling to undergo surgery, medical treatment including a mineralocorticoid receptor (MR) antagonist is recommended. If an aldosterone to renin ratio (ARR)-positive patient is unwilling or unable to undergo further investigations, medical treatment including an MR antagonist is recommended.</li> <li>• In patients with primary aldosteronism due to bilateral adrenal disease, medical treatment with an MR antagonist is recommended; spironolactone is suggested as the primary agent, with eplerenone as an alternative</li> <li>• In patients with glucocorticoid remediable aldosteronism (GRA), administering the lowest dose of glucocorticoid to lower adrenocorticotropic hormone (ACTH) and thus normalize BP and potassium levels is recommended as the first-line treatment. In addition, if BP fails to normalize with glucocorticoid alone, an MR antagonist may be added. For children, the glucocorticoid dosage should be adjusted for age and body weight, and BP targets should be determined from age- and gender-specific published normative data.</li> </ul>

\*Agent not available in the United States.

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the mineralocorticoid (aldosterone) receptor antagonists are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Mineralocorticoid (Aldosterone) Receptor Antagonists<sup>1-7</sup>**

Indication(s)	Single Entity Agents			Combination Products
	Eplerenone	Finerenone	Spironolactone	Spironolactone and HCTZ
<b>Edematous Conditions</b>				
Maintenance therapy together with bed rest and the restriction of fluid and sodium in patients with cirrhosis of the liver accompanied by edema and/or ascites			✓	✓
Management of edema and sodium retention when the patient is only partially responsive to, or is intolerant of, other therapeutic measures			✓	✓
Nephrotic patients when treatment of the underlying disease, restriction of fluid and sodium intake, and the use of other diuretics do not provide an adequate response			✓	✓
Patients with congestive heart failure taking digitalis when other therapies are considered inappropriate			✓	✓
<b>Heart Failure</b>				
Increase survival and reduce the need for hospitalization for heart failure when used in addition to standard therapy in patients with severe heart failure (New York Heart			✓	

Indication(s)	Single Entity Agents			Combination Products
	Eplerenone	Finerenone	Spirolactone	Spirolactone and HCTZ
Association functional class III-IV)				
<b>Hypertension</b>				
Essential hypertension				✓ *
Hypertension	✓ †		✓ ‡	
<b>Hypokalemia</b>				
Prophylaxis of hypokalemia in patients taking digitalis when other measures are considered inadequate or inappropriate			✓	
Treatment of a diuretic-induced hypokalemia in patients with congestive heart failure when other measures are considered inappropriate				✓
Treatment of diuretic-induced hypokalemia in patients with hypertension when other measures are considered inappropriate				✓
Treatment of patients with hypokalemia when other measures are considered inappropriate or inadequate			✓	
<b>Myocardial Infarction</b>				
To improve survival of stable patients with left ventricular systolic dysfunction (ejection fraction ≤40%) and clinical evidence of congestive heart failure after an acute myocardial infarction	✓			
<b>Primary Hyperaldosteronism</b>				
Establish the diagnosis of primary hyperaldosteronism by therapeutic trial			✓	
Short-term preoperative treatment of patients with primary hyperaldosteronism			✓	
Long-term maintenance therapy for patients with discrete aldosterone-producing adrenal adenomas who are judged to be poor operative risks or who decline surgery			✓	
Long-term maintenance therapy for patients with bilateral micro or macronodular adrenal hyperplasia (idiopathic hyperaldosteronism)			✓	
<b>Diabetic Kidney Disease</b>				
To reduce the risk of sustained eGFR decline, end stage kidney disease, cardiovascular death, non-fatal myocardial infarction, and hospitalization for heart failure in adult patients with chronic kidney disease associated with type 2 diabetes		✓		

\*In patients in whom other measures are considered inadequate or inappropriate.

†Alone or in combination with other antihypertensive agents.

‡Usually in combination with other drugs, in patients who cannot be treated adequately with other agents or for whom other agents are considered inappropriate.

eGFR=estimated glomerular filtration rate, HCTZ=hydrochlorothiazide

#### IV. Pharmacokinetics

The pharmacokinetic parameters of the mineralocorticoid (aldosterone) receptor antagonists are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Mineralocorticoid (Aldosterone) Receptor Antagonists<sup>2</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Eplerenone	69	50	Liver, extensive (% not reported)	Renal (67) Feces (32)	3 to 6
Finerenone	44	92	Liver (% not reported)	Renal (80) Feces (20)	2 to 3
Spironolactone	73	90	Liver (% not reported) Renal (% not reported)	Renal (47 to 57) Feces (35 to 41)	1.3 to 1.4
<b>Combination Products</b>					
Spironolactone and HCTZ	73/ 60 to 80	90/40	Liver (% not reported) Renal (% not reported)	Feces (35 to 41) Renal (47 to 57)/ Renal (50 to 70)	1.3 to 1.4/ 4 to 5

HCTZ=hydrochlorothiazide

#### V. Drug Interactions

Major drug interactions with the mineralocorticoid (aldosterone) receptor antagonists are listed in Table 5.

**Table 5. Major Drug Interactions with the Mineralocorticoid (Aldosterone) Receptor Antagonists<sup>2</sup>**

Generic Name(s)	Interaction	Mechanism
Mineralocorticoid receptor antagonists (eplerenone, spironolactone)	ACE inhibitors	Serious hyperkalemia, possibly with cardiac arrhythmias or arrest, may occur with the combination of aldosterone blockers and ACE inhibitors. Potassium sparing effects are additive when combining ACE inhibitors with aldosterone blockers. Aldosterone acts in the renal cortical collecting ducts by inducing synthesis of proteins that constitute the Na <sup>+</sup> , K <sup>+</sup> -ATPase pump. The pump acts to reabsorb sodium and water in exchange for potassium, which is then eliminated in the urine. Aldosterone antagonism can cause hyperkalemia.
Mineralocorticoid receptor antagonists (eplerenone)	Amiloride	Aldosterone blockers and amiloride may exert additive pharmacologic effects. Hyperkalemia with the potential for cardiac arrhythmias may result. Aldosterone blockers and amiloride may cause additive adverse effects when co-administered.
Mineralocorticoid receptor antagonists (eplerenone, spironolactone)	Potassium Preparations	Potassium preparations will increase serum potassium concentrations. This may increase the potential for clinically important hyperkalemia, especially when used concomitantly with aldosterone blockers.
Mineralocorticoid receptor antagonists (eplerenone, spironolactone)	Triamterene	Eplerenone and triamterene may exert additive pharmacologic effects. Hyperkalemia with the potential for cardiac arrhythmias may result.
Mineralocorticoid receptor antagonists (eplerenone)	HIV Protease Inhibitors	Inhibition of CYP3A4 isoenzymes by HIV protease inhibitors may decrease the metabolic elimination of aldosterone blockers. HIV protease inhibitors may increase plasma concentrations and pharmacologic or toxic effects of aldosterone blockers.
Mineralocorticoid receptor antagonists	Imidazoles	Certain azole antifungal agents may decrease the elimination of eplerenone by inhibiting its hepatic metabolism via



Generic Name(s)	Interaction	Mechanism
(eplerenone)		CYP3A4 isoenzyme resulting in increased concentration and consequently increased pharmacologic and toxic (hyperkalemia associated with potentially fatal arrhythmias) effects of eplerenone.
Mineralocorticoid receptor antagonists (eplerenone)	Macrolides	Macrolides may decrease the elimination of eplerenone by inhibiting its hepatic metabolism via CYP3A4 isoenzyme resulting in increased concentration and consequently increased pharmacologic and toxic (hyperkalemia associated with potentially fatal arrhythmias) effects of eplerenone.
Mineralocorticoid receptor antagonists (eplerenone)	Nefazodone	Nefazodone may decrease the elimination of eplerenone by inhibiting its hepatic metabolism via CYP3A4 isoenzyme resulting in increased concentration and consequently increased pharmacologic and toxic (hyperkalemia associated with potentially fatal arrhythmias) effects of eplerenone. Coadministration of eplerenone with nefazodone is contraindicated.
Mineralocorticoid receptor antagonists (eplerenone)	Spironolactone	Eplerenone and spironolactone may exert additive pharmacologic effects. Hyperkalemia with the potential for cardiac arrhythmias may result.
Mineralocorticoid receptor antagonists (eplerenone)	Verapamil	Inhibition of CYP3A4 isoenzymes by verapamil may decrease the metabolic elimination of eplerenone. Verapamil may increase plasma concentrations and pharmacologic or toxic effects of eplerenone.
Mineralocorticoid receptor antagonists (spironolactone)	Angiotensin II Receptor Antagonists	Decreased aldosterone activity by angiotensin II receptor antagonists may function synergistically with potassium conservation by spironolactone to produce substantial hyperkalemia. The risk of hyperkalemia may be increased when spironolactone is co-administered with angiotensin II receptor antagonists.
Mineralocorticoid receptor antagonists (spironolactone)	Eplerenone	Concurrent use of eplerenone and spironolactone may result in increased risk of hyperkalemia.
Mineralocorticoid receptor antagonists (spironolactone)	Digoxin	Concurrent use of digoxin and spironolactone may result in increased digoxin exposure.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics may induce hypokalemia which may increase the risk of torsades de pointes. The coadministration of dofetilide with a thiazide diuretic is contraindicated.
Mineralocorticoid receptor antagonists (spironolactone)	Lithium	Concurrent use of lithium and spironolactone may result in increased lithium concentrations and lithium toxicity (weakness, tremor, excessive thirst, confusion).
Mineralocorticoid receptor antagonists (finerenone)	Strong CYP3A4 inhibitors	Concurrent use of finerenone and strong CYP3A4 inhibitors may result in increased finerenone exposure.
Mineralocorticoid receptor antagonists (finerenone)	Moderate or weak CYP3A4 inhibitors	Concurrent use of finerenone and moderate or weak CYP3A4 inhibitors may result in increased finerenone exposure.
Mineralocorticoid receptor antagonists (finerenone)	Strong or moderate CYP3A4 inducers	Concurrent use of finerenone and strong or moderate CYP3A4 inducers may result in decreased finerenone exposure.
Thiazide diuretics (HCTZ)	Diazoxide	The combination of diazoxide with a thiazide diuretic may lead to hyperglycemia through an unknown mechanism; therefore the combination should be avoided. When used together, blood and urine glucose levels should be frequently monitored, and dosage reductions may be required.
Thiazide diuretics	Digitalis glycosides	Thiazide diuretics may induce electrolyte disturbances which

Generic Name(s)	Interaction	Mechanism
(HCTZ)		may predispose patients to digitalis-induced arrhythmias. Measure plasma levels of potassium and magnesium, supplement low levels, and use dietary sodium restriction or potassium-sparing diuretics to prevent further losses.

ACE inhibitors=angiotensin converting enzyme inhibitors, CYP=cytochrome P450 isoenzyme, HCTZ=hydrochlorothiazide, HIV=human immunodeficiency virus

## VI. Adverse Drug Events

The most common adverse drug events reported with the mineralocorticoid (aldosterone) receptor antagonists are listed in Table 6.

**Table 6. Adverse Drug Events (%) Reported with the Mineralocorticoid (Aldosterone) Receptor Antagonists<sup>1-7</sup>**

Adverse Events	Single Entity Agents			Combination Products
	Eplerenone	Finerenone	Spirolactone	Spirolactone and HCTZ
<b>Cardiovascular</b>				
Hypotension	-	4.8	-	-
Orthostatic hypotension	-	-	-	✓
<b>Central Nervous System</b>				
Ataxia	-	-	✓	✓
Confusion	-	-	✓	✓
Dizziness	3	-	-	✓
Drowsiness	-	-	✓	✓
Fatigue	2	-	✓	✓
Fever	-	-	✓	✓
Headache	-	-	✓	✓
Insomnia	-	-	-	✓
Lethargy	-	-	✓	✓
Restlessness	-	-	-	✓
Vertigo	-	-	-	✓
<b>Dermatological</b>				
Alopecia	-	-	-	✓
Cutaneous vasculitis	-	-	-	✓
Erythema multiforme	-	-	-	✓
Exfoliative dermatitis	-	-	-	✓
Maculopapular eruptions	-	-	-	✓
Necrotizing angitis	-	-	-	✓
Photosensitivity	-	-	-	✓
Pruritus	-	-	-	✓
Purpura	-	-	-	✓
Rash	<1	-	✓	✓
Stevens-Johnson syndrome	-	-	-	✓
Toxic epidermal necrolysis	-	-	-	✓
Urticaria	-	-	✓	✓
<b>Endocrine and Metabolic</b>				
Amenorrhea	-	-	✓	✓
Breast cancer	-	-	✓	✓
Deepening of the voice	-	-	✓	✓
Dehydration	-	-	✓	✓
Gynecomastia	≤1	-	9	9
Hyperchloremic metabolic acidosis	-	-	✓	✓



Adverse Events	Single Entity Agents			Combination Products
	Eplerenone	Finerenone	Spirolactone	Spirolactone and HCTZ
Irregular menses	-		✓	✓
Mastodynia	≤1		2	2
Postmenopausal bleeding	-		✓	✓
<b>Gastrointestinal</b>				
Abdominal pain	1		-	✓
Anorexia	-		✓	✓
Cholestatic toxicity	-		✓	✓
Constipation	-		-	✓
Cramping	-		✓	✓
Diarrhea	2		✓	✓
Gastric bleeding	-		✓	✓
Gastritis	-		✓	✓
Nausea	-		✓	✓
Pancreatitis	-		-	✓
Sialoadenitis	-		-	✓
Ulceration	-		✓	✓
Vomiting	-		✓	✓
Xerostomia	-		✓	✓
<b>Genitourinary</b>				
Abnormal vaginal bleeding	≤2		-	-
Albuminuria	1		-	-
Glucosuria	-		-	✓
Impotence	-		✓	✓
Interstitial nephritis	-		-	✓
Renal dysfunction	-		✓	✓
Renal failure	-		✓	✓
<b>Hematologic</b>				
Agranulocytosis	-		✓	✓
Aplastic anemia	-		-	✓
Eosinophilia	-		✓	✓
Hemolytic anemia	-		-	✓
Leukopenia	-		-	✓
Thrombocytopenia	-		-	✓
<b>Laboratory Test Abnormalities</b>				
Blood urea nitrogen increased	<1		✓	✓
Creatinine increased	6		-	-
Hypercholesterolemia	≤1		-	-
Hyperglycemia	-		-	✓
Hyperkalemia	≤32	18.3	≤40	≤40
Hypertriglyceridemia	<15		-	-
Hyponatremia	2	1.4	✓	✓
Hyperuricemia	<1		-	✓
Liver function tests increased	<1		-	-
<b>Respiratory</b>				
Cough	2		-	-
Respiratory distress	-		-	✓
<b>Other</b>				
Anaphylaxis	-		✓	✓
Angioneurotic edema	<1		-	-
Blurred vision	-		-	✓
Flu-like syndrome	2		-	-

Adverse Events	Single Entity Agents			Combination Products
	Eplerenone	Finerenone	Spironolactone	Spironolactone and HCTZ
Hepatocellular toxicity	-	-	✓	✓
Jaundice	-	-	-	✓
Muscle cramps	-	-	-	✓
Vasculitis	-	-	✓	✓
Weakness	-	-	-	✓
Xanthopsia	-	-	-	✓

✓ Percent not specified  
-Event not reported

## VII. Dosing and Administration

The usual dosing regimens for the mineralocorticoid (aldosterone) receptor antagonists are listed in Table 7.

**Table 7. Usual Dosing Regimens for the Mineralocorticoid (Aldosterone) Receptor Antagonists<sup>1-7</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability	
<b>Single Entity Agents</b>				
Eplerenone	<p><u>Heart Failure:</u> Tablet: initial, 25 mg once daily for four weeks; maintenance, 50 mg once daily</p> <p><u>Hypertension:</u> Tablet: initial, 50 mg once daily; maximum, 50 mg twice daily</p>	Safety and efficacy in children have not been established.	Tablet: 25 mg 50 mg	
Finerenone	<p><u>Chronic kidney disease associated with type 2 diabetes:</u> Tablet: initial, 10 to 20 mg once daily based on estimated glomerular filtration rate; target dose is 20 mg once daily</p>	Safety and efficacy in children have not been established.	Tablet: 10 mg 20 mg	
Spironolactone	<p><u>Edema (congestive heart failure, hepatic cirrhosis, nephrotic syndrome):</u> Tablet: initial, 100 mg once daily in a single or divided dose(s); maintenance, 25 to 200 mg once daily</p> <p><u>Edema caused by cirrhosis:</u> Suspension: initiate therapy in a hospital setting and titrate slowly; initial, 75 mg (15 mL) per day in single or divided doses; in patients requiring titration above 100 mg, use another formulation</p> <p><u>Heart failure (Severe NYHA function class III to IV)</u> Suspension: initial, 20 mg (4 mL) once daily; maintenance, 37.5 mg (7.5 mL) once daily</p> <p>Tablet: initial, 25 mg once daily; maintenance, 25 mg every other day to 50 mg once daily</p> <p><u>Hypertension</u> Suspension: 20 mg (4 mL) to 75 mg (15 mL) per day in single or divided doses</p> <p>Tablet: initial, 50 to 100 mg once daily in a single or divided dose(s); maintenance, 25 to 200 mg once daily;</p>	Safety and efficacy in children have not been established.	Suspension: 25 mg/5 mL	Tablet: 25 mg 50 mg 100 mg

Mineralocorticoid (Aldosterone) Receptor Antagonists  
AHFS Class 243220

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<p>maximum, 400 mg/day</p> <p><u>Hypokalemia:</u> Tablet: 25 to 100 mg once daily</p> <p><u>Primary hyperaldosteronism (diagnosis):</u> Tablet (long test): 400 mg/day for three to four weeks</p> <p>Tablet (short test): 400 mg daily for four days</p> <p><u>Primary hyperaldosteronism (short-term preoperative therapy):</u> Tablet: 100 to 400 mg/day prior to surgery</p> <p><u>Primary hyperaldosteronism (long-term maintenance therapy):</u> Tablet: initial, 100 to 400 mg/day; maximum, 400 mg/day</p>		
<b>Combination Products</b>			
<p>Spironolactone and HCTZ</p>	<p><u>Edema (congestive heart failure, hepatic cirrhosis, nephrotic syndrome):</u> Tablet: maintenance, 100-100 mg/day in a single or divided dose(s); maintenance, 25-25 to 200-200 mg/day</p> <p><u>Hypertension:</u> Tablet: maintenance, 50-50 to 100-100 mg/day in a single or divided dose(s)</p>	<p>Safety and efficacy in children have not been established.</p>	<p>Tablet: 25-25 mg 50-50 mg</p>

HCTZ=hydrochlorothiazide

**VIII. Effectiveness**

Clinical studies evaluating the safety and efficacy of the mineralocorticoid (aldosterone) receptor antagonists are summarized in Table 8.

**Table 8. Comparative Clinical Trials with the Mineralocorticoid (Aldosterone) Receptor Antagonists**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Diabetes/Diabetic Nephropathy/Renal Disease</b>				
<p>Bianchi et al.<sup>31</sup> (2006)</p> <p>Spirolactone 25 mg/day</p> <p>vs</p> <p>placebo</p> <p>All patients were receiving conventional therapy (ACE inhibitor and/or ARB).</p>	<p>OL, PC, PRO, RCT</p> <p>Adult patients with chronic kidney disease</p>	<p>N=165</p> <p>1 year</p>	<p>Primary: Change in proteinuria, eGFR, blood pressure, and serum potassium</p> <p>Secondary: Not reported</p>	<p>Primary: While there was a significant reduction in proteinuria from baseline among spironolactone-treated patients (P&lt;0.001), there was no difference in placebo-treated patients (P&gt;0.05).</p> <p>At one year, there was no significant difference between spironolactone- and placebo-treated patients in eGFR (P value not reported). However, spironolactone-treated patients exhibited a lower monthly rate of decrease in eGFR from baseline compared to conventional therapy-treated patients (P&lt;0.01). Patients whose baseline eGFR was &lt;60 mL/min experienced a greater decline in eGFR compared to patients with baseline eGFR &gt;60 mL/min (P&lt;0.01).</p> <p>At one year of therapy, spironolactone-treated patients experienced a reduction in blood pressure from baseline (P&lt;0.05). In contrast, placebo-treated patients did not exhibit blood pressure reduction from baseline (P value not reported).</p> <p>While there was a significant increase in serum potassium from baseline among spironolactone-treated patients (P&lt;0.001), there was no difference in placebo-treated (P&gt;0.05).</p> <p>Secondary: Not reported</p>
<p>Bianchi et al.<sup>32</sup> (2010)</p> <p>Spirolactone 25 mg, ramipril 10 mg, irbesartan 300 mg, and atorvastatin 10</p>	<p>RCT, OL</p> <p>Patients with a clinical diagnosis of idiopathic chronic glomerulonephritis and urine</p>	<p>N=128</p> <p>36 months</p>	<p>Primary: Changes over time in proteinuria and eGFR</p> <p>Secondary: Adverse events,</p>	<p>Primary: SBP decreased more in the intensive-therapy group (from 156.6 to 113.5 mm Hg) than in the conventional therapy group (from 155.7 to 122.7 mm Hg; P&lt;0.01).</p> <p>Urine protein excretion decreased from 2.65 to 0.45 g/g creatinine with intensive therapy (P&lt;0.001). With conventional therapy, urine protein</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD (intensive therapy)</p> <p>vs</p> <p>ramipril 10 mg and atorvastatin 10 mg QD (conventional therapy)</p> <p>The addition of diuretics, calcium antagonists, <math>\beta</math>-blockers or <math>\alpha</math>1-receptor antagonists were added to achieve blood pressure &lt;130/80 mm Hg</p>	<p>protein-creatinine ratio &gt;1 g/g</p>		<p>drop outs</p>	<p>excretion decreased from 2.60 to 1.23 g/g creatinine (P&lt;0.001).</p> <p>With intensive therapy, eGFR did not significantly change over time (64.6 vs 62.9 mL/min/1.73 m<sup>2</sup>). With conventional therapy, eGFR decreased from 62.5 to 55.8 mL/min/1.73 m<sup>2</sup> (P&lt;0.01).</p> <p>Secondary: In the conventional therapy group, eight patients discontinued the study due to hyperkalemia, cough, and rapid deterioration in kidney function. In the intensive therapy group, 15 dropped out due to hyperkalemia, cough, and hypotension. Nine patients in the intensive therapy group developed gynecomastia. Twelve patients on conventional and 31 on intensive therapy had to interrupt the study temporarily because of low blood pressure. No patient developed an increase in creatine kinase, alanine aminotransferase, and alkaline phosphatase levels during the study.</p>
<p>Ogawa et al.<sup>33</sup> (2006)</p> <p>Spirolactone 25 mg/day plus imidapril* 5 mg/day</p> <p>vs</p> <p>furosemide 20 mg/day plus imidapril* 5 mg/day</p> <p>All patients were pre-treated with imidapril* for 1 year prior to trial onset.</p>	<p>PRO, RCT</p> <p>Adult patients with HTN and type 2 diabetes, with a urine albumin/creatinine ratio &gt;30 mg/g creatinine, and plasma BNP levels &gt;100 pg/mL (suggestive of mild heart failure)</p>	<p>N=30</p> <p>24 months</p>	<p>Primary: Change in BNP, urine albumin/creatinine ratio and blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: At 12 months, spironolactone-treated patients exhibited a significant reduction in BNP level from baseline compared to furosemide-treated patients (P&lt;0.05).</p> <p>At 12 months, spironolactone-treated patients exhibited a significant reduction in urine albumin/creatinine ratio from baseline compared to furosemide-treated patients (P&lt;0.05).</p> <p>Both treatments exhibited similar reductions in blood pressure from baseline (P value not reported).</p> <p>No adverse events were reported in this trial.</p> <p>Secondary: Not reported</p>
<p>Chrysostomou et</p>	<p>DB, PC, RCT</p>	<p>N=41</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>al.<sup>34</sup> (2006)</p> <p>Spironolactone 25 mg/day plus irbesartan 150 mg/day and ramipril 5 mg/day</p> <p>vs</p> <p>ramipril 5 mg/day plus spironolactone 25 mg/day and placebo</p> <p>vs</p> <p>ramipril 5 mg/day plus irbesartan 150 mg/day and placebo</p> <p>vs</p> <p>ramipril 5 mg/day plus placebo and placebo</p>	<p>Patients 18 to 75 years of age, with a 24 hour urinary protein excretion &gt;1.5 g/24 hours on ≥2 occasions ≥3 months apart, serum creatinine level ≤200 μmol/L with &lt;20% variability in the preceding 3 months and treatment with an ACE inhibitor ≥6 months</p>	<p>6 months</p>	<p>Change in 24 hour urinary protein excretion at three months</p> <p>Secondary: Change in 24 hour urinary protein excretion at six months, change in blood pressure and creatinine clearance, adverse effects</p>	<p>Compared to ramipril-treated patients, the 24 hour urinary protein excretion reduction at three months was significantly greater in ramipril plus spironolactone-treated patients (P=0.004).</p> <p>Ramipril-, irbesartan- and spironolactone-treated patients exhibited a significant reduction in 24 hour urinary protein excretion compared to ramipril-treated patients (P&lt;0.001).</p> <p>There was no significant difference in 24 hour urinary protein excretion with ramipril- and ramipril plus irbesartan-treated patients (P=1.00).</p> <p>At three months, spironolactone-treated patients exhibited a significant reduction in proteinuria from baseline (P≤0.001). In contrast, non-spironolactone-treated patients did not experience a significant reduction in proteinuria from baseline (P=0.840).</p> <p>Secondary: At six months, spironolactone-treated patients exhibited the greatest reduction in proteinuria compared to the other treatments (P&lt;0.05).</p> <p>At six months, DBP was higher among ramipril monotherapy-treated patients compared to the other treatments (P=0.046). There was no difference in SBP among the treatments (P value not reported).</p> <p>There were no differences in creatinine clearance among the treatments (P&gt;0.05).</p> <p>Gynecomastia was not observed with any of the treatments.</p>
<p>Furumatsu et al.<sup>35</sup> (2008)</p> <p>Spironolactone 25 mg/day (triple blockade group)</p> <p>vs</p>	<p>MC, OL, PRO, RCT</p> <p>Patients 20 to 70 years of age, with controlled blood pressure &lt;130/80 mm Hg, chronic nephropathy (defined by serum</p>	<p>N=32</p> <p>12 months</p>	<p>Primary: Reduction in proteinuria, urinary type IV collagen, SBP, DBP, mean blood pressure, creatinine, creatinine clearance,</p>	<p>Primary: At one year of therapy, patients randomized to the triple blockage group experienced a statistically significant 58% reduction in urinary protein level from baseline (P&lt;0.05), while there was no difference in the control group. Compared to the control group, the triple blockade group experienced a significant reduction in proteinuria at one year of therapy (P&lt;0.05).</p> <p>At one year of therapy, patients randomized to the triple blockage group</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>trichlormethiazide* 1 mg/day or furosemide 10 mg/day (control group)</p> <p>Study medications were added to ongoing therapy consisting of enalapril 5 mg/day and losartan 50 mg/day.</p>	<p>creatinine level &lt;3 mg/dL or calculated creatinine concentration &lt;30 mL/min), daily treatment with enalapril 5 mg and losartan 50 mg for at least 12 weeks, and persistent proteinuria (urinary protein excretion &gt;0.5 g/day)</p>		<p>potassium, urinary aldosterone</p> <p>Secondary: Not reported</p>	<p>experienced a statistically significant 40% reduction in urinary type IV collagen from baseline (P&lt;0.05); while there was no difference in the control group. However there was no statistically significant difference in the change of urinary type IV collagen from baseline between the two study groups.</p> <p>There were no statistically significant differences between the two study groups in the following outcome measures: SBP, DBP, mean blood pressure, creatinine, creatinine clearance, potassium, and urinary aldosterone.</p> <p>Secondary: Not reported</p>
<p>van den Meiracker et al.<sup>36</sup> (2006)</p> <p>Spironolactone 25 mg BID</p> <p>vs</p> <p>placebo</p> <p>All patients were also receiving their ongoing antihypertensive therapy.</p>	<p>DB, PC, PG, RCT</p> <p>Adult patients with type 2 diabetes, macroalbuminuria (24 hour urinary albumin excretion &gt;300 mg or urinary albumin to creatinine ratio &gt;20 mg/mmol) despite use of an ACE inhibitor or ARB in recommended dosages for ≥1 year</p>	<p>N=59</p> <p>1 year</p>	<p>Primary: Change in albuminuria, DBP and SBP, GFR, aldosterone level, plasma renin activity and serum potassium</p> <p>Secondary: Not reported</p>	<p>Primary: Compared to placebo-treated patients, spironolactone-treated patients exhibited a significant 40.6% reduction in albuminuria from baseline (P=0.002).</p> <p>Compared to placebo, spironolactone-treated patients exhibited a significant reduction in SBP from baseline (P=0.04), with a comparable reduction in DBP (P value not reported).</p> <p>Both treatments exhibited comparable changes in GFR from baseline (P value not reported).</p> <p>Compared to placebo, spironolactone-treated patients exhibited a significant increase in aldosterone level and plasma renin activity from baseline (P&lt;0.05).</p> <p>There was a significant increase in serum potassium level in spironolactone-treated patients compared to placebo (P=0.02).</p> <p>Secondary: Not reported</p>
<p>Schjoedt et al.<sup>37</sup> (2006)</p>	<p>DB, RCT, XO</p> <p>Patients with</p>	<p>N=20</p> <p>2 weeks</p>	<p>Primary: Change in proteinuria,</p>	<p>Primary: Compared to placebo, spironolactone therapy was associated with a significant 32% reduction in proteinuria from baseline (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Spirolactone 25 mg/day</p> <p>vs</p> <p>placebo</p> <p>Study medications were added to ongoing antihypertensive therapy (ACE inhibitor or ARB).</p>	<p>diabetic nephropathy and nephrotic range albuminuria (&gt;2,500 mg/24 hour) despite recommended antihypertensive treatment</p>		<p>ambulatory DBP and SBP, GFR, fractional albumin clearance, aldosterone level, plasma renin activity, and serum potassium</p> <p>Secondary: Not reported</p>	<p>Compared to placebo, spironolactone therapy was associated with a significant reduction in systolic and diastolic ambulatory 24-hr blood pressures from baseline (P=0.004, P=0.001, respectively).</p> <p>Both groups exhibited comparable changes in GFR from baseline (P=0.13).</p> <p>Compared to placebo, spironolactone therapy was associated with a significant 31% reduction in fractional albumin clearance from baseline (P&lt;0.001).</p> <p>Compared to placebo, spironolactone therapy was associated with significant increases in aldosterone level and plasma renin activity from baseline, 80 and 91%, respectively (P&lt;0.005).</p> <p>There was a trend towards an increase in the serum potassium level with spironolactone therapy compared to placebo (P=0.054).</p> <p>Secondary: Not reported</p>
<p>Davidson et al.<sup>38</sup> (2008)</p> <p>Spirolactone 25 mg added to existing ACE inhibitor therapy</p>	<p>OL</p> <p>Patients ≥18 years of age with type 2 diabetes on an ACE inhibitor for &gt;1 month with a urinary albumin to creatinine ratio &gt;100 mg/g</p>	<p>N=24</p> <p>12 weeks</p>	<p>Primary: Change in urinary albumin excretion</p> <p>Secondary: Changes in serum creatinine, serum potassium, and SBP</p>	<p>Primary: Urinary albumin excretion decreased 25.7% from a 404.6 mg/day to 302.7 mg/day (P&lt;0.001). Urinary albumin excretion decreased 27.2% in the microalbuminuria group (P=0.05) and 24.3% in the macroalbuminuria group (P=0.02).</p> <p>Secondary: There were no significant changes in serum sodium, potassium, creatinine, or glucose.</p> <p>There was a significant decrease in SBP with the addition of spironolactone (141.2 to 132.5 mm Hg; P=0.002).</p>
<p>Saklayen et al.<sup>39</sup> (2008)</p> <p>Spirolactone 25</p>	<p>DB, PC, RCT, XO</p> <p>Patients with diabetic nephropathy</p>	<p>N=30</p> <p>7 months</p>	<p>Primary: Blood pressure, serum creatinine, and spot urine</p>	<p>Primary: With spironolactone, the mean SBP at the beginning of the treatment period was 153.64 mm Hg and 141.60 at the end (P=0.01). DBP was 79.56 mm Hg at baseline and 76.68 at study endpoint (P=0.25). The mean SBP</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
to 50 mg/day  vs  placebo  Study medications were added to existing ACE inhibitor or ARB therapy.	with any level of proteinuria who were already being treated with ACE inhibitor (lisinopril) or ARB (losartan) at moderate to maximum dose		protein/creatinine  Secondary: Not reported	with placebo was 154.52 mm Hg at the beginning of the treatment period and 148.82 mm Hg at the end of the study period (P=0.34). DBP was 79.74 mm Hg at baseline and 77.91 at study endpoint (P=0.49).  The urine protein/creatinine increased from 1.24 to 1.57 (24%) with placebo (P=0.35) and decreased from 1.80 to 0.79 (57%) with spironolactone (P=0.004).  Serum creatinine increased from 1.43 to 1.50 on placebo (P=0.19) and from 1.35 to 1.56 on spironolactone (P=0.006).  Secondary: Not reported
Sengul et al. <sup>40</sup> (2009)  Spironolactone 25 mg QD  Study medication was added to existing ACE inhibitor or ARB therapy.	PRO  Patients with overt proteinuria (>300 mg/day) despite the regular use of ACE inhibitors and/or ARBs for ≥6 months	N=33  8 weeks	Primary: Proteinuria, blood pressure  Secondary: Not reported	Primary: At week four, there was a 25.4% reduction in proteinuria with spironolactone (P=0.003). SBP and DBP were significantly reduced (P=0.013 and P=0.040, respectively). Serum potassium level increased 0.28 mEq/L (P<0.001).  At week eight, the 24-hr median urinary protein excretion decreased from 1,428 to 743 mg/day (47.9%) with spironolactone. SBP and DBP were significantly reduced (P<0.004 and P<0.001, respectively). Serum potassium level increased 0.55 mEq/L (P<0.001). There was no difference in creatinine clearance or serum creatinine levels. Serum albumin increased from 3.88 to 4.01 g/dL (P=0.003).  Secondary: Not reported
Tylicki et al. <sup>41</sup> (2008)  Spironolactone 25 mg QD plus background therapy for 8 weeks (triple RAAS blockade)	OL, RCT, XO  Patients with chronic nondiabetic proteinuric kidney diseases	N=18  16 weeks	Primary: 24-hr urine excretion of protein, blood pressure, serum creatinine, serum potassium, plasma renin activity	Primary: A total of 17 patients achieved blood pressure goal of <130/80 mm Hg. There was no difference in ambulatory SBP and DBP between the treatments (P=0.9 and P=0.1).  Serum creatinine and eGFR remained stable during the study periods (P=0.6 and P=0.9, respectively).  A significant increase in plasma renin activity was observed after

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>background therapy for 8 weeks (double RAAS blockade)</p> <p>Background therapy included HCTZ, telmisartan, cilazapril (ACE inhibitor).</p>			<p>Secondary: Not reported</p>	<p>treatment with triple RAAS blockade compared to double RAAS blockade (P=0.02).</p> <p>Triple RAAS therapy provided an additional 55.37% decrease in proteinuria compared to double RAAS blockade (P=0.01). The decrease in proteinuria was shown in 16 of 18 patients. Changes in proteinuria did not correlate with changes in SBP, DBP, or plasma renin activity.</p> <p>There was a significant increase in potassium levels after triple RAAS blockade compared to baseline (P=0.02).</p> <p>Secondary: Not reported</p>
<b>Heart Failure</b>				
<p>Pitt et al.<sup>42</sup> (2003) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE inhibitors, ARBs, diuretics, <math>\beta</math>-blockers, coronary reperfusion therapy)</p>	<p>DB, MC, RCT</p> <p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq</math>40%) and heart failure (patients with diabetes were not required to have heart failure)</p>	<p>N=6,632</p> <p>16 months (mean follow-up)</p>	<p>Primary: Death from any cause, composite of death from cardiovascular causes or hospitalization for a cardiovascular event (heart failure, recurrent acute MI, stroke or ventricular arrhythmia)</p> <p>Secondary: Death from any cause or any hospitalization, death from cardiovascular causes, any hospitalization, hospitalization for</p>	<p>Primary: Significantly fewer eplerenone-treated patients died from any cause compared to placebo-treated patients (478 vs 554; RR, 0.85; 95% CI, 0.75 to 0.96; P=0.008).</p> <p>Significantly fewer eplerenone-treated patients died from or required hospitalization for cardiovascular events compared to placebo-treated patients (885 vs 993; RR, 0.87; 95% CI, 0.79 to 0.95; P=0.002).</p> <p>Secondary: Significantly fewer eplerenone-treated patients died from any cause or required hospitalization (1,730 vs 1,829; RR, 0.92; 95% CI, 0.86 to 0.98; P=0.02).</p> <p>Death from cardiovascular causes was 12.3 and 14.6% in eplerenone- and placebo-treated patients (RR, 0.83; 95% CI, 0.72 to 0.94; P=0.005).</p> <p>Fewer eplerenone-treated patients required hospitalization (1,493 vs 1,526; RR, 0.95; 95% CI, 0.89 to 1.02; P=0.2); however, the difference was not significant.</p> <p>Fewer eplerenone-treated patients required hospitalization due to a cardiovascular event (606 vs 649; RR, 0.91; 95% CI, 0.81 to 1.01;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			cardiovascular causes, hospitalization for heart failure, adverse events	<p>P=0.09); however, the difference was not significant.</p> <p>There was a RR of 15% in the risk of hospitalization for heart failure in the eplerenone-treated patients (RR, 0.85; P=0.03) and 23% fewer episodes of hospitalization for heart failure were reported in these patients (RR, 0.77; P=0.002).</p> <p>Serious hyperkalemia (serum potassium <math>\geq 6.0</math> mmol/L) occurred in 5.5 and 3.9% of eplerenone- and placebo-treated patients (P=0.002). The incidence of hyperkalemia was higher among patients with a lower baseline creatinine clearance (P&lt;0.001).</p> <p>At one year, the serum creatinine concentration had increased by 0.02 and 0.06 mg/dL in placebo- and eplerenone-treated patients (P&lt;0.001).</p> <p>There were no significant differences between eplerenone- and placebo-treated patients in the incidence of sex hormone-related adverse events, including gynecomastia, impotence, breast pain and abnormal vaginal bleeding (P&gt;0.05).</p>
<p>Pitt et al.<sup>43</sup> (2005) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE</p>	<p>Subanalysis of EPHESUS</p> <p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq 40\%</math>) and heart failure (patients with diabetes were not required to have heart failure)</p>	<p>N=6,632</p> <p>30 days post randomization</p>	<p>Primary: Death from any cause, composite of death from cardiovascular causes or hospitalization for a cardiovascular event at 30 days</p> <p>Secondary: Death from cardiovascular causes, sudden cardiac death, fatal or nonfatal heart failure hospitalization,</p>	<p>Primary: A significantly lower percentage of eplerenone-treated patients died from any cause (3.2 vs 4.6%; P=0.004).</p> <p>A lower percentage of eplerenone-treated patients died from or required hospitalization for cardiovascular events (8.6 vs 9.9%; P=0.074); however, the difference was not significant.</p> <p>Secondary: A significantly lower percentage of eplerenone-treated patients died from cardiovascular cause (3.0 vs 4.4%; P=0.003).</p> <p>A lower incidence of sudden cardiac death was noted among eplerenone-treated patients (0.9 vs 1.4%; P=0.051); however, the difference was not significant.</p> <p>A lower percentage of eplerenone-treated patients required hospitalization for fatal/nonfatal heart failure (3.4 vs 4.2%; P=0.106); however, the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
inhibitors, ARBs, diuretics, $\beta$ -blockers, coronary reperfusion therapy)			adverse events	<p>difference was not significant.</p> <p>There was no significant difference between the two treatments in the number of patients experiencing at least one adverse event during 30 days of therapy (P=0.29).</p> <p>At 30 days, the serum potassium concentration had increased by 0.17 and by 0.24 mmol/L in placebo- and eplerenone-treated patients (P&lt;0.001).</p>
<p>Pitt et al.<sup>44</sup> (2006) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE inhibitors, ARBs, diuretics, <math>\beta</math>-blockers, coronary reperfusion therapy)</p>	<p>Subanalysis of EPHESUS evaluating effects of eplerenone in patients with LVEF <math>\leq</math>30%</p> <p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq</math>40%) and heart failure (patients with diabetes were not required to have heart failure)</p>	<p>N=2,106</p> <p>16 months</p>	<p>Primary: Death from any cause, composite of death from cardiovascular causes or hospitalization for a cardiovascular event</p> <p>Secondary: Death from cardiovascular causes, sudden cardiac death, composite of heart failure death and heart failure hospitalizations</p>	<p>Primary: Eplerenone therapy was associated with a significant 21% reduction in the risk of all-cause mortality compared to placebo (P=0.012).</p> <p>Eplerenone therapy was associated with a significant reduction in the risk of the composite endpoint of death from cardiovascular causes or hospitalization for a cardiovascular event compared to placebo (P=0.001).</p> <p>Secondary: Eplerenone therapy was associated with a significant 23% reduction in the risk of cardiovascular mortality compared to placebo (P=0.008).</p> <p>The RR of sudden cardiac death was reduced by 33% (P=0.01) and the heart failure mortality/heart failure hospitalization composite endpoint was reduced by 25% (P=0.005) in eplerenone-treated patients compared to placebo-treated patients.</p> <p>At 30 days, eplerenone therapy was associated with RRRs of 43 (P=0.002), 29 (P=0.006) and 58% (P=0.008) for all-cause mortality, the cardiovascular mortality/cardiovascular hospitalization composite endpoint for sudden cardiac death.</p>
<p>O'Keefe et al.<sup>45</sup> (2007) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by</p>	<p>Subanalysis of EPHESUS evaluating effects of eplerenone in patients with diabetes</p>	<p>N=1,483</p> <p>16 months</p>	<p>Primary: Death from any cause, composite of death from cardiovascular causes or hospitalization for</p>	<p>Primary: Eplerenone therapy was not associated with a significant reduction in the risk of all-cause mortality compared to placebo (P=0.131).</p> <p>Eplerenone therapy in diabetic patients was associated with a significant 17% reduction in the risk of death from cardiovascular causes or hospitalization for a cardiovascular event compared to placebo (P=0.031).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE inhibitors, ARBs, diuretics, <math>\beta</math>-blockers, coronary reperfusion therapy)</p>	<p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq 40\%</math>) and heart failure (patients with diabetes were not required to have heart failure)</p>		<p>a cardiovascular event</p> <p>Secondary: Death from cardiovascular causes, sudden cardiac death, hyperkalemia</p>	<p>Secondary: Eplerenone therapy was not associated with a significant reduction in the risk of cardiovascular mortality compared to placebo (P=0.128).</p> <p>Eplerenone therapy was not associated with a significant reduction in the risk of sudden cardiac death compared to placebo (P=0.533).</p> <p>Eplerenone therapy was associated with a greater incidence of hyperkalemia compared to placebo (5.6 vs 3.0%; P=0.015).</p>
<p>Gheorghide et al.<sup>46</sup> (2009) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE inhibitors, ARBs, diuretics, <math>\beta</math>-blockers, coronary reperfusion therapy)</p>	<p>Subanalysis of EPHESUS evaluating effects of eplerenone on length of stay and total days of heart failure hospitalization</p> <p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq 40\%</math>) and heart failure (patients with diabetes were not required to have heart failure)</p>	<p>N=828</p> <p>16 months</p>	<p>Primary: Mean length of stay/episode of heart failure hospitalization, total number of days of heart failure hospitalizations following the index hospitalization during the subsequent follow up period</p> <p>Secondary: Determine the difference between the five regions in the mean length of stay and the total number of days for</p>	<p>Primary: Over a mean follow up of 16 months, eplerenone therapy was associated with a significant reduction in the mean length of hospital stay/episode of heart failure hospitalization of 1.6 days (9.2 vs 10.8 days; P=0.019).</p> <p>Eplerenone-treated patients achieved a reduction in the total number of days of heart failure hospitalization/patient of 3.6 days (13.3 vs 16.9 days; P=0.0006).</p> <p>Secondary: The length of stay/heart failure hospitalization episode and total number of days of heart failure hospitalization/patient were consistently and similarly reduced in eplerenone-treated patients in all geographic regions as demonstrated by the nonsignificant interaction of study region on treatment effect (P=0.63 for length of stay/episode and P=0.45 for total hospitalization days for heart failure, respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			heart failure hospitalization	
<p>Adamopoulos et al.<sup>47</sup> (2010) EPHESUS</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by titration to 50 mg/day</p> <p>vs</p> <p>placebo</p> <p>Patients were allowed to receive optimal medical therapy (ACE inhibitors, ARBs, diuretics, <math>\beta</math>-blockers, coronary reperfusion therapy)</p>	<p>Subanalysis of EPHESUS evaluating the differential effects of time-to-eplerenone initiation vs placebo</p> <p>Patients with acute MI, left ventricular dysfunction (ejection fraction <math>\leq 40\%</math>) and heart failure (patients with diabetes were not required to have heart failure)</p>	<p>N=6,632</p> <p>16 months</p>	<p>Primary: Death from any cause, composite of death from cardiovascular causes or hospitalization for a cardiovascular event (heart failure, recurrent acute MI, stroke or ventricular arrhythmia)</p> <p>Secondary: Sudden cardiac death</p>	<p>Primary: “Earlier” eplerenone-treated patients had significantly lower event rates when compared to “earlier” placebo-treated patients for all-cause mortality (11.5 vs 16.1%) and the composite of cardiovascular hospitalization or death (24.0 vs 30.3%). No significant differences were found between “later” eplerenone- and placebo-treated patients.</p> <p>“Earlier” eplerenone therapy significantly reduced the risk for all-cause mortality (HR, 0.72; 95% CI, 0.58 to 0.89; P=0.002) and cardiovascular hospitalization or death (HR, 0.78; 95% CI, 0.67 to 0.90; P=0.001).</p> <p>Secondary: “Earlier” eplerenone-treated patients had significantly lower event rates when compared to “earlier” placebo-treated patients for sudden cardiac death (3.7 vs 6.9%). No significant differences were found between “later” eplerenone- and placebo-treated patients.</p> <p>“Earlier” eplerenone therapy significantly reduced the risk for sudden cardiac death (HR, 0.54; 95% CI, 0.38 to 0.77; P=0.001).</p> <p>In a head-to-head comparison between the two eplerenone treatment groups, “earlier” therapy was associated with significantly lower risk with respect to all endpoints. No significant difference was found in a direct comparison between the two placebo treatment groups.</p>
<p>Udelson et al.<sup>48</sup> (2010)</p> <p>Eplerenone 25 mg/day for 4 weeks, followed by 50 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients <math>\geq 21</math> years of age with current symptoms consistent of mild to moderate heart failure (NYHA Class II and III) who had LVEF <math>\leq 35\%</math> and were on therapy</p>	<p>N=226</p> <p>36 weeks</p>	<p>Primary: Change in left ventricular end-diastolic volume index</p> <p>Secondary: Changes in left ventricular end-systolic volume index and LVEF,</p>	<p>Primary: Over 36 weeks, there was no evidence of an effect of eplerenone therapy on left ventricular end-diastolic volume index compared to placebo (P value not reported).</p> <p>Secondary: Over 36 weeks, there was no evidence of an effect of eplerenone therapy on left ventricular end-systolic volume index compared to placebo (P value not reported).</p> <p>Over 36 weeks, there was no evidence of an effect of eplerenone therapy</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	with an ACE inhibitor and/or ARB and $\beta$ -blocker for $\geq 3$ months and at a dose that has not been adjusted within the previous 4 weeks		markers of collagen turnover	on LVEF compared to placebo (P value not reported).  During the course of treatment, eplerenone-treated patients exhibited a greater reduction in PINP and BNP compared to placebo-treated patients (P=0.01 and P=0.04, respectively). No difference between the two treatments was observed in the change from baseline to week 36 in PIIINP (P value not reported).
<p>Zannad et al.<sup>49</sup> (2011) EMPHASIS-HF</p> <p>Eplerenone 25 mg QD for 4 weeks, followed by 50 mg QD</p> <p>vs</p> <p>placebo</p> <p>Randomization occurred within 6 months after hospitalization for a cardiovascular reason.</p> <p>Patients who had not been hospitalized for a cardiovascular reason within 6 months of the screening visit</p>	<p>DB, RCT</p> <p>Patients <math>\geq 55</math> years of age with NYHA Class II symptoms, and ejection fraction <math>\leq 30\%</math> and treatment with an ACE inhibitor, ARB or both and a <math>\beta</math>-blocker at the recommended dose or maximal tolerated dose</p>	<p>N=2,737</p> <p>21 months (median follow up)</p>	<p>Primary: Composite of death from cardiovascular causes or a first hospitalization for heart failure</p> <p>Secondary: Hospitalization for heart failure or death from any cause, death from any cause, death from cardiovascular causes, hospitalization for any reason, hospitalization for heart failure</p>	<p>Primary: The primary composite endpoint occurred in 18.3 and 25.9% of eplerenone- and placebo-treated patients (HR, 0.63; 95% CI, 0.54 to 0.74; P&lt;0.001).</p> <p>Secondary: Death from any cause or hospitalization for heart failure occurred in 19.8 and 27.4% of eplerenone- and placebo-treated patients (HR, 0.65; 95% CI, 0.55 to 0.76; P&lt;0.001).</p> <p>A total of 12.5 and 15.5% of eplerenone- and placebo-treated patients died (HR, 0.76; 95% CI, 0.62 to 0.93; P=0.008).</p> <p>A total of 10.8 and 13.5% of deaths were attributed to cardiovascular causes in eplerenone- and placebo-treated patients (HR, 0.76; 95% CI, 0.61 to 0.94; P=0.01).</p> <p>A total of 29.9 and 35.8% of eplerenone- and placebo-treated patients were hospitalized for any reason (HR, 0.77; 95% CI, 0.67 to 0.88; P&lt;0.001).</p> <p>Of the hospitalized patients, 12.0 vs 18.4% of eplerenone- and placebo-treated patients were hospitalized for heart failure (HR, 0.58; 95% CI, 0.47 to 0.70; P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
could be enrolled if the plasma BNP was $\geq 250$ pg/mL or if the plasma N-terminal pro-BNP was $\geq 500$ pg/mL in men and $\geq 750$ pg/mL in women.				
<p>Eschalier et al.<sup>50</sup> (2013) EMPHASIS-HF</p> <p>Eplerenone 25 mg QD for 4 weeks, followed by 50 mg QD</p> <p>vs</p> <p>placebo</p>	<p>Subgroup analysis of EMPHASIS-HF</p> <p>Patients included in the EMPHASIS-HF trial aged <math>\geq 75</math> years with diabetes, CKD, and SBP &lt;median (123 mm Hg)</p>	<p>N=2,737</p> <p>21 months (median follow up)</p>	<p>Primary: Hospitalization for HF or death from cardiovascular causes</p> <p>Secondary: Serum potassium, hyperkalemia leading to study drug discontinuation, hospitalization for hyperkalemia and hospitalization for worsening renal function (WRF), change in eGFR</p>	<p>Primary: Eplerenone was effective at reducing the risk of cardiovascular death or HF hospitalization in the high-risk subgroups, which is consistent with result in the overall EMPHASIS-HF study population (HR, 0.63; 95% CI, 0.54 to 0.74; <math>P &lt; 0.001</math>). The HR for the primary outcome in the eplerenone group as compared with the placebo group was 0.66 (95% CI, 0.49 to 0.88; <math>P = 0.005</math>) in patients <math>\geq 75</math> years of age, 0.54 (95% CI, 0.42 to 0.70; <math>P &lt; 0.0001</math>) in patients with diabetes, 0.62 (95% CI, 0.49 to 0.79; <math>P = 0.0001</math>) in patients with CKD, and 0.63 (95% CI, 0.51 to 0.79; <math>P &lt; 0.0001</math>) in patients with SBP &lt; median.</p> <p>Secondary: The number of patients with study drug stopped due to adverse events was evenly distributed within and among the study high-risk subgroups in patients age <math>\geq 75</math> years (18.2% in eplerenone vs 19.0% in placebo), in patients with SBP &lt;123 mm Hg (16.6% in eplerenone vs 18.0% in placebo), in patients with CKD (16.1% in eplerenone vs 22.3% in placebo), and in patients with diabetes mellitus (15.1% in eplerenone vs 18.1% in placebo). In patients with CKD (eGFR &lt;60 ml/min/1.73 m<sup>2</sup>), there were fewer patients in eplerenone group who had their treatment stopped due to an adverse event or due to any other reason than in placebo group.</p>
<p>Krum et al.<sup>51</sup> (2013) EMPHASIS-HF</p> <p>Eplerenone 25 mg QD for 4 weeks, followed by 50 mg</p>	<p>Subgroup analysis of EMPHASIS-HF</p> <p>Patients included in the EMPHASIS-HF trial analyzed according to the use</p>	<p>N=2,737</p> <p>21 months (median follow up)</p>	<p>Primary: Hospitalization for HF or death from cardiovascular causes</p> <p>Secondary:</p>	<p>Primary: The beneficial clinical effects of eplerenone (as observed in the main study) were preserved for the EMPHASIS-HF primary end point in patients receiving higher doses of ACE Inhibitor or ARB, <math>\beta</math>-blocker, or both. P values for interaction between high and low doses for the EMPHASIS-HF primary end point were not significant.</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
QD vs placebo	and dose of ACE inhibitors/ARBs, $\beta$ -blockers, or both		Not reported	Secondary: Not reported
Girerd et al. <sup>52</sup> (2015) EMPHASIS-HF  Eplerenone 25 mg QD for 4 weeks, followed by 50 mg QD  vs placebo	Subgroup analysis of EMPHASIS-HF  Patients included in the EMPHASIS-HF trial analyzed according to whether the treatment was initiated <42 or 42+ days after qualifying CV hospitalization	N=2,338  21 months (median follow up)	Primary: Composite of cardiovascular mortality or hospitalization for HF  Secondary: Adverse effects	Primary: The relative rate reductions in CV death/hospitalization for HF, hospitalization for HF, and all-cause mortality were similar (P for interaction=0.65, 0.44, and 0.40, respectively) whether the treatment was initiated <42 or 42+ days after qualifying CV hospitalization. Absolute rate reductions were -5.61 (95% CI, -8.67 to -2.55) events per 100 patient-years in the <42 days group and -3.58 (CI, -6.37 to -0.79) in the 42+ days group. Regardless of the event considered, cumulative incidences of events in patients treated with eplerenone were lower than the rates observed in patients allocated to placebo in both the <42 and the 42+ days groups.  Secondary: The adverse effects of eplerenone were also unaffected by the time from the qualifying CV hospitalization.
Pitt et al. <sup>53</sup> (1999) RALES  Spironolactone 25 mg/day; in the absence of hyperkalemia, the dose could be increased to 50 mg/day after 8 weeks; if hyperkalemia developed the dose could be decreased to 25 mg every other day	DB, MC, RCT  Patients with NYHA class 4 heart failure within 6 months and with NYHA class 3 to 4 at study onset, diagnosed with CHF $\geq$ 6 weeks, treated with an ACE inhibitor and a loop diuretic, with a LVEF $\leq$ 35%	N=1,663  24 months (mean follow-up)	Primary: Death from any cause  Secondary: Death from cardiac causes, hospitalization for cardiac causes, combined incidence of death or hospitalization for cardiac causes, combined end point of death or hospitalizations from any cause,	Primary: There were 386 and 284 deaths from any cause in placebo- and spironolactone-treated patients (RR, 0.70; 95% CI, 0.60 to 0.82; P<0.001).  Secondary: There were 314 and 226 deaths in placebo- and spironolactone-treated patients that were attributed to cardiac causes (RR, 0.69; 95% CI, 0.58 to 0.82; P<0.001).  There were 753 and 515 hospitalizations for cardiac causes in placebo- and spironolactone-treated patients (RR, 0.70; 95% CI, 0.59 to 0.82; P<0.001).  The combined end point of death from cardiac causes or hospitalizations from cardiac causes showed a 32% reduction in risk among spironolactone-treated patients compared to placebo-treated patients (RR, 0.68; 95% CI, 0.59 to 0.78; P<0.001).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			combined end point of death from any cause or hospitalizations from cardiac causes, change in the NYHA class, adverse events	<p>The combined end point of death or hospitalizations from any cause showed a 23% reduction in risk among spironolactone-treated patients compared to placebo-treated patients (RR, 0.77; 95% CI, 0.68 to 0.86; P&lt;0.001).</p> <p>The combined end point of death from any cause or hospitalizations from cardiac causes showed a 32% reduction in risk among spironolactone-treated patients as compared to placebo-treated patients (RR, 0.68; 95% CI, 0.60 to 0.77; P&lt;0.001).</p> <p>A significantly greater percentage of spironolactone-treated patients experienced improvement in the NYHA class compared to placebo-treated patients (41 vs 33%; P&lt;0.001).</p> <p>Gynecomastia or breast pain was reported in 10 and 1% of spironolactone- and placebo-treated men (P&lt;0.001). The incidence of hyperkalemia was minimal with both treatments.</p>
Vardeny et al. <sup>54</sup> (2012) RALES  Spironolactone 25 or 50 mg/day  vs placebo	Post-hoc analysis  Patients with NYHA class III or IV heart failure with an ejection fraction <35%	N=1,658  24 months (mean follow-up)	Primary: Death from any cause  Secondary: Death from cardiac causes, hospitalization for cardiac causes, combined incidence of death or hospitalization for cardiac causes, combined end point of death or hospitalizations from any cause, combined end point of death from	<p>Primary: Patients with reduced baseline eGFR exhibited similar RR reductions in all cause mortality and the composite of death or hospital stays for heart failure compared to patients with a baseline eGFR &gt;60 mL/min/1.73 m<sup>2</sup>, and a greater absolute risk reduction compared to patients with a higher baseline eGFR (10.3 vs 6.4%).</p> <p>Worsening renal failure (17 vs 7%; P&lt;0.001) was associated with an increased adjusted risk of death with placebo (HR, 1.9; 95% CI, 1.3 to 2.6) but not with spironolactone (HR, 1.1; 95% CI, 0.79 to 1.5; P=0.009).</p> <p>The risk of hyperkalemia and renal failure was higher in patients with worse baseline renal function and patients with worsening renal failure, particularly with spironolactone.</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			any cause or hospitalizations from cardiac causes, change in the NYHA class, adverse events	
<p>Vizzard et al.<sup>55</sup> (2010)</p> <p>Spirolactone 25 mg QD, followed by up-titration every 2 weeks to 50 or 100 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, RCT</p> <p>Patients with clinical evidence of heart failure, NHYA class 1 to 2 severity of symptoms at the time of enrollment and receiving optimal medical treatment maintained at stable doses for <math>\geq 6</math> months</p>	<p>N=158</p> <p>6 months</p>	<p>Primary: Change in LVEF, left ventricular end-diastolic and -systolic volumes, left ventricular mass and laboratory examinations</p> <p>Secondary: Not reported</p>	<p>Primary: After six months, LVEF increased (<math>P &lt; 0.001</math>) and left ventricular end-diastolic and -systolic volumes decreased (<math>P &lt; 0.001</math> for both) significantly in spironolactone-treated patients compared to placebo-treated patients.</p> <p>After six months, left ventricular mass decreased significantly in spironolactone-treated patients compared to placebo-treated patients (from <math>269 \pm 74</math> to <math>243 \pm 67</math> g vs <math>250 \pm 43</math> to <math>247 \pm 38</math> g; <math>P &lt; 0.05</math>).</p> <p>Serum potassium increased in spironolactone-treated patients from <math>4.2 \pm 0</math> to <math>4.6 \pm 0.3</math> mmol/L (<math>P &lt; 0.001</math>). Serum aldosterone and renin levels increased, respectively, from <math>157.1 \pm 1.03</math> to <math>205 \pm 56.5</math> pg/mL (<math>P = 0.08</math>) and from <math>3.7 \pm 10.5</math> to <math>6.2 \pm 2.8</math> ng/mL/hr (<math>P = 0.03</math>) in these patients. No significant changes were found in serum creatinine, serum urea nitrogen and uric acid.</p> <p>Secondary: Not reported</p>
<p>Chan et al.<sup>56</sup> (2007)</p> <p>Spirolactone 25 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patients received candesartan 8</p>	<p>DB, PC, RCT</p> <p>Patients with LVEF <math>&lt; 40\%</math> on ACE inhibitors for <math>&gt; 6</math> months</p>	<p>N=48</p> <p>1 year</p>	<p>Primary: Change in LVEF, left ventricular end-diastolic volume index, end-systolic volume index, left ventricular mass index, SBP, quality of life</p> <p>Secondary: Not reported</p>	<p>Primary: At one year, combination therapy was associated with a significant improvement in LVEF from baseline (<math>P &lt; 0.01</math>).</p> <p>At one year, combination therapy was associated with a significant reduction in left ventricular end-diastolic volume index from baseline (<math>P &lt; 0.001</math>).</p> <p>At one year, combination therapy was associated with a significant reduction in end-systolic volume index from baseline (<math>P &lt; 0.0005</math>).</p> <p>At one year, combination therapy was associated with a significant reduction in left ventricular mass index from baseline (<math>P = 0.002</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day.				<p>At one year, combination therapy was associated with a significant reduction in SBP from baseline (P&lt;0.05).</p> <p>The control group was not associated with significant improvements in any of the above primary outcome measures.</p> <p>The quality of life score improved in both study groups.</p> <p>Secondary: Not reported</p>
<p>Edelmann et al.<sup>57</sup> (2013) Aldo-DHF</p> <p>Spirolactone 25 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, PRO, RCT</p> <p>Patients with chronic NYHA class II or III heart failure, preserved LVEF ≥50%, and evidence of diastolic dysfunction</p>	<p>N=422</p> <p>12 months</p>	<p>Primary: Change in diastolic function and maximal exercise capacity</p> <p>Secondary: Left ventricular mass index, neuroendocrine activation, symptoms of heart failure, QOL, 6-minute walking distance</p>	<p>Primary: Diastolic function decreased from 12.7±3.6 to 12.1±3.7 with spironolactone and increased from 12.8±4.4 to 13.6±4.3 with placebo (adjusted mean difference, -1.5; 95% CI, -2.0 to -0.9; P&lt;0.001).</p> <p>With regards to exercise capacity, peak VO<sub>2</sub> did not significantly change with spironolactone vs placebo (from 16.3±3.6 to 16.8±4.6 vs from 16.4±3.5 to 16.9±4.4 mL/min/kg, respectively; adjusted mean difference, 0.1 mL/min/kg; 95% CI, -0.6 to 0.8; P=0.81).</p> <p>Secondary: Compared to placebo, treatment with spironolactone induced reverse remodeling (left ventricular mass index declined; adjusted mean difference, -6 g/m<sup>2</sup>; 95% CI, -10 to -1; P=0.009) and improved neuroendocrine activation (N-terminal pro-brain-type natriuretic peptide geometric mean ratio, 0.86; 95% CI, 0.75 to 0.99; P=0.03).</p> <p>Compared to placebo, spironolactone did not improve heart failure symptoms or QOL.</p> <p>Compared to placebo, spironolactone slightly reduced 6-minute walking distance (-15 m; 95% CI, -27 to -2; P=0.03).</p>
<p>Pitt et al.<sup>58</sup> (2014) TOPCAT</p>	<p>DB, MC, RCT</p> <p>Patients ≥50 years of age with at least 1</p>	<p>N=3445</p> <p>3 years</p>	<p>Primary: Composite of death from cardiovascular</p>	<p>Primary: 18.6% of patients in the spironolactone group and 20.4% of patients in the placebo group had at least one confirmed primary-outcome event (P=0.14).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Spirolactone vs placebo</p>	<p>sign and 1 symptom of HF, a LVEF <math>\geq 45\%</math>, controlled BP, and potassium <math>&lt; 5.0</math> mmol/L who were hospitalized in the previous year</p>		<p>causes, aborted cardiac arrest, or hospitalization for the management of heart failure</p> <p>Secondary: Death from any cause, hospitalization for any cause, hyperkalemia (K <math>\geq 5.5</math> mmol/L), hypokalemia (K <math>&lt; 3.5</math> mmol/L), an elevated serum creatinine level (<math>\geq 2</math> times the baseline value and above the upper limit of normal), and a serum creatinine level <math>\geq 3.0</math> mg/deciliter</p>	<p>Secondary: There were no significant differences between study groups in time to death from any cause or first hospitalization for any reason. Frequency of hospitalization for any reason (including recurrent hospitalization) did not differ significantly according to study group (36.8 hospitalizations per 100 person-years in the spironolactone group and 36.3 per 100 person-years in the placebo group; <math>P=0.71</math>).</p> <p>The spironolactone group had a higher rate of hyperkalemia (18.7 vs 9.1% in the placebo group) and a lower rate of hypokalemia (16.2 vs 22.9%). The spironolactone group was more likely to have a doubling of the serum creatinine level to a value above the upper limit of the normal range (10.2 vs 7.0%). However, there were no significant between-group differences in the proportion of patients with a serum creatinine level of 3.0 mg per deciliter or higher or who required dialysis.</p>
<p>Levy et al.<sup>59</sup> (1977)  Spirolactone and HCTZ 25-25 mg/day (fixed-dose combination product) for 16 weeks following 8 weeks of furosemide monotherapy</p>	<p>DB, RCT  Patients 27 to 79 years of age with arteriosclerotic heart disease, hypertensive heart disease, or rheumatic heart disease classes 1 to 3, and congestive heart failure requiring diuretic</p>	<p>N=32  24 weeks</p>	<p>Primary: Change in heart failure symptoms, glucose, renin concentration, calcium, blood urea nitrogen, uric acid, creatinine, aldosterone, serum potassium level, adverse effects</p> <p>Secondary:</p>	<p>Primary: The combination therapy group and furosemide monotherapy group exhibited comparable control of heart failure symptoms.</p> <p>The combination therapy group was associated with a significant decrease in glucose and an increase in plasma renin concentration compared to furosemide monotherapy group (<math>P&lt;0.01</math>).</p> <p>There were no significant differences in calcium, blood urea nitrogen, uric acid, or creatinine between the study groups.</p> <p>There was a significant increase in aldosterone secretion among patients randomized to the spironolactone and HCTZ group compared to the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  furosemide 25 mg daily for 24 weeks	therapy		Not reported	furosemide group (P<0.01).  There was no significant difference in serum potassium level between treatment groups.  No serious adverse effects were observed in either of the study groups.  Secondary: Not reported
Lee et al. <sup>60</sup> (2013)  Patients receiving spironolactone  vs  patients not receiving spironolactone	OBS  Patients with newly diagnosed HF with documented LVEF of <40% who had no aldosterone receptor antagonist use in the 12 months before study entry	N=2358  Median follow-up of 2.5 years	Primary: all-cause mortality, all-cause hospitalization, severe hyperkalemia, and acute kidney injury  Secondary: Not reported	Primary: Incident spironolactone use was associated with lower crude rates of mortality (5.5 vs 9.8 per 100 person-years; P<0.01) and all-cause hospitalization (49.4 vs 56.1 per 100 person-years, P<0.05) compared with nonuse. After adjustment for differences in patient characteristics and concurrent use of other HF therapies, use of spironolactone was not significantly associated with either death (adjusted HR, 0.93; 95% CI, 0.60 to 1.44) or all-cause hospitalization (adjusted HR, 0.91; 95% CI, 0.77 to 1.08).  Crude rates of acute kidney injury were also significantly lower during periods of spironolactone use (7.2 per 100 person-years) compared with periods of nonuse (16.2 per 100 person-years). Conversely, spironolactone use was associated with higher crude rates of severe hyperkalemia (4.8 per 100 person-years) compared with nonuse (1.6 per 100 person-years, P<0.001). After adjustment for potential confounders, incident spironolactone use was associated with a higher adjusted rate of severe hyperkalemia (adjusted HR, 3.46; 95% CI, 1.97 to 6.06) but not with acute kidney injury (adjusted HR, 0.66; 95% CI, 0.42 to 1.05).  Secondary: Not reported
Inampudi et al. <sup>61</sup> (2014)  Spironolactone on discharge	OBS  hospitalized Medicare beneficiaries with HFrEF (EF <45%)	N=1140  1 year post-discharge	Primary: 30-day all-cause readmission  Secondary: 30-day all-cause	Primary: Within 30 days postdischarge, unadjusted all-cause readmissions rates were 30% and 25% for patients receiving and not receiving spironolactone, respectively. Propensity score (PS)-adjusted HR (95% CI) associated with spironolactone use was 1.41 (1.04 to 1.90).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  No spironolactone on discharge	and advanced CKD		mortality, HF readmissions, and combined end point of all-cause mortality or all-cause readmission	Secondary: The risk of all-cause readmission (PS-adjusted HR, 1.36; 95% CI, 1.13 to 1.63) and the combined end point of all-cause readmission or all-cause mortality (PS-adjusted HR, 1.30; 95% CI, 1.09 to 1.54) during 1 year postdischarge were higher among patients in the spironolactone group.
Maisel et al. <sup>62</sup> (2014) COACH  Spironolactone-treated patients  vs  patients not treated with spironolactone	Secondary analysis  Patients enrolled in the COACH biomarker substudy	N=534  30 days	Primary: 30-day mortality and HF-related rehospitalization  Secondary: Biomarker levels (NT-proBNP, ST2, Gal-3, and creatinine)	Primary: Spironolactone significantly reduced the 30-day composite of mortality and HF-related rehospitalization (HR, 0.538; 95% CI, 0.299 to 0.968; P=0.039).  Secondary: Elevated NT-proBNP, creatinine, and ST2 were associated with increased 30-day mortality and HF-related hospitalizations. Spironolactone treatment was significantly beneficial in groups with elevations of Gal-3, ST-2, NT-proBNP, or creatinine (P=0.037, 0.007, 0.035, and 0.009, respectively). In contrast, spironolactone treatment effects were not significant for groups with lower levels of any biomarker.
<b>Hyperaldosteronism</b>				
Karagiannis et al. <sup>63</sup> (2008)  Eplerenone 25 mg BID, titrated up to 200 mg/day if blood pressure remained $\geq$ 140/90 mm Hg  vs  spironolactone 25 mg BID, titrated up to 400 mg/day if blood pressure remained $\geq$ 140/90 mm Hg	OL, PRO, RCT  Patients with bilateral hyperaldosteronism	N=34  16 weeks	Primary: Percentage of patients whose blood pressure <140/90 mm Hg at week 16  Secondary: Adverse events	Primary: At 16 weeks, 76.5 and 82.4% of spironolactone- and eplerenone-treated patients, respectively, exhibited reductions in blood pressure to <140/90 mm Hg (P=1.00).  Secondary: Serum potassium levels were normalized with both treatments after four weeks of therapy (P value not reported). Mild hyperkalemia was noted in two spironolactone 400 mg-treated patients and in three eplerenone 150 mg-treated patients.  Two spironolactone-treated patients reported bilateral gynecomastia at week 16 (P value not reported). Switching from spironolactone 400 mg/day to eplerenone 150 mg/day was effective in resolving gynecomastia symptoms without disrupting blood pressure control.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 12.5 mg was added to the study regimen if blood pressure remained uncontrolled at week 16.</p>				
<p>Karashima et al.<sup>64</sup> (2015)</p> <p>Eplerenone 50 mg, titrated up to 100 mg/day if blood pressure remained <math>\geq 140/90</math> mm Hg</p> <p>vs</p> <p>spironolactone 25 mg, titrated up to 100 mg/day if blood pressure remained <math>\geq 140/90</math> mm Hg</p> <p>Calcium channel blocker was added to the study regimen if blood pressure remained uncontrolled at week eight.</p>	<p>OL</p> <p>Patients <math>\geq 18</math> years of age with HTN and primary aldosteronism</p>	<p>N=54</p> <p>12 months</p>	<p>Primary: Blood pressure</p> <p>Secondary: Metabolic factors</p>	<p>Primary: Treatment with spironolactone or eplerenone significantly decreased systolic BP and diastolic BP from baseline (<math>P &lt; 0.001</math>). The BP-lowering effects between the two agents did not differ.</p> <p>Secondary: Urinary albumin excretion was significantly improved by treatment (<math>P = 0.024</math>). Spironolactone significantly increased plasma aldosterone concentration compared with eplerenone (<math>P = 0.007</math>). Plasma renin activity, serum potassium, eGFR and urinary albumin excretion did not differ between the two groups. The metabolic factors did not significantly differ between the two groups. Body weight, BMI, waist circumference, visceral and subcutaneous adipose tissue area were not different between the two groups. Although BMI and visceral adipose tissue area were significantly decreased in all patients (<math>P &lt; 0.05</math>), no significant differences in BMI, visceral adipose tissue and subcutaneous adipose tissue area were observed between the two groups. Two patients treated with spironolactone experienced gynecomastia. No patients treated with eplerenone showed gynecomastia.</p>
<b>Hypertension</b>				
<p>Kohvakka et al.<sup>65</sup> (1979)</p> <p>Amiloride 5 mg QD</p> <p>vs</p>	<p>PC, RCT, XO</p> <p>Patients 41 to 70 years of age with uncomplicated HTN, previously treated</p>	<p>N=31</p> <p>3 months</p>	<p>Primary: Changes in blood pressure, serum potassium, sodium, creatinine, urate and total body</p>	<p>Primary: No significant changes in blood pressure were observed with any of the treatments (P values not reported).</p> <p>Mean serum potassium was reduced with all treatments except with spironolactone. KCl supplementation was least effective in elevating</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>triamterene 75 mg QD</p> <p>vs</p> <p>KCl 1,500 mg QD</p> <p>vs</p> <p>spironolactone 50 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patients were also receiving HCTZ 50 mg QD.</p>	<p>with antihypertensive agents for 1 to 6 years</p>		<p>potassium</p> <p>Secondary: Not reported</p>	<p>serum potassium. Total body potassium remained constant throughout treatment (P values not reported).</p> <p>Serum sodium remained within normal limits with all treatments (P values not reported).</p> <p>There were no significant changes in mean serum creatinine with any of the treatments (P values not reported).</p> <p>Serum urate concentration increased significantly with all treatments, including HCTZ monotherapy (P values not reported).</p> <p>Secondary: Not reported</p>
<p>Dahlöf et al.<sup>66</sup> (1991) Hypertension (STOP)</p> <p>Atenolol 50 mg QD, HCTZ 25 mg QD plus amiloride 2.5 mg QD, metoprolol 100 mg QD, or pindolol 5 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, RCT</p> <p>Swedish men and women 70 to 84 years old with treated or untreated essential HTN defined as SBP <math>\geq</math>180 mm Hg with a DBP of <math>\geq</math>90 mm Hg, or DBP &gt;105 mm Hg irrespective of the SBP measured on 3 separate occasions during a 1-month placebo run-in phase in previously</p>	<p>N=1,627</p> <p>25 months</p>	<p>Primary: Frequency of stroke, MI, and other cardiovascular death</p> <p>Secondary: Not reported</p>	<p>Primary: The active treatments significantly reduced the number of all primary endpoints (94 vs 58; RR, 0.60; 95% CI, 0.43 to 0.85; P=0.0031), frequency of stroke (53 vs 29; RR, 0.53; 95% CI, 0.33 to 0.86; P=0.0081) and frequency of other cardiovascular deaths (13 vs 4; RR, 0.30; 95% CI, 0.07 to 0.97) compared to placebo.</p> <p>There was not a statistically significant decrease observed in the rate of MI between the active treatments and placebo (28 vs 25; RR, 0.87; 95% CI, 0.49 to 1.56).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	untreated patients			
White et al. <sup>67</sup> (2003)  Eplerenone 25 mg QD  vs  eplerenone 50 mg QD  vs  eplerenone 100 mg QD  vs  eplerenone 200 mg BID  vs  placebo	DB, MC, RCT  Adult patients with untreated HTN and seated SBP <180 mm Hg, DBP between 95 to 110 mm Hg, and the 24 hour mean DBP ≥85 mm Hg	N=400  12 weeks	Primary: Mean change from baseline in seated DBP at 12 weeks  Secondary: Change from baseline in SBP, 24 hour SBP and DBP, heart rate, adverse events	Primary: Eplerenone 50, 100 and 200 mg-treated patients experienced significant mean reductions in DBP from baseline compared to placebo (P≤0.01). The reduction in BP in eplerenone 25 mg-treated patients failed to meet significance (P=0.10).  Secondary: Eplerenone 50, 100 and 200 mg-treated patients experienced significant mean reductions in SBP from baseline compared to placebo (P≤0.01).  All eplerenone-treated patients experienced significant reductions in 24 hour ambulatory blood pressure measurements compared to placebo (P<0.006 for SBP and P<0.005 for DBP).  There were no significant differences from baseline in 24 hour mean heart rate with any eplerenone-treated patient compared to placebo (P value not reported).  Treatment emergent adverse events were reported in 48 and 49% of eplerenone- and placebo-treated patients. None of the adverse events were significantly different between the treatments (P value not reported). Two cases of impotence, gynecomastia, menstrual abnormalities and female breast pain were reported during the trial; one case occurred in a placebo-treated patient and the other in an eplerenone 100 mg/day-treated patient.
Krum et al. <sup>68</sup> (2002)  Eplerenone 50 to 100 mg/day  vs  placebo  All patients were receiving	DB, MC, PG, RCT  Patients 18 to 85 years of age taking an ACE inhibitor or an ARB for mild to moderate HTN (DBP ≥95 but <110 mm Hg and SBP <180 mm Hg), with potassium >3 mEq/L but ≤5 mEq/L	N=341  8 weeks	Primary: Mean change from baseline in trough cuff seated DBP and SBP at week eight  Secondary: Percentage of responders (DBP <90 mm Hg or exhibited a ≥10	Primary: Eplerenone-treated patients exhibited a significant mean reduction from baseline in SBP compared to placebo-treated patients at eight weeks of therapy (P≤0.05), regardless of concurrent ACE inhibitor or ARB use.  While eplerenone plus ARB-treated patients exhibited a significant mean reduction from baseline in DBP compared to ARB-treated patients at week eight (P≤0.05), eplerenone plus ACE inhibitor-treated patients experienced a reduction in baseline DBP similar to ACE inhibitor-treated patients (P value not reported).  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
background ACE inhibitor or ARB monotherapy.			mm Hg reduction from baseline), adverse events	<p>A significantly greater percentage of eplerenone plus ARB-treated patients exhibited a positive response to therapy compared to ARB-treated patients (P=0.003). No significant differences in response rate were observed between eplerenone plus ACE inhibitor- and ACE inhibitor-treated patients (P value not reported).</p> <p>Adverse effects were mild to moderate and were similar in eplerenone- and placebo-treated groups (P value not reported).</p>
<p>Weinberger et al.<sup>69</sup> (2002)</p> <p>Eplerenone 50 mg QD</p> <p>vs</p> <p>eplerenone 25 mg BID</p> <p>vs</p> <p>eplerenone 100 mg QD</p> <p>vs</p> <p>eplerenone 50 mg BID</p> <p>vs</p> <p>eplerenone 400 mg QD</p> <p>vs</p> <p>eplerenone 200 mg</p>	<p>DB, MC, PG, RCT</p> <p>Patients 21 to 80 years of age, with seated, cuff-assessed DBP <math>\geq</math>95 but &lt;114 mm Hg, a 24 hour mean DBP &gt;85 mm Hg</p>	<p>N=409</p> <p>8 weeks</p>	<p>Primary: Mean change in seated DBP from baseline</p> <p>Secondary: Mean change from baseline in SBP, 24 hour SBP and DBP, renin, aldosterone levels</p>	<p>Primary: Eplerenone therapy, across all doses studied, was associated with a significant reduction from baseline in seated and standing DBP compared to placebo (P&lt;0.05).</p> <p>The eplerenone 50 mg BID regimen was associated with a significant reduction in baseline seated and standing DBP compared to the eplerenone 100 mg QD regimen (P&lt;0.05). However, there were no differences in DBP reduction between any of the other QD and BID eplerenone regimens (P value not reported).</p> <p>Compared to placebo, spironolactone therapy was associated with significant reductions in DBP (P<math>\leq</math>0.001).</p> <p>The eplerenone 50 mg BID and 100 mg QD regimens were associated with DBP reductions comparable to 50 to 75% of the effect observed with the spironolactone 50 mg BID regimen (P value not reported).</p> <p>Secondary: Eplerenone therapy, across all doses studied, was associated with a significant reduction from baseline in seated and standing SBP compared to placebo therapy (P&lt;0.05).</p> <p>The eplerenone 200 mg BID regimen was associated with a significant reduction in baseline seated and standing SBP compared to the eplerenone 400 mg QD regimen (P&lt;0.05). However, there were no differences in SBP reduction between any of the other QD and BID eplerenone regimens (P value not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>BID</p> <p>vs</p> <p>spironolactone 50 mg BID</p> <p>vs</p> <p>placebo</p>				<p>Eplerenone therapy, across all doses studied, was associated with a significant reduction in ambulatory SBP and DBP compared to placebo therapy, as observed during a 24 hour monitoring (P&lt;0.05).</p> <p>Compared to placebo, spironolactone was associated with a significant reduction in SBP (P≤0.001).</p> <p>The eplerenone 50 mg BID and 100 mg QD regimens were associated with SBP reductions comparable to 50 to 75% of the effect observed with the spironolactone 50 mg BID regimen (P value not reported).</p> <p>The incidence of adverse events in eplerenone-treated patients was similar to placebo-treated patients (P value not reported). Additionally, the incidence of adverse events was comparable with eplerenone- and spironolactone-treated patients (P value not reported).</p> <p>The spironolactone 50 mg BID regimen was associated with a significant increase from baseline in serum potassium level compared to the eplerenone 50 mg QD and 100 mg QD regimens, regardless of QD or BID dosing (P&lt;0.05).</p> <p>Eplerenone therapy was not associated with an increased incidence of gynecomastia or impotence compared to placebo therapy. There were no treatment-related menstrual abnormalities reported with eplerenone therapy, while one spironolactone-treated patient reporting treatment related intermenstrual bleeding.</p>
<p>Hollenberg et al.<sup>70</sup> (2003)</p> <p>Eplerenone 50 mg/day</p> <p>vs</p> <p>amlodipine 2.5 mg/day</p>	<p>RCT</p> <p>Patients ≥50 years of age, with untreated SBP between 140 to 190 mm Hg</p>	<p>N=269</p> <p>24 weeks</p>	<p>Primary: Change in SBP and DBP, discontinuation rate, symptom distress index, SF-36 Health Survey</p> <p>Secondary: Not reported</p>	<p>Primary: Both treatments exhibited similar reductions in SBP and DBP from baseline (P=0.01).</p> <p>The dropout rate was 50% greater in amlodipine-treated patients compared to eplerenone-treated patients (P value not reported).</p> <p>Symptom distress (technique used to assess the influence of drug treatment on quality of life) index was assessed and results favored eplerenone therapy (P=0.03).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Both medications were titrated to a maximum of 200 (eplerenone) or 10 (amlodipine) mg/day to achieve a SBP&lt;140 mm Hg.</p>				<p>SF-36 Health Survey showed no significant difference between the two treatments (P value not reported).</p> <p>Both treatments experienced similar incidences of adverse effects (P value not reported). Eplerenone-treated patients did not experience breast pain/tenderness, breast enlargement, changes in menstruation, gynecomastia or loss of libido.</p> <p>Secondary: Not reported</p>
<p>White et al.<sup>71</sup> (2003)</p> <p>Eplerenone 50 mg/day</p> <p>vs</p> <p>amlodipine 2.5 mg/day</p> <p>Both medications were titrated to a maximum of 200 (eplerenone) or 10 (amlodipine) mg/day to achieve a SBP&lt;140 mm Hg.</p>	<p>AC, DB, MC, RCT</p> <p>Patients ≥50 years of age with systolic HTN (seated clinic SBP 150 to 165 mm Hg with a pulse pressure ≥70 mm Hg or 165 to 200 mm Hg with a DBP ≤95 mm Hg)</p>	<p>N=269</p> <p>24 weeks</p>	<p>Primary: Mean change from baseline in SBP, DBP, 24 hour ambulatory BP, pulse pressure, and heart rate at week 24; urine albumin/creatinine ratio; adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: Mean reduction in SBP from baseline was comparable in eplerenone- and amlodipine-treated patients (P=0.83).</p> <p>Eplerenone-treated patients exhibited significant reductions in DBP from baseline at 24 weeks of therapy compared to amlodipine-treated patients (P=0.014).</p> <p>The two treatments exhibited comparable decreases in 24 hour ambulatory BP, pulse pressure and heart rate after 24 weeks of therapy (P&gt;0.05).</p> <p>Eplerenone-treated patients exhibited a significant reduction from baseline in the urine albumin/creatinine ratio compared to amlodipine-treated patients (P=0.002).</p> <p>Treatment-emergent adverse events were reported in 64 and 70% of eplerenone- and amlodipine-treated patients. The only adverse event that was significant between the two treatments was the incidence of edema (3.7 vs 25.5%; P&lt;0.05). There were no reports of gynecomastia, breast tenderness or menstrual irregularities with either treatment.</p> <p>Secondary: Not reported</p>
<p>Williams et al.<sup>72</sup> (2004)</p> <p>Eplerenone 50 mg</p>	<p>AC, DB, MC, PG, RCT</p> <p>Patients ≥18 years of</p>	<p>N=499</p> <p>12 months</p>	<p>Primary: Change in seated trough DBP at 6 months</p>	<p>Primary: At six months, both treatments exhibited comparable reductions in DBP from baseline (P=0.91).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD vs enalapril 10 mg QD</p> <p>Both medications were titrated to 200 (eplerenone) or 40 (enalapril) mg/day if needed for optimal blood pressure control (DBP &lt; 90 mm Hg).</p>	<p>age with stage 1 to 2 HTN (seated DBP ≥90 but &lt;110 mm Hg, with a seated SBP &lt;190 mm Hg)</p>		<p>Secondary: Change in seated trough SBP at 6 months, reduction in SBP and DBP at 12 months, reduction in urine albumin/ creatinine ratio, adverse events</p>	<p>Secondary: At six months, both treatments exhibited comparable reductions in SBP from baseline (P=0.20).</p> <p>At 12 months, both treatments exhibited comparable reductions in SBP and DBP from baseline (P=0.25 and P=0.33).</p> <p>Eplerenone-treated patients exhibited a significant reduction from baseline in urine albumin/creatinine ratio compared to enalapril-treated patients (61.5 vs 25.7%; P=0.01).</p> <p>There were no significant differences in overall treatment-emergent adverse events between the two treatments (P value not reported). There were no sex hormone related adverse events in eplerenone-treated patients. There were no clinically significant differences between the two treatments in any of the laboratory tests assessed. There were two eplerenone- and enalapril-treated patients that experienced hyperkalemia of ≥5.5 mmol/L.</p>
<p>Flack et al.<sup>73</sup> (2003)</p> <p>Eplerenone 50 mg QD vs losartan 50 mg QD vs placebo</p> <p>Doses were increased if blood pressure remained uncontrolled.</p>	<p>DB, MC, PG, RCT</p> <p>Men and women ≥18 years old, with mild to moderate HTN, with SBP &lt;180 mm Hg and DBP 95 to 109 mm Hg (off medication) or if patients were receiving antihypertensive therapy their blood pressure was &lt;140/90 mm Hg</p>	<p>N=551</p> <p>16 weeks</p>	<p>Primary: Mean change from baseline in DBP at 16 weeks</p> <p>Secondary: Mean change from baseline at 16 weeks in SBP, SBP and DBP within and between racial groups, response rate (defined as the percentage of patients with DBP &lt;90 mm Hg or DBP ≥90 mm Hg but ≥10 mm Hg below baseline),</p>	<p>Primary: At 16 weeks, patients randomized to eplerenone exhibited significantly greater mean changes in DBP from baseline compared to either losartan- or placebo-treated groups (P&lt;0.001).</p> <p>Secondary: At 16 weeks, patients randomized to eplerenone exhibited significantly greater mean changes in SBP from baseline compared to either losartan- or placebo-treated groups (P&lt;0.001).</p> <p>At 16 weeks, African American patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the placebo-treated African American patients (P&lt;0.001).</p> <p>At 16 weeks, African American patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the losartan-treated African American patients (P≤0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>urinary albumin/creatinine ratio, effect of eplerenone in patients with various baseline renin and aldosterone levels, adverse effects</p>	<p>At 16 weeks, white patients randomized to eplerenone exhibited significantly greater mean changes in SBP and DBP from baseline compared to the placebo-treated white patients (P=0.001). However, the difference in SBP- and DBP-lowering effects was not significant different between the eplerenone ad losartan groups (P=0.126, P=0.068, respectively).</p> <p>Significantly greater percentage of patients randomized to eplerenone exhibited a positive response to therapy compared to either placebo (64.5 vs 41.2%; P&lt;0.001) or losartan group (64.5 vs 48.3%; P=0.003).</p> <p>The eplerenone group (regardless of race) exhibited statistically significant improvement in urinary albumin/creatinine ratio from baseline compared to placebo (P=0.003). However, the difference in urinary albumin/creatinine ratio change from baseline was not significantly different between the eplerenone and losartan groups (P=0.652).</p> <p>Compared to losartan, eplerenone was more effective in lowering SBP and DBP in patients with low-moderate baseline renin levels (P&lt;0.05). However, the difference was not statistically significant in patients with high baseline renin levels.</p> <p>Compared to losartan, eplerenone was more effective in lowering SBP in patients with low or high baseline aldosterone levels (P&lt;0.05). However, the difference was not statistically significant in patients with moderate baseline aldosterone levels.</p> <p>Compared to losartan, eplerenone was more effective in lowering DBP in patients with low baseline aldosterone levels (P&lt;0.05). However, the difference was not statistically significant in patients with moderate-high baseline aldosterone levels.</p> <p>There were no significant differences in the incidence of adverse events noted in eplerenone, placebo or losartan groups. The reported incidence of gynecomastia, breast pain, menstrual abnormalities, impotence, hyperkalemia and decreased libido with eplerenone was low and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				comparable to losartan and placebo.
Hanazawa et al. <sup>74</sup> (abstract) (2011)  Spironolactone 12.5 or 25 mg/day  In addition to existing antihypertensive regimens (monotherapy with a calcium channel blocker, ACE inhibitor, or ARB).	PRO  Patients with uncontrolled HTN	N=86  Not reported	Primary: Change in baseline blood pressure  Secondary: Not reported	Primary: Morning home SBP/DBP reduction was similar among patients not controlled on a calcium channel blocker (n=30, -8.2/-2.6 mmHg), ACE inhibitor (n=22, -13.0/-4.7 mmHg), and ARB (n=34, -11.5/-5.1 mmHg).  An increase in serum potassium correlated positively with the decline in morning SBP.  Secondary: Not reported
Schersten et al. <sup>75</sup> (2002)  Spironolactone 50 mg/day  vs  spironolactone 100 mg/day  vs  spironolactone 200 mg/day  vs  placebo	RCT, SB, XO  Patients <75 years of age, with DBP 105 to 135 mm Hg, after 10 to 15 minutes of supine rest	N=45  11 months	Primary: Change from baseline in DBP and SBP, adverse effects  Secondary: Not reported	Primary: All spironolactone-treated patients exhibited a significantly reduced BP level from baseline as compared to placebo (P<0.001).  While spironolactone 200 mg/day-treated patients exhibited a significantly greater lowered mean supine SBP compared to spironolactone 50 mg/day-treated patients (P<0.05), the difference between spironolactone 50 mg- and 100 mg/day-treated patients was not significant (P value not reported).  Spironolactone 200 mg/day-treated patients exhibited a significant reduction in mean upright SBP from baseline compared to spironolactone 100 mg/day- and 50 mg/day-treated patients (P<0.01).  The difference in the lowering of DBP from baseline was not significantly different among any of the spironolactone-treated patients (P value not reported).  Spironolactone 100 mg/day-treated patients exhibited a significant increase in baseline potassium and serum creatinine concentrations (P<0.05). However, spironolactone 50 mg/day-treated patients did not exhibit a change in potassium level from baseline (P value not reported).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Oxlund et al. <sup>76</sup> (2013)  Spironolactone 25 mg (option to titrate to twice daily)  vs  placebo	DB, MC, RCT  Patients aged 30 to 74 years with blood pressure at or above 130/80 mmHg despite triple antihypertensive therapy	N=119  16 weeks	Primary: Reduction of mean SBP and DBPs  Secondary: Glycemic control, urinary albumin excretion, adverse effects	Primary: All measures of BP by office as well as ambulatory monitoring showed marginal, insignificant reductions in the placebo group and significant reductions in the spironolactone group. Maximum reduction of office BP (11.3/5.3 mmHg) was found at 8 weeks of treatment (P<0.0001), after which no further reduction was found. Mean daytime SBP/DBP at 16 weeks of follow-up was 137 (13)/75 (8) mmHg in the spironolactone group and 145 (12)/79 (7) mmHg in the placebo group, P=0.0001 for difference of systolic measures and P=0.0038 for diastolic.  Secondary: Urinary albumin/creatinine ratio was reduced significantly in the spironolactone group (P=0.001), but not in the placebo group. There was a nonsignificant decrease of eGFR in the spironolactone group. Hb1Ac did not change during intervention. The frequency of adverse events was comparable in the two groups.
Václavík et al. <sup>77</sup> (2014) ASPIRANT-EXT  Spironolactone 25 mg  vs  placebo	DB, MC, RCT  Patients with office SBP >140 mm Hg or DBP >90 mm Hg despite treatment with at least 3 antihypertensive drugs, including a diuretic	N=150  8 weeks	Primary: Comparison of the fall of daytime systolic and diastolic pressure on ABPM between the spironolactone and placebo groups after 8 weeks of treatment  Secondary: Nighttime BP, serum sodium, potassium, and creatinine, change in body weight	Primary: At 8 weeks, BP values were decreased more by spironolactone, with differences in mean fall of SBP of -9.8, -13.0, -10.5, and -9.9 mm Hg (P<0.001 for all) in daytime, nighttime, and 24-hour ambulatory BP monitoring and in the office. The respective DBP differences were -3.2, -6.4, -3.5, and -3.0 mm Hg (P=0.013, P<0.001, P=0.005, and P=0.003).  Secondary: A small comparable weight gain was observed in both study groups. With spironolactone treatment, serum sodium decreased by a median of 1.0 mmol/L, and serum potassium increased by a median 0.4 mmol/L. The mean serum potassium increased during the 8 weeks of spironolactone treatment from 4.10 to 4.49 mmol/L, the highest reached serum potassium value at 8 weeks was 5.6 mmol/L.
Li et al. <sup>78</sup>	DB, MC, PC, RCT	N=304	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2010)</p> <p><u>Phase A</u> Spironolactone 25 mg QD (low-dose), 25 mg BID (middle-dose), 50 mg BID (high-dose) for 6 weeks</p> <p><u>Phase B</u> Spironolactone 25 mg QD, 25 mg BID, 50 mg BID for 4 weeks</p> <p>vs</p> <p>placebo</p>	<p>Children 4 to 16 years of age with SBP <math>\geq</math>95th percentile</p>		<p>Change in SBP during phase B</p> <p>Secondary: Change in DBP, safety</p>	<p>Change in SBP from baseline of phase B to the end of the study (differences from placebo) were -2.61, 2.32, and -2.76 mm Hg for the low-, middle-, and high-dose groups, respectively (P value not significant, P value not significant, P=0.048, respectively).</p> <p>Secondary: There were no significant effects of eplerenone on change in DBP from baseline of phase B to end of study compared to placebo.</p> <p>During phase A, adverse events were reported by 40.2% of subjects in the high-dose group, 30.6% of those in the middle-dose group, and 37.9% of those in the low-dose group. In phase B, there were no differences in adverse event frequencies between active therapy and placebo (high-dose: 38.4 vs 45.2%; middle-dose: 50.0 vs 25.0%; low-dose 26.9 vs 34.6%, eplerenone vs, placebo, respectively).</p> <p>Serious adverse events in phase A included diarrhea, sleep apnea, syncope, pericarditis, arthritis, pneumonia, sepsis, and pleural effusion. In phase B, serious adverse events included sleep apnea, abdominal pain, and fever.</p>
<p>Hood et al.<sup>79</sup> (2007) SALT</p> <p>Spironolactone 50 mg/day</p> <p>vs</p> <p>spironolactone 100 mg/day</p> <p>vs</p> <p>amiloride 20 mg/day</p> <p>vs</p>	<p>DB, RCT, XO</p> <p>Adult patients with seated blood pressure of 140/90 to 170/110 mm Hg, plasma renin of <math>\leq</math>12 mU/L, plasma aldosterone-renin ratio <math>&gt;</math>750, previous fall in SBP <math>\geq</math>20 mm Hg after 1 month of OL treatment with spironolactone 50 mg/day</p>	<p>N=57</p> <p>42 weeks</p>	<p>Primary: Change in blood pressure and plasma renin from baseline between spironolactone 100 mg/day and bendroflumethiazide 5 mg/day</p> <p>Secondary: Change in blood pressure and plasma renin from baseline between amiloride and other diuretics and</p>	<p>Primary: Spironolactone 100 mg/day- and bendroflumethiazide 5 mg/day-treated patients did not exhibit a significant difference in BP reduction from baseline (P value not reported).</p> <p>Secondary: Spironolactone 50 mg/day-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (P&lt;0.01).</p> <p>Losartan 100 mg-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (P&lt;0.05).</p> <p>High-dose bendroflumethiazide- and amiloride-treated patients exhibited significantly greater reductions in blood pressure compared to the lower doses (P&lt;0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>amiloride 40 mg/day</p> <p>vs</p> <p>bendroflumethiazide* 2.5 mg/day</p> <p>vs</p> <p>bendroflumethiazide* 5 mg/day</p> <p>vs</p> <p>losartan 100 mg/day</p> <p>vs</p> <p>placebo</p>			<p>between lower and higher doses of each diuretic</p>	<p>Spirolactone-treated patients exhibited a four-fold increase in baseline renin level compared to a two-fold increase observed in bendroflumethiazide-treated patients (P=0.003).</p>
<p>Nash et al.<sup>80</sup> (1977)</p> <p>Spirolactone 50 mg BID</p> <p>vs</p> <p>spironolactone 100 mg BID</p> <p>vs</p>	<p>DB, RCT</p> <p>Male outpatients between the ages of 21 to 65 years, with essential HTN, DBP between 90 to 114 mm Hg</p>	<p>N=79</p> <p>12 weeks</p>	<p>Primary: Change in SBP, DBP, blood urea nitrogen, serum potassium, gynecomastia</p> <p>Secondary: Not reported</p>	<p>Primary: At week 12, all study groups exhibited significant reductions in SBP and DBP from baseline (P&lt;0.05).</p> <p>At week 12, all three spironolactone monotherapy groups exhibited statistically significant increases in blood urea nitrogen from baseline (P&lt;0.05).</p> <p>At week 12, the HCTZ monotherapy group was associated with a statistically significant decrease in serum potassium levels (P&lt;0.001).</p> <p>At week 12, all three spironolactone monotherapy groups exhibited statistically significant increases in serum potassium levels from baseline</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
spironolactone 200 mg BID  vs  HCTZ 50 mg BID  vs  spironolactone and HCTZ 25-25 mg BID (fixed-dose combination product)				(P<0.05).  At week 12, the spironolactone and HCTZ combination group was not associated with statistically significant increases in serum potassium levels from baseline.  A dose-related risk of gynecomastia was observed in the spironolactone-treated patients. Among patients treated with spironolactone 50, 100, or 200 mg BID; 5.5, 11.8, and 40% reported gynecomastia symptoms. Of the patients randomized to spironolactone and HCTZ combination product, 7.7% reported gynecomastia symptoms.  Secondary: Not reported
Schrijver et al. <sup>81</sup> (1979)  Spironolactone 50 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IA)  vs  spironolactone 50 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks	DB  Patients, between 24 to 63 years of age, with DBP between 90 to 114 mm Hg	N=49  20 weeks (4-week placebo run-in, 8-week single drug therapy, 4-week two-drug therapy, 4-week recovery)	Primary: Change in MABP, serum potassium, uric acid level, blood glucose, blood urea nitrogen, creatinine, plasma renin activity, aldosterone, side effects  Secondary: Not reported	Primary: Following eight weeks of therapy with a single drug, all study groups exhibited a statistically significant reduction in MABP from baseline (P<0.01). There were no significant differences in MABP reduction among the study groups.  The addition of a second drug to the antihypertensive regimen was not associated with a significant improvement in MABP. At the end of the two-drug treatment period, there were no differences in MABP among any of the study groups.  Spironolactone therapy was associated with a significant decrease in serum potassium concentration from baseline (P<0.001).  Spironolactone regimens were not associated with a significant change in potassium levels from baseline.  Following eight weeks of therapy with a single drug, HCTZ-treated patients experienced a statistically significant increase in uric acid from baseline (P<0.001). Groups IIA and IIB also experienced a significant but smaller increase in uric acid level from baseline (P<0.05) with no change in groups I and IV.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(group IB)</p> <p>vs</p> <p>spironolactone 100 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IIA)</p> <p>vs</p> <p>spironolactone 100 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IIB)</p> <p>vs</p> <p>spironolactone 200 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IIIA)</p>				<p>During the single-drug treatment phase, patients randomized to group I experienced a significant increase in blood glucose from baseline (P&lt;0.05).</p> <p>During the single-drug treatment phase, all patients except those randomized to group I experienced a significant increase in blood urea nitrogen from baseline (P&lt;0.05).</p> <p>During the single-drug treatment phase, patients randomized to groups I and II experienced a significant increase in serum creatinine from baseline (P&lt;0.05).</p> <p>During the single-drug treatment phase, all treatment groups experienced a significant increase in plasma renin activity from baseline (P&lt;0.01). The addition of HCTZ in the two-drug study phase was associated with a rise in plasma renin activity in all study groups (P&lt;0.05).</p> <p>All treatment groups experienced a significant increase in plasma aldosterone from baseline (P&lt;0.05).</p> <p>Gynecomastia was reported only by patients randomized to the higher-dose spironolactone groups.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>spironolactone 200 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IIIB)</p> <p>vs</p> <p>HCTZ 50 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IVA)</p> <p>vs</p> <p>HCTZ 50 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IVB)</p>				
<p>Wray et al.<sup>82</sup> (2010)</p>	<p>DB, RCT</p> <p>Patients ≥60 years of</p>	<p>N=36</p> <p>6 months</p>	<p>Primary: Blood pressure, sympathetic</p>	<p>Primary: Arterial blood pressure decreased significantly with spironolactone (SBP: 160 to 134 mm Hg and DBP: 77 to 68 mm Hg) and with HCTZ (SBP: 161</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Spirolactone 25 to 100 mg QD</p> <p>vs</p> <p>HCTZ 12.5 to 50 mg QD</p> <p>Patients also received potassium 0 to 40 mEq to maintain blinding.</p>	<p>age with stage 1 HTN</p>		<p>nervous system activity</p> <p>Secondary: Not reported</p>	<p>to 145 mm Hg and 78 to 73 mm Hg). There was no significant difference between the groups.</p> <p>Sympathetic nervous system activity was significantly reduced after spironolactone (plasma norepinephrine: 378 to 335 pg/mL; P=0.04; [3H]-norepinephrine release rate: 2.74 to 1.97 µg/min/m<sup>2</sup>; P=0.04), but not with HCTZ (plasma norepinephrine: 368 to 349 pg/mL; P=0.47; [3H]-norepinephrine release rate: 2.63 to 2.11 µg/min/m<sup>2</sup>; P=0.21).</p> <p>There were no instances of hyperkalemia, and no other adverse effects were reported.</p> <p>Secondary: Not reported</p>
<p>Krieger et al.<sup>83</sup> (2018) ReHOT</p> <p>Spirolactone 12.5 mg QD (could be titrated to 25 or 50 mg/ day)</p> <p>vs</p> <p>clonidine 0.1 mg BID (could be titrated to 0.2 or 0.3 mg BID)</p>	<p>OL, RCT</p> <p>Patients with resistant hypertension (no office and ambulatory BP monitoring control, despite treatment with 3 drugs, including a diuretic, for 12 weeks)</p>	<p>N=162</p> <p>12 weeks</p>	<p>Primary: BP control (determined by office BP&lt;140/90 and ambulatory 24-hour mean BP &lt;130/80)</p> <p>Secondary: BP control by each evaluation method, absolute BP reduction</p>	<p>Primary: Compared with the spironolactone group, the clonidine group presented similar rates of achieving the primary end point (20.5 vs 20.8%, respectively; RR, 1.01; 95% CI, 0.55 to 1.88; P=1.00).</p> <p>Secondary: Secondary end point analysis showed similar office BP (33.3 vs 29.3%) and ambulatory BP monitoring (44 vs 46.2%) control for spironolactone and clonidine, respectively. However, spironolactone promoted greater decrease in 24-hour systolic and diastolic BP and diastolic daytime ambulatory BP than clonidine.</p>
<p>Bomback et al.<sup>84</sup> (2009)</p> <p>Spirolactone 12.5 mg QD for 4 weeks in addition to ACE inhibitor therapy</p>	<p>OL</p> <p>Patients with obesity, longstanding hypertension and evidence of target organ damage who</p>	<p>N=21</p> <p>8 weeks</p>	<p>Primary: Change in 24-hour ambulatory blood pressure, changes in office blood pressure, nocturnal blood pressure, and</p>	<p>Primary: Mean office, 24-hr ambulatory, and nocturnal ambulatory blood pressures declined significantly during the four weeks of spironolactone therapy from 110.6 to 105.0 mm Hg (office P=0.004), 100.6 to 95.5 mm Hg (24-hr P=0.03) and 95.3 to 87.5 mm Hg (nocturnal P=0.004).</p> <p>The mean urine albumin: creatinine ratio dropped from 13.8 to 8.5 mg/g (P=0.002) during spironolactone therapy and returned to 13.2 mg/g after</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	were treated with ACE inhibitors		urine albumin: creatinine ratio  Secondary: Not reported	the drug was withdrawn.  Serum potassium was not significantly affected by spironolactone therapy. There was a significant increase in serum creatinine from 0.95 before therapy to 1.03 mg/dl after spironolactone. The eGFR decreased from 81.9 to 76.8 mL/min/1.73m <sup>2</sup> .  Secondary: Not reported
Williams et al. <sup>85</sup> (2015) PATHWAY-2  Twelve weeks of once daily treatment with each of spironolactone (25 to 50 mg), bisoprolol (5 to 10 mg), doxazosin modified release (4 to 8 mg), and placebo, in addition to their baseline blood pressure drugs	DB, PC, XO  Patients 18 to 79 years of age with seated clinic SBP $\geq$ 140 mmHg (or $\geq$ 135 mmHg for patients with diabetes) and home SBP (18 readings over four days) $\geq$ 130 mmHg, despite treatment for at least three months with maximally tolerated doses of three drugs (an ACE or ARB, a CCB, and a diuretic)	N=335  12 months	Primary: Average home SBP, recorded in the morning and the evening in triplicate, on four consecutive days before study visits  Secondary: Clinic SBP, BP control rates, adverse events	Primary: The average reduction in home SBP by spironolactone was significantly greater compared to placebo (-8.70 mmHg; 95% CI, -9.72 to -7.69; P<0.0001), to the mean of the other two active treatments (doxazosin and bisoprolol; -4.26; 95% CI, -5.13 to -3.38; P<0.0001), and to the individual treatments; versus doxazosin (-4.03; 95% CI, -5.04 to -3.02; P<0.0001) and versus bisoprolol (-4.48; 95% CI, -5.50 to -3.46; P<0.0001).  Secondary: The results for seated clinic SBP largely mirror those seen with home SBP except that there was a large placebo effect on clinic BP that was not seen with home BP measurement.  Overall 219 (68.9%; 95% CI, 63.6 to 73.8) of 314 patients achieved target home SBP of <135 mmHg. 58% of patients had their BP controlled with spironolactone, which was significantly greater than rates for other treatments (P<0.001 when compared to doxazosin, bisoprolol, and placebo). Most patients who were controlled by doxazosin or bisoprolol had a still greater fall in blood pressure on spironolactone, which was consequently the most effective treatment in almost 60% of patients. This was at least three times the proportion in whom doxazosin or bisoprolol were the most effective.  All active treatments were well tolerated with similar low rates of adverse events and withdrawals due to adverse events.
Chapman et al. <sup>86</sup> (2007)	Subanalysis of ASCOT-BPLA	N=1,411	Primary: Change in DBP	Primary: Spironolactone-treated patients lead to a significant 21.9 mm Hg reduction



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>ASCOT-BPLA</p> <p>Atenolol 50 to 100 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); bendroflumethiazide* plus potassium 1.25 to 2.5 mg plus doxazosin were added for additional blood pressure control; if blood pressure remained elevated on the 3 above drugs, spironolactone 25 mg was added to the regimen</p> <p>vs</p> <p>amlodipine 5 to 10 mg titrated to target blood pressure &lt;140/90 mm Hg (or &lt;130/90 mm Hg in diabetic patients); perindopril 4 to 8 mg and doxazosin were added for additional control; if blood pressure remained elevated</p>	<p>evaluating effects of spironolactone on treatment-resistant HTN</p> <p>Patients 40 to 79 years of age with HTN and <math>\geq 3</math> cardiovascular risk factors, with SBP <math>\geq 160</math> mm Hg and/or DBP <math>\geq 100</math> mm Hg (not on antihypertensive therapy) or SBP <math>\geq 140</math> mm Hg and/or DBP <math>\geq 90</math> mm Hg (on antihypertensive therapy)</p>	<p>1.3 years</p>	<p>and SBP, adverse effects</p> <p>Secondary: Not reported</p>	<p>in SBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 20.8 to 23.0 mm Hg; <math>P &lt; 0.001</math>).</p> <p>Spironolactone-treated patients lead to a significant 9.5 mm Hg reduction in DBP among patients whose blood pressure was previously uncontrolled on at least three other antihypertensive drugs (95% CI, 9.0 to 10.1; <math>P &lt; 0.001</math>).</p> <p>Spironolactone-treated patients exhibited small but significant decreases in sodium, LDL-C and TC as well as increases in potassium, glucose, creatinine and HDL-C (<math>P &lt; 0.05</math>).</p> <p>The most common adverse effect reported in the trial was gynecomastia in men (<math>P</math> value not reported).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
on the 3 above drugs, spironolactone 25 mg was added to the regimen				
<b>Diabetic Kidney Disease</b>				
Filippatos et al. <sup>87</sup> (2021) FIDELIO-DKD Finerenone 10 or 20 mg QD vs placebo	DB, MC, PC, PG, RCT  Patients ≥18 years of age with a clinical diagnosis of T2D and moderately elevated albuminuria and an eGFR ≥25 to <75mL/min/1.73m <sup>2</sup>	N=5,674  Median follow-up 2.6 years (range, 2.0 to 3.4 years)	Primary: Composite cardiovascular outcome included time to first onset of cardiovascular death, nonfatal myocardial infarction, nonfatal stroke or hospitalization for heart failure  Secondary: Composite kidney outcome included time to first onset of kidney failure, a sustained ≥40% decrease in eGFR from baseline over at least four weeks, or renal death	Primary: Over a median follow-up of 2.6 years, finerenone reduced the risk of the composite cardiovascular outcome compared with placebo (HR, 0.86; 95% CI, 0.75 to 0.99; P=0.034), with no significant interaction between patients with and without CVD (HR, 0.85; 95% CI, 0.71 to 1.01 in patients with a history of CVD; HR, 0.86; 95% CI, 0.68 to 1.08 in patients without a history of CVD; P value for interaction=0.85).  Secondary: The composite kidney outcome was lower with finerenone versus placebo; however, the effects were more pronounced in patients with a history of CVD than those without it (P value for interaction=0.016). In patients with a history of CVD, the composite kidney outcome occurred in 200 (15.3%) patients in the finerenone group and 267 (20.5%) patients in the placebo group (incidence rate per 100 patient-years, 6.6 and 9.06, respectively; HR, 0.70; 95% CI, 0.58 to 0.84). In patients without a history of CVD, the composite kidney outcome occurred in 304 (19.9%) patients in the finerenone group and 333 (21.6%) patients in the placebo group (incidence rate per 100 patient-years, 8.42 and 9.1, respectively; HR, 0.94; 95% CI, 0.81 to 1.10).
Pitt et al. <sup>88</sup> (2021) FIGARO-DKD Finerenone 10 mg or 20 mg QD vs	DB, MC, PC, RCT  Patients ≥18 years of age with T2D and CKD treated with RAS inhibitor at the maximum dose on the manufacturer's	N=7,437  Median follow-up 3.4 year	Primary: Composite of death from cardiovascular causes, nonfatal myocardial infarction, nonfatal stroke or	Primary: The incidence of death from cardiovascular causes, nonfatal myocardial infarction, nonfatal stroke, or hospitalization for heart failure (the primary composite outcome) was significantly lower in the finerenone group than in the placebo group (458 of 3,686 patients [12.4%] vs. 519 of 3,666 patients [14.2%]; HR, 0.87; 95% CI, 0.76 to 0.98; P=0.03).  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>placebo</p>	<p>label that di not cause unacceptable side effects</p>		<p>hospitalization for heart failure</p> <p>Secondary: Composite of the first occurrence of kidney failure, a sustained decrease from baseline of at least 40% in the eGFR for a period of at least four weeks or death from renal causes</p>	<p>There was no significant between-group difference in the incidence of the first secondary composite outcome of failure, a sustained decrease from baseline of at least 40% in the eGFR for a period of at least four weeks or death from renal causes (350 patients [9.5%] in the finerenone group and 395 [10.8%] in the placebo group; HR, 0.87; 95% CI, 0.76 to 1.01).</p>
<b>Miscellaneous</b>				
<p>Pitt et al.<sup>89</sup> (2003) 4E-Left Ventricular Hypertrophy Study Eplerenone 200 mg QD vs enalapril 40 mg QD vs enalapril 10 mg plus eplerenone 200 mg  If the blood pressure was uncontrolled on study medication at</p>	<p>AC, DB, PGd, RCT  Patients with left ventricular hypertrophy, a history of HTN and predominantly in sinus rhythm</p>	<p>N=153  9 months</p>	<p>Primary: Change in left ventricular mass as assessed by MRI</p> <p>Secondary: Reduction in SBP and DBP, response rate (DBP &lt;90 mm Hg), change in urine albumin creatinine ratio</p>	<p>Primary: Both treatments were associated with a significant reduction in left ventricular mass from baseline (P&lt;0.001). The difference in left ventricular mass reduction from baseline between the two treatments was not significant (P=0.258).</p> <p>While enalapril plus eplerenone therapy demonstrated a significantly greater reduction in left ventricular mass from baseline compared to eplerenone therapy (P=0.007); the effect was not statistically different from that observed with enalapril therapy (P=0.107).</p> <p>Secondary: The SBP was reduced significantly more in enalapril plus eplerenone-treated patients compared to eplerenone-treated patients (P=0.048). The other treatment groups exhibited statistically comparable reductions from baseline in mean SBP and DBP (P value not reported).</p> <p>While 70.0% of eplerenone-treated patients responded to therapy, 40.7% of enalapril-treated patients responded (P=0.003). In addition, 79.6% of enalapril plus eplerenone-treated patients responded to therapy compared to 40.7% enalapril-treated patients (P=0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>week 8, OL HCTZ 12.5 to 25 mg/day and/or amlodipine 10 mg/day were allowed.</p>				<p>Enalapril plus eplerenone therapy was associated with a significant reduction in urine albumin creatinine ratio compared to either eplerenone or enalapril therapy (P&lt;0.05).</p> <p>Adverse events were reported with similar incidence among all treatment groups (P value not reported). Cough was significant in enalapril-treated patients compared to eplerenone-treated patients (P=0.033). Two cases of gynecomastia were reported (one eplerenone- and one enalapril plus eplerenone-treated patients). Four patients (three enalapril- and one enalapril plus eplerenone-treated patients) experienced impotence during the trial. Seven eplerenone-, two enalapril- and three enalapril plus eplerenone-treated patients experienced serious hyperkalemia (<math>\geq 6.0</math> mmol/L).</p>
<p>Taniguchi et al.<sup>90</sup> (2006)</p> <p>Candesartan 8 mg in addition to spironolactone 25 mg QD for 6 months, after 6 months of candesartan monotherapy (combination group)</p> <p>vs</p> <p>candesartan 8 mg daily for 12 months</p>	<p>DB, RCT, XO</p> <p>Patients, 67 years of age on average, with essential HTN and left ventricular hypertrophy</p>	<p>N=97</p> <p>1 year</p>	<p>Primary: Change in blood pressure and relative wall thickness</p> <p>Secondary: Not reported</p>	<p>Primary: Both study groups experienced a statistically significant reduction in blood pressure from baseline (P&lt;0.05).</p> <p>While candesartan was associated with a significant reduction in relative wall thickness among patients with concentric left ventricular remodeling or hypertrophy (P&lt;0.05), the addition of spironolactone did not provide additional benefit.</p> <p>Secondary: Not reported</p>
<p>Edwards et al.<sup>91</sup> (2009)</p> <p>Spironolactone 25 mg QD</p>	<p>DB, PC, RCT</p> <p>Patients 18 to 80 years of age with stage 2 and 3 chronic kidney</p>	<p>N=115</p> <p>36 weeks</p>	<p>Primary: Change in left ventricular mass and arterial stiffness measured</p>	<p>Primary: Treatment with spironolactone resulted in significant reductions in left ventricular mass and left ventricular mass index. The prevalence of left ventricular hypertrophy decreased by 50% with spironolactone, but was unchanged with placebo. Spironolactone did not affect left ventricular volumes or ejection fraction.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  placebo  Study medications were added to existing ACE inhibitor or ARB therapy.	disease with controlled blood pressure (mean daytime ambulatory blood pressure <130/85 mm Hg) on and ACE inhibitors or ARB for 6 months		Secondary: Aortic distensibility, Aug AIx, blood pressure, and albuminuria	Secondary: Treatment with spironolactone resulted in a significant decrease in pulse wave velocity, central aortic pressure augmentation, Aug Ix, and Aug Ix 75. Aortic distensibility increased with the use of spironolactone compared to placebo.  Treatment with spironolactone resulted in a significant decrease in office systolic blood pressure (-11 vs -5 mm Hg, P<0.05), office pulse pressure (-5 mm Hg vs -1 mm Hg, P<0.05), central systolic blood pressure (-12 mm Hg vs -4 mm Hg, P<0.01), central mean arterial pressure (-8 mm Hg vs -4 mm Hg, P<0.05), and central pulse pressure (-5 mm Hg vs -1 mm Hg, P<0.01). Office, central, and ambulatory diastolic pressures were not different between treatment groups.  Treatment with spironolactone was not associated with a significant decrease in eGFR compared to placebo (-3 vs -1 mL/min/1.73 m <sup>2</sup> , respectively; P value not significant). Treatment with spironolactone reduced albuminuria by -21 mg/mmol compared to -8 mg/mmol with placebo (P<0.05).

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, QD=once daily

Study regimen abbreviations: AC=active comparator, BE=blinded endpoint, DB=double blind, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, SB=single blind, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor blocker, BNP=brain natriuretic peptide, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, eGFR=estimated glomerular filtration rate, GFR=glomerular filtration rate, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HR=hazard ratio, HTN=hypertension, KCl=potassium chloride, LDL-C=low-density lipoprotein cholesterol, LVEF=left ventricular ejection fraction, MABP=mean arterial blood pressure, MI=myocardial infarction, MRI=magnetic resonance imaging, NYHA=New York Heart Association, OR=odds ratio, PINP=procollagen type 1 N-terminal peptide, QOL=quality of life, RAAS=renin-angiotensin-aldosterone system, RR=relative risk, SBP=systolic blood pressure, T2D=type 2 diabetes, TC=total cholesterol

**Additional Evidence**

Dose Simplification

Ludbrook et al. evaluated the differences in blood pressure control and adverse events with spironolactone 300 to 400 mg administered either once daily or in in divided doses. Both administration schedules were associated with comparable systolic and diastolic blood pressure reductions. None of the regimens reduced the incidence of adverse effects (85% in both groups).<sup>92</sup>

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 9. Relative Cost of the Mineralocorticoid (Aldosterone) Receptor Antagonists**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Eplerenone	tablet	Inspra®*	\$\$\$\$\$	\$\$
Finerenone	tablet	Kerendia®	\$\$\$\$\$	N/A
Spironolactone	suspension, tablet	Aldactone®*, Carospir®	\$\$\$\$\$	\$
<b>Combination Products</b>				
Spironolactone and HCTZ	tablet	Aldactazide®*	\$\$\$\$	\$\$\$\$

\*Generic is available in at least one dosage form or strength.  
HCTZ=hydrochlorothiazide

**X. Conclusions**

The mineralocorticoid (aldosterone) receptor antagonists, eplerenone and spironolactone, are approved for the treatment of hypertension.<sup>3-6</sup> Eplerenone is also indicated to improve survival in patients with left ventricular systolic dysfunction (ejection fraction ≤40%) and clinical evidence of congestive heart failure after an acute

myocardial infarction.<sup>5</sup> Spironolactone is approved for the management of hyperaldosteronism, hypokalemia, and edema associated with congestive heart failure, cirrhosis, or the nephrotic syndrome. It is also indicated for patients with severe heart failure (NYHA class III to IV) to increase survival, and to reduce the need for hospitalization for heart failure when used in addition to standard therapy.<sup>3</sup> Spironolactone is now available as an oral suspension. Of note, Carospir<sup>®</sup> is not therapeutically equivalent to Aldactone<sup>®</sup>.<sup>4</sup> Spironolactone is available as single entity agents, as well as in combination with hydrochlorothiazide as a fixed-dose combination product.

Finerenone is a novel, non-steroidal, selective mineralocorticoid receptor antagonist indicated to reduce the risk of sustained eGFR decline, end stage kidney disease, cardiovascular death, non-fatal myocardial infarction, and hospitalization for heart failure in adult patients with CKD associated with T2D.<sup>7</sup> All of the mineralocorticoid (aldosterone) receptor antagonist products are available in a generic formulation except for finerenone.

There are several national and international guidelines that provide recommendations regarding the use of the mineralocorticoid (aldosterone) receptor antagonists.<sup>10-30</sup> For the treatment of heart failure, a mineralocorticoid (aldosterone) receptor antagonist is routinely recommended in addition to standard therapy (ACE inhibitor or ARB, and  $\beta$ -blocker) in patients with symptoms and an LVEF  $\leq 35\%$ . A mineralocorticoid (aldosterone) receptor antagonist is also recommended following a myocardial infarction in patients with an LVEF  $\leq 40\%$  who also have either diabetes or heart failure. Once again, therapy should be in addition to standard heart failure therapy (ACE inhibitor or ARB, and  $\beta$ -blocker).<sup>18-19</sup> There are several national and international organizations that have published guidelines on the treatment of hypertension. Most of the guidelines do not address the use of the mineralocorticoid (aldosterone) receptor antagonists. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>20-26</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>20</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>20-26</sup> Most patients will require more than one antihypertensive agent to achieve blood pressure goals.<sup>20-26</sup>

For the treatment of cirrhosis and ascites, spironolactone is recommended as first line therapy in addition to sodium restriction.<sup>29</sup> Spironolactone is also recommended for the treatment of patients with unilateral primary aldosteronism (in lieu of surgery) and in those with bilateral adrenal disease. Eplerenone is considered an alternative treatment option, especially in men who experience erectile dysfunction and gynecomastia with spironolactone therapy.<sup>30</sup>

For diabetic kidney disease, guidelines recommend the use of ACEIs or ARBs in patients who have CKD and albuminuria and have more recently recommended the addition of SGLT2 inhibitor. The ADA guidelines recommend a nonsteroidal mineralocorticoid receptor antagonist (finerenone) to reduce CKD progression and cardiovascular events in patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a SGLT2 inhibitor.<sup>28</sup>

Eplerenone and spironolactone have been shown to reduce cardiovascular morbidity and mortality in patients with heart failure when added to standard therapy.<sup>42-47,49-54</sup> These agents have also been shown to effectively lower blood pressure.<sup>65-86</sup> Only one trial in hypertensive patients included both eplerenone and spironolactone. Both products significantly decreased blood pressure compared to placebo; however, statistical analyses were not performed among the two agents. The authors noted that there was a greater reduction in blood pressure with spironolactone 50 mg twice daily compared to eplerenone 50 mg twice daily. This information suggests that eplerenone may only be 50 to 75% as potent as spironolactone.<sup>69</sup> Most patients will require more than one antihypertensive agent to achieve blood pressure goals.<sup>20-27</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>22-23,26,90</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations. Several studies in diabetic and non-diabetic patients with renal disease have demonstrated a reduction in proteinuria with the addition of spironolactone to existing ACE inhibitor and/or ARB therapy.<sup>31-41</sup> Studies have demonstrated efficacy of finerenone compared to placebo in reducing incidence of composite cardiovascular outcome including time to cardiovascular death, myocardial infarction, stroke, or hospitalization for heart failure, among patients with CKD and T2D.<sup>87,88</sup>

In general, adverse events are similar with the mineralocorticoid (aldosterone) receptor antagonists and both agents can increase serum potassium levels. While eplerenone is a selective aldosterone receptor antagonist, spironolactone may also antagonize glucocorticoid, progesterone, and androgen receptors. Consequently, there is an increased risk of steroid-related adverse effects with spironolactone (e.g., gynecomastia, impotence, menstrual abnormalities).<sup>1-6</sup> Finerenone is a novel, non-steroidal, selective mineralocorticoid (aldosterone) receptor antagonists that appears to be more selective for the mineralocorticoid receptor. Finerenone has shown to reduce albuminuria in T2D patients with CKD, while revealing only a low risk of hyperkalemia.<sup>7-9</sup>

There is insufficient evidence to support that one brand mineralocorticoid (aldosterone) receptor antagonist is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand mineralocorticoid (aldosterone) receptor antagonists within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand mineralocorticoid (aldosterone) receptor antagonist is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.



## XII. References

1. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
2. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
3. Aldactone® [package insert]. New York (NY): Pfizer Inc; 2021 Feb.
4. Carospir® [package insert]. Farmville (NC): CMP Pharma, Inc. 2021 Jul.
5. Inspra® [package insert]. New York (NY): Pfizer Inc; 2020 Aug.
6. Aldactazide® [package insert]. New York (NY): Pfizer Inc; 2021 Mar.
7. Kerendia® [prescribing information]. Whippany (NJ): Bayer HealthCare Pharmaceuticals Inc.; 2021 Jul.
8. Bakris GL, Agarwal R, Anker SD, Pitt B, Ruilope LM, Nowack C et al. Design and Baseline Characteristics of the Finerenone in Reducing Kidney Failure and Disease Progression in Diabetic Kidney Disease Trial. *Am J Nephrol*. 2019;50(5):333-344.
9. Perkovic V, Badve S, Bakris GL. Treatment of diabetic kidney disease. Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2022 Jan]. Available from: <http://www.uptodate.com/utd/index.do>.
10. Fraker T, Fihn S, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of chronic stable angina: a report of the American College of Cardiology/American Heart Association task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *Circulation*. 2007 Dec 4;116(23):2762-72.
11. Task Force Members, Montalescot G, Sechtem U, Achenbach S, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2019 Aug;41(3):407-477. doi:10.1093/eurheartj/ehz425.
12. Qaseem A, Fihn SD, Dallas P, Williams S, Owens DK, Shekelle P, et al. Management of Stable Ischemic Heart Disease: Summary of a Clinical Practice Guideline From the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. 2012;157:735-743. doi:10.7326/0003-4819-157-10-201211200-00011.
13. Amsterdam EA, Wenger NK, Brindis RG, Casey Jr DE, Ganiats TG, Holmes Jr DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ, 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes, *Journal of the American College of Cardiology* (2014), doi: 10.1016/j.jacc.2014.09.017.
14. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al, ESC Scientific Document Group, 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC), *European Heart Journal*, Volume 42, Issue 14, 7 April 2021, Pages 1289–1367, <https://doi.org/10.1093/eurheartj/ehaa575>.
15. O’Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *J Am Coll Cardiol*. 2012. doi:10.1016/j.jacc.2012.11.019.
16. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno Het al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2017;39:119-177.
17. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019 Mar 17. pii: S0735-1097(19)33877-X. doi: 10.1016/j.jacc.2019.03.010. [Epub ahead of print].
18. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
19. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by

- the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
20. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
  21. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, Ramirez A, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
  22. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, Butalia S, Leung AA, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
  23. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
  24. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
  25. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
  26. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
  27. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
  28. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl.1).
  29. Biggins SW, Angeli P, Garcia-Tsao G, Ginès P, Ling SC, Nadim MK, Wong F, Kim WR. Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology*. 2021 Aug;74(2):1014-1048. doi: 10.1002/hep.31884.
  30. Funder JW, Carey RM, Mantero F, Murad MH, Reincke M, Shibata H, et al. The Management of Primary Aldosteronism: Case Detection, Diagnosis, and Treatment: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2016 May;101(5):1889-916. doi: 10.1210/jc.2015-4061.
  31. Bianchi S, Bigazzi R, Campese VM. Long-term effects of spironolactone on proteinuria and kidney function in patients with chronic kidney disease. *Kidney Int*. 2006 Dec; 70(12):2116-23.
  32. Bianchi S, Bigazzi R, Campese VM. Intensive versus conventional therapy to slow the progression of idiopathic glomerular diseases. *Am J Kidney Dis* 2010;55:671-81.
  33. Ogawa S, Takeuchi K, Mori T, Nako K, Ito S. Spironolactone further reduces urinary albumin excretion and plasma B-type natriuretic peptide levels in hypertensive type II diabetes treated with angiotensin-converting enzyme inhibitor. *Clin Exp Pharmacol Physiol*. 2006 May-Jun;33(5-6):477-9.
  34. Chrysostomou A, Pedagogos E, MacGregor L, Becker GJ. Double-blind, placebo-controlled study on the effect of the aldosterone receptor antagonist spironolactone in patients who have persistent proteinuria and are on long-term angiotensin-converting enzyme inhibitor therapy, with or without an angiotensin II receptor blocker. *Clin J Am Soc Nephrol*. 2006 Mar;1(2):256-62.
  35. Furumatsu Y, Nagasawa Y, Tomida K, Mikami S, Kaneko T, Okada N, Tsubakihara Y, Imai E, Shoji T. Effect of renin-angiotensin-aldosterone system triple blockade on non-diabetic renal disease: addition of an aldosterone blocker, spironolactone, to combination treatment with an angiotensin-converting enzyme inhibitor and angiotensin II receptor blocker. *Hypertens Res*. 2008 Jan;31(1):59-67.
  36. van den Meiracker AH, Baggen RG, Pauli S, Lindemans A, Vulto AG, Poldermans D, Boomsma F. Spironolactone in type 2 diabetic nephropathy: Effects on proteinuria, blood pressure and renal function. *J Hypertens*. 2006 Nov;24(11):2285-92.

37. Schjoedt KJ, Rossing K, Juhl TR, Boomsma F, Tarnow L, Rossing P, Parving HH. Beneficial impact of spironolactone on nephrotic range albuminuria in diabetic nephropathy. *Kidney Int.* 2006 Aug;70(3):536-42.
38. Davidson MB, Wong A, Hamrahian AH, et al. Effect of spironolactone therapy on albuminuria in patients with type 2 diabetes treated with angiotensin-converting enzyme inhibitors. *Endocr Pract* 2008;14:985-92.
39. Saklayen MG, Gyebi LK, Tasosa J, et al. Effects of additive therapy with spironolactone on proteinuria in diabetic patients already on ACE inhibitor or ARB therapy: results of a randomized, placebo-controlled, double-blind, crossover trial. *J Investig Med* 2008;56:714-9.
40. Sengul E, Sahin T, Sevin E, et al. Effect of spironolactone on urinary protein excretion in patients with chronic kidney disease. *Ren Fail* 2009;31:928-32.
41. Tylicki L, Rutkowski P, Renke M, et al. Triple pharmacological blockade of the renin-angiotensin-aldosterone system in nondiabetic CKD: an open-label crossover randomized controlled trial. *Am J Kidney Dis* 2008;52:486-93.
42. Pitt B, Remme W, Zannad F et al. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med.* 2003;348(14):1309-21.
43. Pitt B, White H, Nicolau J et al. Eplerenone reduces mortality 30 days after randomization following acute myocardial infarction in patients with left ventricular systolic dysfunction and heart failure. *J Am Coll Cardiol.* 2005;46(3):425-31.
44. Pitt B, Gheorghide M, Zannad F, Anderson JL, van Veldhuisen DJ, Parkhomenko A, Corbalan R, Klug EQ, Mukherjee R, Solomon H; EPHEUS Investigators. Evaluation of eplerenone in the subgroup of EPHEUS patients with baseline left ventricular ejection fraction  $\leq 30\%$ . *Eur J Heart Fail* 2006;8:295-301.
45. O'Keefe JH, Abuissa H, Pitt B. Eplerenone improves prognosis in postmyocardial infarction diabetic patients with heart failure: results from EPHEUS. *Diabetes Obes Metab.* 2007 May 8; [Epub ahead of print].
46. Gheorghide M, Khan S, Blair JE, et al. The effects of eplerenone on length of stay and total days of heart failure hospitalization after myocardial infarction in patients with left ventricular systolic dysfunction (EPHEUS). *Am Heart J* 2009;158:437-43.
47. Adamopoulos C, Ahmed A, Fay R, et al. Timing of eplerenone initiation and outcomes in patients with heart failure after acute myocardial infarction complicated by left ventricular systolic dysfunction: insights from the EPHEUS trial. *Eur J Heart Fail* 2009;11:1099-105.
48. Udelson JE, Feldman AM, Greenberg B, et al. Randomized, double-blind, multicenter, placebo-controlled study evaluating the effect of aldosterone antagonism with eplerenone on ventricular remodeling in patients with mild-to-moderate heart failure and left ventricular systolic dysfunction. *Circ Heart Fail* 2010;3:347-53.
49. Zannad F, McMurray JJV, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med.* 2011;364:11-21.
50. Eschaliel R, McMurray JJ, Swedberg K, et al. Safety and efficacy of eplerenone in patients at high risk for hyperkalemia and/or worsening renal function: analyses of the EMPHASIS-HF study subgroups (Eplerenone in Mild Patients Hospitalization And Survival Study in Heart Failure). *J Am Coll Cardiol.* 2013 Oct 22;62(17):1585-93.
51. Krum H, Shi H, Pitt B, et al. Clinical benefit of eplerenone in patients with mild symptoms of systolic heart failure already receiving optimal best practice background drug therapy: analysis of the EMPHASIS-HF study. *Circ Heart Fail.* 2013 Jul;6(4):711-8.
52. Girerd N, Collier T, Pocock S, Krum H, McMurray JJ, Swedberg K, et al. Clinical benefits of eplerenone in patients with systolic heart failure and mild symptoms when initiated shortly after hospital discharge: analysis from the EMPHASIS-HF trial.
53. Pitt B, Zannad F, Remme WJ et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med.* 1999;341(10):709-17.
54. Vardeny O, Wu DH, Desai A, Rossignol P, Zannad F, Pitt B, et al. Influence of baseline and worsening renal function on efficacy of spironolactone in patients with severe heart failure. *J Am Coll Cardiol.* 2012;60(20):2089-9.
55. Vizzardi E, D'Aloia A, Giubbini R, Bordonali T, Bugatti S, Pezzali N, et al. Effect of spironolactone on left ventricular ejection fraction and volumes in patients with class I or II heart failure. *Am J Cardiol.* 2010;106:1292-6.
56. Chan AK, Sanderson JE, Wang T, Lam W, Yip G, Wang M, Lam YY, Zhang Y, Yeung L, Wu EB, Chan WW, Wong JT, So N, Yu CM. Aldosterone receptor antagonism induces reverse remodeling when added to angiotensin receptor blockade in chronic heart failure. *J Am Coll Cardiol.* 2007 Aug 14;50(7):591-6.
57. Edelmann F, Wachter R, Schmidt AG, Kraigher-Krainer E, Colantonio C, Kamke W, et al. Effect of spironolactone on diastolic function and exercise capacity in patients with heart failure with preserved ejection fraction. *JAMA.* 2013;309(8):781-791.

58. Pitt B, Pfeffer MA, Assmann SF, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med.* 2014 Apr 10;370(15):1383-92.
59. Levy B. The efficacy of safety of furosemide and a combination of spironolactone and hydrochlorothiazide in congestive heart failure. *J Clin Pharmacol.* 1977;17(7):420-30.
60. Lee KK, Shilane D, Hlatky MA, Yang J, Steimle AE, and Go AS. Effectiveness and safety of spironolactone for systolic heart failure. *Am J Cardiol.* 2013 Nov 1;112(9):1427-32.
61. Inampudi C, Parvataneni S, Morgan CJ, et al. Spironolactone use and higher hospital readmission for Medicare beneficiaries with heart failure, left ventricular ejection fraction <45%, and estimated glomerular filtration rate <45 ml/min/1.73 m<sup>2</sup>. *Am J Cardiol.* 2014 Jul 1;114(1):79-82.
62. Maisel A, Xue Y, van Veldhuisen DJ, et al. Effect of spironolactone on 30-day death and heart failure rehospitalization (from the COACH Study). *Am J Cardiol.* 2014 Sep 1;114(5):737-42.
63. Karagiannis A, Tziomalos K, Papageorgiou A, Kakafika AI, Pagourelis ED, Anagnostis P, Athyros VG, Mikhailidis DP. Spironolactone versus eplerenone for the treatment of idiopathic hyperaldosteronism. *Expert Opin Pharmacother.* 2008 Mar;9(4):509-15.
64. Karashima S, Yoneda T, Kometani M, Ohe M, Mori S, Sawamura T, et al. Comparison of eplerenone and spironolactone for the treatment of primary aldosteronism. *Hypertens Res.* 2016 Mar;39(3):133-7.
65. Kohvakka A, Eisalo A, Manninen V. Maintenance of potassium balance during diuretic therapy. *Acta Med Scand.* 1979;205(4):319-24.
66. Dahlöf B, Lindholm LH, Hansson L, Scherstén B, Ekblom T, Wester PO. Morbidity and mortality in the Swedish Trial in Old Patients with Hypertension (STOP-Hypertension). *Lancet.* 1991 Nov 23;338(8778):1281-5.
67. White WB, Carr AA, Krause S et al. Assessment of the novel selective aldosterone blocker eplerenone using ambulatory and clinical blood pressure in patients with systemic hypertension. *Am J Cardiol.* 2003;92(1):38-42.
68. Krum, Nolly H, Workman D et al. Efficacy of eplerenone added to renin-angiotensin blockade in hypertensive patients. *Hypertension* 2002;40(2):117-23.
69. Weinberger MH, Roniker B, Krause SL et al. Eplerenone, a Selective Aldosterone Blocker, in Mild-to-Moderate Hypertension. *Am Journal Hypertens.* 2002;15(8):709-16.
70. Hollenberg HK, Williams GH, Anderson H et al. Symptoms and the distress they cause: comparison of an aldosterone antagonist and a calcium-channel blocking agent in patients with systolic hypertension. *Arch Intern Med.* 2003;163(13):1543-8.
71. White WB, Duprez D, St Hillaire R et al. Effects of the selective aldosterone blocker eplerenone versus the calcium antagonist amlodipine in systolic hypertension. *Hypertension* 2003;41(5):1021-6.
72. Williams GH, Burgess E, Kolloch RE et al. Efficacy of eplerenone versus enalapril as monotherapy in systemic hypertension. *Am J of Cardiol.* 2004;93(8):990-6.
73. Flack JM, Oparil S, Pratt JH et al. Efficacy and tolerability of eplerenone and losartan in hypertensive black and white patients. *Journal of the American College of Cardiology* 2003;41(7):1148-55.
74. Hanazawa T, Obara T, Ogasawara K, Shinki T, Katada S, Inaue R, et al. Low-dose and very low-dose spironolactone in combination therapy for essential hypertension: evaluation by self-measurement of blood pressure at home (abstract). *Clin Exp Hypertens.* 2011;33(7):427-36.
75. Schersten B, Thulin T, Kuylensstierna J et al. Clinical and biochemical effects of spironolactone administered once daily in primary hypertension, Multicenter Sweden study. *Hypertension* 1980;2(5):672-79.
76. Oxlund CS, Henriksen JE, Tarnow L, Schousboe K, Gram J, and Jacobsen IA. Low dose spironolactone reduces blood pressure in patients with resistant hypertension and type 2 diabetes mellitus: a double blind randomized clinical trial. *J Hypertens.* 2013 Oct;31(10):2094-102.
77. Václavík J, Sedlák R, Jarkovský J, Kociánová E, and Táborský M. Effect of spironolactone in resistant arterial hypertension: a randomized, double-blind, placebo-controlled trial (ASPIRANT-EXT). *Medicine (Baltimore).* 2014 Dec;93(27):e162.
78. Li JS, Flynn JT, Portman R, et al. The efficacy and safety of the novel aldosterone antagonist eplerenone in children with hypertension: a randomized, double-blind, dose-response study. *J Pediatr* 2010;157:282-7.
79. Hood SJ, Taylor KP, Ashby MJ, Brown MJ. The spironolactone, amiloride, losartan, and thiazide (SALT) double-blind crossover trial in patients with low-renin hypertension and elevated aldosterone-renin ratio. *Circulation.* 2007 Jul 17;116(3):268-75.
80. Nash DT. Antihypertensive effect and serum potassium homeostasis: comparison of hydrochlorothiazide and spironolactone alone and in combination. *J Med.* 1977;8(5):367-77.
81. Schrijver G, Weinberger MH. Hydrochlorothiazide and spironolactone in hypertension. *Clin Pharmacol Ther.* 1979;25(1):33-42.

82. Wray DW, Supiano MA. Impact of aldosterone receptor blockade compared with thiazide therapy on sympathetic nervous system function in geriatric hypertension. *Hypertension* 2010;55:1217-23.
83. Krieger EM, Drager LF, Giorgi DMA, Pereira AC, Barreto-Filho JAS, Nogueira AR, et al. Spironolactone Versus Clonidine as a Fourth-Drug Therapy for Resistant Hypertension: The ReHOT Randomized Study (Resistant Hypertension Optimal Treatment). *Hypertension*. 2018 Apr;71(4):681-690.
84. Bomback AS, Muskala P, Bald E, et al. Low-dose spironolactone, added to long-term ACE inhibitor therapy, reduces blood pressure and urinary albumin excretion in obese patients with hypertensive target organ damage. *Clin Nephrol* 2009;72:449-56.
85. Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G, et al. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial.
86. Chapman N, Dobson J, Wilson S, Dahlöf B, Sever PS, Wedel H, Poulter NR; Anglo-Scandinavian Cardiac Outcomes Trial Investigators. Effect of spironolactone on blood pressure in subjects with resistant hypertension. *Hypertension*. 2007 Apr;49(4):839-45.
87. Filippatos G, Anker SD, Agarwal R, Pitt B, Ruilope LM, Rossing P, et al. Finerenone and Cardiovascular Outcomes in Patients With Chronic Kidney Disease and Type 2 Diabetes. *Circulation*. 2021 Feb 9;143(6):540-552.
88. Pitt B, Filippatos G, Agarwal R, Anker SD, Bakris GL, Rossing P et al. Cardiovascular Events with Finerenone in Kidney Disease and Type 2 Diabetes. *N Engl J Med*. 2021 Dec 9;385(24):2252-2263. doi: 10.1056/NEJMoa2110956. Epub 2021 Aug 28.
89. Pitt B, Reichek N, Willenbrock R et al. Effects of eplerenone, enalapril, and eplerenone/enalapril in patients with essential hypertension and left ventricular hypertrophy: the 4E-left ventricular hypertrophy study. *Circulation* 2003;108(15):1831-8.
90. Taniguchi I, Kawai M, Date T, Yoshida S, Seki S, Taniguchi M, Shimizu M, Mochizuki S. Effects of spironolactone during an angiotensin II receptor blocker treatment on the left ventricular mass reduction in hypertensive patients with concentric left ventricular hypertrophy. *Circ J*. 2006 Aug; 70(8):995-1000.
91. Edwards NC, Steeds RP, Stewart PM, et al. Effect of spironolactone on left ventricular mass and aortic stiffness in early-stage chronic kidney disease: a randomized controlled trial. *J Am Coll Cardiol* 2009;54:505-12.
92. Ludbrook A, Dyonon M, Mendelsohn FA, Louis WJ. Comparison of a single-dose and twice-a-day spironolactone therapy in mild hypertension. *Med J Aust*. 1980 Feb 9;1(3):124-5.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Renin Inhibitors  
AHFS Class 243240  
May 4, 2022**

**I. Overview**

The renin-angiotensin-aldosterone system (RAAS) is the most important component in the homeostatic regulation of blood pressure.<sup>1</sup> Excessive activity of the RAAS may lead to hypertension, as well as fluid and electrolyte disorders. Renin catalyzes the conversion of angiotensinogen to angiotensin I, which is the first and rate-limiting step of the RAAS.<sup>1-3</sup> Angiotensin I is then cleaved to angiotensin II by angiotensin-converting enzyme (ACE). Angiotensin II may also be generated through other pathways (angiotensin I convertase). Through a negative feedback mechanism, angiotensin II inhibits renin release. Angiotensin II can increase blood pressure by direct vasoconstriction, as well as through actions on the brain and autonomic nervous system. In addition, angiotensin II induces aldosterone synthesis from the adrenal cortex, leading to sodium and water reabsorption. Angiotensin II exerts other detrimental effects, including ventricular hypertrophy, remodeling and myocyte apoptosis.<sup>4,5</sup>

Aliskiren is the only renin inhibitor that is currently available and it is approved for the treatment of hypertension. It decreases plasma renin activity and inhibits the conversion of angiotensinogen to angiotensin I. It is unknown if aliskiren affects other RAAS components, such as ACE or non-ACE pathways. Aliskiren is available as a single entity product, as well as in combination with hydrochlorothiazide.<sup>6-7</sup> Previously, aliskiren was available in combination with valsartan under the name Valtorna®; however, this agent was removed from the market in 2012 due to evidence suggesting an increased risk of renal impairment, hypotension, and hyperkalemia in patients taking aliskiren and ACE inhibitors or ARBs concomitantly.<sup>8</sup> Combination products containing aliskiren and amlodipine, with or without hydrochlorothiazide, are also no longer available.

The renin inhibitors that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. Aliskiren is available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Renin Inhibitors Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Aliskiren	tablet	Tekturna®*	aliskiren
<b>Combination Products</b>			
Aliskiren and hydrochlorothiazide	tablet	Tekturna HCT®	none

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the renin inhibitors are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Renin Inhibitors**

Clinical Guideline	Recommendations
Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b> <sup>9</sup>	<ul style="list-style-type: none"> <li>Pharmacologic treatment should be initiated in patients <math>\geq 60</math> years of age to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 150</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>In patients <math>&lt; 60</math> years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure <math>\geq 90</math> mm Hg to a goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 150</math> mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>10</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment               <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment               <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy               <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>11</sup></p>	<p><b>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</b></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk</li> </ul>



Clinical Guideline	Recommendations
	<p>factors.</p> <ul style="list-style-type: none"> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq 130</math> mmHg, intensive management to target a SBP <math>&lt; 120</math> mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt; 60</math> years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in</li> </ul>

Clinical Guideline	Recommendations
	<p>combination therapy.</p> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt;40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (<math>&lt;40\%</math>) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium <math>&lt;5.2</math> mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul>

Clinical Guideline	Recommendations
	<p><b>Treatment of hypertension in association with stroke</b></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours)               <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke               <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours)               <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><b>Treatment of hypertension in association with LVH</b></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><b>Treatment of hypertension in association with nondiabetic chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><b>Treatment of hypertension in association with renovascular disease</b></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amenable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><b>Treatment of hypertension in association with diabetes mellitus</b></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>12</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt; 150</math> mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or</li> </ul>

Clinical Guideline	Recommendations
	<p>mineralocorticoid receptor antagonist (MRA) if required.</p> <ul style="list-style-type: none"> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>13</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> </ul>



Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on “Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence”.</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African–Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person’s medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in</b></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however,</li> </ul>

Clinical Guideline	Recommendations
<p><b>Blacks (2010)<sup>14</sup></b></p>	<p>the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</p> <ul style="list-style-type: none"> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p><b>Kidney Disease Improving Clinical Outcomes Group; KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)<sup>15</sup></b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130</li> </ul>

Clinical Guideline	Recommendations
	<p>mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</p> <ul style="list-style-type: none"> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>16</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HF<sub>r</sub>EF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HF<sub>r</sub>EF.</li> <li>• In adults with HF<sub>p</sub>EF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HF<sub>p</sub>EF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be</li> </ul>

Clinical Guideline	Recommendations
	<p>prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</p> <ul style="list-style-type: none"> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> <li>• For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of</li> </ul>

Clinical Guideline	Recommendations
	<p>comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</p> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP <math>\geq</math>180 mmHg or DBP <math>\geq</math>110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>17</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq</math>140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq</math>180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq</math>15%), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure</li> </ul>

Clinical Guideline	Recommendations
	<p>target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</p> <ul style="list-style-type: none"> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq</math>300 mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq</math>300 mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq</math>300 mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> or <math>\geq</math>300 mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> </ul>

Clinical Guideline	Recommendations
	<ul style="list-style-type: none"> <li>Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the renin inhibitors are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Renin Inhibitors<sup>4,7</sup>**

Indication(s)	Single Entity Agents	Combination Products
	Aliskiren	Aliskiren and HCTZ
<b>Hypertension</b>		
Treatment of hypertension	✓	✓

HCTZ=hydrochlorothiazide

### IV. Pharmacokinetics

The pharmacokinetic parameters of the renin inhibitors are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Renin Inhibitors<sup>5</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Aliskiren	2.5	47 to 51	Liver, minor (%) not reported)	Feces (91) Renal (<1)	40



Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Combination Products</b>					
Aliskiren and HCTZ	2.5/50 to 75	47 to 51/ 40 to 68	Liver, minor (% not reported)/ Not reported	Feces (91) Renal (<1)/ Renal (>95)	40/6 to 15

HCTZ=hydrochlorothiazide

## V. Drug Interactions

Major drug interactions with the renin inhibitors are listed in Table 5.

**Table 5. Major Drug Interactions with the Renin Inhibitors<sup>5</sup>**

Generic Name(s)	Interaction	Mechanism
Renin Inhibitors (aliskiren)	ACE inhibitors	Aliskiren is contraindicated in patients with diabetes who are receiving ARBs or ACEIs because of the increased risk of renal impairment, hyperkalemia, and hypotension. In general, avoid combined use of aliskiren with ACE inhibitors or ARBs, particularly in patients with creatinine clearance (CrCl) <60 mL/min.
Renin Inhibitors (aliskiren)	ARBs	Aliskiren is contraindicated in patients with diabetes who are receiving ARBs or ACEIs because of the increased risk of renal impairment, hyperkalemia, and hypotension. In general, avoid combined use of aliskiren with ACE inhibitors or ARBs, particularly in patients with CrCl <60 mL/min.
Renin Inhibitors (aliskiren)	Azole antifungals	Increased absorption of aliskiren resulting from inhibition of P-gp expression by certain azole antifungal agents may occur. In addition, azole antifungal agents may inhibit aliskiren metabolism (CYP3A4).
Renin Inhibitors (aliskiren)	Cyclosporine	Concurrent use of aliskiren and cyclosporine may result in increased aliskiren exposure and plasma concentrations.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics may induce hypokalemia and increase the risk of torsades de pointes.
Thiazide diuretics (HCTZ)	Lithium	Decreased lithium clearance may occur with thiazide use, which may lead to increased serum lithium levels and possibly lithium toxicity.
Thiazide diuretics (HCTZ)	Diazoxide	The combination of diazoxide with a thiazide diuretic may lead to hyperglycemia, hyperuricemia, and hypotension.
Thiazide diuretics (HCTZ)	Digitalis glycosides	Thiazide diuretics may induce electrolyte disturbances which may predispose patients to digitalis-induced arrhythmias.

ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor antagonist

## VI. Adverse Drug Events

The most common adverse drug events reported with the renin inhibitors are listed in Table 6. The boxed warning for aliskiren-containing products is listed in Table 7.

**Table 6. Adverse Drug Events (%) Reported with the Renin Inhibitors<sup>4-7</sup>**

Adverse Events	Single Entity Agents	Combination Products
	Aliskiren	Aliskiren and HCTZ
<b>Cardiovascular</b>		
Hypotension	<1	<1
Peripheral edema	✓	✓
<b>Central Nervous System</b>		
Dizziness	>1	2
Fatigue	>1	>1
Headache	>1	>1
Restlessness	-	✓
Seizure	✓	✓
Vertigo	1	1
<b>Dermatologic</b>		
Erythema multiforme	-	✓
Exfoliative dermatitis	-	✓
Photosensitivity	-	✓
Rash	1	1
Stevens-Johnson syndrome	-	✓
Toxic epidermal necrolysis	-	✓
Urticaria	-	✓
<b>Endocrine and Metabolic</b>		
Gout	<1	<1
<b>Gastrointestinal</b>		
Abdominal pain	✓	✓
Cramping	-	✓
Diarrhea	2	2
Dyspepsia	✓	✓
Gastric irritation	-	✓
Gastroesophageal reflux	✓	✓
<b>Genitourinary</b>		
Glycosuria	-	✓
<b>Hematologic</b>		
Agranulocytosis	-	✓
ALT increased	-	1
Anemia	✓	-
Aplastic anemia	-	✓
Hematocrit decreased	✓	✓
Hemoglobin decreased	✓	✓
Hemolytic anemia	-	✓
Leukopenia	-	✓
Thrombocytopenia	-	✓
<b>Laboratory Test Abnormalities</b>		
Alanine aminotransaminase increased	-	1
Blood urea nitrogen increased	7	12
Creatine kinase increased	1	-
Hyperglycemia	-	✓

Adverse Events	Single Entity Agents	Combination Products
	Aliskiren	Aliskiren and HCTZ
Hyperkalemia	1	1
Hypokalemia	-	2
Serum creatinine increased	7	1
Uric acid increased	<1	<1
<b>Musculoskeletal</b>		
Arthralgia	-	1
Asthenia	-	1
Back pain	>1	>1
Muscle cramps	-	-
Muscle spasm	-	✓
Myositis	<1	-
Rhabdomyolysis	<1	-
Weakness	-	✓
<b>Renal</b>		
Interstitial nephritis	-	✓
Renal dysfunction	-	✓
Renal failure	-	✓
Renal stones	<1	<1
<b>Respiratory</b>		
Cough	1	1
Influenza	-	2
Nasopharyngitis	✓	>1
Respiratory distress	-	✓
Sinusitis	-	-
Upper respiratory infection	>1	>1
<b>Other</b>		
Allergic reaction	-	-
Angioedema	✓	✓
Blurred vision	-	✓
Edema (face, hands, or whole body)	<1	<1
Fever	-	✓
Jaundice	-	✓
Necrotizing angitis	-	✓
Pancreatitis	-	✓
Periorbital edema	✓	✓
Purpura	-	✓
Xanthopsia	-	✓

✓ Percent not specified.

- Event not reported.

**Table 7. Boxed Warning for Aliskiren Products<sup>4</sup>**

<b>WARNING</b>
When pregnancy is detected, discontinue aliskiren as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

## VII. Dosing and Administration

The usual dosing regimens for the renin inhibitors are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Renin Inhibitors<sup>4-7</sup>**

<b>Generic Name(s)</b>	<b>Usual Adult Dose</b>	<b>Usual Pediatric Dose</b>	<b>Availability</b>
<b>Single Entity Agents</b>			
Aliskiren	<u>Hypertension:</u> Tablet: initial, 150 mg once daily; maintenance, titrate as needed; maximum, 300 mg/day	<u>Hypertension in pediatric patients 6 to 17 years of age:</u> Tablet: 20 to 50 kg, initial, 75 mg once daily; maximum, 150 mg daily; ≥50 kg, follow adult dosing	Tablet: 150 mg 300 mg
<b>Combination Products</b>			
Aliskiren and HCTZ	<u>Hypertension:</u> Tablet: initial, 150-12.5 mg once daily; maximum, 300-25 mg/day	Safety and efficacy in children have not been established.	Tablet: 150-12.5 mg 150-25 mg 300-12.5 mg 300-25 mg

HCTZ=hydrochlorothiazide

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the renin inhibitors are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Renin Inhibitors**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Cardiovascular Risk Reduction</b>				
Parving et al. <sup>18</sup> (2012) ALTITUDE  Aliskiren  vs  placebo  Both in addition to standard treatment	DB, PC, RCT  Patients ≥35 years of age with type 2 diabetes and evidence of microalbuminuria, macroalbuminuria, or cardiovascular disease	N=8,561  Median of 32.9 months	Primary: Composite of death from cardiovascular causes or the first occurrence of cardiac arrest with resuscitation; nonfatal MI; nonfatal stroke; unplanned hospitalization for HF; end-stage renal disease, death attributable to kidney failure, or the need for renal- replacement therapy with no dialysis or transplantation available or initiated; or a serum creatinine value that was at least double the baseline value and that exceeded the upper limit of the normal range  Secondary: Composite of cardiovascular	The independent data and safety monitoring committee recommended termination of the study medication, on the basis of their assessment that the excess risk of adverse events in the aliskiren group could not be offset by a reduction in major cardiovascular and renal events.  Primary: The primary outcome occurred in 783 participants in the aliskiren group (18.3%) and 732 in the placebo group (17.1%). The hazard ratio for this outcome in the aliskiren group as compared with the placebo group was 1.08 (95% CI, 0.98 to 1.20; P=0.12).  Secondary: The secondary cardiovascular composite outcome occurred in 590 participants in the aliskiren group (13.8%) and 539 in the placebo group (12.6%); the HR in the aliskiren group was 1.11 (95% CI, 0.99 to 1.25; P=0.09).  The secondary renal composite outcome occurred in 257 participants in the aliskiren group (6.0%) and 251 in the placebo group (5.9%); the HR in the aliskiren group was 1.03 (95% CI, 0.87 to 1.23; P=0.74).  The number of deaths from any cause did not differ significantly between the study groups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			components and of renal components of the primary composite end point	
<p>McMurray et al.<sup>19</sup> (2016) ATMOSPHERE</p> <p>Enalapril 5 or 10 mg BID</p> <p>vs</p> <p>aliskiren 150 mg QD</p> <p>vs</p> <p>combination of aliskiren 150 mg QD and enalapril 5 or 10 mg BID</p>	<p>DB, DD, RCT</p> <p>Patients with CHF (NYHA class II to IV) and EF <math>\leq</math>35% receiving stable doses of an ACE inhibitor (equivalent to at least 10 mg of enalapril daily) and of a <math>\beta</math>-blocker at the time of enrollment</p>	<p>N=7,016</p> <p>Median of 36.6 months</p>	<p>Primary: Composite of death from cardiovascular causes or hospitalization for heart failure</p> <p>Secondary: Change from baseline to 12 months in the Kansas City Cardiomyopathy Questionnaire (KCCQ) clinical summary score</p>	<p>Primary: Overall, the primary outcome occurred in 770 patients (32.9%) in the combination-therapy group (11.7 events per 100 person-years), in 791 patients (33.8%) in the aliskiren group (12.1 events per 100 person-years), and in 808 patients (34.6%) in the enalapril group (12.4 events per 100 person-years). The hazard ratio in the combination-therapy group, as compared with the enalapril group, was 0.93 (95% CI, 0.85 to 1.03; P=0.17); the hazard ratio in the aliskiren group, as compared with the enalapril group, was 0.99 (95% CI, 0.90 to 1.10; P=0.91 for superiority). Although the noninferiority margin of 1.104 was met with the use of the 95% confidence interval, the one-sided P value of 0.0184 did not fulfill the prespecified requirement of a P value of 0.0123 or less. A sensitivity analysis that included only patients who received the assigned trial regimen gave consistent results, as did an analysis in which data that were collected after regulatory censoring were included.</p> <p>Secondary: There were no significant between-group differences in the secondary outcome. The exploratory composite renal outcome (the composite of death from renal causes, end-stage renal disease, or doubling of the serum creatinine level) occurred significantly more frequently in the combination-therapy group than in the enalapril group.</p>
<b>Hypertension</b>				
<p>Tocci et al.<sup>20</sup> (abstract) 2012</p> <p>Aliskiren 150 to 300 mg/day</p>	<p>MC, OL, OS, PRO</p> <p>Patient with HTN not adequately controlled on <math>\geq</math>2 other antihypertensive agents</p>	<p>N=1,186</p> <p>12 months</p>	<p>Primary: Efficacy, safety</p> <p>Secondary: Not reported</p>	<p>Primary: SBP and DBP was 141.1/82.4, 134.9/79.8, and 133.6/78.9 mmHg at one, six and 12 month follow-up visits, respectively (P&lt;0.0001 vs baseline for all comparisons). These effects were consistent in all predefined subgroups, including those with left ventricular hypertrophy, renal disease, diabetes mellitus, CAD, or cerebrovascular disease.</p> <p>Reduced levels of microalbuminuria were reported, without affecting other renal and electrolyte parameters.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Oh et al. <sup>21</sup> (2007)  Aliskiren 150, 300, or 600 mg QD  vs  placebo	DB, MC, PC, PG, RCT  Men and women ≥18 years (mean age 53 years) with mild-to-moderate essential HTN (DBP ≥95 and <110 mm Hg)	N=672  8 weeks	Primary: Change in mean sitting DBP  Secondary: Change in mean sitting SBP, 24-hour ABPM, proportion achieving a successful treatment response (DBP <90 mm Hg or ≥10 mm Hg pressure reduction from baseline) or blood pressure control (<140/90 mm Hg), plasma renin activity and concentration, safety and tolerability	Primary: All three doses investigated provided significantly greater reductions in mean sitting DBP from baseline compared to placebo (P<0.0001 for all). The mean sitting DBP reductions were 10.3 mm Hg with 150 mg, 11.1 mm Hg with 300 mg and 12.5 mm Hg with 600 mg compared to 4.9 mm Hg with placebo.  Secondary: All three doses provided significantly greater reductions in mean sitting SBP from baseline compared to placebo (P<0.0001 for all). The mean sitting SBP reductions were 13.0 mm Hg with 150 mg, 14.7 mm Hg with 300 mg and 15.8 mm Hg with 600 mg compared to 3.8 mm Hg with placebo.  Reduction in the 24-hour ABPM was significantly greater in all doses of aliskiren compared to placebo (n=216; P<0.0001 for all). Reductions in mean ambulatory DBP and SBP were consistent across the 24-hour dosing interval with all aliskiren doses.  The proportion of patients achieving a successful treatment response was 59.3% with aliskiren 150 mg, 63.3% with 300 mg and 69.3% with 600 mg compared to 36.2% with placebo (P<0.0001 for all).  The proportion of patients achieving blood pressure control was 35.9% with 150 mg, 41.6% with 300 mg and 46.4% with 600 mg compared to 20.3% with placebo (P<0.0001 for all).  Plasma renin activity decreased 79.5% with 150 mg, 81.1% with 300 mg and 75.0% with 600 mg compared to an increase of 19.5% with placebo. Aliskiren resulted in dose-dependent increases from baseline in renin concentrations (51.5%, 101.6%, and 228.5% for 150, 300 and 600 mg, respectively). Renin concentrations were almost unchanged with placebo.
Kushiro et al. <sup>22</sup> (2006)	DB, MC, PC, PG, RCT	N=455  8 weeks	Primary: Change in mean sitting DBP	Primary: All three aliskiren doses provided significantly greater reductions in mean sitting DBP from baseline compared to placebo. The placebo-corrected

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Aliskiren 75, 150, or 300 mg QD</p> <p>vs</p> <p>placebo</p>	<p>Japanese men and women between the ages of 20 and 80 years with essential HTN (mean sitting DBP of <math>\geq 90</math> mm Hg and <math>&lt; 110</math> mm Hg during the run-in period and <math>\geq 95</math> mm Hg and <math>&lt; 110</math> mm Hg at baseline)</p>		<p>Secondary: Change in mean trough sitting SBP, proportion of patients responding to treatment (mean sitting DBP <math>&lt; 90</math> mm Hg and/or <math>\geq 10</math> mm Hg decrease in mean sitting DBP from baseline), dose-response relationship, safety</p>	<p>reductions in mean sitting DBP were 4.0 mm Hg with 75 mg aliskiren, 4.5 mm Hg with 150 mg and 7.5 mm Hg with 300 mg (<math>P &lt; 0.0005</math> for all).</p> <p>Secondary: The mean sitting SBP reductions were significantly lower with all aliskiren doses when compared to placebo. The placebo-corrected reductions in mean sitting SBP were 5.7 mm Hg with 75 mg aliskiren, 5.9 mm Hg with 150 mg and 11.2 mm Hg with 300 mg (<math>P &lt; 0.001</math> for all).</p> <p>The proportion of responders at study end point was 47.8% with aliskiren 75 mg, 48.2% with 150 mg and 63.7% with 300 mg compared to 27.8% with placebo (<math>P &lt; 0.005</math> for all).</p> <p>Dose-response analysis showed that the relationship between reductions in mean sitting DBP and SBP and aliskiren dose was almost linear. However, further analyses revealed that a pattern of similar reductions with aliskiren 75 and 150 mg and greater reductions with aliskiren 300 mg was a better fit for both mean sitting DBP and SBP.</p> <p>The incidence of drug-related adverse events was comparable between aliskiren (53 to 55%) and placebo (50%). There was no evidence of a dose-dependent increase in the incidence of all-causality adverse events at the aliskiren doses evaluated in this study.</p>
<p>Musini et al.<sup>23</sup> (2009)</p> <p>Aliskiren (variable doses)</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients <math>\geq 18</math> years of age with mild to moderate essential HTN (defined as mean sitting DBP <math>\leq 95</math> mm Hg and <math>\leq 110</math> mm Hg at baseline)</p>	<p>N=3,694 (6 trials)</p> <p>8 weeks</p>	<p>Primary: Changes in dose-related SBP and DBP</p> <p>Secondary: Variability of blood pressure, pulse pressure, heart rate, withdrawals due to adverse effects, and rates of specific adverse effects</p>	<p>Primary: Aliskiren monotherapy was more effective than placebo in lowering mean sitting SBP. The additional magnitude of blood pressure lowering minus the placebo effect: aliskiren 75 mg vs placebo -2.94 (95% CI, -4.56 to -1.31); aliskiren 150 mg vs placebo -5.45 (95% CI, -6.46 to -4.43); aliskiren 300 mg vs placebo -8.66 (95% CI, -9.68 to 7.64); aliskiren 600 mg vs placebo -11.36 (95% CI, -13.53 to -9.19).</p> <p>Aliskiren monotherapy was more effective than placebo in lowering mean sitting DBP. The additional magnitude of blood pressure lowering minus the placebo effect: aliskiren 75 mg vs placebo -2.29 (95% CI, -3.31 to -1.26); aliskiren 150 mg vs placebo -3.00 (95% CI, -3.65 to -2.34); aliskiren 300 mg vs placebo -4.97 (95% CI, -5.62 to -4.31); aliskiren 600 mg vs placebo -6.57 (95% CI, -7.92 to -5.23).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Secondary: No trials reported on pulse pressure at baseline or end point. Two trials recorded baseline heart rate, but no data were provided at week eight.</p> <p>There were no significant differences in withdrawals between placebo and aliskiren at any dose. The relative risk for aliskiren 75 mg vs placebo was 0.97 (95% CI, 0.49 to 1.89); for aliskiren 150 mg vs placebo was 1.01 (95% CI, 0.61 to 1.69); for aliskiren 300 mg vs placebo was 0.91 (95% CI, 0.57 to 1.47) and for aliskiren 600 mg vs placebo was 0.63 (9% CI, 0.21 to 1.89).</p> <p>One trial reported on the incidence of dry cough: placebo (1.1%); aliskiren 75 mg (1.1%); aliskiren 150 mg (2.8%); aliskiren 300 mg (0.6%). No trials reported angioedema.</p> <p>The blood pressure lowering efficacy of aliskiren 150 mg vs 75 mg, as well as aliskiren 600 mg vs 300 mg was not significantly different. Aliskiren 300 mg significantly lowered both SBP and DBP as compared to 150 mg (SBP: -2.97; 95% CI, -3.99 to -1.95; DBP: -1.66; 95% CI, -2.32 to -1.0).</p>
<p>Braun-Dullaes et al.<sup>24</sup> (2012)</p> <p>Aliskiren 150 mg QD, up titrated to 300 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patients received amlodipine 5 mg/day, up titrated</p>	<p>DB, MC, RCT</p> <p>Patients with HTN (mean sitting SBP <math>\geq</math>160 to &lt;200 mm Hg)</p>	<p>N=485</p> <p>8 weeks</p>	<p>Primary: Change in baseline mean sitting SBP and DBP, blood pressure control rate (&lt;140/90 mm Hg)</p> <p>Secondary: Safety</p>	<p>Primary: After eight weeks, add-on treatment with aliskiren resulted in significantly greater reductions in mean sitting SBP and DBP compared to placebo (-37.7/-16.1 vs -30.6/-12.3 mm Hg; P&lt;0.0001).</p> <p>After eight weeks, significantly more patients receiving aliskiren add-on therapy achieved blood pressure control compared to placebo (67.0 vs 49.1%; P=0.0001).</p> <p>Secondary: The overall incidence of adverse events was similar between both treatments. The most commonly reported adverse event was peripheral edema, with a higher incidence occurring in patients receiving placebo (18.3 vs 14.4%).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
up to 10 mg/day.				
Weinberger et al. <sup>25</sup> (abstract) 2012 ACCESS  Aliskiren 150 mg/day, up titrated to 300 mg/day  vs  placebo  All patients were receiving amlodipine 5 mg/day, up titrated to 10 mg/day	DB, RCT  African American patients with stage 2 HTN (mean sitting SBP 160 to 199 mm Hg) with obesity or metabolic syndrome	N=489  8 weeks	Primary: Change in baseline mean sitting SBP  Secondary: Not reported	Primary: LSM reductions in mean sitting SBP were significantly higher with add-on aliskiren compared to placebo in both obese (-33.7 vs -27.9 mm Hg; P<0.001) and metabolic syndrome patients (-36.4 vs -28.5 mm Hg; P<0.001).  Secondary: Not reported
Teo et al. <sup>26</sup> (2014) APOLLO  Aliskiren 300 mg daily  vs  placebo  and  amlodipine 5 mg daily  or	DB, PC, RCT  Patients ≥65 years with SBP between 130 and 159 mm Hg with either CVD or one additional CV risk factor	N=11,000  5 years  Study was terminated by sponsor after 1759 subjects were randomized and followed for 0.6 year due to non-scientific reasons without any knowledge	Primary: Original endpoints (risks of the composite of CV death, non-fatal MI, non-fatal stroke, and clinically significant heart failure) could not be assessed, so tolerability and effects on BP lowering were reported  Secondary: Not reported	Primary: Postrandomization, aliskiren reduced adjusted mean SBP by 3.5 (SE [standard error] 0.5) mmHg, (P<0.001), and DBP by 1.7 (SE 0.3) mmHg (P<0.001) compared with placebo (first co-primary outcome), HCTZ or amlodipine by 6.8 (SE 0.5) mmHg, (P<0.001) for SBP and 3.3 (SE 0.3) mmHg (P<0.001) for DBP. The reduction in SBP in the double therapy compared with double placebo (second co-primary outcome) was 10.3 (SE 0.8) mmHg (P<0.001) for SBP, and 5.0 (SE 0.5) mmHg, P<0.001 in mean DBP.  There were few serious adverse events, both during run-in and after randomization, with no excess associated with any treatment group.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
HCTZ 25 mg daily		of blinded trial data and despite objections of the Steering Committee		
<p>Schmieder et al.<sup>27</sup> (2009)</p> <p>Aliskiren 150 to 300 mg QD (with optional addition of amlodipine 5 to 10 mg QD)</p> <p>vs</p> <p>HCTZ 12.5 to 25 mg QD (with optional addition of amlodipine 5 to 10 mg QD)</p> <p>vs</p> <p>placebo for 6 weeks, then randomized to either aliskiren 300 mg QD or HCTZ 25 mg QD</p>	<p>AC, DB, RCT</p> <p>Adults with essential HTN</p>	<p>N=1,124</p> <p>12 months</p>	<p>Primary: Safety and change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP</p>	<p>Primary: The proportion of patients who experienced adverse events during the six week placebo-controlled period was similar in the aliskiren monotherapy, HCTZ monotherapy, and placebo groups (26.4, 24.5, and 28.5%, respectively).</p> <p>During the 52 week double-blind treatment period, adverse events were reported by a similar proportion of patients receiving the aliskiren and hydrochlorothiazide regimens. Most adverse events were mild or moderate in intensity.</p> <p>At week 26, the aliskiren regimen provided significantly greater reductions from baseline in DBP compared to HCTZ (-14.2 and -13.0 mm Hg, respectively; P&lt;0.05). The greater reduction in DBP with the aliskiren regimen compared to the HCTZ regimen was maintained at week 52 (-16.0 and -15.0 mm Hg, respectively; P&lt;0.05).</p> <p>Secondary: At week 26, the aliskiren regimen provided significantly greater reductions from baseline in SBP compared to HCTZ (-20.3 and -18.6 mm Hg, respectively; P&lt;0.05). Reductions in SBP at week 52 were not inferior to those of HCTZ (-22.1 and -21.2 mm Hg, respectively; P&lt;0.0001 for non-inferiority).</p>
<p>Schmieder et al.<sup>28</sup> (2009)</p> <p>Aliskiren 150 mg QD, followed by 300 mg QD after 3</p>	<p>Subgroup analysis of obese patients in Schmieder et al.<sup>25</sup></p> <p>Patients 18 years of age and older with</p>	<p>N=1,124</p> <p>52 weeks</p>	<p>Primary: Mean sitting DBP</p> <p>Secondary: Mean sitting SBP at week 26, mean</p>	<p>Primary: The LSM DBP and SBP reductions at week 12 were significantly greater with aliskiren compared to HCTZ (P&lt;0.0001 and P=0.001 respectively).</p> <p>Secondary: At week 52, aliskiren resulted in significantly greater mean sitting DBP</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>weeks</p> <p>vs</p> <p>HCTZ 12.5 mg QD, followed by 25 mg QD after 3 weeks</p> <p>vs</p> <p>placebo, followed by aliskiren 300 mg QD or HCTZ 25 mg QD after 6 weeks</p>	<p>essential HTN, a mean sitting DBP <math>\geq 90</math> and <math>&lt; 110</math> mm Hg; at randomization, patients had to have a mean sitting DBP <math>\geq 95</math> and <math>&lt; 110</math> mm Hg and show a difference of <math>\leq 10</math> mm Hg since the previous visit</p>		<p>sitting DBP and SBP at week 52, proportion of patients with response to treatment, blood pressure control at weeks 26 and 52, and safety</p>	<p>reductions compared to HCTZ (<math>P &lt; 0.001</math>).</p> <p>Blood pressure response rates were significantly greater with aliskiren compared to HCTZ at both week 12 and week 52 (<math>P &lt; 0.05</math>).</p> <p>Significantly more obese patients achieved blood pressure control with aliskiren compared to HCTZ at week 12 (<math>P = 0.0013</math>). Blood pressure control rates were similar between groups at week 52 (<math>P</math> value not reported).</p>
<p>Littlejohn et al.<sup>29</sup> (2009)</p> <p>Aliskiren 150 to 300 mg and amlodipine 5 to 10 mg QD</p> <p>HCTZ may be added if additional blood pressure control was required.</p>	<p>OL, MC</p> <p>Patients <math>\geq 18</math> years of age with essential HTN (mean sitting DBP <math>\geq 90</math> mm Hg and <math>&lt; 110</math> mm Hg)</p>	<p>N=556</p> <p>12 months</p>	<p>Primary: Safety and tolerability</p> <p>Secondary: Blood pressure-lowering efficacy</p>	<p>Primary: Long-term treatment with aliskiren and amlodipine was generally well tolerated. In total, 76.3% of patients reported at least one adverse event. The majority were mild or moderate in severity and transient. The most frequently reported adverse events were peripheral edema, upper respiratory tract infection, headache, and bronchitis.</p> <p>Peripheral edema was reported in 20.5% of patients who received aliskiren and amlodipine and in 14.0% of patients who received aliskiren and amlodipine and HCTZ.</p> <p>Edema was reported as mild in 59.5%, moderate in 33.3% and severe in 7.1% of patients.</p> <p>Secondary: At week two, treatment with aliskiren/amlodipine led to a mean reduction in blood pressure of 13.5/8.3 mm Hg. At week 10, there was a mean reduction in blood pressure of 23.5/15.1 mm Hg. Blood pressure reductions were sustained from week 10 until the end of the study. At week 54, aliskiren and amlodipine decreased mean blood pressure from 153.5/97.6 mm Hg at baseline to 129.4/82.2 mm Hg (<math>P &lt; 0.001</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				The BP control rate was 74.3% with aliskiren/amlodipine at week 54.
<p>Drummond et al.<sup>30</sup> (2007)</p> <p>Aliskiren and amlodipine 150-5 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amlodipine 5 mg QD</p> <p>vs</p> <p>amlodipine 10 mg QD</p> <p>Patients not responding to amlodipine 5 mg QD at the end of 4 week single-blind run-in period received combination therapy, continuation of amlodipine 5 mg QD or titration to amlodipine 10 mg QD.</p>	<p>AC, DB, MC, PG, RCT</p> <p>Patients 18 years of age and older with mild to moderate HTN</p>	<p>N=545</p> <p>6 weeks</p>	<p>Primary: Change in DBP at 6 weeks</p> <p>Secondary: SBP, comparison of SBP and DBP reductions between combination therapy group and amlodipine 10 mg group, proportion of patients responding to treatment, and proportion of patients achieving blood pressure control</p>	<p>Primary: DBP reduction was significantly greater in the combination therapy group compared to those in the amlodipine 5 mg group (P&lt;0.0001).</p> <p>Secondary: SBP reduction was significantly greater in the combination therapy group compared to those in the amlodipine 5 mg group (P&lt;0.0001).</p> <p>No significant differences were observed in DBP or SBP reduction between the combination therapy group and the amlodipine 10 mg group (P=0.6167 and P=0.2666 respectively).</p> <p>The proportion of patients responding to treatment was significantly higher in the combination therapy group compared to the amlodipine 5 mg group (P&lt;0.0001). No significant difference was observed between the combination therapy group and the amlodipine 10 mg group (P value not reported).</p> <p>The proportion of patients achieving blood pressure control was significantly higher in the combination therapy group compared to the amlodipine 5 mg group (P&lt;0.0001). No significant difference was observed between the combination therapy group and the amlodipine 10 mg group (P=0.5229).</p>
Villamil et al. <sup>31</sup> (2007)	DB, MC, PC, RCT	N=2,776	Primary: Change in mean	Primary: Aliskiren monotherapy significantly reduced mean sitting DBP (P=0.0002).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Aliskiren 75 to 300 mg QD</p> <p>vs</p> <p>HCTZ 6.25 to 25 mg QD</p> <p>vs</p> <p>aliskiren and HCTZ (every dose combination except aliskiren 300 mg and HCTZ 6.25 mg) QD</p> <p>vs</p> <p>placebo</p>	<p>Men and women ≥18 years with mild-to-moderate essential HTN</p>	<p>8 weeks</p>	<p>sitting DBP</p> <p>Secondary: Change in mean sitting SBP, dose-response efficacy for all treatment groups, proportion achieving a successful response (DBP &lt;90 mm Hg or ≥10 mm Hg), proportion achieving blood pressure control (&lt;140/90 mm Hg), plasma renin activity, renin concentrations, safety</p>	<p>Only the aliskiren 150 and 300 mg doses were more effective than placebo (P=0.09 for aliskiren 75 mg). HCTZ monotherapy significantly reduced DBP from baseline (P&lt;0.01 for all vs placebo).</p> <p>All combinations were more effective than placebo (P&lt;0.0001) with reductions in DBP ranging from 10.4 to 14.3 mm Hg. Most combination regimens were more effective than monotherapy with the individual components (exceptions were aliskiren 150 mg and HCTZ 6.25 mg vs monotherapy, and aliskiren 75 mg and HCTZ 12.5 mg vs HCTZ monotherapy).</p> <p>Secondary: After eight weeks of therapy, aliskiren 150 and 300 mg regimens (both P&lt;0.0001) were more effective than placebo in lowering mean sitting SBP, but the 75 mg dose was not (P=0.151).</p> <p>Combination therapy was consistently more effective in reducing SBP than monotherapy with the individual components, with the exception of aliskiren 75 mg plus HCTZ 12.5 vs HCTZ monotherapy. Reductions in SBP with combination therapy ranged from 14.3 to 21.2 mm Hg.</p> <p>Blood pressure reductions were related to the doses of both aliskiren and HCTZ.</p> <p>Responder rates were significantly higher with aliskiren 300 mg (63.9%; P=0.0005), HCTZ 12.5 and 25 mg (60.6 and 59.0%, respectively; both P&lt;0.02) and all combination doses (58.4 to 80.6%; all P&lt;0.05) than placebo (45.8%). Responder rates for all combinations of aliskiren and HCTZ 25 mg, and aliskiren 300 mg and HCTZ 12.5 mg were higher than both monotherapies (P&lt;0.05), while aliskiren 75 mg and HCTZ 12.5 mg and aliskiren 150 mg and HCTZ 12.5 mg were more effective than their respective aliskiren monotherapies (P&lt;0.05).</p> <p>In the aliskiren and HCTZ monotherapy groups, only aliskiren 300 mg led to statistically significantly greater control rates than placebo (46.7 vs 28.1%; P=0.0001). Control rates for all combinations, with the exception of aliskiren 75 mg and HCTZ 6.25 mg, were higher than placebo (all P&lt;0.02).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>There was a trend towards improved control rates with combination therapy (37.4 to 59.5%) compared to aliskiren monotherapy (29.0 to 46.7%) or HCTZ monotherapy (32.5 to 37.8%). Combinations utilizing the higher doses of one or both drugs (aliskiren 75 to 300 mg with HCTZ 25 mg or aliskiren 150 to 300 mg with HCTZ 12.5 mg) yielded control rates that were significantly higher than monotherapy with either component.</p> <p>While all doses of aliskiren decreased plasma renin activity and all doses of HCTZ increased plasma renin activity, combination therapy resulted in decreased plasma renin activity of 46.1 to 63.5%. Renin concentrations increased in all monotherapy and combination regimens with the exception of HCTZ 6.25 and 12.5 mg.</p> <p>All active treatments were well tolerated with 37.3 to 39.2% of patients experiencing adverse events with aliskiren monotherapy, 38.7 to 42.0% with HCTZ monotherapy, 34.6 to 45.3% with aliskiren and HCTZ, and 44% with placebo. Hypokalemia (serum potassium &lt;3.5 mmol/L) occurred with the highest frequency with HCTZ 12.5 and 25 mg (3.9 and 5.2%, respectively). When administered in combination with aliskiren, the frequency of hypokalemia was 0.7 to 2.0% with HCTZ 12.5 mg and 2.2% to 3.4% with HCTZ 25 mg.</p>
<p>Maddury et al.<sup>32</sup> (2013)</p> <p>aliskiren</p> <p>vs</p> <p>aliskiren + hydrochlorothiazide (HCT) single-pill combination</p>	<p>MC, OBS, OL, PRO</p> <p>Patients ≥18 years of age with a diagnosis of hypertension for which aliskiren or aliskiren HCT therapy had been prescribed by the treating physician</p>	<p>N=4,826</p> <p>26 ± 8 weeks</p>	<p>Primary: Proportion of patients who achieved therapeutic goal, defined as a target BP &lt;140/90 mmHg</p> <p>Secondary: Absolute change from baseline to end of study in mean sitting SBP (msSBP) and mean sitting DBP (msDBP), and the</p>	<p>Primary: The proportion of patients who reached the defined therapeutic BP goal at week 26 was 49.2% overall, 49.5% for aliskiren, and 48.3% for aliskiren HCT.</p> <p>Secondary: At week 26, the proportion of aliskiren-treated patients achieving the predefined response for SBP and DBP was 83.6 and 84.4%, respectively; the corresponding BP response rates in patients receiving aliskiren HCT were 84.4 and 86.5%. Treatment with aliskiren and aliskiren HCT was also associated with significant reductions from baseline to the end of study in msSBP and msDBP (P&lt;0.001 vs baseline for both treatments).</p> <p>Adverse effects occurred in a total of 101 (2.1%) patients. The most common AEs included headache, onset of diabetes mellitus, abdominal discomfort, and dizziness.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			proportion of patients achieving a BP response, safety	
<p>Fukutomi et al.<sup>33</sup> (2014)</p> <p>Aliskiren 150 mg/ amlodipine 5 mg group (AL/AM), after 8 weeks aliskiren dose was doubled to 300 mg for another 8 weeks</p> <p>vs</p> <p>high-dose amlodipine 10 mg (AM) group</p>	<p>MC, OL, RCT</p> <p>Hypertensive patients who were untreated or being treated with 5mg amlodipine</p> <p>During a 4-week run-in period, untreated patients started 5 mg amlodipine monotherapy and treated patients continued their medication. At the end of the run-in period, patients with BP <math>\geq</math>140mm Hg and/or diastolic BP (DBP) <math>\geq</math>90mm Hg were considered eligible for the study</p>	<p>N=87</p> <p>4-week run-in plus 16 weeks of treatment</p>	<p>Primary: Brachial flow-mediated vasodilation (FMD) and nitroglycerin-mediated vasodilation (NMD)</p> <p>Secondary: Not reported</p>	<p>Primary: FMD significantly improved in the AL/AM group but significantly decreased in the AM group. At the end of the study, FMD was significantly higher in the AL/AM group than in the AM group (<math>3.7 \pm 1.9\%</math> vs. <math>2.3 \pm 1.1\%</math>; <math>P &lt; 0.001</math>). NMD did not change after the treatment period in either group.</p> <p>Secondary: Not reported</p>
<p>Jordan et al.<sup>34</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p>	<p>DB, DD, MC, PG, RCT</p> <p>Obese men and women (BMI <math>\geq</math>30 kg/m<sup>2</sup>) <math>\geq</math>18 years with essential HTN (mean sitting DBP 95 to 109 mm Hg)</p>	<p>N=489</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting DBP with aliskiren 300 mg plus HCTZ vs HCTZ alone at 8 weeks</p> <p>Secondary:</p>	<p>Primary: Aliskiren 300 mg added to HCTZ 25 mg significantly reduced mean sitting DBP compared to HCTZ alone at week eight (mean difference, -4.0; <math>P &lt; 0.0001</math>).</p> <p>Secondary: Aliskiren 300 mg added to HCTZ caused numerically larger reductions in mean sitting DBP and SBP compared to amlodipine 10 mg plus HCTZ and irbesartan 300 mg plus HCTZ at week eight, but there were no statistically</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>amlodipine 5 to 10 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>irbesartan 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>HCTZ 25 mg QD (existing therapy)</p>	<p>and SBP &lt;180 mm Hg) who had not responded to 4 weeks of treatment with HCTZ 25 mg</p>		<p>Comparisons of mean sitting DBP and SBP with aliskiren plus HCTZ vs the other treatment groups, percentage of responders (mean sitting DBP &lt;90 mm Hg or <math>\geq</math>10 mm Hg reduction from baseline), proportion of patients achieving blood pressure control (mean sitting blood pressure &lt;140/90 mm Hg), plasma renin activity, safety and tolerability</p>	<p>significant differences between treatment groups (P&gt;0.05).</p> <p>Responder rates were significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0193) and week 12 (P=0.004) but comparable to responder rates observed with amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>The proportion of patients achieving blood pressure control was significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0005) and week 12 (P=0.0001) but not statistically different than amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>Plasma renin activity significantly increased (P&lt;0.05) during four weeks of HCTZ monotherapy. Combination with aliskiren neutralized this increase and led to an overall significant reduction in plasma renin activity compared to pretreatment baseline (P&lt;0.05) whereas amlodipine and irbesartan led to further significant increases (P&lt;0.05).</p> <p>All of the study treatments were generally well tolerated. Amlodipine plus HCTZ (45.2%) was associated with a higher incidence of adverse events than the other treatment groups (36.1 to 39.3%), largely due to a higher rate of peripheral edema (11.1 vs 0.8 to 1.6%).</p>
<p>Nickenig et al.<sup>35</sup> (2008)</p> <p>Aliskiren and HCTZ 300-25 mg QD (fixed-dose combination)</p> <p>vs</p> <p>aliskiren and HCTZ 300-12.5 mg QD (fixed-dose combination)</p>	<p>DB, MC, RCT</p> <p>Patients with HTN and an inadequate response to aliskiren (mean sitting DBP &gt;90 and <math>\leq</math>110 mm Hg following 4 weeks of aliskiren 300 mg)</p>	<p>N=880</p> <p>8 weeks</p>	<p>Primary: Changes in mean sitting SBP and DBP, rates of blood pressure control (&lt;140/90 mm Hg)</p> <p>Secondary: Not reported</p>	<p>Primary: Treatment with aliskiren and HCTZ 300-25 mg and 300-12.5 mg led to significantly greater reductions in mean sitting SBP/DBP from baseline (15.9/11.0 mm Hg and 13.5/10.5 mm Hg, respectively) compared to aliskiren 300 mg (8.0/7.4 mm Hg; both P&lt;0.001).</p> <p>Rates of blood pressure control were significantly higher with aliskiren and HCTZ 300-25 mg (60.2%) and 300-12.5 mg (57.9%) compared to aliskiren 300 mg (40.9%; both P&lt;0.001).</p> <p>Patients treated with aliskiren and HCTZ or aliskiren monotherapy demonstrated similar tolerability.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  aliskiren 300 mg QD (existing therapy)				
Blumenstein et al. <sup>36</sup> (2009)  Aliskiren and HCTZ 300-25 mg QD (fixed-dose combination product)  vs  aliskiren and HCTZ 150-25 mg QD (fixed-dose combination product)  vs  HCTZ 25 mg (existing therapy)	DB, MC, RCT  Patients with HTN and an inadequate response to HCTZ (mean sitting DBP >90 and ≤110 mm Hg following 4 weeks of HCTZ 25 mg)	N=722  8 weeks	Primary: Changes in mean sitting SBP/DBP, proportion of patients achieving blood pressure control (mean sitting blood pressure <140/90 mm Hg), and blood pressure response rates (msDBP <90 mm Hg or a ≥10 mm Hg decrease from baseline)  Secondary: Not reported	Primary: The mean reductions in mean sitting SBP/DBP from baseline with aliskiren and HCTZ 300-25 and 150-25 mg were significantly greater compared to those achieved with HCTZ monotherapy (P<0.001 for all).  Rates of blood pressure control were significantly higher with aliskiren and HCTZ 300-25 and 150-25 mg compared to HCTZ monotherapy (P<0.001 for both).  Aliskiren and HCTZ 300-25 mg provided significantly greater reductions in mean sitting SBP/DBP and rates of blood pressure control compared to aliskiren and HCTZ 150-25 mg dose (P<0.05 for all).  Blood pressure response rates were significantly higher with aliskiren and HCTZ 300-25mg (78.5%) and aliskiren and HCTZ 150-25 mg (67.4%) compared to HCTZ monotherapy (47.1%; P<0.001 for both comparisons).  All treatments were generally well-tolerated and the proportion of patients experiencing adverse events was similar across treatment groups. The majority of adverse events were mild and transient. Adverse events reported in >2% of patients were nasopharyngitis, dizziness, back pain, and vertigo.  The proportion of patients with serum potassium <3.5 mmol/L was lower with aliskiren and HCTZ (1.3 to 2.2%) compared to HCTZ monotherapy (3.4%). Hyperkalemia (serum potassium >5.5 mmol/L) was observed in only one patient receiving aliskiren and HCTZ and two patients in the HCTZ monotherapy group. No patient had increases in serum creatinine above the pre-specified clinically significant threshold.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Lacourciere et al.<sup>37</sup> (2012)</p> <p>Aliskiren and amlodipine and HCTZ 150-5-12.5 mg /day (fixed-dose combination product), up titrated to double the initial dose</p> <p>vs</p> <p>aliskiren and amlodipine 150-5 mg/day (fixed-dose combination product) , up titrated to double the initial dose</p> <p>vs</p> <p>aliskiren and HCTZ 150-12.5 mg/day (fixed-dose combination product) , up titrated to double the initial dose</p> <p>vs</p> <p>amlodipine and HCTZ 5-12.5 mg/day (fixed-</p>	<p>DB, RCT</p> <p>Patients ≥18 years of age with moderate to severe HTN</p>	<p>N=1,191</p> <p>8 weeks</p>	<p>Primary: Change in baseline mean sitting SBP and DBP, blood pressure control rate (&lt;140/90 mm Hg)</p> <p>Secondary: Safety</p>	<p>Primary: Treatment with aliskiren and amlodipine and HCTZ resulted in significant LSM reductions in mean sitting SBP/DBP (week 4: -30.7/-15.9 mm Hg; week 8: -37.9/-20.6 mm Hg) compared to any combination therapy (P&lt;0.001 for all). Significant reductions with triple therapy were observed as early as two weeks compared to dual therapies (P&lt;0.05).</p> <p>Significantly more patients receiving aliskiren and amlodipine and HCTZ achieved blood pressure control compared to dual therapies with moderate to severe (62.3%) and severe (57.5%) HTN.</p> <p>Secondary: The majority of adverse events were mild or moderate in nature. The overall incidence of events was comparable among treatments (36.2 vs 33.4 vs 32.3 vs 33.6%). Peripheral edema was the most commonly reported adverse event.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
dose combination product)* , up titrated to double the initial dose				
Ferdinand et al. <sup>38</sup> (abstract) (2012)  Aliskiren and amlodipine and HCTZ 300-10-25 mg /day (fixed-dose combination product)  vs  aliskiren and amlodipine 150-5 mg/day (fixed-dose combination product)	Subgroup analysis  Patients with HTN and any of the following: diabetes, cardiometabolic syndrome, obesity, or black patients	N=not reported  8 weeks	Primary: Change in baseline mean sitting SBP  Secondary: Not reported	Primary: LSM reductions in mean sitting SBP, across all subgroups ranged from 35 to 37 mm Hg with aliskiren and amlodipine and HCTZ compared to 28 to 30 mm Hg with aliskiren and amlodipine (P<0.01).  Secondary: Not reported
Gradman et al. <sup>39</sup> (2005)  Aliskiren 150 to 600 mg QD  vs  irbesartan 150 mg QD  vs  placebo	DB, MC, PC, PG, RCT  Men and women, age 18 years or older, with mild-to-moderate essential HTN (mean sitting DBP ≥95 mm Hg and <110 mm Hg)	N=652  8 weeks	Primary: Change in mean sitting DBP and SBP  Secondary: Proportion of patients achieving blood pressure control (<140/90 mm Hg), safety	Primary: Decreases in mean sitting DBP at eight weeks were significantly greater with all doses of aliskiren compared to placebo (P<0.001). The least-squares mean reductions in trough DBP for aliskiren 150, 300, and 600 mg were 9.3, 11.8, and 11.5 mm Hg, respectively, vs 6.3 mm Hg for placebo.  Decreases in mean sitting SBP at eight weeks were significantly greater with all doses of aliskiren compared to placebo (P<0.001). The least-squares mean reductions in trough SBP for aliskiren 150, 300, and 600 mg were 11.4, 15.8, and 15.7 mm Hg, respectively, vs 5.3 mm Hg for placebo.  The antihypertensive effect of aliskiren 150 mg was comparable to irbesartan 150 mg with reductions of 8.9 and 12.5 mm Hg for mean sitting DBP and SBP, respectively. Aliskiren 300 and 600 mg produced significantly greater mean sitting DBP reductions than irbesartan 150 mg

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>(P&lt;0.05). While the reductions in mean sitting SBP were greater with aliskiren 300 and 600 mg than irbesartan 150 mg, these differences were not statistically significant).</p> <p>Secondary: The percentage of patients achieving blood pressure control was significantly greater with all doses of aliskiren (37.8%-150 mg, 50.0%-300 mg, 45.7%-600 mg) and irbesartan (33.8%) compared to placebo (20.8%; P&lt;0.05). More patients on aliskiren 300 and 600 mg achieved blood pressure control compared to irbesartan (P&lt;0.05).</p> <p>Drug-related adverse events for both aliskiren and irbesartan were comparable to placebo and the most commonly reported adverse events were headache, dizziness, and diarrhea. The number of patients discontinuing therapy was similar in all groups.</p>
<p>O'Brien et al.<sup>40</sup> (2007)</p> <p>Aliskiren 150 mg QD for 3 weeks, then HCTZ 25 mg QD was added for an additional 3 weeks (if ABPM remained <math>\geq</math>135/85 mm Hg)</p> <p>vs</p> <p>irbesartan 150 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p>	<p>3 OL studies</p> <p>Men and women 18 to 80 years with ambulatory SBP <math>\geq</math>140 and <math>\leq</math>180 mm Hg without treatment</p>	<p>N=67</p> <p>6 to 9 weeks</p>	<p>Primary: Change in daytime systolic ABPM with combination therapy compared to monotherapy</p> <p>Secondary: Change in daytime diastolic ABPM, nighttime systolic and diastolic ABPM, daytime and nighttime heart rates, plasma renin activity</p>	<p>Primary: Aliskiren coadministered with HCTZ (P=0.0007) or ramipril (P=0.03) led to significantly greater reductions in daytime systolic ABPM compared to monotherapy. There was a trend for a reduction in daytime systolic ABPM with the addition of aliskiren to irbesartan; however, this trend was not statistically significant.</p> <p>Secondary: Aliskiren plus HCTZ significantly lowered daytime diastolic ABPM compared to aliskiren monotherapy (P=0.0006). Changes in nighttime systolic and diastolic ABPM followed similar trends but did not achieve statistical significance (P=0.06 and P=0.09, respectively). No changes in heart rate were observed with either aliskiren regimen.</p> <p>Aliskiren added to irbesartan did not significantly change diastolic ABPM compared to irbesartan monotherapy; however, nighttime systolic and diastolic ABPM were significantly reduced (P&lt;0.05 for all). No changes in heart rate were observed with either irbesartan regimen.</p> <p>Mean diastolic ABPM was significantly decreased with the addition of aliskiren 150 mg (P&lt;0.05) but not aliskiren 75 mg to ramipril monotherapy. Both aliskiren doses significantly decreased nighttime systolic and diastolic</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>ramipril 5 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p>				<p>ABPM (P&lt;0.05 for all). No changes in heart rate were observed with either ramipril regimen.</p> <p>Aliskiren alone significantly inhibited plasma renin activity by 65% (P&lt;0.0001), while ramipril and irbesartan monotherapy increased renin activity by 90 and 175%, respectively. When aliskiren was coadministered with HCTZ, ramipril or irbesartan, plasma renin activity remained similar to baseline levels or decreased.</p>
<p>Strasser et al.<sup>41</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>lisinopril 20 to 40 mg QD</p> <p>HCTZ may be added if additional blood pressure control was required.</p>	<p>AC, DB, DD, MC, PG, RCT</p> <p>Men and women with uncomplicated severe HTN (mean sitting DBP 105 to 119 mm Hg)</p>	<p>N=183</p> <p>8 weeks</p>	<p>Primary: Safety</p> <p>Secondary: Change in mean sitting DBP and SBP, percentage of responders</p>	<p>Primary: Both active treatments were well tolerated with an incidence of adverse events of 32.8% for aliskiren and 29.3% for lisinopril. The proportion of patients discontinuing treatment due to adverse events was 3.2% for aliskiren and 3.4% for lisinopril. The most frequently reported adverse events in both groups were headache, nasopharyngitis and dizziness.</p> <p>Secondary: Aliskiren showed similar reductions from baseline to lisinopril in mean sitting DBP (-18.5 vs -20.1 mm Hg) and SBP (-20.0 and -22.3 mm Hg).</p> <p>Responder rates were 81.5% with aliskiren and 87.9% with lisinopril. Approximately half of patients required the addition of HCTZ to achieve blood pressure control (53.6% for aliskiren and 44.8% for lisinopril).</p>
<p>Stanton et al.<sup>42</sup> (2003)</p> <p>Aliskiren 37.5 to 300 mg QD</p> <p>vs</p> <p>losartan 100 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Men and women 21 to 70 years of age with mild-to-moderate HTN (SBP ≥140 mm Hg)</p>	<p>N=226</p> <p>4 weeks</p>	<p>Primary: Change in daytime ambulatory SBP</p> <p>Secondary: Changes in clinic SBP and DBP, plasma renin activity, plasma aliskiren levels,</p>	<p>Primary: A dose-dependent reduction in daytime ambulatory SBP was observed with increasing aliskiren doses (with mean changes of -0.40 mm Hg with aliskiren 37.5 mg, -5.3 mm Hg with aliskiren 75 mg, -8.0 mm Hg with aliskiren 150 mg, and -11 mm Hg with aliskiren 300 mg; P=0.0002). The change in daytime SBP with losartan 100 mg (-10.9 mm Hg) was significantly different than aliskiren 37.5 mg, but not the other higher aliskiren dosages).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			adverse events	<p>Clinic SBP and DBP, both in the sitting and standing positions, decreased with aliskiren in a dose-dependent manner, whereas heart rate was unaltered. The decreases in clinic blood pressures were similar for losartan 100 mg and aliskiren 150 and 300 mg.</p> <p>Dose-dependent reductions in plasma renin activity were also observed (median change -55, -60, -77, and -83% with 37.5, 75, 150, and 300 mg aliskiren, respectively; P=0.0008). By contrast, plasma renin activity increased by 110% with losartan 100 mg.</p> <p>Rate of adverse events was 22% with aliskiren 37.5 mg, 35% with aliskiren 75 mg, 25% with aliskiren 150 mg, 23% with aliskiren 300 mg, and 32% with losartan 100 mg. There was no increase in the number of adverse events when increasing the dose of aliskiren.</p>
<p>Uresin et al.<sup>43</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and ramipril 5 to 10 mg QD</p>	<p>DB, MC, RCT</p> <p>Patients ≥18 years of age with type 1 or type 2 diabetes mellitus and stage 1 to 2 HTN (mean sitting DBP &gt;95 and &lt;110 mm Hg)</p>	<p>N=837</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, proportion of patients with a successful response to treatment (trough mean sitting DBP &lt;90 mm Hg and/or ≥10 mm Hg reduction from baseline), rates of blood pressure control (blood pressure &lt;130/80 mm Hg), changes from baseline in 24-hour ABPM measurements, and</p>	<p>Primary: Aliskiren monotherapy, ramipril monotherapy, and aliskiren and ramipril combination therapy lowered mean sitting DBP by 11.3, 10.7, and 12.8 mm Hg, respectively. Treatment with aliskiren and ramipril combination therapy produced significantly greater reductions from baseline in mean sitting DBP compared to either aliskiren monotherapy (P=0.043) or ramipril monotherapy (P=0.004). Aliskiren 300 mg was statistically non-inferior (P=0.0002) to ramipril 10 mg for the change in mean sitting DBP.</p> <p>Secondary: Aliskiren monotherapy, ramipril monotherapy, and aliskiren and ramipril combination therapy lowered mean sitting SBP by 14.7, 12.0, and 16.6 mm Hg, respectively. Treatment with aliskiren and ramipril combination therapy produced significantly greater reductions from baseline in mean sitting SBP compared to ramipril monotherapy (P&lt;0.0001), but not aliskiren monotherapy (P=0.088). Aliskiren monotherapy was statistically superior to ramipril for the change in mean sitting SBP (P=0.021).</p> <p>The proportion of patients with a successful response to therapy was similar for aliskiren and ramipril combination therapy (74.1%) and aliskiren monotherapy (73.1%). The responder rates in both groups were significantly higher (P&lt;0.05) compared to ramipril monotherapy (65.8%).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>changes in biomarkers (plasma renin concentration, plasma renin activity, aldosterone)</p>	<p>Rates of blood pressure control with aliskiren and ramipril combination pressure (13.1%) were not significantly different compared to aliskiren monotherapy (8.2%) or ramipril monotherapy (8.4%).</p> <p>All treatments significantly lowered mean 24-hour ambulatory blood pressure. Aliskiren and ramipril combination therapy was significantly more effective compared to ramipril monotherapy in lowering 24-hour mean ambulatory DBP (P=0.034). There was no significant difference in 24-hour ambulatory SBP compared to ramipril monotherapy.</p> <p>Aliskiren significantly reduced plasma renin activity from baseline as monotherapy (by 66%, P&lt;0.0001) or in combination with ramipril (by 48%, P&lt;0.0001).</p>
<p>Duprez et al.<sup>44</sup> (2010) AGELESS</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>The addition of HCTZ was allowed at week 12 and amlodipine was allowed at week 22 in patients not achieving adequate blood pressure control.</p>	<p>AC, DB, MC, RCT</p> <p>Patients ≥65 years of age with essential HTN (mean sitting SBP ≥140 and &lt;180 mm Hg and mean sitting DBP &lt;110mm Hg)</p>	<p>N=901</p> <p>36 weeks</p>	<p>Primary: Change in mean seated SBP at week 12</p> <p>Secondary: Change in mean sitting SBP at week 36, change in mean sitting DBP at week 12 and week 36, percentage of patients who achieved blood pressure control (mean sitting SBP/DBP &lt;140/90 mm Hg in non-diabetic patients and &lt;130/80 mm Hg in diabetic patients) at week 12 and week 36, percentage of</p>	<p>Primary: At week 12, aliskiren lowered mean sitting SBP by 14 mm Hg and ramipril decreased mean sitting SBP by 11.6 mm Hg (difference, -2.3 mm Hg; 95% CI, -4.3 to -0.3). Aliskiren monotherapy showed statistically non-inferior (P&lt;0.001) and statistically superior (P=0.02) reductions in mean sitting SBP compared to ramipril monotherapy.</p> <p>Secondary: At week 22, aliskiren decreased mean sitting SBP by 19.6 mm Hg and ramipril decreased mean sitting SBP by 17 mm Hg (difference, -2.4 mm Hg; 95% CI, -4.5 to -0.3; P=0.03).</p> <p>At week 36, aliskiren decreased mean sitting SBP by 20 mm Hg and ramipril decreased mean sitting SBP by 18.1 mm Hg (difference, -1.9 mm Hg; 95% CI, -4.0 to 0.2; P=0.07).</p> <p>At week 12, aliskiren decreased mean sitting DBP by 5.1 mm Hg and ramipril decreased mean sitting DBP by 3.6 mm Hg (difference, -1.5 mm Hg; 95% CI, -2.6 to -0.5; P&lt;0.01).</p> <p>At week 22, aliskiren decreased mean sitting DBP by 8.2 mm Hg and ramipril decreased mean sitting DBP by 7.3 mm Hg (difference, -0.8 mm Hg; 95% CI, -2.0 to 0.3; P=0.14).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			patients who required add-on therapy	<p>At week 36, aliskiren decreased mean sitting DBP by 8.2 mm Hg and ramipril decreased mean sitting DBP by 7.0 mm Hg (difference, -1.2 mm Hg; 95% CI, -2.3 to -0.1; P=0.03).</p> <p>The percentage of patients achieving blood pressure control was significantly greater with aliskiren (42%) compared to ramipril (33%) at week 12 (P&lt;0.01). At week 22, a significantly greater proportion of patients achieved blood pressure control with aliskiren (62%) compared to ramipril (50%; P&lt;0.001). At week 36, similar blood pressure control rates were achieved with aliskiren (59%) and ramipril (51%; P=0.01).</p> <p>By week 36, a significantly greater percentage of patients receiving ramipril compared to aliskiren required additional HCTZ (56 vs 46%; P&lt;0.01).</p> <p>By week 36, a greater percentage of patients receiving ramipril (16%) compared to aliskiren (12%) required add-on therapy with both HCTZ and amlodipine (P=0.048).</p> <p>More patients receiving aliskiren were receiving monotherapy (42%) than patients receiving ramipril (29%) at week 36.</p>
<p>Anderson et al.<sup>45</sup> (2008)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>ramipril 5 to 10 mg QD</p> <p>The addition of HCTZ was allowed in patients not achieving adequate blood</p>	<p>AC, DB, MC, PC, RCT</p> <p>Men and women ≥18 years with essential HTN (mean sitting DBP 90 to 109 mm Hg)</p>	<p>N=842</p> <p>26 weeks</p>	<p>Primary: Change in mean sitting DBP at week 26</p> <p>Secondary: Change in mean sitting SBP at week 26, change in mean sitting SBP and DBP at week 6 and 12 (comparing aliskiren and ramipril monotherapy), proportion</p>	<p>Primary: Reductions in mean sitting DBP at week 26 were significantly greater with aliskiren-based therapies (-13.2 mm Hg) compared to ramipril-based therapies (-12.0 mm Hg; P=0.0250).</p> <p>Secondary: Reductions in mean sitting SBP at week 26 were significantly greater with aliskiren-based therapies (-17.9 mm Hg) compared to ramipril-based therapies (-15.2 mm Hg; P=0.0036).</p> <p>Mean changes in sitting SBP were significantly greater with aliskiren (-12.9 and -14.0 mm Hg, respectively) compared to ramipril (-10.5 and -11.3, respectively) at weeks six and 12 (P=0.0041 and P=0.0027, respectively).</p> <p>Mean changes in sitting DBP were not significantly greater with aliskiren</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>pressure control.</p> <p>The study did not specifically analyze the effects of HCTZ on either treatment regimen.</p>			<p>achieving blood pressure control (&lt;140/90 mm Hg), proportion achieving SBP control (&lt;140 mm Hg), safety</p>	<p>(-10.5 and -11.3 mm Hg, respectively) compared to ramipril (-9.5 and -9.7, respectively) at week six, but were significantly greater at week 12 (P=0.0689 and P=0.0056, respectively).</p> <p>The proportion of patients achieving overall blood pressure control (&lt;140/90 mm Hg) was significantly higher with aliskiren-based therapy (61.4%) compared to ramipril-based therapy (53.1%; P=0.0205) at week 26. Also, the proportion of patients achieving SBP control (&lt;140 mm Hg) was significantly higher with aliskiren-based therapy (72.5%) compared to ramipril-based therapy (64.1%; P=0.0075) at week 26.</p> <p>The majority of adverse events reported during the active treatment period were mild or moderate in intensity and transient. Most events occurred at a similar incidence in the two groups with the exception of cough which was considered treatment-related in 5.5% of patients receiving ramipril vs 2.1% of patients receiving aliskiren.</p>
<p>Oparil et al.<sup>46</sup> (2007)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>valsartan 160 to 320 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and valsartan 160 to 320 mg QD (single entity products)</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Men and women aged 18 years or over with stage 1-2 essential HTN (mean sitting DBP 95 to 109 mm Hg and 8-hr ambulatory DBP ≥90 mm Hg)</p>	<p>N=1,797</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP, proportion of patients achieving a successful response to treatment (mean sitting DBP &lt;90 mm Hg and/or ≥10 mm Hg reduction from baseline) or achieving blood pressure control (mean sitting SBP/DBP &lt;140/90 mm Hg), change in 24-hr ABPM,</p>	<p>Primary: The combination of aliskiren 300 mg and valsartan 320 mg lowered mean sitting DBP from baseline by 12.2 mm Hg, significantly more than either monotherapy with aliskiren 300 mg (-9.0 mm Hg; P&lt;0.0001), valsartan 320 mg (-9.7 mm Hg; P&lt;0.0001) or with placebo (-4.1 mm Hg; P&lt;0.0001). Monotherapy with aliskiren or valsartan provided significantly greater reductions in mean sitting DBP than did placebo at week 8 (P&lt;0.0001 for all).</p> <p>Secondary: The combination of aliskiren 300 mg and valsartan 320 mg lowered mean sitting SBP from baseline by 17.2 mm Hg, significantly more than either monotherapy with aliskiren 300 mg (-13.0 mm Hg; P&lt;0.0001), valsartan 320 mg (-12.8 mm Hg; P&lt;0.0001), or with placebo (-4.6 mm Hg; P&lt;0.0001). Monotherapy with aliskiren or valsartan provided significantly greater reductions in mean sitting SBP than did placebo at week eight end point (all P&lt;0.0001).</p> <p>The proportion of patients achieving a successful response to treatment at week eight was significantly higher with the combination of aliskiren and valsartan (66%) than with aliskiren alone (53%; P=0.0003) or valsartan</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
placebo			change in biomarkers, safety	<p>alone (55%; P=0.0010). All active treatments were associated with significantly greater responder rates than placebo (30%; P&lt;0.0001 for all).</p> <p>The proportion of patients achieving blood pressure control was significantly greater in the combination group (49%) than in the aliskiren (37%; P=0.0005) or valsartan (34%; P&lt;0.0001) monotherapy groups. All active treatments were associated with significantly greater control rates than placebo (16%; P&lt;0.0001 for all).</p> <p>The combination of aliskiren and valsartan was significantly more effective in lowering mean 24-hr ambulatory SBP and DBP than was either agent alone (P&lt;0.0001 for all). The greater reductions in ambulatory blood pressure with aliskiren plus valsartan were maintained throughout the entire 24-hour dosing interval.</p> <p>Aliskiren and valsartan (P&lt;0.0001) and monotherapy with aliskiren (P&lt;0.0001) or valsartan (P=0.0002) provided significant increases in plasma renin concentrations versus placebo. Increases in plasma renin concentrations were significantly greater for the combination than aliskiren (P=0.0014) or valsartan (P&lt;0.0001) monotherapy.</p> <p>Valsartan monotherapy produced significantly greater increases in plasma renin activity than placebo (160 vs 18%; P=0.0003). By contrast, aliskiren alone significantly reduced plasma renin activity by 73% (P&lt;0.0001 vs placebo), while the combination of aliskiren plus valsartan led to a reduction in plasma renin activity of 44% (P&lt;0.0001 vs placebo).</p> <p>The combination of aliskiren and valsartan (-31%; P&lt;0.0001) and valsartan monotherapy (-25%; P=0.0007) provided significantly greater reductions in plasma aldosterone concentration than did placebo (7%), while aliskiren monotherapy had no significant effect (-5.9%; P=0.1059).</p> <p>Rates of adverse events and laboratory abnormalities were similar in all groups.</p>
Yarows et al. <sup>47</sup> (2008)	Post-hoc analysis of patients with stage 2 HTN from Oparil et	N=1,797 8 weeks	Primary: Change in mean sitting DBP	<p>Primary: In patients with stage 2 HTN, significantly greater reductions in DBP were demonstrated in the aliskiren and valsartan 300-320 mg group compared to</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Aliskiren 150 mg QD for 4 weeks, followed by 300 mg QD for 4 weeks</p> <p>vs</p> <p>valsartan 160 mg QD for 4 weeks, followed by 320 mg QD for 4 weeks</p> <p>vs</p> <p>aliskiren and valsartan 150-160 mg QD for 4 weeks, followed by 300-320 mg QD for 4 weeks (fixed-dose combination products)</p> <p>vs</p> <p>placebo</p>	<p>al.<sup>48</sup></p> <p>Men and women ≥18 years of age with stage 1 to 2 essential HTN (mean sitting DBP 95 to 109 mm Hg and 8-hour ambulatory DBP ≥90 mm Hg)</p>		<p>Secondary:</p> <p>Change in mean sitting SBP, proportion of patients achieving a successful response to treatment (mean sitting DBP &lt;90 mm Hg and/or ≥10 mm Hg reduction from baseline) or achieving blood pressure control (mean sitting SBP/DBP &lt;140/90 mm Hg)</p>	<p>either higher-dose monotherapy group (P&lt;0.05) and placebo (P&lt;0.0001).</p> <p>Secondary:</p> <p>In patients with stage 2 HTN, significantly greater reductions in SBP were demonstrated in the aliskiren and valsartan 300-320 mg group compared to either higher-dose monotherapy group (P&lt;0.05) and placebo (P&lt;0.0001).</p> <p>DBP and SBP reductions in both monotherapy groups were significantly greater compared to placebo (P&lt;0.0001).</p> <p>The proportion of patients with stage 2 HTN achieving blood pressure control at week eight was significantly greater in the aliskiren and valsartan 300-320 mg group compared to both monotherapy groups and placebo (P≤0.044).</p> <p>Blood pressure control rates in the aliskiren group were significantly greater than placebo (P&lt;0.001). No significant difference was observed between the valsartan monotherapy and placebo groups.</p>
<p>Bakris et al.<sup>48</sup> (2013) VIVID</p> <p>Therapy with aliskiren/valsartan 150/160 mg titrated to 300/320</p>	<p>AC, DB, RCT</p> <p>Hypertensive adults with type 2 diabetes and stage 1 or 2 chronic kidney disease (CKD)</p>	<p>N=1143</p> <p>8 weeks</p>	<p>Primary: ABP</p> <p>Secondary: Safety</p>	<p>Primary:</p> <p>Both treatments produced significant reductions from baseline to week 8 in all ABPM measures (P&lt;0.0001). The addition of aliskiren to valsartan was associated with an incremental benefit of 4.0 mm Hg of lowering in 24-hour mean ambulatory (ma)SBP and 2.4 mm Hg of lowering in 24-hour maDBP (both P&lt;0.001).</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg vs valsartan monotherapy 160 mg titrated to 320 mg				Adverse events were experienced by 202 participants (35.2%) in the combination aliskiren/valsartan group and by 182 participants (32.2%) in the valsartan group.
Pool et al. <sup>49</sup> (2007)  Aliskiren 75 to 300 mg QD  vs  valsartan 80 to 320 mg  vs  aliskiren 75 to 300 mg and valsartan 80 to 320 mg  vs  valsartan and HCTZ 160-12.5 mg QD (fixed-dose combination)  vs  placebo	DB, MC, PC, PG, RCT  Men and women ≥18 years with mild-to-moderate essential HTN (mean sitting DBP ≥95 mm Hg after a 3- to 4-week single-blind placebo run-in period)	N=1,123  8 weeks	Primary: Change in mean sitting DBP  Secondary: Change in mean sitting SBP, safety	Primary: Aliskiren 300 mg significantly (P<0.0001) lowered mean sitting DBP compared to placebo. Reductions in mean sitting DBP for aliskiren 75 and 150 mg compared to placebo failed to reach statistical significance (P=0.052 and P=0.051, respectively).  Secondary: Aliskiren 300 mg significantly (P<0.0001) lowered mean sitting SBP compared to placebo.  A statistically significant linear dose relationship was observed for the effect of aliskiren (75 to 300 mg) on mean sitting DBP (P=0.0002) and mean sitting SBP (P=0.0005). The effects of aliskiren monotherapy on mean sitting DBP and SBP across the 75 to 300 mg dose range were similar to the effects of valsartan 80 to 320 mg.  Coadministration of aliskiren and valsartan produced a greater antihypertensive effect than either drug alone. Reductions in mean sitting DBP and SBP obtained with aliskiren 150 mg plus valsartan 160 mg and aliskiren 300 mg plus valsartan 320 mg were not significantly different from those observed with valsartan 160 mg plus HCTZ 12.5 mg. Responder rates were significantly greater than placebo for all 3 aliskiren monotherapy groups and for all aliskiren plus valsartan combinations. The proportion of responders with aliskiren 75 mg plus valsartan 80 mg was significantly greater than either component monotherapy (P<0.05). There was no significant difference between the proportion of responders to aliskiren 150 mg plus valsartan 160 mg or aliskiren 300 mg plus valsartan 320 mg compared to valsartan 160 mg plus HCTZ 12.5 mg.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Control rates were higher with aliskiren 300 mg compared to placebo and with valsartan 160 mg plus HCTZ 12.5 mg compared to aliskiren 150 mg plus valsartan 160 mg, but there were no significant differences between aliskiren plus valsartan combinations and the respective monotherapies.</p> <p>Aliskiren and valsartan were generally well tolerated either as monotherapy or in combination. The overall incidence of adverse events and rate of discontinuations because of adverse events were similar to placebo in all active treatment groups.</p>
<p>Geiger et al.<sup>50</sup> (2009)</p> <p>Aliskiren 150 to 300 mg QD, added to existing HCTZ therapy</p> <p>vs</p> <p>valsartan 160 to 320 mg QD, added to existing HCTZ therapy</p> <p>vs</p> <p>aliskiren 150 to 300 mg and valsartan 160 to 320 mg QD, added to existing HCTZ therapy</p> <p>vs</p> <p>HCTZ 25 mg QD</p>	<p>AC, DB, RCT</p> <p>Patients ≥18 years of age with mild to moderate essential HTN who were taking HCTZ for 4 weeks with a DBP ≥95 mm Hg</p>	<p>N=641</p> <p>8 weeks</p>	<p>Primary: Change in DBP at week 8</p> <p>Secondary: Change SBP at week 8, change in DBP and SBP at week 4, proportion of patients achieving blood pressure control (SBP/DBP &lt;140/90 mm Hg), change in plasma renin activity, plasma renin concentration</p>	<p>Primary: After eight weeks of therapy, the triple therapy showed significantly greater reductions in SBP and DBP compared to the other groups. The additional SBP and DBP reductions were 7 and 5 mm Hg, respectively compared to aliskiren and HCTZ (P&lt;0.0001), 3 and 2 mm Hg compared to valsartan and HCTZ (P&lt;0.01), and 15 and 10 mm Hg compared to HCTZ monotherapy (P&lt;0.001).</p> <p>Aliskiren and HCTZ and valsartan and HCTZ combination therapies were more effective compared to HCTZ monotherapy. Valsartan and HCTZ was more effective than aliskiren and HCTZ. SBP and DBP were reduced by 15 and 11 mm Hg, respectively in the aliskiren and HCTZ group. SBP and DBP were reduced by 18 and 14 mm Hg, respectively, in the valsartan and HCTZ group.</p> <p>Secondary: Blood pressure control rate was significantly higher with triple therapy compared to aliskiren and HCTZ (40.9%, P&lt;0.001), valsartan and HCTZ (48.7%, P&lt;0.001), and HCTZ monotherapy (20.5%, P&lt;0.001).</p> <p>At week four, a significantly greater blood pressure control rate was observed for the triple therapy group at lower doses (150-160-25 mg) compared to the respective doses of the other groups: aliskiren and valsartan and HCTZ (300-320-25 mg) group (56%) compared to aliskiren and HCTZ (36.6%, P&lt;0.05), valsartan and HCTZ (42.2%, P&lt;0.05), and HCTZ monotherapy (19.9%, P&lt;0.01).</p> <p>At week eight, plasma renin concentration was unchanged in the HCTZ</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>group, but was significantly increased in other groups. A significant decrease in plasma renin activity from baseline was observed in the aliskiren and HCTZ group (P&lt;0.001) and a significant increase was observed in the valsartan and HCTZ (P&lt;0.001). In the HCTZ and triple therapy groups, there was no change in plasma renin activity (both P&gt;0.75).</p>
<p>Dietz et al.<sup>51</sup> (2008)</p> <p>Aliskiren 150 to 300 mg QD</p> <p>vs</p> <p>atenolol 50 to 100 mg QD</p> <p>vs</p> <p>aliskiren 150 to 300 mg and atenolol 50 to 100 mg QD</p>	<p>RCT, DB, MC</p> <p>Patients ≥18 years of age with HTN (mean sitting DBP ≥95 and &lt;110 mm Hg)</p>	<p>N=694</p> <p>12 weeks</p>	<p>Primary: Changes in mean sitting SBP and mean sitting DBP, rates of blood pressure control (&lt;140/90 mm Hg), pulse pressure and pulse rate, plasma renin concentration, plasma renin activity</p> <p>Secondary: Not reported</p>	<p>Primary: Treatment with aliskiren and atenolol combination therapy led to a significantly greater reduction in mean sitting SBP by 17.3 mm Hg compared to aliskiren monotherapy (difference, -2.9 mm Hg; P=0.039) or atenolol monotherapy (difference, -3.0 mm Hg; P=0.034). There was no difference between mean sitting SBP reductions with aliskiren and atenolol monotherapy (difference, -0.1 mm Hg; P=0.954).</p> <p>Treatment with aliskiren and atenolol combination therapy led to a significantly greater reduction in mean sitting DBP by 14.1 mm Hg compared to aliskiren monotherapy (difference, -2.9 mm Hg; P&lt;0.001), but not atenolol monotherapy (difference, -0.5 mm Hg; P=0.545). Reductions in mean sitting DBP with atenolol were larger compared to those observed with aliskiren (difference, 2.4 mm Hg; P=0.003).</p> <p>Rates of blood pressure control were higher with aliskiren and atenolol combination therapy (51.3%) compared to aliskiren monotherapy (36.1%, P&lt;0.001) or atenolol monotherapy (42.2%, P=0.009). There was no significant difference in blood pressure control rates between aliskiren and atenolol monotherapy (P=0.388).</p> <p>Mean pulse pressure was reduced by 3.0 mm Hg with aliskiren and atenolol combination therapy and aliskiren monotherapy. Atenolol monotherapy did not affect pulse pressure. Aliskiren monotherapy did not affect pulse rate. Significant mean reductions in pulse rate of &gt;10 bpm were observed with atenolol monotherapy and the aliskiren and atenolol combination (P&lt;0.001 vs aliskiren monotherapy for both).</p> <p>Aliskiren monotherapy increased plasma renin concentration by 241% and aliskiren/atenolol increased plasma renin concentration by 85% (P=0.010 vs aliskiren). Atenolol monotherapy decreased plasma renin concentration by 24% (P&lt;0.001 vs aliskiren and aliskiren/atenolol). Aliskiren, atenolol</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				and aliskiren/atenolol reduced plasma renin activity by 65, 52, and 61%, respectively.  Secondary: Not reported
Stanton et al. <sup>52</sup> (2010)  Aliskiren 300 mg QD  vs  irbesartan, losartan, valsartan, ramipril, HCTZ, placebo	MA  Adults with mild to moderate essential HTN	N=4,877 (8 trials)  4 to 12 weeks	Primary: Paradoxical blood pressure rises, as well as the percentage of patients with SBP increases (>10 or >20 mm Hg) or DBP increases (>5 or >10 mm Hg) from baseline  Secondary: Not reported	Primary: There were no significant differences among the pooled aliskiren, irbesartan, losartan, valsartan, ramipril, and HCTZ groups in the incidence of SBP increases >10 mm Hg (P=0.30) and >20 mm Hg (P=0.28) or DBP increases >5 mm Hg (P=0.65) and >10 mm Hg (P=0.5).  Increases in SBP and DBP occurred significantly more frequently in the pooled placebo group than the aliskiren group (P<0.001).  Secondary: Not reported
Wysong et al. <sup>53</sup> (2007)  Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)  vs  β-blockers (atenolol, metoprolol, oxprenolol*, or	MA  13 RCTs evaluating patients ≥18 years of age with HTN	N=91,561  Duration varied	Primary: All-cause mortality  Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions	Primary: There was not a significant difference observed in all-cause mortality between β-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with β-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).  Secondary: There was a significant decrease in stroke observed with β-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
propranolol)				<p>CHD risk was not significantly different between <math>\beta</math>-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07], diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with <math>\beta</math>-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of <math>\beta</math>-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with <math>\beta</math>-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>
<p>Baguet et al.<sup>54</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*,</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the <math>\beta</math>-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).</p> <p>The weighted average reduction of SBP and DBP for each drug class were as follows: Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>enalapril, ramipril, trandolapril, and aliskiren)</p> <p>Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>β-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively.            Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively.            ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.            ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively.            Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary: Not reported</p>
<b>Diabetes/Diabetic Nephropathy/Renal Dysfunction</b>				
<p>Persson et al.<sup>55</sup> (2009)</p> <p>Aliskiren 300 mg QD</p> <p>vs</p> <p>irbesartan 300 mg QD</p> <p>vs</p> <p>aliskiren 300 mg QD and irbesartan</p>	<p>DB, RCT, XO</p> <p>Adults with type 2 diabetes, HTN, and albuminuria</p>	<p>N=26</p> <p>Four 2-month treatment periods</p>	<p>Primary: Albuminuria (urinary albumin excretion rate)</p> <p>Secondary: 24-hour blood pressure, GFR</p>	<p>Primary: Treatment with aliskiren led to a significant reduction in albuminuria by 48% compared to placebo (P&lt;0.001). Treatment with irbesartan led to a significant reduction in albuminuria by 58% compared to placebo (P&lt;0.001). There was no significant difference in albuminuria between aliskiren and irbesartan (P value not reported). The combination of aliskiren and irbesartan significantly reduced albuminuria by 71% compared to placebo (P&lt;0.001), which was also significantly better than with monotherapy (P&lt;0.001 for aliskiren and P=0.028 for irbesartan).</p> <p>Secondary: SBP and DBP 24-hr blood pressure were reduced by 3 and 4 mm Hg, respectively by aliskiren (P value not significant and P=0.009, respectively), 12 and 5 mm Hg, respectively by irbesartan (P&lt;0.001 and P=0.002, respectively), and 10 and 6 mm Hg, respectively with the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
300 mg QD  vs  placebo				combination (P=0.001 and P <0.001, respectively) compared to placebo. There was no significant change in 24-hr blood pressure with irbesartan compared to combination therapy.  GFR was significantly reduced 4.6 mL/min/1.73 m <sup>2</sup> with aliskiren (P=0.037), 8.0 mL/min/1.73 m <sup>2</sup> with irbesartan (P<0.001), and 11.7 mL/min/1.73 m <sup>2</sup> with the combination (P<0.001) compared to placebo.
Parving et al. <sup>56</sup> (2008) AVOID  Aliskiren 150 mg QD for 3 months, followed by 300 mg QD for 3 months  vs  placebo  Study medications were added to losartan 100 mg and other pre-existing antihypertensive treatments.	DB, MC, PC, RCT  Hypertensive patients who were 18 to 85 years of age who had type 2 diabetes and nephropathy	N=599  6 months	Primary: Reduction in albumin:creatinine ratio at 6 months  Secondary: Blood pressure reductions, adverse events	Primary: Treatment with aliskiren 300 mg/day as compared to placebo reduced the mean urinary albumin:creatinine ratio by 20% (95% CI, 9 to 30; P<0.001), with a reduction of 50% or more in 24.7% of the patients who received aliskiren as compared to 12.5% of those who received placebo (P<0.001).  Secondary: A small difference in blood pressure was seen between the treatment groups by the end of the study period with SBP and DBP pressures 2 and 1 mm Hg lower, respectively, in the aliskiren group (P=0.07 and P=0.08, respectively).  The total numbers of adverse and serious adverse events were similar in the groups.
<b>Miscellaneous</b>				
Solomon et al. <sup>57</sup> (2009) ALLAY  Aliskiren 300 mg QD  vs	AC, RCT  Adults with HTN and increased left ventricular wall thickness	N=465  9 months	Primary: Change in left ventricular mass  Secondary: Not reported	Primary: There were reductions in left ventricular mass from baseline in all treatment groups, with 4.9 g/m <sup>2</sup> (5.4%), 4.8 g/m <sup>2</sup> (4.7%), and 5.8 g/m <sup>2</sup> (6.4%) reductions in the aliskiren, losartan, and combination arms, respectively (P<0.0001 for all treatment groups).  The reduction in left ventricular mass in the combination group was not significantly different from that with losartan alone (P=0.52).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
losartan 100 mg QD  vs  aliskiren 300 mg and losartan 100 mg QD				The difference in left ventricular mass regression between the aliskiren and losartan arms was within the prespecified non-inferiority margin, suggesting that aliskiren was as effective as losartan in reducing left ventricular hypertrophy (P<0.0001 for non-inferiority).  Secondary: Not reported
McMurray et al. <sup>58</sup> (2008) ALOFT  Aliskiren 150 mg QD  vs  placebo	DB, MC, PC, RCT  Patients ≥18 years of age with NYHA class II to IV heart failure, current or past history of NTH, and plasma brain natriuretic peptide concentration >100 pg/mL who had been treated with an ACE inhibitor (or angiotensin receptor blocker) and β-blocker	N=302  3 months	Primary: N-terminal pro-brain natriuretic peptide, brain natriuretic peptide, aldosterone, signs and symptoms of heart failure echocardiographic measures of cardiac size and ventricular function, blood pressure, heart rate variability, quality of life, neurohumoral and inflammatory biomarkers, and glycemic measures  Secondary: Not reported	Primary: Plasma N-terminal pro- brain natriuretic peptide increased by 762 pg/mL with placebo and decreased by 244 pg/mL with aliskiren (P=0.0106).  Brain natriuretic peptide decreased by a mean of 12.2 pg/mL in the placebo group and by 61.0 pg/mL in the aliskiren group (P=0.0160).  Plasma aldosterone did not differ between groups. Urinary aldosterone decreased with aliskiren by 9.24 nmol/day and by 6.96 nmol/day with placebo (P=0.0150).  Plasma renin activity decreased 5.71 ng·mL <sup>-1</sup> ·h <sup>-1</sup> with aliskiren compared to a decrease of 0.97 ng·mL <sup>-1</sup> ·h <sup>-1</sup> with placebo (P<0.0001).  There was no difference between treatments for change in signs or symptoms of heart failure, echocardiographic measurements of wall thickness, chamber volumes, or LVEF.  The mean decrease in seated systolic blood pressure was 1.7 mm Hg in the placebo group and 4.1 mm Hg in the aliskiren group (P=0.2257). The mean decrease in seated diastolic blood pressure was 0.2 mm Hg in the placebo group and 2.9 mm Hg in the aliskiren group (P=0.0599). The mean increase in seated heart rate was 0.2 bpm in the placebo group and 1.1 bpm in the aliskiren group (P=0.6774).  Mean standing systolic blood pressure decreased by 1.7 mm Hg in the placebo group and by 3.5 mm Hg in the aliskiren group (P=0.497). The mean standing diastolic blood pressure increased by 0.7 mm Hg with

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>placebo and decreased by 3.5 mm Hg with aliskiren (P=0.0045). The mean standing heart rate decreased by 0.3 bpm in the placebo group and increased by 0.7 bpm in the aliskiren group (P=0.466).</p> <p>There were no differences between treatments in any of the other prespecified comparisons, including autonomic measurements, the Kansas City Cardiomyopathy questionnaire, inflammatory and other plasma and urinary biomarkers (including urinary protein excretion), or measurements of glucose/insulin metabolism.</p> <p>Secondary: Not reported</p>

\*Agent not available in the United States.

Drug regimen abbreviations: QD=once daily, SR=sustained-release

Study design abbreviations: AC=active comparator, DB=double blind, DD=double dummy, MA=meta-analysis, MC=multicenter, OL=open-label, OS=observational, PC=placebo-controlled, PG=parallel-group, PRO=prospective, RCT=randomized-controlled trial, XO=cross-over

Miscellaneous abbreviations: ABPM=ambulatory blood pressure monitoring, ACE inhibitors=angiotensin converting enzyme inhibitors, ARB=angiotensin II receptor blocker, BMI=body mass index, CAD=coronary artery disease, CHD=coronary heart disease, CI=confidence interval, DBP=diastolic blood pressure, GFR=glomerular filtration rate, HCTZ=hydrochlorothiazide, HTN=hypertension, LVEF=left ventricular ejection fraction, LSM=least squares mean, NYHA=New York Heart Association, RR=relative risk, SBP=systolic blood pressure

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Renin Inhibitors**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Aliskiren	tablet	Tekturna <sup>®</sup> *	\$\$\$\$\$	\$\$\$\$\$
<b>Combination Products</b>				
Aliskiren and HCTZ	tablet	Tekturna HCT <sup>®</sup>	\$\$\$\$\$	\$\$\$\$\$

\*Generic is available in at least one dosage form or strength.  
HCTZ=hydrochlorothiazide, N/A=not available

**X. Conclusions**

Aliskiren is the only renin inhibitor in this class and it is approved for the treatment of hypertension.<sup>6-7</sup> It is available as a single entity product, as well as in combination with hydrochlorothiazide. Aliskiren is available generically.

There are several national and international organizations that have published guidelines on the treatment of hypertension. Most of the guidelines do not address the use of the renin inhibitors, with the exception of European Society of Hypertension and European Society of Cardiology which state that evidence is available to justify the use of aliskiren for the management of hypertension, particularly in combination with other antihypertensive agents.<sup>9-17</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension. According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National

Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>9</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>9-17</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>9-17</sup>

Several clinical trials have demonstrated that renin inhibitors effectively lower blood pressure. The reduction in blood pressure with aliskiren monotherapy was similar to monotherapy with ACE inhibitors, angiotensin II receptor blockers,  $\beta$ -blockers, and dihydropyridines. In clinical trials comparing combination therapy to monotherapy, the more aggressive treatment regimen lowered blood pressure to a greater extent than the less-intensive treatment regimen.<sup>20-58</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>9-17</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>11,12,14</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations. Aliskiren is not recommended for use in combination with ACE inhibitors or ARBs, largely due to the findings of the ALITITUDE trial in which the risk of renal impairment, hypotension, and hyperkalemia increased in patients with GFR <60 mL/min and patients with diabetes.<sup>6-8,18</sup> Aliskiren has been shown to have positive effects on surrogate markers of cardiovascular and renal damage in patients with type 2 diabetes and nephropathy, heart failure and left ventricular hypertrophy.<sup>55-58</sup> However, the effects of aliskiren on hard cardiovascular and renal endpoints have not been established.

At this time, there is insufficient evidence to conclude that the renin inhibitors offer a significant clinical advantage over other alternatives in general use. Therefore, all brand renin inhibitors within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand renin inhibitor is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. *Pharmacotherapy: a pathophysiologic approach*. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
2. Aliskiren (Tekturna) for hypertension. *Med Lett Drugs Ther*. 2007 Apr 9;49(1258):29-31.
3. Van Tassell BW, Munger MA. Aliskiren for renin inhibition: a new class of antihypertensives. *Ann Pharmacother*. 2007 Mar;41:456-64.
4. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
5. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
6. Tekturna® [package insert]. Orlando (FL): Noden Pharma USA; 2021 Mar.
7. Tekturna HCT® [package insert]. Orlando (FL): Noden Pharma USA; 2020 Sep.
8. FDA Drug Safety Communication: New Warning and Contraindication for blood pressure medicines containing aliskiren (Tekturna) [press release on the Internet]. Rockville (MD): Food and Drug Administration (US); 2013 Mar 15 [cited 2022 Jan 25]. Available from: <http://www.fda.gov/Drugs/DrugSafety/ucm300889.htm>.
9. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
10. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
11. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
12. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
13. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
14. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010;56:780-800.
15. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
16. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324
17. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl. 1).
18. Parving HH, Brenner BM, McMurray JJV, et al. Cardiorenal End Points in a Trial of Aliskiren for Type 2 Diabetes. *N Engl J Med* 2012; 367:2204-2213.
19. McMurray JJ, Krum H, Abraham WT, Dickstein K, Køber LV, Desai AS, et al. Aliskiren, Enalapril, or Aliskiren and Enalapril in Heart Failure. *N Engl J Med*. 2016 Apr 21;374(16):1521-32.
20. Tocci G, Aimo G, Caputo D, De Matteis C, Di Napoli T, Granantelli A, et al. An observational, prospective, open-label multicentre evaluation of aliskiren in treated, uncontrolled patients: a real-life, long-term, follow-up, clinical practice in Italy. *High Blood Pressure Cardiovasc Prev*. 2012 Jun 1;19(2):78-83.
21. Oh BH, Mitchell J, Herron JR, et al. Aliskiren, an oral renin inhibitor, provides dose-dependent efficacy and sustained 24-hour blood pressure control in patients with hypertension. *J Am Coll Cardiol*. 2007 Mar 20;49(11):1157-63.
22. Kushiro T, Itakura H, Abo Y, et al. Aliskiren, a novel oral renin inhibitor, provides dose-dependent efficacy and placebo-like tolerability in Japanese patients with hypertension. *Hypertens Res*. 2006;29(12):997-1005.



23. Musini VM, Fortin PM, Bassett K, Wright JM. Blood pressure lowering efficacy of renin inhibitors for primary hypertension: a Cochrane systematic review. *J Hum Hypertens* 2009;23:495-502.
24. Braun-Dullaeus RC, Shustov SB, Alvarez C, Rogelio GG, Zhang J, Hristoskova S, et al. Treatment with aliskiren/amlodipine combination in patients with moderate-to-severe hypertension: a randomised, double-blind, active comparator trial. *Int J Clin Pract*. 2012;66(9):834-842.
25. Weinberger MH, Izzo JL Jr, Purkayastha D, Weitzman R, Black HR. Comparative efficacy and safety of combination aliskiren/amlodipine and amlodipine monotherapy in African American patients with stage 2 hypertension and obesity or metabolic syndrome. *J Am Soc Hypertens*. 2011 Nov-Dec;5(6):489-97.
26. Teo KK, Pfeffer M, Mancia G, et al. Aliskiren alone or with other antihypertensives in the elderly with borderline and stage 1 hypertension: the APOLLO trial. *Eur Heart J*. 2014 Jul;35(26):1743-51.
27. Schmieder RE, Philipp T, Guerediaga J, et al. Long-term antihypertensive efficacy and safety of the oral direct renin inhibitor aliskiren: a 12-month randomized, double-blind comparator trial with hydrochlorothiazide. *Circulation* 2009;119:417-25.
28. Schmieder R, Philipp T, Guerediaga J, Gorostidi M, Bush C, Keefe D. Aliskiren-based therapy lowers blood pressure more effectively than hydrochlorothiazide-based therapy in obese patients with hypertension: sub-analysis of a 52-week, randomized, double-blind trial. *J Hypertens*. 2009;27:1493-1501.
29. Littlejohn T, Trenkwalder P, Hollanders G, et al. Long-term safety, tolerability and efficacy of combination therapy with aliskiren and amlodipine in patients with hypertension. *Curr Med Res Opin* 2009;25:951-959.
30. Drummond W, Munger M, Essop M, Maboudian M, Khan M, Keefe D. Antihypertensive efficacy of the oral direct renin inhibitor aliskiren as add-on therapy in patients not responding to amlodipine monotherapy. *J Clin Hypertens*. 2007;9:742-50.
31. Villamil A, Chrysant SG, Calhoun D, et al. Renin inhibition with aliskiren provides additive antihypertensive efficacy when used in combination with hydrochlorothiazide. *J Hypertens*. 2007 Jan;25(1):217-26.
32. Maddury SR, Pande A, Haque KM, et al. Effectiveness and safety of aliskiren and aliskiren hydrochlorothiazide (HCT) in a multiethnic, real-world setting. *Adv Ther*. 2013 Feb;30(2):176-89.
33. Fukutomi M, Hoshida S, Mizuno H, and Kario K. Differential effects of aliskiren/amlodipine combination and high-dose amlodipine monotherapy on endothelial function in elderly hypertensive patients. *Am J Hypertens*. 2014 Jan;27(1):14-20.
34. Jordan J, Engeli S, Boye S, et al. Direct renin inhibition with aliskiren in obese patients with arterial hypertension. *Hypertension*. 2007 May;49:1047-55.
35. Nickenig G, Simanenkova V, Lembo G, et al. Efficacy of aliskiren/hydrochlorothiazide single-pill combinations in aliskiren non-responders. *Blood Press Suppl* 2008;2:31-40.
36. Blumenstein M, Romaszko J, Calderón A, et al. Antihypertensive efficacy and tolerability of aliskiren/hydrochlorothiazide (HCT) single-pill combinations in patients who are non-responsive to HCT 25 mg alone. *Curr Med Res Opin* 2009;25:903-910.
37. Lacourciere Y, Taddei S, Konis G, Fang H, Severin T, Zhang J. Clinical and ambulatory blood pressure lowering effect of aliskiren/amlodipine/hydrochlorothiazide combination in patients with moderate-to-severe hypertension: a randomized active-controlled trial. *J Hypertens*. 2012;30:2047-2055.
38. Ferdinand KC, Weitzman R, Purkayastha D, Sridharan K, Jaimes EA. Aliskiren-based dual- and triple-combination therapies in high-risk US minority patients with stage 2 hypertension (abstract). *J Am Soc Hypertens*. 2012 May-Jun;6(3):219-27.
39. Gradman AH, Schmieder RE, Lins RL, et al. Aliskiren, a novel orally effective renin inhibitor, provides dose-dependent antihypertensive efficacy and placebo-like tolerability in hypertensive patients. *Circulation*. 2005 Mar 1;111(8):1012-8.
40. O'Brien E, Barton J, Nussberger J, et al. Aliskiren reduces blood pressure and suppresses plasma renin activity in combination with a thiazide diuretic, an angiotensin-converting enzyme inhibitor, or an angiotensin receptor blocker. *Hypertension*. 2007 Feb;49(2):276-84.
41. Strasser RH, Puig JG, Farsang C, et al. A comparison of the tolerability of the direct renin inhibitor aliskiren and lisinopril in patients with severe hypertension. *J Hum Hypertens*. 2007 Oct;21(10):780-7.
42. Stanton A, Jensen C, Nussberger J, et al. Blood pressure lowering in essential hypertension with an oral renin inhibitor, aliskiren. *Hypertension*. 2003 Dec;42:1137-43.
43. Uresin Y, Taylor AA, Kilo C, et al. Efficacy and safety of the direct renin inhibitor aliskiren and ramipril alone or in combination in patients with diabetes and hypertension. *J Renin Angiotensin Aldosterone Syst* 2007;8:190-8.
44. Duprez D, Munger M, Botha J, et al. Aliskiren for geriatric lowering of systolic hypertension: a randomized controlled trial. *J Hum Hypertens* 2010;24:600-608.

45. Andersen K, Weinberger MH, Egan B, et al. Comparative efficacy and safety of aliskiren, an oral direct renin inhibitor, and ramipril in hypertension: a 6-month, randomized, double-blind trial. *J Hypertens*. 2008;26(3):589-99.
46. Oparil S, Yarows S, Patel S, et al. Efficacy and safety of combined use of aliskiren and valsartan in patients with hypertension: a randomized, double-blind trial. *Lancet*. 2007;370:221-9.
47. Yarows S, Oparil S, Patel S, Fang H, Zhang J. Aliskiren and valsartan in stage 2 hypertension; subgroup analysis of a randomized, double-blind study. *Adv Ther*. 2008;25(12):1288-302.
48. Bakris GL, Oparil S, Purkayastha D, Yadao AM, Alessi T, and Sowers JR. Randomized study of antihypertensive efficacy and safety of combination aliskiren/valsartan vs valsartan monotherapy in hypertensive participants with type 2 diabetes mellitus. *J Clin Hypertens (Greenwich)*. 2013 Feb;15(2):92-100.
49. Pool JL, Schmieder RE, Azizi M, et al. Aliskiren, an orally effective renin inhibitor, provides antihypertensive efficacy alone and in combination with valsartan. *Am J Hypertens* 2007;20(1):11-20.
50. Geiger H, Barranco E, Gorostidi M, et al. Combination therapy with various combinations of aliskiren, valsartan, and hydrochlorothiazide in hypertensive patients not adequately responsive to hydrochlorothiazide alone. *J Clin Hypertens (Greenwich)* 2009;11:324-32.
51. Dietz R, Dechend R, Yu C, et al. Effects of the direct renin inhibitor aliskiren and atenolol alone or in combination in patients with hypertension. *J Renin Angiotensin Aldosterone Syst* 2008;9:163-175.
52. Stanton AV, Gradman AH, Schmieder RE, et al. Aliskiren monotherapy does not cause paradoxical blood pressure rises. Meta-analysis of data from 8 clinical trials. *Hypertension* 2010;55:54-60.
53. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
54. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig*. 2007;27(11):735-53.
55. Persson F, Rossing P, Reinhard H, et al. Renal effects of aliskiren compared with and in combination with irbesartan in patients with type 2 diabetes, hypertension, and albuminuria. *Diabetes Care* 2009;32:1873-9.
56. Parving HH, Persson F, Lewis JB, et al; for the AVOID Study Investigators. Aliskiren combined with losartan in type 2 diabetes and nephropathy. *N Engl J Med*. 2008 Jun 5;358(23):2433-46.
57. Solomon SD, Appelbaum E, Manning WJ et al. Effect of the direct renin inhibitor aliskiren, the angiotensin receptor blocker losartan, or both on left ventricular mass in patients with hypertension and left ventricular hypertrophy. *Circulation* 2009;119:530-7.  
McMurray JJ, Pitt B, Latini R, et al; Aliskiren Observation of Heart Failure Treatment (ALOFT) Investigators. Effects of the oral direct renin inhibitor aliskiren in patients with symptomatic heart failure. *Circ Heart Fail* 2008;1:17-24.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Loop Diuretics  
AHFS Class 402808  
May 4, 2022**

**I. Overview**

Diuretics are commonly used for the treatment of hypertension, heart failure, and various edematous conditions.<sup>1,2</sup> These agents act at different sites within the nephron, which leads to the increased urinary excretion of sodium, chloride and water.<sup>2</sup> The diuretics are categorized into several different AHFS classes, including loop diuretics, potassium-sparing diuretics, thiazide diuretics, thiazide-like diuretics, vasopressin antagonists, and miscellaneous diuretics. The agents which make up these classes differ with regards to their Food and Drug Administration (FDA)-approved indications, mechanism of action, efficacy, safety profiles, tolerability, and ease of use.

The loop diuretics are approved for the treatment of edema and hypertension.<sup>3-6</sup> They primarily act in the thick ascending limb of the loop of Henle to increase the urinary excretion of sodium, chloride, and water. Furosemide and ethacrynic acid also inhibit the absorption of sodium and chloride in the proximal and distal tubules. Bumetanide may also have an additional action in the proximal tubule. The loop diuretics are considered to be the most potent diuretics.<sup>3-7</sup> When given at their maximum dosages, they can lead to the excretion of up to 20% to 25% of the filtered sodium. As renal function declines (glomerular filtration rate <30 mL/minute), a loop diuretic should be considered rather than a thiazide diuretic. Loop diuretics do not possess the added property of arterial vasodilation, as seen with the thiazide diuretics.<sup>1,2</sup> Some studies have suggested that hydrochlorothiazide (a thiazide diuretic) is more effective in lowering blood pressure than the loop diuretics.<sup>8</sup>

The loop diuretics that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All agents are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Loop Diuretics Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Bumetanide	injection, tablet	N/A	bumetanide
Ethacrynate sodium	injection^	Sodium Edocrin <sup>®*</sup>	none
Ethacrynic acid	tablet	Edocrin <sup>®*</sup>	ethacrynic acid
Furosemide	injection, solution, tablet	Lasix <sup>®*</sup>	furosemide
Torsemide	tablet	N/A	torsemide

\*Generic is available in at least one dosage form or strength.

^Product is primarily administered in an institution.

PDL=Preferred Drug List

N/A=Not available

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the loop diuretics are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Loop Diuretics**

Clinical Guideline	Recommendation(s)
American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure	<b>Treatment of Stage A heart failure (HF)</b> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding</li> </ul>

Clinical Guideline	Recommendation(s)
(2022) <sup>9</sup>	<p>smoking are helpful to reduce future risk of HF. (LoE: B)</p> <ul style="list-style-type: none"> <li>● Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><u>Treatment of Stage B heart failure</u></p> <ul style="list-style-type: none"> <li>● In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>● In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>● In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>● In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>● In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>● In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>● In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>● In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><u>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</u></p> <ul style="list-style-type: none"> <li>● For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>● In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>● For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>● In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>● In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>● In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>● In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>● ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>● ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>● In patients with HFrEF, with current or previous symptoms, use of one of the</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>three <math>\beta</math>-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is <math>&gt;30</math> mL/min/1.73 m<sup>2</sup> and serum potassium is <math>&lt;5.0</math> mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at <math>&lt;5.5</math> mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq 35\%</math>) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq 70</math> beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>10</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>prognosis.</p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>11</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>blood pressure <math>\geq 140</math> mm Hg or diastolic blood pressure <math>\geq 90</math> mm Hg and to a goal systolic blood pressure <math>&lt; 140</math> mm Hg and goal diastolic blood pressure <math>&lt; 90</math> mm Hg.</p> <ul style="list-style-type: none"> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>12</sup></p>	<p><u>Lifestyle modifications</u></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio <math>&lt; 0.5</math> is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment           <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment           <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>• Core drug-treatment strategy           <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>13</sup></p>	<p><b>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</b></p> <ul style="list-style-type: none"> <li>• Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>• Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>• For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>selection for intensive management is recommended and caution should be taken in certain high-risk groups.</p> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients &lt;60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt; 40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (<math>&lt; 40\%</math>) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium <math>&lt; 5.2</math> mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours)</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> <li>● BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>● BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>● Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>● The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>● Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>● For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>● In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>● The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>● Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>● Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>● Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>● Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>● Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>● For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>ACE inhibitor or an ARB is recommended as initial therapy.</p> <ul style="list-style-type: none"> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>≥140 mmHg or DBP measurements of ≥90 mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</p> <ul style="list-style-type: none"> <li>• Women with severe hypertension with SBP ≥160 mmHg or DBP ≥110 mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>14</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), β-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</p> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p>



Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP <math>&lt; 220</math> mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to <math>&lt; 180</math> mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at <math>&lt; 180/105</math> mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</p> <ul style="list-style-type: none"> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>15</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>bendroflumethiazide or hydrochlorothiazide.</p> <ul style="list-style-type: none"> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on “Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence”.</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African–Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person’s medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in</b></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however,</li> </ul>

Clinical Guideline	Recommendation(s)
<p><b>Blacks (2010)<sup>16</sup></b></p>	<p>the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</p> <ul style="list-style-type: none"> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p><b>Kidney Disease Improving Clinical Outcomes Group: KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)<sup>17</sup></b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</p> <ul style="list-style-type: none"> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><b>BP management in children with CKD</b></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>18</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</p> <ul style="list-style-type: none"> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> <li>• For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</p> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP <math>\geq</math>180 mmHg or DBP <math>\geq</math>110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>19</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq</math>140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq</math>180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq</math>15%), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</p> <ul style="list-style-type: none"> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq</math>300 mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq</math>300 mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq</math>300 mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> or <math>\geq</math>300 mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Association for the Study of Liver Diseases: <b>Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance (2021)</b><sup>20</sup></p>	<p><u>Treatment of ascites</u></p> <ul style="list-style-type: none"> <li>• Moderate sodium restriction (2 g or 90 mmol/day) and diuretics (spironolactone with or without furosemide) are the first-line treatment in patients with cirrhosis and grade 2 ascites.</li> <li>• After ascites is adequately mobilized, attempts should be made to taper the diuretics to the lowest dose necessary to maintain minimal or no ascites to prevent the development of adverse effects.</li> <li>• Fluid restriction is not necessary unless serum sodium is <math>&lt; 125</math> mmol/L.</li> <li>• In patients receiving diuretics, body weight and serum creatinine and sodium should be regularly monitored to assess response and to detect the development of adverse effects.</li> <li>• Human albumin solution (20 to 40 g/week) or baclofen administration (10 mg/day, with a weekly increase of 10 mg/day, up to 30 mg/day) can be considered in cases of severe muscle cramps.</li> <li>• Large-volume paracentesis is the first-line treatment of grade 3 ascites. After paracentesis, sodium restriction and diuretics should be started.</li> <li>• Referral for liver transplant evaluation should be considered in patients with grade 2 or 3 ascites.</li> <li>• Nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers should be avoided in patients with cirrhosis and ascites.</li> <li>• Aminoglycosides should be avoided whenever possible in the treatment of bacterial infections.</li> <li>• For patients with cirrhosis and diuretic-responsive ascites, controversial data suggest potential benefits of long-term infusion of human albumin solution. At present, no recommendation can be made for its use in routine clinical practice.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the loop diuretics are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Loop Diuretics<sup>3-6</sup>**

Indication	Bumetanide	Ethacrynic Acid	Furosemide	Torsemide
<b>Edema</b>				
Adjunctive therapy in acute pulmonary edema			✓* (injection)	
Treatment of edema associated with congestive heart failure, hepatic disease, and renal disease	✓ †	✓ ‡	✓ §	✓
Rapid onset of diuresis is desired (e.g., in acute pulmonary edema) or when gastrointestinal absorption is impaired or oral therapy is not practical				✓ (injection)
Short-term management of ascites due to malignancy, idiopathic edema, and lymphedema		✓		
Short-term management of hospitalized pediatric patients, other than infants, with congenital heart disease or nephrotic syndrome		✓		
<b>Hypertension</b>				
Treatment of hypertension			✓    ¶	✓

\*The intravenous administration of furosemide is indicated when a rapid onset of diuresis is desired.

†If impaired gastrointestinal absorption is suspected or oral administration is not practical, bumetanide should be given by the intramuscular or intravenous route.

‡Treatment of edema when an agent with greater diuretic potential than those commonly employed is required.

§If impaired gastrointestinal absorption is suspected or oral administration is not practical, furosemide should be given by the intramuscular or intravenous route.

|| Alone or in combination with other antihypertensive agents.

¶If impaired gastrointestinal absorption is suspected or oral administration is not practical, furosemide should be given by the intramuscular or intravenous route.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the loop diuretics are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Loop Diuretics<sup>7</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
Bumetanide	80 to 95	90 to 99	Liver, partial (% not reported)	Bile (2) Feces (10 to 20) Renal (50 to 81)	1 to 1.5
Ethacrynic acid	100	90	Liver (% not reported)	Renal (66)	1 to 4
Furosemide	47 to 70	91 to 99	Liver (10)	Feces (7 to 9) Renal (60 to 90)	0.5 to 2
Torsemide	80 to 90	99	Liver (80)	Renal (69)	3 to 6

### V. Drug Interactions

Major drug interactions with the loop diuretics are listed in Table 5.

**Table 5. Major Drug Interactions with the Loop Diuretics<sup>7</sup>**

Generic Name(s)	Interaction	Mechanism
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Desmopressin	Concomitant use of desmopressin nasal spray and a loop diuretic is contraindicated due to an increased risk of severe hyponatremia.
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Cisapride	Possible additive prolongation of the QT interval due to electrolyte loss increases the risk of life-threatening cardiac arrhythmias, including torsades de pointes.
Loop diuretics (bumetanide, ethacrynic acid, furosemide)	Digitalis Glycosides	Diuretic-induced electrolyte disturbances may predispose patients to digitalis-induced arrhythmias.
Loop diuretics (ethacrynic acid, furosemide, torsemide)	Aminoglycosides	Auditory toxicity may be increased by possible synergistic activity. The mechanism is unknown.
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Cisplatin	The combination of loop diuretics and cisplatin may cause additive ototoxicity through an unknown mechanism.
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Lithium	Increased plasma lithium concentrations increase risk of toxicity. The mechanism is unknown.
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Thiazide diuretics	The two classes of agents exhibit their diuretic action at different sites in the renal tubules and have synergistic effects that may result in profound diuresis and serious electrolyte abnormalities.
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Dofetilide	Concurrent use of dofetilide and diuretics may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Loop diuretics (bumetanide, ethacrynic acid, furosemide, torsemide)	Sotalol	Concurrent use of sotalol and diuretics may result in an increased risk of cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest).
Loop diuretics (bumetanide)	Indomethacin	Concurrent use of bumetanide and indomethacin may result in reduced diuretic effectiveness and possible nephrotoxicity.

## VI. Adverse Drug Events

The most common adverse drug events reported with the loop diuretics are listed in Table 6. The boxed warning for the loop diuretics is listed in Table 7.

**Table 6. Adverse Drug Events (%) Reported with the Loop Diuretics<sup>3-7</sup>**

Adverse Events	Bumetanide	Ethacrynic Acid	Furosemide	Torsemide
<b>Cardiovascular</b>				
Atrial Fibrillation	-	-	-	✓
Chest pain	<1	-	-	1
Edema	-	-	-	1
Electrocardiogram changes	<1	-	-	2
Hypotension	<1	-	✓	✓
Hypovolemia	-	✓	-	✓
Myalgia	-	-	-	2
Orthostatic hypotension	<1	-	✓	✓
Shunt thrombosis	-	-	-	✓
Syncope	-	-	-	✓
Ventricular tachycardia	-	-	-	✓
<b>Central Nervous System</b>				
Apprehension	-	✓	-	-

Adverse Events	Bumetanide	Ethacrynic Acid	Furosemide	Torsemide
Asterixis	<1	-	-	-
Asthenia	-	-	✓	2
Confusion	-	✓	-	-
Dizziness	1	-	✓	3
Fatigue	<1	✓	-	✓
Headache	<1	✓	✓	7
Insomnia	-	-	-	1
Nervousness	-	-	-	1
Paresthesia	-	-	✓	-
Restlessness	-	-	✓	-
Vertigo	<1	✓	✓	-
Xanthopsia	-	-	✓	-
<b>Dermatologic</b>				
Erythema multiforme	-	-	✓	-
Exfoliative dermatitis	-	-	✓	-
Hives	<1	-	-	-
Itching	<1	-	-	-
Pruritus	<1	-	✓	✓
Rash	<1	✓	✓	✓
Photosensitivity	-	-	✓	-
Purpura	-	-	✓	-
Scaling eczema	-	-	✓	-
Stevens-Johnson Syndrome	-	-	✓	-
Urticaria	-	-	✓	-
<b>Endocrine and Metabolic</b>				
Acute gout	-	✓	-	✓
Dehydration	<1	-	-	-
Electrolyte imbalance	-	-	✓	✓
Nipple tenderness	<1	-	-	-
<b>Gastrointestinal</b>				
Abdominal discomfort/pain	<1	✓	-	-
Anorexia	-	✓	✓	-
Constipation	-	-	✓	2
Diarrhea	<1	✓	✓	2
Dry mouth	<1	-	-	-
Dyspepsia	<1	-	✓	2
Dysphagia	-	✓	-	✓
Gastrointestinal bleed	-	✓	-	✓
Loss of appetite	-	-	✓	-
Malaise	-	✓	-	-
Nausea	<1	✓	✓	2
Pancreatitis	-	✓	✓	-
Polydipsia	-	-	-	✓
Vomiting	<1	✓	✓	✓
<b>Genitourinary</b>				
Difficulty maintaining an erection	<1	-	-	-
Premature ejaculation	<1	-	-	-
<b>Hematologic</b>				
Agranulocytosis	-	✓	✓	-
Anemia	-	-	✓	-
Aplastic anemia	-	-	✓	-
Deviations in differential counts	<1	-	-	-
Deviations in hematocrit	<1	-	-	-

Adverse Events	Bumetanide	Ethacrynic Acid	Furosemide	Torsemide
Deviations in hemoglobin	<1	-	-	-
Deviations in prothrombin time	<1	-	-	-
Deviations in white blood cell count	<1	-	-	-
Hemolytic anemia	-	-	✓	-
Henoch-Schönlein purpura	-	✓	-	-
Leukopenia	-	-	✓	-
Neutropenia	-	✓	-	-
Thrombocytopenia	<1	✓	✓	-
<b>Hepatic</b>				
Abnormal liver enzymes	✓	✓	-	-
Encephalopathy	<1	-	-	-
Jaundice	-	✓	✓	-
<b>Laboratory Test Abnormalities</b>				
Azotemia	11	-	-	-
Changes in alkaline phosphatase	<1	-	-	-
Changes in cholesterol	<1	-	-	-
Changes in serum proteins	<1	-	-	-
Changes in total serum bilirubin	<1	-	-	-
Hyperlipidemia	✓	✓	✓	✓
Hyperglycemia	7	✓	✓	✓
Hyperuricemia	18	✓	✓	✓
Hypernatremia	<1	-	-	-
Hypocalcemia	✓	✓	✓	✓
Hypochloremia	15	-	-	-
Hypoglycemia	-	✓	-	-
Hypokalemia	15	✓	✓	✓
Hypomagnesemia	✓	✓	✓	✓
Hyponatremia	9	-	-	-
Serum creatinine increased	7	-	-	-
Variations in bicarbonate	3	-	-	-
Variations in calcium	2	-	-	-
Variation in CO <sub>2</sub> content	4	-	-	-
Variations in phosphorus	5	-	-	-
<b>Musculoskeletal</b>				
Arthralgia	-	-	-	2
Arthritic pain	<1	-	-	✓
Muscle cramps	1	-	✓	✓
Musculoskeletal pain	<1	-	-	-
Spasticity	-	-	✓	-
<b>Renal</b>				
Changes in creatinine clearance	<1	-	-	-
Glycosuria	<1	-	✓	-
Hematuria	-	✓	-	-
Interstitial nephritis	-	-	✓	-
Polyuria	-	-	-	7
Proteinuria	<1	-	-	-
Renal Failure	<1	-	-	-
<b>Respiratory</b>				
Cough	-	-	-	2
Hyperventilation	<1	-	-	-
Rhinitis	-	-	-	3
<b>Special Senses</b>				
Blurred vision	-	✓	✓	-

Adverse Events	Bumetanide	Ethacrynic Acid	Furosemide	Torsemide
Deafness	-	✓	-	-
Ear discomfort	<1	-	-	-
Fullness of ears	-	✓	-	-
Impaired hearing	<1	✓	✓	-
Ototoxicity	✓	✓	✓	✓
Tinnitus	-	✓	✓	-
<b>Other</b>				
Angioedema	-	-	-	✓
Chills	-	✓	-	-
Fever	-	✓	✓	-
Necrotizing angitis	-	-	✓	-
Systemic vasculitis	-	-	✓	-
Sore Throat	-	-	-	2
Sweating	<1	-	-	-
Thrombophlebitis	-	-	✓	-
Weakness	<1	-	✓	✓

✓ Percent not specified  
- Event not reported

**Table 7. Boxed Warning for Bumetanide and Furosemide<sup>6</sup>**

<b>WARNING</b>
Loop diuretics are potent diuretics which, if given in excessive amounts, can lead to a profound diuresis with water and electrolyte depletion. Therefore, careful medical supervision is required and dose and dosage schedule have to be adjusted to the individual patient's needs.

## VII. Dosing and Administration

The usual dosing regimens for the loop diuretics are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Loop Diuretics<sup>3-7</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Bumetanide	<u>Edema:</u> Injection: 0.5 to 1 mg over one minute; maximum, 10 mg/day  Tablet: 0.5 to 2 mg/day; maximum, 10 mg/day	Safety and efficacy in children have not been established.	Injection: 0.25 mg/mL  Tablet: 0.5 mg 1 mg 2 mg
Ethacrynic acid	<u>Edema:</u> Tablet: 50 to 200 mg/day	<u>Edema (13 months and older):</u> Tablet: initial, 25 mg	Tablet: 25 mg
Furosemide	<u>Edema:</u> Injection (acute pulmonary edema): 40 mg intravenously over 1 to 2 minutes; maintenance, may increase to 80 mg intravenously  Injection: 20 to 40 mg as a single intravenous or intramuscular injection; maintenance, may repeat in two hours or increased by 20 mg until desired response	<u>Edema:</u> Injection: initial, 1 mg/kg; maintenance, may increase by 1 mg/kg not sooner than two hours after the previous dose; maximum, 6 mg/kg per dose  Solution, tablet: 2 mg/kg as a single dose; maximum, 6 mg/kg per dose	Injection: 10 mg/mL  Solution: 10 mg/mL 40 mg/5 mL  Tablet: 20 mg 40 mg 80 mg

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
	<p>Oral: 20 to 80 mg/days; maximum, 600 mg/day</p> <p><u>Hypertension:</u> Injection, solution, tablet: 80 mg/day</p>		
Toremide	<p><u>Edema:</u> Injection, tablet (chronic renal failure): initial, 20 mg once daily; maintenance, 200 mg/day</p> <p>Injection, tablet (congestive heart failure): initial, 10 to 20 mg once daily; maximum, 200 mg/day</p> <p>Injection, tablet (hepatic cirrhosis): initial, 5 to 10 mg once daily; maximum, 40 mg/day</p> <p><u>Hypertension:</u> Injection, tablet: initial, 5 to 10 mg/day; maximum, 10 mg/day</p>	Safety and efficacy in children have not been established.	Tablet: 5 mg 10 mg 20 mg 100 mg



## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the loop diuretics are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Loop Diuretics**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Cirrhosis</b>				
Laffi et al. <sup>21</sup> (1991)  Furosemide 25 mg/day  vs  torasemide* 10 mg/day	DB, RCT  Nonazotemic cirrhotic patients with ascites	N=24  3 days	Primary: Percent increase in natriuresis, body weight loss, percent increase in diuresis, plasma aldosterone concentration, plasma renin activity  Secondary: Not reported	Primary: Treatment with torasemide led to significantly greater natriuresis than furosemide (P<0.02). There was a greater percentage increase in basal values (day 1: 130 vs 50%; day 2: 104 vs 42%; and day 3: 65 vs 26%, respectively).  Body weight loss was significantly higher with torasemide (2.5 kg) than with furosemide (1.3 kg; P<0.02).  There was no significant difference (P=0.08) in the percent increase in diuresis among the treatment groups (day 1: 60 vs 26%; day 2: 35 vs 27%; day 3: 31 vs 24%).  Plasma aldosterone concentrations (ng/mL) with torasemide were 0.79 and 0.94 at baseline and day three, respectively. Plasma aldosterone concentrations with furosemide were 0.54 and 0.52 at baseline and day three, respectively.  Plasma renin activity (ng/mL/hr) with torasemide were 5.8 and 9.4 at baseline and day three, respectively. Plasma renin activity with furosemide were 4.2 and 5.4 at baseline and day three, respectively.  Secondary: Not reported
Gerbes et al. <sup>22</sup> (1993)  Furosemide 80 mg as a single dose  vs	DB, RCT, XO  Patients with cirrhosis and ascites	N=28  24 hours	Primary: Urine volume, urine sodium volume, urine potassium volume, plasma aldosterone concentration,	Primary: Treatment with torasemide led to greater cumulative 24 hour diuresis than furosemide (2,863 vs 2,111; P<0.05).  There was no difference in cumulative 0 to 6 hour sodium excretion with torasemide or furosemide (95.7 vs 92.1 mmol, respectively; P value not significant). There was greater cumulative 6 to 24 hour sodium excretion

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>torasemide* 20 mg as a single dose</p>			<p>plasma renin activity</p> <p>Secondary: Not reported</p>	<p>with torasemide compared to furosemide (38.4 vs 16.6 mmol; P&lt;0.05). There was no difference in cumulative 0 to 24 hour sodium excretion with torasemide or furosemide (134.0 vs 108.5 mmol, respectively; P value not significant).</p> <p>There was no difference in cumulative 0 to 6 hour potassium excretion with torasemide or furosemide (57.5 vs 39.9 mmol, respectively; P value not significant). There was greater cumulative 6 to 24 hour potassium excretion with torasemide compared to furosemide (36.0 vs 27.6 mmol; P&lt;0.05). There was no difference in cumulative 0 to 24 hour potassium excretion with torasemide or furosemide (88.3 vs 68.0 mmol, respectively; P value not significant).</p> <p>Plasma aldosterone concentrations (ng/100 mL) with torasemide were 111.9 and 132 at baseline and 24 hours, respectively. Plasma aldosterone concentrations with furosemide were 105.7 and 131 at baseline and 24 hours, respectively.</p> <p>Plasma renin activity (ng/mL/hr) with torasemide were 29.9 and 30.6 at baseline and 24 hours, respectively. Plasma renin activity with furosemide were 34.7 and 36.8 at baseline and 24 hours, respectively.</p> <p>Secondary: Not reported</p>
<p>Fiaccadori et al.<sup>23</sup> (1993)</p> <p>Furosemide 50 mg/day</p> <p>vs</p> <p>torasemide* 20 mg/day</p> <p>Patients also received</p>	<p>DB, RCT</p> <p>Nonazotemic cirrhotic patients with controlled ascites</p>	<p>N=28</p> <p>10 weeks</p>	<p>Primary: Excretion of phosphate, free water, sodium, potassium, calcium, and uric acid</p> <p>Secondary: Not reported</p>	<p>Primary: Furosemide produced more excretion of phosphates (P&lt;0.001) and magnesium (P&lt;0.05) compared to torasemide.</p> <p>Torasemide produced more excretion of free water (P&lt;0.02).</p> <p>There was no difference in the excretion of sodium, potassium, calcium, or uric acid among the treatment groups.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
spironolactone 200 mg/day				
Abecasis et al. <sup>24</sup> (2001)  Frusemide† 40 mg/day  vs  torsemide 20 mg/day  Patients also received spironolactone 200 mg/day	OL, RCT  Cirrhotic patients with ascites	N=46  11 to 12 days	Primary: Resolution of ascites, weight loss, diuretic dosage, diuretic response  Secondary: Not reported	Primary: There was no difference in the percentages of patients with resolution of ascites with torsemide compared to frusemide (73 vs 75%; P value not significant).  There was no difference in weight loss with torsemide compared to frusemide (8 vs 8.5 kg; P value not significant).  More patients receiving frusemide required an increase in diuretic dosage (37.5%) than with torsemide (9%; P<0.05).  Torsemide produced a greater diuretic response in 24 hours than frusemide (P<0.007).
<b>Heart Failure/Edema</b>				
Galløe et al. <sup>25</sup> (2006)  Bumetanide 0.5 mg (0, 1, 2 or 4 tablets BID) plus trandolapril 0.5 mg (0, 1, 2 or 4 tablets QD)  Treatment was combined to achieve 16 different dosage combinations.	DB, DD, RCT, multiple XO  Patients with previous MI ≥3 years ago, had medical treatment for heart failure and ejection fraction between 0.36 and 0.54 estimated by echocardiography	N=16  14 days	Primary: Patient reported quality of life  Secondary: Effects on kidney function, left ventricular function and blood pressure	Primary: Bumetanide 0.5 mg-treated patients experienced a 12% increase in well-being, but higher doses of bumetanide decreased patient's well-being by 12% compared to placebo (P<0.002). Increasing doses of bumetanide tended to increase tiredness (P=0.072). There were no significant effects of bumetanide therapy on the patients' opinion of their health, degree of dyspnea, appetite or work capacity.  Secondary: Bumetanide therapy increased 24 hour urine production in a straight dose-dependent manner (P<0.0001), while trandolapril therapy had no effect (P=0.53). Bumetanide and trandolapril therapy did not alter the 24 hour creatinine excretion and creatinine clearance (P=0.33, P=0.11 and P=0.53, P=0.97, respectively).  Bumetanide therapy decreased left ventricular function and increased heart rate in a dose-dependent manner (P<0.001). Left ventricular function was also nonsignificantly decreased with trandolapril therapy (P>0.062).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Trandolapril therapy significantly reduced SBP by maximally of 7.6 mm Hg (5.8%) with the lowest dose of 0.5 mg/day (P=0.007). Bumetanide therapy had no significant effect on DBP (P=0.23).
Hutcheon et al. <sup>26</sup> (1981)  Bumetanide 1 to 2 mg/day  vs  furosemide 80 mg/day	DB, PG  Patients with severe edema associated with CHF	N=20  3 days	Primary: Edema, symptoms of heart failure, safety and tolerability  Secondary: Not reported	Primary: Each agent was effective in decreasing the edema and relieving the symptoms of heart failure.  Side effects were not severe and were similar in both treatment groups. Muscle cramps and abdominal pain were deemed not severe. Electrolyte shifts indicative of hypochloremic alkalosis and hyponatremia were seen in two patients in the bumetanide group.  Secondary: Not reported
Konecke et al. <sup>27</sup> (1981)  Bumetanide  vs  furosemide  No dose or frequency reported.	OL, PG, RCT  Men and women with clinically detectable edema and signs and symptoms of CHF (e.g., rales, gallop rhythm, orthopnea, dyspnea, engorged neck veins, paroxysmal nocturnal dyspnea, congested liver, etc.)	N=42  6 months	Primary: Changes in weight, blood pressure, pulse, signs and symptoms of CHF, electrolytes and functional capacity, safety and tolerability  Secondary: Not reported	Primary: There were no statistical differences in changes in body weight, blood pressure, edema, abdominal girth, and hepatomegaly and other signs and symptoms of CHF in patients receiving bumetanide vs furosemide.  There were variable minor changes in serum sodium, potassium, chloride, and uric acid in both groups throughout the treatment. Changes remained within normal limits and reached significance for chloride at weeks eight and 16 in the bumetanide group.  Functional capacity improved slightly or remained unchanged throughout treatment in both treatment groups.  There were no major side effects that were medication related in both treatment groups.  Secondary: Not reported
Nicholson et al. <sup>28</sup> (1977)  Bumetanide 1 mg/day alternating	RCT, XO  Patients with cirrhosis and fluid overload	N=10  6 months	Primary: Ascites and edema  Secondary: Adverse events	Primary: Bumetanide and frusemide were both effective in controlling ascites and edema, with nine out of 10 patients showing a satisfactory response.  Secondary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>with 3 mg/day for 3 months</p> <p>vs</p> <p>frusemide† 40 mg/day alternating with 160 mg/day for 3 months</p>				<p>Side effects were reported in six patients. The most common side effects were urinary frequency and nocturia, which occurred in four patients taking bumetanide and 1 patient taking frusemide. There was one patient on bumetanide and one patient on frusemide who developed symptoms of postural hypotension.</p>
<p>Eshaghian et al.<sup>29</sup> (2006)</p> <p>Furosemide 0 to 40 mg/day (group 1)</p> <p>vs</p> <p>furosemide 41 to 80 mg/day (group 2)</p> <p>vs</p> <p>furosemide 81 to 160 mg/day (group 3)</p> <p>vs</p> <p>furosemide &gt;160 mg/day (group 4)</p>	<p>Cohort</p> <p>Men and women with advanced systolic heart failure referred to a single university medical center for heart failure management and/or transplant evaluation from 1985 to 2004</p>	<p>N=1,354</p> <p>2 years</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Composite endpoint of death or urgent transplant</p>	<p>Primary: There were 269 deaths during the two year follow-up, with 182 deaths by year one and 87 deaths during year two. Of the 269 deaths, 91 deaths were due to progressive heart failure, 72 deaths were sudden, eight deaths were secondary to myocardial infarction and 101 were unknown.</p> <p>Survival estimates at one year were 91, 88, 80, and 69% for groups 1, 2, 3, and 4, respectively (P&lt;0.0001). Survival estimates at two years were 83, 81, 68, and 53%, respectively (P&lt;0.0001).</p> <p>Secondary: There were a total of 431 patients who received heart transplants by the end of the two year follow up: 223 urgent and 208 elective.</p> <p>The HRs for death from any cause, death and urgent transplantation, death from progressive heart failure, and sudden death for group 4 compared with group 1 were similar.</p> <p>On univariate analysis, compared with group 1, increasing loop diuretic dose were associated with a progressive increase in mortality (group 2: HR, 1.2; 95% CI, 0.8 to 1.7, group 3: HR, 2.1; 95% CI, 1.5 to 2.9, and group 4: HR, 3.4; 95% CI, 2.4 to 4.7).</p>
<p>Mentz et al.<sup>30</sup> (2016)</p> <p>ASCEND-HF</p> <p>Furosemide</p>	<p>Cohort analysis of ASCEND-HF (diuretic choice not randomized)</p> <p>Patients enrolled in</p>	<p>N=4,177</p>	<p>Primary: All-cause mortality or HF hospitalization through 30-days after discharge</p>	<p>Primary &amp; Secondary: Of the 4,177 patients in the outcomes analysis cohort, 87% (n=3,620) received furosemide and 13% (n=557) received torsemide. Torsemide was associated with similar outcomes on unadjusted analysis (30-day mortality/HF hospitalization OR, 1.03; 95% CI, 0.73 to 1.45, P=0.88 and 180-day mortality HR, 0.97; 95% CI, 0.73 to 1.29; P=0.83). On inverse</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs torsemide	the ASCEND-HF Trial (patients with acute HF) discharged alive on either furosemide or torsemide		Secondary: 30-day all-cause mortality, 30-day HF hospitalization and 180-day all-cause mortality post-discharge	propensity weighted-adjusted analysis, torsemide use was associated with nominally lower 30-day mortality or HF hospitalization (OR, 0.89; 95% CI, 0.62 to 1.29; P=0.55), 30-day mortality (OR, 0.89; 95% CI, 0.40 to 1.97; P=0.78), 30-day HF hospitalization (OR, 0.87; 95% CI, 0.58 to 1.30; P=0.49) and 180-day mortality (HR, 0.86; 95% CI, 0.63 to 1.19; P=0.37) compared with furosemide.
Murray et al. <sup>31</sup> (2001)  Furosemide  vs  torsemide	OL, RCT  Patients with CHF	N=234  12 months	Primary: Readmission to the hospital for heart failure  Secondary: Readmission for all cardiovascular causes and for all causes, numbers of hospital days, health-related quality of life	Primary: Patients receiving torsemide were less likely to need readmission for heart failure (32%) compared to furosemide (17%; P<0.01).  Secondary: Patients receiving torsemide were less likely to need readmission for all cardiovascular causes (59%) compared to furosemide (44%; P=0.03).  There was no difference in the rate of admissions for all causes among the treatment groups (76 vs 71%; P=0.36).  Patients treated with torsemide had significantly fewer hospital days for heart failure (106 vs 296 days; P=0.02).  Improvements in fatigue scores from baseline were significantly greater among patients treated with torsemide compared to furosemide at months 2, 8, and 12 (P<0.05).
Cosín et al. <sup>32</sup> (2002)  Furosemide 40 mg/day orally or other diuretics  vs  torasemide* 10 mg/day orally	OL  Patients with NYHA functional class II to III heart failure	N=1,377  12 months	Primary: Mortality, morbidity, functional class and serum potassium levels (<3.5 or >5 mEq/L)  Secondary: Not reported	Primary: Total mortality was significantly lower in the torasemide group (2.2%) compared to the furosemide/other diuretics group (4.5%; P<0.05).  Cardiac mortality was lower in patients receiving torasemide (1.4%) than in those receiving furosemide/other diuretics (3.5%; P<0.05).  NYHA improvement in at least 1 class occurred in more patients who received torasemide (45.8%) than those who received furosemide/other diuretics (37.2%; P=0.00017).  Abnormal potassium levels were observed in fewer torasemide patients

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				(12.9%) than furosemide/other diuretics patients (17.9%; P=0.013).
Muller et al. <sup>33</sup> (2003)  Furosemide  vs  torasemide*	R, OL  Patients with NYHA functional class II-IV congestive heart failure	N=237  9 months	Primary: Clinical improvement in heart failure, quality of life, hospitalizations, safety and tolerability  Secondary: Not reported	Primary: Clinical improvement in chronic heart failure was seen in both groups, but the trend to improve by at least one NYHA class was significant with torasemide (P=0.014) compared to furosemide-treated patients.  There were no differences in adverse reactions and hospitalizations due to CHF.  Secondary: Not reported
Kasama et al. <sup>34</sup> (2006)  Furosemide 20 to 40 mg/day  vs  torasemide* 4 to 8 mg/day	RCT  Patients with non-ischemic CHF (LVEF <45%) also being treated with an ACE inhibitor	N=40  6 months	Primary: Effect on cardiac sympathetic nerve activity (delayed heart to mediastinum count ratio, delayed total defect score, washout rate)  Secondary: Effect on left ventricular remodeling (left ventricular end diastolic volume, left ventricular end systolic volume)	Primary: In the furosemide group at the end of treatment, mean heart to mediastinum count ratio increased from 1.68±0.18 to 1.71±0.19 (P value not significant), mean total defect score decreased from 42±11 to 40±12 (P value not significant), and mean washout rate decreased from 50±8% to 47±12% (P value not significant).  In the torasemide group at the end of treatment, mean heart to mediastinum count ratio increased from 1.61±0.19 to 1.77±0.24 (P<0.001), mean total defect score decreased from 44±8 to 36±8 (P<0.001), and mean washout rate decreased from 52±12 to 41±14% (P=0.001).  Secondary: In the furosemide group left ventricular end diastolic volume decreased from 174±24 to 165±34 mL (P value not significant), left ventricular end systolic volume decreased from 120±15 to 109±33 mL (P value not significant), and LVEF increased from 31±7 to 32±7% (P value not significant).  In the torasemide group left ventricular end diastolic volume decreased from 173±22 to 147±30 mL (P<0.01), left ventricular end systolic volume decreased from 117±19 to 95±25 mL (P<0.001), and LVE increased from 31±7 to 34±7% (P value not significant).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Levy et al.<sup>35</sup> (1977)</p> <p>Furosemide 25 mg daily for 24 weeks</p> <p>vs</p> <p>spironolactone and HCTZ 25-25 mg/day (fixed-dose combination product) for 16 weeks following 8 weeks of furosemide monotherapy</p>	<p>DB, RCT</p> <p>Patients 27 to 79 years of age with arteriosclerotic heart disease, hypertensive heart disease, or rheumatic heart disease classes 1 to 3, and congestive heart failure requiring diuretic therapy</p>	<p>N=32</p> <p>24 weeks</p>	<p>Primary: Change in heart failure symptoms, glucose, renin concentration, calcium, blood urea nitrogen, uric acid, creatinine, aldosterone, serum potassium level, adverse effects</p> <p>Secondary: Not reported</p>	<p>Primary: The combination therapy group and furosemide monotherapy group exhibited comparable control of heart failure symptoms.</p> <p>The combination therapy group was associated with a significant decrease in glucose and an increase in plasma renin concentration compared to furosemide monotherapy group (P&lt;0.01).</p> <p>There were no significant differences in calcium, blood urea nitrogen, uric acid, or creatinine between the study groups.</p> <p>There was a significant increase in aldosterone secretion among patients randomized to the spironolactone and HCTZ group compared to the furosemide group (P&lt;0.01).</p> <p>There was no significant difference in serum potassium level between treatment groups.</p> <p>No serious adverse effects were observed in either of the study groups.</p> <p>Secondary: Not reported</p>
<p>Austin et al.<sup>36</sup> (1976)</p> <p>Furosemide 40 to 60 mg infused through a pulmonary artery catheter</p> <p>vs</p> <p>ethacrynic acid 25 to 50 mg infused through a pulmonary artery</p>	<p>OS</p> <p>Men and women who underwent diagnostic right and transeptal left heart catheterization with chronic postcapillary pulmonary HTN with heart failure NYHA class II to IV</p>	<p>N=27</p> <p>1 hour</p>	<p>Primary: Hemodynamic response (in the control state and at 20, 40, and 60 minutes after diuretic administration) including cardiac index, pulmonary artery, left atrial and systemic artery mean pressures, plasma volume, PBV and PEV</p>	<p>Primary: The hemodynamic response with each medication was similar. When compared to control state, the reductions in pulmonary artery mean pressure at 20, 40, and 60 minutes after diuretic infusion with either ethacrynic acid or furosemide were significant (P&lt;0.001).</p> <p>The average left atrial mean pressure also decreased from 22 mm Hg during the control period to 18 mm Hg at 20 minutes and to 15 mm Hg at 60 minutes post diuretic infusion (ethacrynic acid or furosemide; P&lt;0.001).</p> <p>The mean cardiac index decreased significantly at 20, 40, and 60 minutes compared to the control state after diuretic infusion with either ethacrynic acid or furosemide (P&lt;0.001).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
catheter			Secondary: Not reported	There was a significant decrease in plasma volume at 60 minutes post drug infusions (ethacrynic acid or furosemide; $P < 0.001$ ).  In contrast, there was no significant change in PBV, PEV, PEV/PBV, and systemic arterial pressure throughout the study period with ethacrynic acid or furosemide.  Secondary: Not reported
Patterson et al. <sup>37</sup> (1994)  Torsemide 5 mg QD  vs  torsemide 10 mg QD  vs  torsemide 20 mg QD  vs  placebo QD	DB, MC, PC, PG  Men and women diagnosed with NYHA class II or III CHF and edema	N=66  7 days	Primary: Change in body weight from baseline  Secondary: Change in urinary sodium, potassium, chloride excretion and urine volume after the first dose of drug	Primary: Patients receiving torsemide 10 and 20 mg had a significant decrease in weight (-1.62 and -1.30 kg, respectively) as compared to placebo.  Torsemide 5 mg did not demonstrated a significant reduction in body weight compared to placebo (-0.60 kg).  Secondary: Severity of edema decreased as the dose of torsemide increased. The adverse events did not increase with higher doses of torsemide.
Senzaki et al. <sup>38</sup> (2008)  Torasemide* (de novo group)  vs  torasemide*	RCT  Pediatric patients (age range from 3 weeks to 17 years) with congested heart failure, patients newly diagnosed with	N=102  3 to 4 weeks	Primary: Clinical signs and symptoms of congestive heart failure  Secondary: Humoral factors, serum potassium	Primary The de novo torasemide group significantly improved the congestive heart failure index from $7.2 \pm 1.6$ to $5.7 \pm 1.4$ ( $P < 0.05$ ); however the replacement group did not. The replacement group baseline value of the congestive heart failure index was $7.4 \pm 2.4$ and after treatment the mean value was $6.8 \pm 2.3$ .  Secondary: The de novo and replacement groups significantly improved brain

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(replacement group) was converted from furosemide dosage using 0.2 mg torasemide* corresponding to 1 mg furosemide	CHF or previously treated with furosemide		levels, and adverse events	natriuretic peptide and aldosterone levels (P<0.05); however, plasma rennin activity was not significantly decreased among both groups.  Serum potassium levels were significantly increased in the replacement group (P<0.05), but not in the de novo group.  The most commonly reported adverse events of torasemide were those associated with loop diuretics in general.
Faris et al. <sup>39</sup> (2006)  Loop diuretics (furosemide, bumetanide), thiazide diuretics (chlorothiazide), or potassium-sparing diuretics (amiloride, triamterene)  vs  placebo or active control (ACE inhibitors, digoxin)	MA  Adult patients with chronic heart failure	N=525 (14 trials)  2 to 52 weeks	Primary: Mortality  Secondary: Effect of diuretic withdrawal on worsening of heart failure and exercise capacity	Primary: Mortality was reported in three of the seven placebo-controlled trials, and this analysis showed that mortality was lower for patients treated with diuretics than with placebo (3/111 [2.7%] vs 12/110 [10.9%], respectively; OR, 0.24; 95% CI, 0.07 to 0.83; P=0.02).  These results showed that patients treated with diuretics had an absolute risk reduction of 8% when compared to placebo and a number needed to treat of 12.5.  Secondary: An analysis of pooled data from two trials showed lower admission rates for worsening heart failure in patients taking diuretics than in patients taking placebo (OR, 0.07; 95% CI, 0.01 to 0.52; P=0.01).  Diuretics were found to improve exercise capacity, with a difference in means of 0.74 (95% CI, 0.37 to 1.11; P<0.0001) and of 0.67 (95% CI, 0.02 to 1.31; P=0.04.), respectively. The combined results of these 4 trials indicated that diuretics improved exercise capacity in participants with chronic heart failure with a difference in means of 0.72 (95% CI, 0.40 to 1.04; P<0.0001).
<b>Hypertension</b>				
Van der Heijden et al. <sup>40</sup> (1998)  Bumetanide 1 mg/day for 6 weeks	DB, PC, XO  Patients with HTN	N=27  24 weeks	Primary: Changes in blood pressure, serum lipid levels, lab values, safety and tolerability	Primary: Bumetanide and furosemide reduced SBP by 8.2% (P<0.0002) and DBP by 4.5% (P<0.002). Overall SBP and DBP measurements were 12 and 4 mm Hg lower, respectively, when receiving bumetanide or furosemide vs placebo.  Both furosemide and bumetanide increased TC by 5.0% (P<0.002), HDL-

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>furosemide 40 mg/day for 6 weeks</p>			<p>Secondary: Not reported</p>	<p>C by 1.7% (P value not significant), LDL-C by 4.8% (P&lt;0.01), and TG by 12.4% (P&lt;0.01).</p> <p>Serum glucose, magnesium, sodium, and potassium levels were unchanged in both treatment groups; whereas serum creatinine tended to increase (3.2%; P=0.09).</p> <p>Side effects were mild in severity with no discontinuation reported. In both bumetanide and furosemide treated patients, four patients reported hypertonic muscles, but was resolved within a couple of days.</p> <p>Secondary: Not reported</p>
<p>De Berrazueta et al.<sup>41</sup> (2007)</p> <p>Furosemide infused in 3 progressive solutions containing 475, 950, and 1,900 nmol/mL for arterial studies and 240, 480, and 960 nmol/mL for venous studies</p> <p>vs</p> <p>torasemide* infused in 3 solutions containing 400, 800, and 1,600 nmol/mL for</p>	<p>RCT</p> <p>Patients with HTN and healthy controls</p>	<p>N=59</p> <p>Single dose</p>	<p>Primary: Dilatory effect on arteries and veins</p> <p>Secondary: Not reported</p>	<p>Primary: There were no significant changes in arterial dilation. Furosemide increased vasodilatation from 0.56±0.09 to 0.88±0.06 (P=0.000) in healthy control subjects and from 0.49±0.10 to 0.75±0.12 (P=0.000) in hypertensive patients.</p> <p>Torsemide increased venodilation from 0.46±0.06 to 0.70±0.11 (P=0.007) in control subjects and from 0.48±0.09 to 0.67±0.12 (P=0.03) in hypertensive patients.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
arterial studies and 200, 400, and 800 nmol/mL for venous studies				
von Dossow et al. <sup>42</sup> (2008)  Furosemide 40 mg IV and 80 mg PO 2 hours after extubation on day 1 after surgery  vs  torasemide* 20 mg IV and 20 mg PO 2 hours after extubation on day 1 after surgery	DB, RCT  Patients with secondary pulmonary HTN scheduled for elective valve replacement and/or coronary artery bypass graft	N=21  Day 1 after surgery	Primary: Cardiac output  Secondary: Endothelin-1 and angiotensin-II	Primary: Cardiac output increased significantly (P=0.03) in the torasemide group compared to the furosemide group.  Secondary: Endothelin-1 and angiotensin-II increased significantly (P=0.031) in the furosemide group compared to the torasemide group.
Vasavada et al. <sup>43</sup> (2003)  <u>Phase 1: Inpatient</u> Furosemide 200 mg/day with sodium-free water (10 mL/kg)  vs  torsemide 100 mg/day with sodium-free water (10 mL/kg)	DB, RCT, two-phase, XO  Patients ≥18 years of age with chronic kidney disease (serum creatinine >1.4 mg/dL) and volume overload	N=14  3 weeks	Primary: <u>Phase 1: Inpatient</u> Change in 24-hour urinary sodium excretion  <u>Phase 2: Outpatient</u> Primary: 24-hour ambulatory SBP  Secondary: Potassium, calcium, protein excretion, diurnal variation of	Primary <u>Phase 1: Inpatient</u> Furosemide and torsemide increased urinary sodium excretion from 199 to 357 mEq/day and 213 to 398 mEq/day, respectively. These differences between the two diuretics were not statistically significant.  <u>Phase 2: Outpatient</u> Both treatments had similar effects in reducing SBP (P=0.43). The SBP was reduced from baseline to post treatment by 9.7 mm Hg for torsemide (P=0.007) and 9.2 mm Hg for furosemide (P=0.021).  Secondary: There were no significant differences in excretion rate profiles between torsemide and furosemide (P>0.17).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p><u>Phase 2:Outpatient</u> Furosemide 80 mg/day</p> <p>vs</p> <p>torsemide 40 mg/day</p>			<p>electrolyte and protein excretion, and glomerular filtration rate</p>	
<p>Pupita et al.<sup>44</sup> (1983)</p> <p>Furosemide 25 mg QD</p> <p>vs</p> <p>chlorthalidone 50 mg QD</p>	<p>RCT, XO</p> <p>Men and women with a mean age of 53.9±9.2 years with mild to moderate HTN</p>	<p>N=36</p> <p>12 months</p>	<p>Primary: Blood pressure</p> <p>Secondary: Plasma electrolytes, adverse events</p>	<p>Primary: Patients taking chlorthalidone had significantly lower SBP at each monthly measurement compared to baseline (P&lt;0.01). However, only DBP values at month five were significant compared to baseline (P&lt;0.05).</p> <p>Patients taking furosemide had significantly lower SBP at months three, four, and five compared to baseline (P&lt;0.05 for month three, and P&lt;0.01 for months four and five). DBP values were significantly lower at all monthly measurements compared to baseline in patients taking furosemide (P&lt;0.01).</p> <p>At month one, SBP decreased by 19.4 mm Hg with chlorthalidone and by 21.2 mm Hg with furosemide (P&lt;0.001). DBP decreased by 11 mm Hg with chlorthalidone and by 12.6 mm Hg with furosemide at month one (P&lt;0.001).</p> <p>Secondary: There were no significant changes in serum sodium levels with either chlorthalidone or furosemide. Patients taking chlorthalidone had significantly lower serum chloride levels compared to baseline at all points (P&lt;0.01), whereas patients taking furosemide had significantly lower levels only at month six (P&lt;0.05). Both chlorthalidone and furosemide significantly reduced serum potassium levels at all points compared to baseline (P&lt;0.01).</p> <p>Patient taking chlorthalidone reported adverse effects including dizziness, transient abdominal disorder, and slight weakness. Patients taking furosemide reported transient early weakness and irritability. The rate of adverse events was not statistically significant in either treatment group.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Valmin K et al.<sup>45</sup> (1975)</p> <p>Furosemide 12.5, 25 or 40 mg BID</p> <p>vs</p> <p>HCTZ 12.5 mg BID</p> <p>vs</p> <p>placebo</p>	<p>DB, RCT, XO, 5 experimental periods each of 4 weeks</p> <p>Men and women with essential HTN</p>	<p>N=34</p> <p>20 weeks</p>	<p>Primary: Blood pressure, urinary output, serum electrolytes, safety and tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: When compared to placebo, there was a significant reduction of blood pressure with HCTZ 12.5 mg BID and furosemide 12.5 mg BID (P&lt;0.05).</p> <p>Paired comparison showed that HCTZ 12.5 mg BID and furosemide 25 and 40 mg BID had a similar hypotensive effect, irrespective of the initial blood pressure (P&gt;0.10).</p> <p>When compared to placebo, the urinary output increased significantly with furosemide 12.5, 25, or 40 mg BID (P&lt;0.05, P&lt;0.01 and P&lt;0.001, respectively) but not with the HCTZ group (P&gt;0.10).</p> <p>Sodium level did not alter during the various treatment periods when compared with the placebo period, or between the individual treatment periods (P&gt;0.10).</p> <p>Potassium level fell significantly during the HCTZ period (P&lt;0.001) and furosemide 25 mg and 40 mg BID period (P&lt;0.01 and P&lt;0.001, respectively). Potassium level was not significantly affected with furosemide 12.5 mg BID (P&gt;0.10).</p> <p>Secondary: Not reported</p>
<p>Araoye et al.<sup>8</sup> (1978)</p> <p>Furosemide 40 mg BID</p> <p>vs</p> <p>HCTZ 50 mg BID</p>	<p>DB, XO</p> <p>Patients with HTN</p>	<p>N=not specified</p> <p>3 months</p>	<p>Primary: Blood Pressure</p> <p>Secondary: Not reported</p>	<p>Primary: Furosemide and HCTZ significantly reduced blood pressure. The decrease in blood pressure was consistently greater in the HCTZ group than with furosemide; however the difference was significant in regards to SBP only.</p> <p>Secondary: Not reported</p>
<p>Ogawa et al.<sup>46</sup> (2006)</p> <p>Furosemide 20 mg/day plus</p>	<p>PRO, RCT</p> <p>Adult patients with HTN and type 2 diabetes, with a</p>	<p>N=30</p> <p>24 months</p>	<p>Primary: Change in BNP, urine albumin/creatinine ratio, and blood pressure</p>	<p>Primary: At 12 months, spironolactone-treated patients exhibited a significant reduction in BNP level from baseline compared to furosemide-treated patients (P&lt;0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>imidapril‡ 5 mg/day</p> <p>vs</p> <p>spironolactone 25 mg/day plus imidapril‡ 5 mg/day</p> <p>All patients were pre-treated with imidapril‡ for 1 year prior to trial onset.</p>	<p>urine albumin/creatinine ratio &gt;30 mg/g creatinine, and plasma BNP levels &gt;100 pg/mL (suggestive of mild heart failure)</p>		<p>Secondary: Not reported</p>	<p>At 12 months, spironolactone-treated patients exhibited a significant reduction in urine albumin/creatinine ratio from baseline compared to furosemide-treated patients (P&lt;0.05).</p> <p>Both treatments exhibited similar reductions in blood pressure from baseline (P value not reported).</p> <p>No adverse events were reported in this trial.</p> <p>Secondary: Not reported</p>
<p>Furumatsu et al.<sup>47</sup> (2008)</p> <p>Spironolactone 25 mg/day (triple blockade group)</p> <p>vs</p> <p>trichlormethiazide * 1 mg/day or furosemide 10 mg/day (control group)</p> <p>Study medications were added to ongoing therapy consisting of enalapril 5 mg/day and losartan 50 mg/day.</p>	<p>MC, OL, PRO, RCT</p> <p>Patients 20 to 70 years of age, with controlled blood pressure &lt;130/80 mm Hg, chronic nephropathy (defined by serum creatinine level &lt;3 mg/dL or calculated creatinine concentration &lt;30 mL/min), daily treatment with enalapril 5 mg and losartan 50 mg for at least 12 weeks, and persistent proteinuria (urinary protein excretion</p>	<p>N=32</p> <p>12 months</p>	<p>Primary: Reduction in proteinuria, urinary type IV collagen, SBP, DBP, mean blood pressure, creatinine, creatinine clearance, potassium, urinary aldosterone</p> <p>Secondary: Not reported</p>	<p>Primary: At one year of therapy, patients randomized to the triple blockage group experienced a statistically significant 58% reduction in urinary protein level from baseline (P&lt;0.05), while there was no difference in the control group. Compared to the control group, the triple blockade group experienced a significant reduction in proteinuria at one year of therapy (P&lt;0.05).</p> <p>At one year of therapy, patients randomized to the triple blockage group experienced a statistically significant 40% reduction in urinary type IV collagen from baseline (P&lt;0.05); while there was no difference in the control group. However there was no statistically significant difference in the change of urinary type IV collagen from baseline between the two study groups.</p> <p>There were no statistically significant differences between the two study groups in the following outcome measures: SBP, DBP, mean blood pressure, creatinine, creatinine clearance, potassium, and urinary aldosterone.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	>0.5 g/day)			
Hansson et al. <sup>48</sup> (2000) NORDIL  Conventional therapy (diuretic, $\beta$ -blocker or both)  vs  diltiazem 180 to 360 mg QD	BE, MC, OL, PRO, RCT  Patients 50 to 74 years of age with DBP $\geq$ 100 mm Hg and previously untreated	N=10,881  4.5 years	Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death  Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI	Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/ $\beta$ -blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).  Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/ $\beta$ -blocker patients (P=0.04).  Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/ $\beta$ -blocker patients (P=0.17).  Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57 and congestive heart failure (P=0.42).
Wysong et al. <sup>49</sup> (2007)  Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)  vs  $\beta$ -blockers (atenolol, metoprolol, oxprenolol*, or propranolol)	MA  13 RCTs evaluating patients $\geq$ 18 years of age with HTN	N=91,561  Duration varied	Primary: All-cause mortality  Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions	Primary: There was not a significant difference observed in all-cause mortality between $\beta$ -blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with $\beta$ -blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).  Secondary: There was a significant decrease in stroke observed with $\beta$ -blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with $\beta$ -blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).  CHD risk was not significantly different between $\beta$ -blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07]), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15)



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with <math>\beta</math>-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of <math>\beta</math>-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with <math>\beta</math>-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>
<b>Miscellaneous</b>				
<p>Bagshaw et al.<sup>50</sup> (2007)</p> <p>Loop diuretics (frusemide†, torasemide*)</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients with acute renal failure</p>	<p>N=555</p> <p>Variable duration</p>	<p>Primary: Mortality, need for renal replacement therapy, and renal recovery</p> <p>Secondary: Urine output, serum potassium level and acid-base status, duration of acute renal failure or renal replacement therapy, length of hospital stay, toxicity</p>	<p>Primary: There was no statistical difference in mortality between loop diuretics compared to placebo (OR, 1.28; 95% CI, 0.89 to 1.84; P=0.18).</p> <p>There was no statistical difference in renal recovery between loop diuretics and control (OR, 0.88; 95% CI, 0.59 to 1.31; P=0.5).</p> <p>Secondary: Loop diuretics were associated with a shorter duration of renal replacement therapy (weighted mean difference of 1.4 days; 95% CI, 0.2 to 2.3; P=0.02), shorter time to spontaneous decline in serum creatinine level (WMD, 2.1 days; 95% CI, 0.4 to 3.7; P=0.01), and a greater increase in urine output from baseline (OR, 2.6; 95% CI, 1.4 to 4.9; P=0.004).</p> <p>There was no data available on acid-base status, hospital status, hospital length of stay or health costs.</p>
<p>Galloe et al.<sup>51</sup> (2006)</p>	<p>DB, PC, RCT, XO</p> <p>Men and women</p>	<p>N=16</p> <p>14 days</p>	<p>Primary: Patient reported quality of life</p>	<p>Primary: Patient's well-being increased 12% with 0.5 mg bumetanide BID but higher doses bumetanide decreased patient's well-being by 12% compared</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Bumetanide 0.5 mg (0, 1, 2, or 4 tablets BID)  vs  trandolapril 0.5 mg (0, 1, 2, or 4 tablets QD)	with previous MI $\geq$ 3 years ago, had medical treatment for heart failure and ejection fraction between 0.36 and 0.54 estimated by echo-cardiography (wall motion index)		Secondary: Effects on the involved organs: kidney function, left ventricular function, blood pressure	<p>to placebo (P&lt;0.002). Increasing doses of bumetanide tended to increase tiredness (P=0.072). There were no statistically significant effects of bumetanide on the patient's opinion of their health, degree of dyspnea, appetite or work capacity.</p> <p>Secondary:            Bumetanide increased 24-hour urine production in a straight dose-dependent manner (P&lt;0.0001) while trandolapril had no effect (P=0.53). Bumetanide and trandolapril did not alter the 24-hour creatinine excretion and creatinine clearance (P=0.33, P=0.11 and P=0.53, P=0.97, respectively).</p> <p>Bumetanide decreased left ventricular function and increased heart rate in a dose dependent manner (P&lt;0.001). Left ventricular function was also decreased with trandolapril but did not reach statistically significant. (P&gt;0.062).</p> <p>Trandolapril significantly reduced SBP by maximally of 7.6 mm Hg (5.8%) with the lowest dose of 0.5 mg/day (P=0.007). Bumetanide had no significant effect on DBP (P=0.23).</p>

\*Synonym for torsemide.

†Synonym for furosemide.

‡Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, IV=intravenous, PO=oral, QD=once daily

Study design abbreviations: BE=blinded endpoint, DB=double blind, DD=double dummy, MA=meta analysis, MC=multicenter, OL=open label, OS=observational, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitors, BNP=brain natriuretic peptide, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HTN=hypertension, HR=hazard ratio, LDL-C=low-density lipoprotein cholesterol, LVEF=left ventricular ejection fraction, MI=myocardial infarction, NYHA=New York Heart Association, PBV=pulmonary blood volume, PEV=pulmonary extravascular fluid volume, RR=relative risk, SBP=systolic blood pressure, TC=total cholesterol, TG=triglycerides

### Additional Evidence

#### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

## IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Loop Diuretics**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Bumetanide	injection, tablet	N/A	N/A	\$\$
Ethacrynic acid	tablet	Edecrin®*	\$\$\$\$\$	\$\$\$\$\$
Furosemide	injection, solution, tablet	Lasix®*	\$\$\$	\$
Torsemide	tablet	N/A	N/A	\$

\*Generic is available in at least one dosage form or strength.

N/A=Not available

## X. Conclusions

All of the loop diuretics are approved for the treatment of edema associated with congestive heart failure, hepatic disease, or renal disease. Furosemide and torsemide are also approved for the treatment of hypertension. Additionally, ethacrynic acid is approved for the short-term treatment of ascites (due to malignancy, idiopathic edema, and lymphedema) and for the short-term treatment of hospitalized pediatric patients with congenital heart disease or the nephrotic syndrome.<sup>3-6</sup> All agents are available in a generic formulation.

Guidelines recommend the use of diuretics and sodium restriction for the management of ascites due to cirrhosis. Spironolactone is recommended as first-line therapy, either as monotherapy or in combination with furosemide.

Amiloride or eplerenone are alternative treatment options in patients experiencing gynecomastia with

**spironolactone**.<sup>20</sup> Several studies have compared furosemide and torsemide in cirrhotic patients with ascites. Although torsemide significantly increased natriuresis and diuresis compared to furosemide, these effects were not consistently demonstrated across the studies. There was no difference in plasma renin or aldosterone concentrations among the treatment groups.<sup>21-24</sup>

For the treatment of chronic heart failure, guidelines recommend the use of diuretics in all patients who have evidence of volume overload. Loop diuretics are generally recommended as initial therapy in patients with left ventricular dysfunction. For those with persistent fluid retention despite treatment with a loop diuretic, a thiazide diuretic or metolazone may be added to the regimen. In patients with normal left ventricular function, either a thiazide diuretic or loop diuretic may be used as initial therapy to manage fluid overload.<sup>9-10</sup> There are relatively few studies that have directly compared the loop diuretics for the treatment of chronic heart failure. In open-label trials, torsemide decreased mortality, hospitalizations and improved NYHA functional class compared to treatment with furosemide. However, due to limitations in the study designs, it is difficult to draw firm conclusions about the results of these studies.<sup>31-33</sup>

There are several published guidelines on the treatment of hypertension. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>11-18</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>11</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>11-18</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>11-18</sup>

In clinical trials, the thiazide diuretics have been shown to effectively lower blood pressure.<sup>40-49</sup> Some studies suggest that hydrochlorothiazide is more effective than a loop diuretic for lowering blood pressure.<sup>8</sup> However, a loop diuretic should be used when the glomerular filtration rate is  $<30$  mL/min.<sup>1,2</sup>

Serious adverse events reported with the loop diuretics include electrolyte abnormalities, hypersensitivity reactions, and ototoxicity. Ethacrynic acid has a higher rate of ototoxicity than other loop diuretics and is less commonly used. Patients allergic to sulfonamides may also show hypersensitivity to bumetanide, furosemide, and torsemide. Ethacrynic acid is the only loop diuretic that is not a sulfonamide derivative and can be safely used in patients with a sulfonamide allergy.<sup>3-7</sup>

There is insufficient evidence to support that one brand loop diuretic is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand loop diuretics within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand loop diuretic is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. *Pharmacotherapy: a pathophysiologic approach*. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
2. Agarwal R. Thiazides versus loop diuretics in the treatment of hypertension. In: *UpToDate*, Bakris GL (Ed), UpToDate, Waltham, MA, 2021.
3. Edecrin® [package insert]. Bridgewater (NJ): Bausch Health US, LLC; 2020 Aug.
4. Lasix® [package insert]. Parsippany (NJ): Validus Pharmaceuticals LLC; 2018 Aug.
5. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
6. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
7. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
8. Araoye MA, Chang MY, Khatri IM, et al. Furosemide compared with hydrochlorothiazide. Long-term treatment of hypertension. *JAMA* 1978;240(17):1863.
9. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
10. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
11. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
12. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
13. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
14. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
15. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
16. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
17. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
18. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
19. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl.1).
20. Biggins SW, Angeli P, Garcia-Tsao G, Ginès P, Ling SC, Nadim MK, Wong F, Kim WR. Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021

- Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology*. 2021 Aug;74(2):1014-1048. doi: 10.1002/hep.31884.
21. Laffi G, Marra F, Buzzelli G, et al. Comparison of the effects of torasemide and furosemide in nonazotemic cirrhotic patients with ascites: A randomized, double-blind study. *Hepatology* 1991;13:1101-5.
  22. Gerbes AL, Bertheau-Reitha U, Falkner C et al. Advantages of the new loop diuretic torasemide over furosemide in patients with cirrhosis and ascites. A randomized, double blind cross-over trial. *J Hepatol* 1993;17:353-8.
  23. Fiaccadori F, Pedretti G, Pasetti G et al. Torsemide versus furosemide in cirrhosis: a long-term double-blind, randomized clinical study. *Clin Investig* 1993;71:579-84.
  24. Abecasis R, Guevera M, Miguez C, et al. Long-term efficacy of torsemide compared to frusemide in cirrhotic patients with ascites. *Scand J Gastroenterol* 2001;3:309-13.
  25. Galløe AM, Skagen K, Christensen NJ, Nielsen SL, Frandsen EK, Bie P, et al. Dosage dependent hormonal counter regulation to combination therapy in patients with left ventricular dysfunction. *J Clin Pharm Ther*. 2006 Apr;31(2):139-47.
  26. Hutcheon D, Vincent ME, Sandhu RS. Clinical use of diuretics in congestive heart failure. *J Clin Pharmacol*. 1981;21(11-12 Pt 2):668-72.
  27. Konecke LL. Clinical trial of bumetanide versus furosemide in patients with congestive heart failure. *J Clin Pharmacol*. 1981;21(11-12 Pt 2):688-90.
  28. Nicholson G. Treatment of fluid retention in cirrhosis: a comparison of bumetanide and furosemide. *Curr Med Res Opin*. 1977;4(9):675-9.
  29. Eshaghian S, Horwich TB, Fonarow GC. Relation of loop diuretic dose to mortality in advanced heart failure. *Am J Cardiol*. 2006 Jun 15;97(12):1759-64. Epub 2006 Apr 27.
  30. Mentz RJ, Hasselblad V, DeVore AD, Metra M, Voors AA, Armstrong PW, et al. Torsemide Versus Furosemide in Patients With Acute Heart Failure (from the ASCEND-HF Trial). *Am J Cardiol*. 2016 Feb 1;117(3):404-11.
  31. Murray MD, Deer MM, Ferguson JA, et al. Open-label randomized trial of torsemide compared with furosemide therapy for patients with heart failure. *Am J Med* 2001;111:513-20.
  32. Cosín J, Díez J; TORIC investigators. Torasemide in chronic heart failure: results of the TORIC study. *Eur J Heart Fail* 2002;4:507-13.
  33. Muller K, Gamba G, Jaquet F, et al. Torsemide vs furosemide in primary care patients with chronic heart failure NYHA II to IV—efficacy and quality of life. *Eur J Heart Fail*. 2003;5(6):793-801.
  34. Kasama S, Toyama T, Hatori T, Sumino H, Kumakura H, Takayama Y, et al. Effects of torasemide on cardiac sympathetic nerve activity and left ventricular remodeling in patients with congestive heart failure. *Heart*. 2006 Oct;92(10):1434-40. Epub 2006 Apr 18.
  35. Levy B. The efficacy of safety of furosemide and a combination of spironolactone and hydrochlorothiazide in congestive heart failure. *J Clin Pharmacol*. 1977;17(7):420-30.
  36. Austin SM, Schreiner BF, Kramer DH, et al. The acute hemodynamic effects of ethacrynic acid and furosemide in patients with chronic postcapillary pulmonary hypertension. *Circulation* 1976;53(2):364-69.
  37. Patterson JH, Adams KF, Applefeld MM, et al. Oral torsemide in patients with chronic congestive heart failure: effects on body weight, edema, and electrolyte excretion. Torsemide Investigators Group. *Pharmacotherapy* 1994;14(5):514-21.
  38. Senzaki H, Kamiyama M Phard, Masutani S, Ishido H, Taketazu M, Kobayashi T, et al. Efficacy and safety of Torasemide in children with heart failure. *Arch Dis Child*. 2008 Mar 12.
  39. Faris R, Flather MD, Purcell H, et al. Diuretics for heart failure. *Cochrane Database of Systematic Reviews* 2006, Issue 1. Art. No.: CD003838. DOI: 10.1002/14651858.CD003838.pub2.
  40. Van der Heijden M, Donders SH, Cleophas TJ, et al. A randomized, placebo-controlled study of loop diuretics in patients with essential hypertension: the bumetanide and furosemide on lipid profile (BUFUL) clinical study report. *J Clin Pharmacol*. 1998;38(7):630-5.
  41. De Berrazuela JR, González JP, de Mier I, Poveda JJ, García-Unzueta MT. Vasodilatory action of loop diuretics: a plethysmography study of endothelial function in forearm arteries and dorsal hand veins in hypertensive patients and controls. *J Cardiovasc Pharmacol*. 2007 Feb;49(2):90-5.
  42. von Dossow V, Spies C, Schenk H, Schlesinger S, von Heymann C. Secondary pulmonary hypertension: haemodynamic effects of torasemide versus furosemide. *Clin Drug Investig*. 2008;28(1):17-26.
  43. Vasavada N, Saha C, Agarwal R. A double-blind randomized crossover trial of two loop diuretics in chronic kidney disease. *Kidney Int*. 2003;64(2):632-40.
  44. Pupita F, Belogi M, Ansuini R, et al. Long-acting and short-acting diuretics in the treatment of hypertension. *Pharmatherapeutica* 1983;3(7):475-81.

45. Valmin K, Hansen T. Treatment of benign essential hypertension: comparison of furosemide and hydrochlorothiazide. *Eur J Clin Pharmacol.* 1975;8(6):393-401.
46. Ogawa S, Takeuchi K, Mori T, Nako K, Ito S. Spironolactone further reduces urinary albumin excretion and plasma B-type natriuretic peptide levels in hypertensive type II diabetes treated with angiotensin-converting enzyme inhibitor. *Clin Exp Pharmacol Physiol.* 2006 May-Jun;33(5-6):477-9.
47. Furumatsu Y, Nagasawa Y, Tomida K, Mikami S, Kaneko T, Okada N, Tsubakihara Y, Imai E, Shoji T. Effect of renin-angiotensin-aldosterone system triple blockade on non-diabetic renal disease: addition of an aldosterone blocker, spironolactone, to combination treatment with an angiotensin-converting enzyme inhibitor and angiotensin II receptor blocker. *Hypertens Res.* 2008 Jan;31(1):59-67.
48. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, et al. Randomized trial of effects of calcium antagonists compared with diuretics and  $\beta$ -blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet.* 2000 Jul 29;356(9227):359-65.
49. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev.* 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.
50. Bagshaw SM, Delaney A, Haase M, Ghali WA, Bellomo R. Loop diuretics in the management of acute renal failure: a systematic review and meta-analysis. *Crit Care Resusc.* 2007 Mar;9(1):60-8.
51. Galløe AM, Skagen K, Christensen NJ, Nielsen SL, Frandsen EK, Bie P, et al. Dosage dependent hormonal counter regulation to combination therapy in patients with left ventricular dysfunction. *J Clin Pharm Ther.* 2006 Apr;31(2):139-47.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Potassium-Sparing Diuretics  
AHFS Class 402816  
May 4, 2022**

## 1. Overview

The potassium-sparing diuretics are approved for the treatment of congestive heart failure, edema, and hypertension.<sup>1-5</sup> They inhibit sodium-potassium ion exchange at the distal convoluted tubule, cortical collecting tubule, and collecting duct. This reduces both potassium and hydrogen secretion and their subsequent excretion.<sup>1-9</sup> When used alone, potassium-sparing diuretics have a weak diuretic and antihypertensive effect and increased risk of hyperkalemia.<sup>5</sup> The potassium-sparing diuretics are generally used in combination with other diuretics to help restore normal serum potassium levels or to prevent the development of hypokalemia.<sup>1-5</sup> Amiloride and triamterene are both available as a fixed-dose combination with hydrochlorothiazide. Hydrochlorothiazide inhibits the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride.<sup>5-8</sup>

The potassium-sparing diuretics that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the products are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Potassium-Sparing Diuretics Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Single Entity Agents</b>			
Amiloride	tablet	N/A	amiloride
Triamterene	capsule	N/A	triamterene
<b>Combination Products</b>			
Amiloride and hydrochlorothiazide	tablet	N/A	amiloride and hydrochlorothiazide
Triamterene and hydrochlorothiazide	capsule, tablet	Maxzide®*	triamterene and hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.

PDL=Preferred Drug List

N/A=Not available

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the potassium-sparing diuretics are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Potassium-Sparing Diuretics**

Clinical Guideline	Recommendation(s)
American Heart Association/American College of Cardiology/Heart Failure Society of America: <b>2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022)</b> <sup>9</sup>	<b>Treatment of Stage A heart failure (HF)</b> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul>



Clinical Guideline	Recommendation(s)
	<p><b>Treatment of Stage B heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>• In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> <li>• In patients with a recent MI and LVEF≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><b>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</b></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three β-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is &gt;30 mL/min/1.73 m<sup>2</sup> and serum potassium</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>is &lt;5.0 mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at &lt;5.5 mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF ≤35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of ≥70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>10</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</p> <ul style="list-style-type: none"> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</li> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults</b> (2014)<sup>11</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients <math>\geq 18</math> years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>12</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio <math>&lt; 0.5</math> is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week. Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality</li> </ul> </li> </ul>



Clinical Guideline	Recommendation(s)
	<p>prevention.</p> <ul style="list-style-type: none"> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> <ul style="list-style-type: none"> <li>● General scheme for drug treatment           <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</li> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> </ul> </li> <li>● Core drug-treatment strategy           <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: 2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)<sup>13</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients &lt;60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> </ul> </li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> </ul>



Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><u>Treatment of hypertension in association with heart failure</u></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction <math>&lt;40\%</math>), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (<math>&lt;40\%</math>) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium <math>&lt;5.2</math> mmol/L, an eGFR <math>\leq 30</math> mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><u>Treatment of hypertension in association with stroke</u></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> <ul style="list-style-type: none"> <li>● BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><u>Treatment of hypertension in association with LVH</u></p> <ul style="list-style-type: none"> <li>● Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>● The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul> <p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>● Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>● For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>● In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>● The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>● Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>● Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>● Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>● Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>● Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>● For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>● For persons with diabetes and hypertension not included in other guidelines in</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</p> <ul style="list-style-type: none"> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95<sup>th</sup> percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral b-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</p> <ul style="list-style-type: none"> <li>• Women with severe hypertension with SBP <math>\geq</math>160 mmHg or DBP <math>\geq</math>110 mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>14</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</li> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is <math>&lt;</math>150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>hypertension), because of the high CV risk of these patients.</p> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (<math>\geq 65</math> years) are treated with the same recommendations in non-older patient population. In very old patients (<math>\geq 80</math> years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of <math>&lt; 80</math> mmHg if tolerated. Treated values of <math>&lt; 130</math> mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• A SBP goal &lt;130 mmHg is recommended in patients with diabetes and &lt;130 mmHg if tolerated, but not &lt;120 mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq 220</math> mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq 140/90</math> mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis and management (2019)</b><sup>15</sup></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq 160</math> mmHg) the same treatment as people with both raised systolic and diastolic blood pressure.</li> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients <math>&lt; 55</math> years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged <math>&lt; 55</math> years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are <math>&gt; 55</math> years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>blood pressure, continue with their treatment.</p> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on “Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence”.</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African–Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person’s medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul> <p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>16</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however,</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</p> <ul style="list-style-type: none"> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease (2021)</b><sup>17</sup></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD) not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>• The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>• The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>• The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>• The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>• The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>• The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>• Treat adult kidney transplant recipients with high BP to a target BP of &lt;130</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</p> <ul style="list-style-type: none"> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><b>BP management in children with CKD</b></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)</b><sup>18</sup></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) <math>\geq 130</math> mmHg or an average diastolic blood pressure (DBP) of <math>\geq 80</math> mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of <math>\geq 10\%</math> and an average SBP of <math>\geq 130</math> mmHg or an average <math>\geq 80</math> mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk <math>&lt; 10\%</math> and an SBP of <math>\geq 140</math> mmHg or a DBP of <math>\geq 90</math> mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of <math>\geq 10\%</math>, a BP target <math>&lt; 130/80</math> mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target <math>&lt; 130/80</math> mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</li> <li>Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP <math>&gt; 20/10</math> mmHg above their BP target.</li> <li>Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal <math>&lt; 130/80</math> mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>In adults with SIHD and hypertension, a BP target <math>&lt; 130/80</math> is recommended.</li> <li>Adults with SIHD and hypertension (BP <math>\geq 130/80</math> mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (<math>\geq 300</math> mg/d, or <math>\geq 300</math> mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</li> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq 220/120</math> mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq 140/90</math> mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal <math>&lt;130/80</math> mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal <math>&lt;130</math> mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP <math>&lt;140</math> mmHg and a DBP <math>&lt;90</math> mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq 130/80</math> mmHg with a treatment goal <math>&lt;130/80</math> mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target <math>&lt;130/80</math> mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP ≥180 mmHg or DBP ≥110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>19</sup></p>	<p><u>Hypertension/blood pressure control</u></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (≥140/90) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure ≥180/110 and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk ≥15%), a blood pressure target of &lt;130/80 mmHg may be appropriate, if it</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>can be safely attained.</p> <ul style="list-style-type: none"> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk &lt;15%), treat to blood pressure target of &lt;140/90.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure &gt;120/80, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>140/90 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq</math>160/100 mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq</math>300 mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul> <p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq</math>300 mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq</math>25 mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq</math>300 mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq</math>25</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>mL/min/1.73 m<sup>2</sup> or ≥300 mg/g, respectively.</p> <ul style="list-style-type: none"> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium–glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (≤30%) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin–to–creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin–to–creatinine ratio ≥300 mg/g creatinine and/or estimated glomerular filtration rate &lt;60 mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin–to–creatinine ratio (&lt;30 mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate &lt;30 mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>

\*Agent is not available in the United States

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the potassium-sparing diuretics are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Potassium-Sparing Diuretics<sup>1-4</sup>**

Indication(s)	Single Entity Agents		Combination Products	
	Amiloride	Triamterene	Amiloride and HCTZ	Triamterene and HCTZ
<b>Congestive Heart Failure (or Edema) and Hypertension</b>				
Help restore normal serum potassium levels in patients who develop hypokalemia on the kaliuretic diuretic	✓ *			
Prevent development of hypokalemia in patients who would be exposed to particular risk if hypokalemia were to develop	✓ *			
Treatment of edema associated with congestive heart failure, cirrhosis of the liver and the nephrotic syndrome; steroid-induced edema,		✓ †		



Indication(s)	Single Entity Agents		Combination Products	
	Amiloride	Triamterene	Amiloride and HCTZ	Triamterene and HCTZ
idiopathic edema and edema due to secondary hyperaldosteronism				
Use in patients who develop hypokalemia when thiazide or other kaliuretic diuretics are used alone, or in whom maintenance of normal serum potassium levels is considered to be clinically important			✓ ‡	
Use in patients who develop hypokalemia on hydrochlorothiazide alone, or in whom require a thiazide diuretic and in whom the development of hypokalemia cannot be risked				✓ ‡

\*As adjunctive treatment with thiazide diuretics or other kaliuretic-diuretic agents.

†May be used alone or with other diuretics, either for its added diuretic effect or its potassium-sparing potential.

‡The fixed combination drug is not indicated for the initial therapy of edema or hypertension except in individuals in whom the development of hypokalemia cannot be risked.

HCTZ=hydrochlorothiazide

#### IV. Pharmacokinetics

The pharmacokinetic parameters of the potassium-sparing diuretics are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Potassium-Sparing Diuretics<sup>6</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
<b>Single Entity Agents</b>					
Amiloride	30 to 90	Not significant (% not reported)	Not metabolized	Feces (40 to 50) Renal (50)	6 to 9
Triamterene	30 to 70	55 to 67	Liver (80)	Renal (21)	1.5 to 2.5
<b>Combination Products</b>					
Amiloride and HCTZ	30 to 90/60 to 80	Not significant (% not reported)/ 10	Not metabolized	Feces (40 to 50) Renal (50)/ Renal (>60)	6 to 9/ 10 to 12
Triamterene and HCTZ	30 to 70/60 to 80	55 to 67/40	Liver (80)/ not reported	Renal (21)/ Renal (>60)	1.5 to 2.5/ 10 to 12

HCTZ=hydrochlorothiazide

#### V. Drug Interactions

Major drug interactions with the potassium-sparing diuretics are listed in Table 5.

**Table 5. Major Drug Interactions with the Potassium-Sparing Diuretics<sup>6</sup>**

Generic Name(s)	Interaction	Mechanism
Potassium-sparing diuretics (amiloride, triamterene)	Aldosterone blockers	Aldosterone blockers and potassium-sparing diuretics may exert additive pharmacologic effects. Hyperkalemia with the potential for cardiac arrhythmias may result.
Potassium-sparing diuretics (amiloride, triamterene)	Potassium preparations	Use of potassium preparations and potassium-sparing diuretics may increase the risk of hyperkalemia. Cardiac arrhythmias or cardiac arrest may occur.
Potassium-sparing diuretics (amiloride, triamterene)	ACE inhibitors	Hyperkalemia, possibly with cardiac arrhythmias or arrest may occur with the combination of amiloride and ACE inhibitors. Decreased aldosterone activity by ACE inhibitors may function synergistically with potassium



Generic Name(s)	Interaction	Mechanism
		conservation by amiloride to produce substantial hyperkalemia.
Potassium-sparing diuretics (amiloride, triamterene)	Indomethacin and derivatives	The combination of indomethacin and derivatives and triamterene may cause a sudden onset of nephrotoxicity.
Potassium-sparing diuretics (amiloride)	ARBs	The risk of hyperkalemia may be increased when amiloride is co-administered with ARBs. Decreased aldosterone activity by ARBs may function synergistically with potassium conservation by amiloride to produce substantial hyperkalemia.
Thiazide diuretics (HCTZ)	Dofetilide	Thiazide diuretics increase potassium excretion. Hypokalemia may occur, increasing the risk of torsades de pointes.
Thiazide diuretics (HCTZ)	Lithium	Thiazide diuretics decrease the renal clearance of lithium which leads to increased serum lithium levels. Lithium toxicity has occurred.
Potassium-sparing diuretics (amiloride)	Aliskiren	Hyperkalemia risk may be increased when aliskiren is coadministered with potassium-sparing diuretics. Decreased aldosterone activity by aliskiren may function synergistically with potassium conservation leading to the development of hyperkalemia.
Potassium-sparing diuretics (amiloride)	Macrolide immunosuppressants	Macrolide immunosuppressives and potassium-sparing diuretics may exert additive effects on potassium leading to hyperkalemia.
Thiazide diuretics (HCTZ)	Diazoxide	Hyperglycemia may occur with symptoms similar to diabetes. The mechanism is unknown.
Thiazide diuretics (HCTZ)	Digitalis glycosides	Diuretic-induced electrolyte disturbances may predispose the patient to digitalis-induced cardiac arrhythmias.

ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor blocker, HCTZ=hydrochlorothiazide

## VI. Adverse Drug Events

The most common adverse drug events reported with the potassium-sparing diuretics are listed in Table 6. The boxed warnings are listed in Tables 7 through 9.

**Table 6. Adverse Drug Events (%) Reported with the Potassium-Sparing Diuretics<sup>1-6</sup>**

Adverse Events	Single Entity Agents		Combination Products	
	Amiloride	Triamterene	Amiloride and HCTZ	Triamterene and HCTZ
<b>Cardiovascular</b>				
Arrhythmia	≤1	-	≤1	✓
Bradycardia	-	-	-	1 to 10
Chest pain	≤1	-	≤1	-
Congestive heart failure	-	-	-	1 to 10
Edema	-	-	-	1 to 10
Hypotension	-	-	1 to 10	1 to 10
Orthostatic hypotension	≤1	-	1 to 10	1 to 10
Palpitations	≤1	-	≤1	-
<b>Central Nervous System</b>				
Dizziness	1 to 10	✓	1 to 10	1 to 10
Fatigue	1 to 10	✓	1 to 10	1 to 10
Headache	1 to 10	✓	1 to 10	1 to 10
<b>Dermatological</b>				
Alopecia	≤1	-	≤1	≤1

Adverse Events	Single Entity Agents		Combination Products	
	Amiloride	Triamterene	Amiloride and HCTZ	Triamterene and HCTZ
Erythema multiforme	-	-	≤1	≤1
Exfoliative dermatitis	-	-	≤1	≤1
Photosensitivity	-	✓	1 to 10	1 to 10
Rash	-	✓	-	1 to 10
Stevens-Johnson syndrome	-	-	<1	<1
Toxic epidermal necrolysis	-	-	<1	<1
<b>Endocrine and Metabolic</b>				
Dehydration	1 to 10	-	1 to 10	<1
Gynecomastia	1 to 10	-	1 to 10	<1
Metabolic acidosis	1 to 10	-	1 to 10	<1
Postmenopausal bleeding	-	-	-	<1
<b>Gastrointestinal</b>				
Abdominal pain	1 to 10	-	1 to 10	✓
Anorexia	-	-	1 to 10	1 to 10
Appetite changes	1 to 10	-	1 to 10	-
Constipation	1 to 10	-	1 to 10	1 to 10
Diarrhea	1 to 10	✓	1 to 10	✓
Dry Mouth	-	✓	-	-
Epigastric distress	-	-	1 to 10	1 to 10
Flatulence	≤1	-	≤1	-
Gastrointestinal bleeding	≤1	-	≤1	-
Nausea	1 to 10	✓	1 to 10	1 to 10
Pancreatitis	-	-	<1	<1
Vomiting	1 to 10	✓	1 to 10	✓
<b>Genitourinary</b>				
Bladder spasms	≤1	-	≤1	-
Dysuria	≤1	-	≤1	-
Impotence	1 to 10	-	1 to 10	<1
Polyuria	≤1	-	≤1	-
Renal dysfunction	-	✓	≤1	≤1
<b>Hematological</b>				
Agranulocytosis	-	-	≤1	≤1
Aplastic anemia	-	-	≤1	≤1
Hemolytic anemia	-	-	<1	<1
Leukopenia	-	-	≤1	≤1
Megaloblastic anemia	-	✓	-	-
Thrombocytopenia	-	✓	≤1	≤1
<b>Laboratory Test Abnormalities</b>				
Hypercalcemia	-	-	<1	<1
Hyperkalemia	<10	✓	-	-
Hypokalemia	-	✓	1 to 10	1 to 10
Hyponatremia	1 to 10	-	1 to 10	1 to 10
<b>Musculoskeletal</b>				
Muscle cramps	1 to 10	-	1 to 10	✓
Weakness	1 to 10	✓	1 to 10	✓
<b>Renal</b>				
Azotemia	-	✓	-	-
Interstitial nephritis	-	-	<1	<1
Renal failure	-	✓	<1	<1
<b>Respiratory</b>				
Cough	1 to 10	-	1 to 10	-
Dyspnea	1 to 10	-	1 to 10	1 to 10

Adverse Events	Single Entity Agents		Combination Products	
	Amiloride	Triamterene	Amiloride and HCTZ	Triamterene and HCTZ
Eosinophilic pneumonitis	-	-	<1	<1
Respiratory distress	-	-	<1	<1
<b>Other</b>				
Allergic myocarditis	-	-	<1	<1
Allergic reactions	-	✓	<1	<1
Hepatic function impairment	-	✓	<1	<1
Increased intraocular pressure	≤1	-	≤1	-
Jaundice	≤1	✓	≤1	-
Tinnitus	≤1	-	≤1	-
Visual disturbance	-	-	≤1	≤1

✓ Percent not specified

- Event not reported

**Table 7. Boxed Warning for Amiloride<sup>5</sup>**

<b>WARNING</b>
Like other potassium-conserving agents, amiloride may cause hyperkalemia (serum potassium levels greater than 5.5 mEq per liter) which, if uncorrected, is potentially fatal. Hyperkalemia occurs commonly (about 10%) when amiloride is used without a kaliuretic diuretic. This incidence is greater in patients with renal impairment, diabetes mellitus (with or without recognized renal insufficiency), and in the elderly. When amiloride is used concomitantly with a thiazide diuretic in patients without these complications, the risk of hyperkalemia is reduced to about 1 to 2%. It is thus essential to monitor serum potassium levels carefully in any patient receiving amiloride, particularly when it is first introduced, at the time of diuretic dosage adjustments, and during any illness that could affect renal function.

**Table 8. Boxed Warning for Triamterene-Hydrochlorothiazide<sup>5</sup>**

<b>WARNING</b>
Abnormal elevation of serum potassium levels (at least 5.5 mEq/L) can occur with all potassium-sparing agents, including triamterene. Hyperkalemia is more likely to occur in patients with renal impairment and diabetes (even without evidence of renal impairment), and in elderly or severely ill patients. Because uncorrected hyperkalemia may be fatal, serum potassium levels must be monitored at frequent intervals especially in patients receiving triamterene, when dosages are changed, or with any illness that may influence renal function.

**Table 9. Boxed Warning for Amiloride and Hydrochlorothiazide<sup>5</sup>**

<b>WARNING</b>
Like other potassium-conserving diuretic combinations, amiloride and hydrochlorothiazide may cause hyperkalemia (serum potassium levels greater than 5.5 mEq per liter). In patients without renal impairment or diabetes mellitus, the risk of hyperkalemia with this combination product is about 1 to 2 percent. This risk is higher in patients with renal impairment or diabetes mellitus (even without recognized diabetic nephropathy). Since hyperkalemia, if uncorrected, is potentially fatal, it is essential to monitor serum potassium levels carefully in any patient receiving amiloride hydrochloride and hydrochlorothiazide, particularly when it is first introduced, at the time of dosage adjustments, and during any illness that could affect renal function.

## VII. Dosing and Administration

The usual dosing regimens for the potassium-sparing diuretics are listed in Table 10.

**Table 10. Usual Dosing Regimens for the Potassium-Sparing Diuretics<sup>1-6</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
-----------------	------------------	----------------------	--------------

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Single Entity Agents</b>			
Amiloride	<u>Congestive heart failure (or edema) and hypertension:</u> Tablet: initial, 5 mg daily; may increase to 10 mg daily if needed; maximum, 20 mg	Safety and efficacy in children have not been established.	Tablet: 5 mg
Triamterene	<u>Treatment of edema:</u> Capsule: initial, 100 mg twice daily; maximum, 300 mg	Safety and efficacy in children have not been established.	Capsule: 50 mg 100 mg
<b>Combination Products</b>			
Amiloride and HCTZ	<u>Congestive heart failure (or edema) and hypertension:</u> Tablet: initial, 5-50 mg once daily; maintenance, 5-50 to 10-100 mg once daily or in divided doses	Safety and efficacy in children have not been established.	Tablet: 5-50 mg
Triamterene and HCTZ	<u>Congestive heart failure (or edema) and hypertension:</u> Capsule, tablet: initial, 37.5-25 mg once daily; maintenance: 37.5-25 to 75-50 mg once daily	Safety and efficacy in children have not been established.	Capsule: 37.5-25 mg  Tablet: 37.5-25 mg 75-50 mg

HCTZ=hydrochlorothiazide

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the potassium-sparing diuretics are summarized in Table 11.

**Table 11. Comparative Clinical Trials with the Potassium-Sparing Diuretics**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Edema/Heart Failure</b>				
Bayliss et al. <sup>20</sup> (1987)  Amiloride 5 mg QD and furosemide 40 mg	OS  Patients with heart failure, 22 to 75 years of age, referred with breathlessness on moderate exertion (NYHA class 2 to 3) who were not previously treated	N=12  1 month	Primary: Average weight, heart rate at rest and maximal exercise, maximal treadmill exercise time, plasma renin, plasma aldosterone, noradrenaline at rest and maximal exercise  Secondary: Not reported	Primary: Average weight was significantly reduced during treatment from 72.4 to 68.5 kg (P=0.0003).  Resting heart rate decreased from 89 to 75 bpm (P=0.03). There was no significant change during exercise.  Maximal treadmill exercise time significantly increased from 9.1 to 17.6 minutes (P=0.007).  Plasma concentrations of renin increased from 1.1 to 4.2 ng/mL/hr at rest and from 2.5 to 11.3 ng/mL/hr upon exercise (P<0.007).  Plasma concentrations of aldosterone increased from 169 to 488 pmol/L at rest and from 223 to 737 pmol/L upon exercise (P<0.007).  Plasma concentrations of noradrenaline were significantly reduced (decreased to within normal ranges) at rest following treatment (P=0.005) but remained abnormally high at maximal exercise following treatment.  Secondary: Not reported
Rengo et al. <sup>21</sup> (1979)  Amiloride 15 mg QD  vs  amiloride and	RCT  Patients 35 to 60 years of age with liver cirrhosis and ascites or CHF	N=30  15 days	Primary: Body weight, 24 hour diuresis, serum sodium, serum potassium, sodium and potassium urinary loss	Primary: All treatment groups had a significant reduction in body weight from baseline (P<0.001 for all). Amiloride and HCTZ-treated patients achieved a significantly greater reduction compared to amiloride-treated patients (P<0.001).  All treatment groups significantly differed from baseline in 24 hour diuresis (P<0.01). Amiloride and HCTZ- and HCTZ-treated patients achieved greater diuresis compared to amiloride-treated patients (P<0.001 for both).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 15-150 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>HCTZ 150 mg QD</p>			<p>Secondary: Not reported</p>	<p>Serum sodium was reduced from baseline in all treatment groups. HCTZ-treated patients had a significantly greater reduction than amiloride- (P&lt;0.01) and amiloride and HCTZ-treated patients (P&lt;0.001). Sodium urinary loss was seen with all treatments at day two, amiloride and HCTZ therapy had maintained the loss at day five (P&lt;0.001 for both).</p> <p>Serum potassium decreased in HCTZ-treated patients but increased in amiloride- and amiloride and HCTZ-treated patients. HCTZ-treated patients had a marked increase in potassium urinary loss (P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Cheitlin et al.<sup>22</sup> (1991)</p> <p>Amiloride 5 or 10 mg QD for 7 days, followed by placebo plus HCTZ 50 or 100 mg QD for 14 days</p> <p>vs</p> <p>placebo for 14 days, followed by amiloride 5 or 10 mg plus HCTZ 50 or 100 mg QD for the next 7 days</p>	<p>DB, PC, RCT, XO</p> <p>Patients with a history CHF and ≥1 episode of pulmonary edema (NYHA class 2 to 3) who were not previously treated</p>	<p>N=11</p> <p>21 days</p>	<p>Primary: Hemodynamic changes at rest and exercise</p> <p>Secondary: Not reported</p>	<p>Primary: At rest, there were no significant differences between placebo- and amiloride-treated patients in right atrial pressure, pulmonary atrial pressure, heart rate, pulmonary artery wedge pressure, systemic arterial pressure, right ventricular stroke work index, left ventricular stroke work index, systemic vascular resistance, cardiac index or stroke volume index (P values not reported).</p> <p>During exercise, there were significant differences between placebo- and amiloride-treated patients at the 50-watt stage in right atrial pressure (15.0 vs 10.5 mm Hg), pulmonary artery wedge pressure (28.6 vs 22.1 mm Hg), pulmonary artery diastolic pressure (32.2 vs 21.6 mm Hg), mean pulmonary artery pressure (44.4 vs 38.9 mm Hg), left ventricular stroke work index (69.5 vs 77.9 g-m/m<sup>2</sup>) and stroke volume index (44.9 vs 46.2 cc/beat/m<sup>2</sup>), respectively (P values not reported).</p> <p>There were no significant differences between placebo and amiloride therapy during exercise in right ventricular stroke work index, heart rate, aortic pressure, cardiac index and total systemic vascular resistance (P values not reported).</p> <p>Secondary: Not reported</p>
<p>Ghosh et al.<sup>23</sup> (1987)</p>	<p>PG, RCT, SB</p>	<p>N=60</p>	<p>Primary: Body weight,</p>	<p>Primary: Body weight was reduced with both treatments (P values not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Amiloride and HCTZ 2.5-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>triamterene and HCTZ 50-25 mg QD (fixed-dose combination product)</p>	<p>Elderly patients with stable, mild to moderate CHF</p>	<p>8 weeks</p>	<p>clinical score, biochemistry</p> <p>Secondary: Not reported</p>	<p>Both treatments resulted in improvements in clinical scores; 95 and 88% of the amiloride/HCTZ- and triamterene/HCTZ-treated patients showed an improvement in heart failure signs with no patient's symptoms becoming worse (P values were not reported).</p> <p>Eighty five and 84% of amiloride/HCTZ- and triamterene/HCTZ-treated patients showed an improvement in heart failure symptoms (P values were not reported).</p> <p>There were no significant differences in serum sodium, potassium or urea between the two treatments (P values were not reported).</p> <p>Secondary: Not reported</p>
<p>Kohvakka.<sup>24</sup> (1998)</p> <p>HCTZ 50 mg BID</p> <p>vs</p> <p>amiloride 5 mg BID plus HCTZ 50 mg BID</p> <p>vs</p> <p>triamterene 75 mg BID plus HCTZ 50 mg BID</p> <p>vs</p> <p>KCl 1,000 mg BID plus HCTZ 50 mg BID</p>	<p>RCT, XO</p> <p>Patients 41 to 69 years of age with CHF (NYHA class 2 to 3) who developed persistent hypokalemia on HCTZ alone</p>	<p>N=25</p> <p>5 months</p>	<p>Primary: Changes in weight, blood pressure, serum sodium, serum potassium and total body potassium</p> <p>Secondary: Percentage with hypokalemia, median days until hypokalemia detection, serum magnesium</p>	<p>Primary: Weight loss was significant in amiloride plus HCTZ- and triamterene plus HCTZ-treated patients (P=0.05 for both), but not in KCl plus HCTZ-treated patients (P value not reported), compared to HCTZ-treated patients.</p> <p>No significant changes in blood pressure were observed (P values not reported).</p> <p>No differences in serum sodium were observed in amiloride plus HCTZ- or triamterene plus HCTZ-treated patients (P values not reported). Serum sodium levels were slightly higher in KCl plus HCTZ-patients compared to HCTZ-treated patients (P=0.01).</p> <p>Serum potassium was found to be significantly higher in all combination treated-patients compared to HCTZ-treated patients (P=0.01 for all comparisons). Total body potassium was significantly higher in amiloride plus HCTZ- and triamterene plus HCTZ-treated patients (P=0.05 for both), but not in KCl plus HCTZ-treated patients (P value not reported), compared to HCTZ-treated patients.</p> <p>Secondary: The percentages of patients that became hypokalemic were 39, 52 and 52% in</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>amiloride plus HCTZ-, triamterene plus HCTZ- and KCl plus HCTZ-treated patients (P values not reported).</p> <p>The median days until hypokalemia detection were 114.0, 75.0 and 51.5 for amiloride plus HCTZ-, triamterene plus HCTZ- and KCl plus HCTZ-treated patients (P values not reported).</p> <p>Serum magnesium was maintained at a significantly higher rate in amiloride plus HCTZ- and triamterene plus HCTZ- patients compared to KCl plus HCTZ-treated patients (P values not reported).</p>
<p>Faris et al.<sup>25</sup> (2006)</p> <p>Potassium-sparing diuretics (amiloride, triamterene), loop diuretics (furosemide, bumetanide), or thiazide diuretics (chlorothiazide)</p> <p>vs</p> <p>placebo or active control (ACE inhibitors, digoxin)</p>	<p>MA (14 trials)</p> <p>Adult patients with chronic heart failure</p>	<p>N=525</p> <p>2 to 52 weeks</p>	<p>Primary: Mortality</p> <p>Secondary: Effect of diuretic withdrawal on worsening of heart failure and exercise capacity</p>	<p>Primary: Pooled data from three PC trials (n=202) reporting on mortality revealed that mortality was lower for diuretic-treated patients compared to placebo-treated patients (2.7 vs 10.9%, respectively; OR, 0.24; 95% CI, 0.07 to 0.83; P=0.02). The difference represents an absolute risk reduction of 8% in mortality in diuretic-treated patients (NNT, 12.5).</p> <p>Secondary: Pooled data from two PC trials (n=169) reporting on the effect of diuretics on worsening heart failure revealed lower admission rates for worsening heart failure in diuretic-treated patients compared to placebo-treated patients (OR, 0.07; 95% CI, 0.01 to 0.52; P=0.01).</p> <p>Pooled data from two parallel RCTs (n=43) reporting on the effect of diuretics on exercise capacity revealed that diuretic therapy improved exercise capacity compared to active control (WMD, 0.74; 95% CI, 0.37 to 1.11; P&lt;0.0001). Pooled data from two XO RCTs (n=48) revealed similar results (WMD, 0.67; 95% CI, 0.02 to 1.31; P=0.04). In total (n=91), diuretic therapy improved exercise capacity in patients with chronic heart failure (WMD, 0.72; 95% CI, 0.40 to 1.04; P&lt;0.0001).</p>
<b>Hypertension</b>				
<p>Heran et al.<sup>26</sup> (2010)</p> <p>Potassium sparing diuretics (amiloride,</p>	<p>SR (6 RCTs)</p> <p>Patients with a baseline office SBP ≥140 mm Hg and/or DBP ≥90</p>	<p>N=496</p> <p>3 to 12 weeks</p>	<p>Primary: Quantify the dose-related SBP and DBP lowering efficacy of potassium sparing</p>	<p>Primary: <i>Blood pressure lowering efficacy of potassium sparing diuretics as a second drug:</i></p> <p>There was no effect on SBP (-0.03 mm Hg; 95% CI, -2.90 to 2.83) and DBP (-0.22 mm Hg; 95% CI, -2.01 to 1.57) when potassium sparing diuretics were initiated at a dose of half the recommended starting dose. Due to the lack of</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>triamterene)</p> <p>Monotherapy vs placebo and as combination therapy with another antihypertensive drug class (ACE inhibitor, ARB, <math>\beta</math>-blocker, calcium channel blocker, centrally-acting drugs, diuretics and renin inhibitors)</p>	<p>mm Hg</p>		<p>diuretics</p> <p>Secondary: Variability of blood pressure, pulse pressure, heart rate, withdrawals due to adverse effects</p>	<p>data, an estimate of the effect of higher doses or whether there was a dose response effect could not be determined.</p> <p>Secondary: <i>Blood pressure lowering efficacy of potassium sparing diuretics as a second drug:</i> The limited data did not suggest any effect of potassium sparing on blood pressure variability.</p> <p>Analysis of six trials assessing amiloride and triamterene did not suggest any effect of potassium sparing diuretics on pulse pressure.</p> <p>Two trials provided heart rate data and did not suggest any effect of potassium sparing diuretics on heart rate.</p> <p>An analysis of withdrawals due to adverse effects during three to 12 weeks of treatment with potassium sparing diuretics was reported in five of trials. The overall estimate showed no significant effect of potassium sparing diuretics on this outcomes (RR, 0.53; 95% CI, 0.19 to 1.51).</p>
<p>Multicenter Diuretic Cooperative Study Group<sup>27</sup> (1981)</p> <p>Amiloride 5 mg QD</p> <p>vs</p> <p>amiloride and HCTZ 5-50 mg QD (fixed-dose combination product)</p> <p>vs</p>	<p>DB, MC, RCT</p> <p>Patients 21 to 69 years of age with mild to moderate essential HTN (supine DBP 95 to 115 mm Hg)</p>	<p>N=179</p> <p>12 weeks</p>	<p>Primary: Change from baseline in average supine SBP and DBP</p> <p>Secondary: Heart rate, body weight, serum potassium</p>	<p>Primary: Baseline vs 12 week average supine blood pressure was 153/101 vs 139/93 for amiloride-, 160/100 vs 137/90 for amiloride and HCTZ- and 154/101 vs 134/89 mm Hg for HCTZ-treated patients. Reductions in supine blood pressure were significant with all treatments (P&lt;0.01). The SBP reduction was significantly greater with amiloride and HCTZ-treated patients compared to amiloride-treated patients at all weeks and HCTZ-treated patients at four and eight weeks (P&lt;0.05, both).</p> <p>Secondary: No significant changes from baseline in heart rate were observed in amiloride and HCTZ-treated patients (P values not reported). An increase in heart rate of 3.3 bpm was observed in these patients (P&lt;0.05).</p> <p>Changes in body weight from baseline were -1.17 kg in amiloride and HCTZ-, -0.72 kg in HCTZ- and 0.045 kg in amiloride-treated patients (P&lt;0.05, for amiloride plus HCTZ only).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
HCTZ 50 mg QD				Changes in serum potassium from baseline were 0.23 in amiloride- (P<0.01), -0.38 in amiloride and HCTZ- (P<0.01) and -0.59 mEq/L in HCTZ-treated patients (P<0.01). The change in HCTZ-treated patients was statistically greater than the change in the amiloride and HCTZ-treated patients (P<0.05). Twenty three, two and zero percent of HCTZ-, amiloride and HCTZ- and amiloride-treated patients experienced hypokalemia.
Salmela et al. <sup>28</sup> (1986)  Amiloride 2.5 mg/day and HCTZ 25 mg/day  vs  HCTZ 25 mg/day daily	DB, MC, PG, RCT  Adult patients with mild to moderate HTN	N=40  12 weeks	Primary: Changes in blood pressure  Secondary: Not reported	Primary: At the end of the first treatment period (four weeks), mean supine SBP and DBP was 161 and 91 mm Hg in amiloride plus HCTZ-treated patients (P<0.01 and P<0.001, respectively).  At the end of the first treatment period (four weeks), mean supine SBP and DBP was 165 and 96 mm Hg in HCTZ-treated patients (P<0.01 for both).  At the end of the second treatment period (eight weeks), mean supine SBP and DBP was 154 and 86 mm Hg in amiloride plus HCTZ-treated patients (P<0.01 and P<0.001).  At the end of the second treatment period (eight weeks), mean supine SBP and DBP was 155 and 90 mm Hg in HCTZ-treated patients (P<0.001 and P<0.001).  There were no significant differences in blood pressure reduction between the two treatments (P value not reported).  Secondary: Not reported
Hood et al. <sup>29</sup> (2007) SALT study  Amiloride 20 mg/day  vs  amiloride 40	DB, RCT, XO  Adult patients with seated blood pressure of 140/90 to 170/110 mm Hg, plasma renin of ≤12 mU/L, plasma aldosterone-renin	N=57  42 weeks	Primary: Change in blood pressure and plasma renin from baseline between spironolactone 100 mg/day and bendroflumethiazide 5 mg/day	Primary: Spironolactone 100 mg/day- and bendroflumethiazide 5 mg/day-treated patients did not exhibit a significant difference in blood pressure reduction from baseline (P value not reported).  Secondary: Spironolactone 50 mg/day-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (P<0.01).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day vs spironolactone 50 mg/day vs spironolactone 100 mg/day vs bendroflumethiazide* 2.5 mg/day vs bendroflumethiazide* 5 mg/day vs losartan 100 mg/day vs placebo	ratio >750, previous fall in SBP $\geq$ 20 mm Hg after 1 month of OL treatment with spironolactone 50 mg/day		Secondary: Change in blood pressure and plasma renin from baseline between amiloride and other diuretics and between lower and higher doses of each diuretic	Losartan 100 mg-treated patients exhibited a significant decrease in blood pressure from baseline compared to bendroflumethiazide 2.5 mg/day-treated patients (P<0.05).  High-dose bendroflumethiazide- and amiloride-treated patients exhibited significantly greater reductions in blood pressure compared to the lower doses (P<0.05).  Spironolactone-treated patients exhibited a four-fold increase in baseline renin level compared to a two-fold increase observed in bendroflumethiazide-treated patients (P=0.003).
Kohvakka et al. <sup>30</sup> (1979)  Amiloride 5 mg QD	PC, RCT, XO  Patients 41 to 70 years of age with uncomplicated	N=31  3 months	Primary: Changes in blood pressure, serum potassium, sodium, creatinine, urate	Primary: No significant changes in blood pressure were observed with any of the treatments (P values not reported).  Mean serum potassium was reduced with all treatments except with

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>triamterene 75 mg QD</p> <p>vs</p> <p>KCl 1,500 mg QD</p> <p>vs</p> <p>spironolactone 50 mg QD</p> <p>vs</p> <p>placebo</p> <p>All patients were also receiving HCTZ 50 mg QD.</p>	<p>HTN, previously treated with antihypertensive agents for 1 to 6 years</p>		<p>and total body potassium</p> <p>Secondary: Not reported</p>	<p>spironolactone. KCl supplementation was least effective in elevating serum potassium. Total body potassium remained constant throughout treatment (P values not reported).</p> <p>Serum sodium remained within normal limits with all treatments (P values not reported).</p> <p>There were no significant changes in mean serum creatinine with any of the treatments (P values not reported).</p> <p>Serum urate concentration increased significantly with all treatments, including HCTZ monotherapy (P values not reported).</p> <p>Secondary: Not reported</p>
<p>Larochelle et al.<sup>31</sup> (1985)</p> <p>Amiloride 5 mg/day and HCTZ 50 mg/day</p> <p>vs</p> <p>HCTZ 50 mg/day</p>	<p>DB, RCT</p> <p>Ambulant patients 18 to 70 years of age with essential HTN who after not being treated for <math>\geq 2</math> weeks prior to the trial had a supine DBP of 95 to 109 mm Hg and a serum potassium level of <math>&gt;3.5</math> mmol/L</p>	<p>N=266</p> <p>8 weeks</p>	<p>Primary: Blood pressure, serum potassium concentration</p> <p>Secondary: Not reported</p>	<p>Primary: At eight weeks, there were no differences between the two treatments in the mean blood pressure reductions (P value not reported).</p> <p>During the eight weeks of treatment, the HCTZ plus amiloride-treated patients experienced a decrease in mean supine blood pressure (159/99 to 138/88 mm Hg) and serum potassium levels (4.23 to 3.91 mmol/L) (P values not reported).</p> <p>During the eight weeks of treatment, HCTZ-treated patients experienced a reduction in mean supine blood pressure (157/99 to 138/87 mm Hg) and serum potassium levels (4.16 to 3.69 mmol/L) (P values not reported).</p> <p>Hypokalemia occurred less frequently in HCTZ plus amiloride-treated patients compared to HCTZ-treated patients (14 and 29%, respectively; P=0.0026). However, the proportions of patients with a potassium level exceeding 4.5</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>mmol/L were similar (4.5 vs 3.9%, respectively; P value not reported).</p> <p>Secondary: Not reported</p>
<p>Dean et al.<sup>32</sup> (1984)</p> <p>Amloride and HCTZ 5-50 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>triamterene and HCTZ 50-25 mg QD (fixed-dose combination product)</p>	<p>RCT, SB, XO</p> <p>Patients with mild to moderate HTN (DBP 95 to 110 mm Hg)</p>	<p>N=20</p> <p>8 weeks</p>	<p>Primary: Blood pressure, hypokalemia, hyperkalemia, renal function tests</p> <p>Secondary: Not reported</p>	<p>Primary: Both treatments produced a comparable effect on blood pressure. The baseline standing and lying blood pressure was 168/105 and 168/104 mm Hg, respectively. After eight weeks, amloride and HCTZ-treated patients had a standing and lying blood pressure of 145/92 and 145/90 mm Hg, respectively (P values not reported). After eight weeks, triamterene and HCTZ-treated patients had a standing and lying blood pressure of 142/93 and 143/91 mm Hg, respectively (P values were not reported).</p> <p>There were no cases of hypokalemia or hyperkalemia and no renal function changes with either treatment (P values were not reported).</p> <p>Secondary: Not reported</p>
<p>Maxwell et al.<sup>33</sup> (1985)</p> <p>Amloride and HCTZ 50-25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>triamterene and HCTZ 5-50 mg QD (fixed-dose combination product)</p>	<p>OL, PRO, RCT</p> <p>Patients with mild to moderate HTN, mean supine DBP &lt;90 or &gt;114 mm Hg at the end of a 3 week placebo, run-in phase</p>	<p>N=84</p> <p>9 weeks</p>	<p>Primary: Mean blood pressure changes</p> <p>Secondary: Not reported</p>	<p>Primary: Seventy three (n=30) and 81% (n=35) of triamterene and HCTZ- and amloride and HCTZ-treated patients were maintained on the initial dosage throughout the trial, with no significant differences between the two treatments (P value not reported).</p> <p>At week nine, mean SBP and DBP was 136.2 and 87.4 mm Hg in triamterene and HCTZ-treated patients (P value not reported). At week nine, mean SBP and DBP was 132.6 and 85.7 mm Hg in amloride and HCTZ-treated patients (P value not reported).</p> <p>At week nine, mean serum potassium levels were 4.13 and 3.98 mEq/L in triamterene and HCTZ- and amloride and HCTZ-treated patients (P&lt;0.05).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>All patients received placebo for 3 weeks prior to the treatment phase.</p> <p>After 2 weeks of treatment, dosage could be doubled.</p>				
<p>Hansson et al.<sup>34</sup> (1999) HYPERTENSION -2 (STOP)</p> <p><u>Conventional drug group</u> Atenolol 50 mg QD, HCTZ 25 mg QD plus amiloride 2.5 mg QD, metoprolol 100 mg QD, or pindolol 5 mg QD</p> <p>vs</p> <p><u>Newer drug group</u> ACE inhibitors (enalapril 10 mg QD or lisinopril 10 mg QD) or calcium channel blockers (felodipine 2.5 mg QD, or isradipine 2 to 5 mg QD)</p>	<p>BE, MC, OL, RCT</p> <p>Swedish men and women between 70 to 84 years old with treated or untreated essential with HTN on 3 separate occasions defined by SBP <math>\geq 180</math> mm Hg, DBP <math>&gt;105</math> mm Hg, or both</p>	<p>N=6,614</p> <p>60 months</p>	<p>Primary: Combined fatal stroke, MI, and other fatal cardiovascular disease; combined fatal and nonfatal stroke, MI, and other cardiovascular Mortality</p> <p>Secondary: Not reported</p>	<p>Primary: The combined fatal mortality endpoints occurred in 221 of the 2,213 patients in the conventional drugs group and in 438 of 4,401 in the newer drugs group (RR, 0.99; 95% CI, 0.84 to 1.16; P=0.89).</p> <p>The combined fatal and nonfatal mortality endpoints occurred in 460 patients taking conventional drugs and in 887 taking newer drugs (RR, 0.96; 95% CI, 0.86 to 1.08; P=0.49).</p> <p>Secondary: Not reported</p>
<p>Williams et al.<sup>35</sup></p>	<p>3 phase, OL</p>	<p>N=156</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1984)</p> <p>Phase 1 (Baseline, 2 weeks): Triamterene and HCTZ 75-50 mg/day (fixed-dose combination product) (Group 1)</p> <p>vs</p> <p>triamterene and HCTZ 150-100 mg/day (fixed-dose combination product) (Group 3)</p> <p>vs</p> <p>no antihypertensive medications (Group 3)</p> <p>Phase 2 (4 weeks): Triamterene and HCTZ 75-50 mg/day (fixed-dose combination product) (Groups 1, 2 and 3)</p> <p>Phase 3 (up to 8 months): Triamterene and HCTZ 75-50</p>	<p>Patients 21 to 70 years of age, with essential HTN</p>	<p>6 to 32 weeks</p>	<p>Blood pressure and weight comparisons between Phase 1 and 2</p> <p>Secondary: Serum potassium concentrations</p>	<p>During Phase 1, mean standing DBP, mean standing SBP and weight for Group 1-treated patients were: 91 mm Hg, 138 mm Hg and 82 kg (P values not reported). During Phase 2, the comparisons in these patients were: 88 mm Hg, 135 mm Hg and 82 kg (P values not reported).</p> <p>During Phase 1, mean standing DBP, mean standing SBP and weight for Group 2-treated patients were: 93 mm Hg, 139 mm Hg and 87 kg (P values not reported). During Phase 2, the comparisons in these patients were: 98 mm Hg, 149 mm Hg and 79 kg (P value not reported).</p> <p>During Phase 1, mean standing DBP, mean standing SBP and weight for Group 3-treated patients were: 98 mm Hg, 149 mm Hg and 80 kg (P values not reported). During Phase 2, the comparisons in these patients were: 94 mm Hg, 136 mm Hg and 78 kg (P value not reported).</p> <p>Of these Phase 1 and 2 comparisons, mean standing DBP and SBP differences were reported to be significant during Phase 2 for Group 1- and Group 3-treated patients (P values not reported).</p> <p>Secondary: When Group 2-treated patients were switched 75/50 mg/day, no patient became hypokalemic (serum potassium concentration &lt;3.5 mEq/L) (P value not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day (fixed-dose combination product) (Groups 1, 2 and 3)				
<p>Hansson et al.<sup>36</sup> (2000) NORDIL</p> <p>Conventional therapy (diuretic, <math>\beta</math>-blocker or both)</p> <p>vs</p> <p>diltiazem 180 to 360 mg QD</p>	<p>BE, MC, OL, PRO, RCT</p> <p>Patients 50 to 74 years of age with DBP <math>\geq</math>100 mm Hg and previously untreated</p>	<p>N=10,881</p> <p>4.5 years</p>	<p>Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death</p> <p>Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI</p>	<p>Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/<math>\beta</math>-blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).</p> <p>Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/<math>\beta</math>-blocker patients (P=0.04).</p> <p>Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/<math>\beta</math>-blocker patients (P=0.17).</p> <p>Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57 and congestive heart failure (P=0.42).</p>
<p>Messerli et al.<sup>37</sup> (1998)</p> <p>Diuretics (amiloride, chlorthalidone, HCTZ, HCTZ and triamterene [fixed-dose combination product], or thiazide)</p> <p>vs</p> <p><math>\beta</math>-blockers (atenolol, metoprolol or pindolol)</p>	<p>MA</p> <p>10 RCTs lasting <math>\geq</math>1 year, which used as first line agents diuretics and/or <math>\beta</math>-blockers and reported morbidity and mortality outcomes in patients <math>\geq</math>60 years of age with HTN</p>	<p>N=16,164</p> <p>1 year</p>	<p>Primary: Cardiovascular morbidity and mortality, all-cause morbidity</p> <p>Secondary: Not reported</p>	<p>Primary: Diuretic treatment significantly reduced the odds for cardiovascular mortality by 25% (OR, 0.75; 95% CI, 0.64 to 0.87), while <math>\beta</math>-blockers did not reduce cardiovascular mortality (OR, 0.98; 95% CI, 0.78 to 1.23; P values not reported).</p> <p>Diuretic treatment significantly reduced the odds for all-cause mortality by 14% (OR, 0.86; 95% CI, 0.77 to 0.96), while <math>\beta</math>-blockers did not reduce all-cause mortality (OR, 1.05; 95% CI, 0.88 to 1.25; P values not reported).</p> <p>Secondary: Not reported</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Lindholm et al.<sup>38</sup> (2005)</p> <p>β-blocker therapy (atenolol, metoprolol, oxprenolol*, pindolol or propranolol)</p> <p>vs</p> <p>other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan or verapamil)</p> <p>or</p> <p>placebo</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a β-blocker as first line treatment (in ≥50% of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary: The RR of stroke was 16% higher with β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non β-blockers (RR, 1.26%; 95% CI, 1.15 to 1.38; P&lt;0.0001).</p> <p>The relative risk of MI was 2% higher for β-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).</p> <p>The RR of all-cause mortality was 3% higher for β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).</p> <p>Secondary: Not reported</p>
<p>Wysong et al.<sup>39</sup> (2007)</p> <p>Other antihypertensive therapies (i.e.,</p>	<p>MA</p> <p>13 RCTs evaluating patients ≥18 years of age with HTN</p>	<p>N=91,561</p> <p>Duration varied</p>	<p>Primary: All-cause mortality</p> <p>Secondary: Stroke, CHD, cardiovascular</p>	<p>Primary: There was not a significant difference observed in all-cause mortality between β-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)</p> <p>vs</p> <p>β-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)</p>			<p>death, total cardiovascular disease, adverse reactions</p>	<p>β-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).</p> <p>Secondary:</p> <p>There was a significant decrease in stroke observed with β-blocker therapy compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between β-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with β-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of β-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with β-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, QD=once daily

Study regimen abbreviations: BE=blinded endpoint, DB=double blind, MA=meta analysis, MC=multicenter, OL=open label, OS=observational, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, SB=single blind, SR=systematic review, XO=cross over

Miscellaneous abbreviations: ACE inhibitor=angiotensin converting enzyme inhibitor, ARB=angiotensin II receptor blocker, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, HCTZ=hydrochlorothiazide, HTN=hypertension, MI=myocardial infarction, NNT=number needed to treat, NYHA=New York Heart Association, OR=odds ratio, RR=relative risk, SBP=systolic blood pressure, WMD=weighted mean difference

### Additional Evidence

#### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

## IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 11. Relative Cost of the Potassium-Sparing Diuretics**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Single Entity Agents</b>				
Amiloride	tablet	N/A	N/A	\$
Triamterene	capsule	N/A	N/A	\$\$\$\$
<b>Combination Products</b>				
Amiloride and HCTZ	tablet	N/A	N/A	\$
Triamterene and HCTZ	capsule, tablet	Maxzide®*	\$\$\$	\$

\*Generic is available in at least one dosage form or strength.

HCTZ=hydrochlorothiazide, N/A=Not available

## X. Conclusions

The potassium-sparing diuretics are approved for the treatment of congestive heart failure, edema, and hypertension.<sup>1-4</sup> When used alone, potassium-sparing diuretics have a weak diuretic and antihypertensive effect and an increased risk of hyperkalemia. The potassium-sparing diuretics are generally used in combination with other diuretics to help restore normal serum potassium levels or to prevent the development of hypokalemia.<sup>1-4</sup> Amiloride and triamterene are available as a fixed-dose combination with hydrochlorothiazide. All of the products are available in a generic formulation.

For the treatment of chronic heart failure, guidelines recommend the use of diuretics in all patients who have evidence of volume overload. The loop diuretics are generally recommended as initial therapy in patients with left ventricular dysfunction. For those with persistent fluid retention despite treatment with a loop diuretic, a thiazide diuretic or metolazone may be added to the regimen. In patients with normal left ventricular function, either a thiazide diuretic or loop diuretic may be used as initial therapy to manage fluid overload.<sup>8-9</sup> As indicated by the FDA-approved indications of the potassium-sparing diuretics, these agents are typically used as adjunctive therapy in patients receiving thiazide diuretics to prevent the development of hypokalemia or to restore normal serum potassium levels.<sup>1-4</sup> Triamterene may be used along or with other diuretics, either for its added diuretic effect or its potassium-sparing potential.<sup>3</sup>

There are several national and international organizations that have published guidelines on the treatment of hypertension.<sup>11-19</sup> Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension. According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another antihypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>11</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>11-19</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>11-19</sup> The use of a fixed-dose combination product may simplify the treatment regimen and improve adherence.<sup>13,14,17</sup> However, there are no prospective, randomized trials that have demonstrated better clinical outcomes with a fixed-dose combination product compared to the coadministration of the individual components as separate formulations.

Amiloride has been shown to be effective for the treatment of edema and hypertension, as well as for the prevention of serum potassium loss in patients taking a thiazide or loop diuretic. Clinical trials have also demonstrated comparable efficacy with the fixed-dose combination of amiloride-hydrochlorothiazide and triamterene-hydrochlorothiazide in patients with hypertension and heart failure.<sup>20-39</sup>

There is insufficient evidence to support that one brand potassium-sparing diuretic is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand potassium-sparing diuretics within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand potassium-sparing diuretic is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Maxzide® [package insert]. Morgantown (WV): Mylan Pharmaceuticals, Inc.; 2018 Jul.
2. Dyazide® [package insert]. Research Triangle Park (NC): GlaxoSmithKline; 2011 Mar.
3. Dyrenium® [package insert]. Dublin (Ireland): Concordia Pharmaceuticals; 2019 Sep.
4. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
5. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
6. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
7. DiPiro JT, Yee GC, Posey LM, Haines ST, Nolin TD, Ellingrod V, editors. Pharmacotherapy: a pathophysiologic approach. 11th edition. New York (NY): McGraw-Hill; 2020. <https://accesspharmacy.mhmedical.com/book.aspx?bookid=2577>. Accessed January 2022.
8. Kaplan NM, Victor RG, Flynn JT. Kaplan's clinical hypertension. 11<sup>th</sup> ed. Philadelphia (PA): Lippincott, Williams, and Wilkins; 2015.
9. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
10. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
11. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
12. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, Ramirez A, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
13. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, Butalia S, Leung AA, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
14. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
15. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
16. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
17. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
18. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
19. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl.1).
20. Bayliss J, Norell M, Canepa-Anson R, Sutton G, Poole-Wilson P. Untreated heart failure: clinical and neuroendocrine effects of introducing diuretics. *Br Heart J*. 1987 Jan;57(1):17-22.
21. Rengo F, Trimarco B, Bonaduce D, Petretta M, Ferrara N, D'Ascia C, et al. Potassium-sparing effect of amiloride in patients receiving diuretics: a quantitative review. *Acta Cardiol*. 1979;34(4)259-67.

22. Cheitlin MD, Byrd R, Benowitz N, Liu E, Modin G. Amiloride improves hemodynamics in patients with chronic congestive heart failure treated with chronic digoxin and diuretics. *Cardiovasc Drugs Ther.* 1991 Aug;5(4):719-25.
23. Ghosh AK, Mankikar G, Strouthidis T, Windsor A, Long C, Glover DR. A single-blind, comparative study of hydrochlorothiazide/amiloride ('Moduret' 25) and hydrochlorothiazide/triamterene ('dyazide') in elderly patients with congestive heart failure. *Curr Med Res Opin.* 1987;10(9):573-9.
24. Kohvakka A. Maintenance of potassium balance during long-term diuretic therapy in chronic heart failure patients with thiazide-induced hypokalemia: comparison of potassium supplementation with potassium chloride and potassium-sparing agents, amiloride and triamterene. *Int J Clin Pharmacol Ther Toxicol.* 1988;26(5):273-7.
25. Faris R, Flather MD, Purcell H, Poole-Wilson PA, Coats AJ. Diuretics for heart failure. *Cochrane Database Syst Rev.* 2006 Jan 25;(1):CD003838.
26. Heran BS, Chen JMH, Wang JJ, Wright JM. Blood pressure lowering efficacy of potassium-sparing diuretics (that block the epithelial sodium channel) for primary hypertension. *Cochrane Database of Systemic Reviews.* 2010, Issue 1. Art. No.:CD008167. DOI:10.1002/14651858.CD008167.pub2.
27. Multiclinic comparison of amiloride, hydrochlorothiazide, and hydrochlorothiazide plus amiloride in essential hypertension. Multicenter Diuretic Cooperative Study Group. *Arch Intern Med.* 1981 Mar;141(4):482-6.
28. Salmela P, Juustila H, Kinnunen O, and Koistinen P. Comparison of low doses of hydrochlorothiazide plus amiloride and hydrochlorothiazide alone in hypertension in elderly patients. *Ann Clin Res.* 1986;18:88-92.
29. Hood SJ, Taylor KP, Ashby MJ, Brown MJ. The spironolactone, amiloride, losartan, and thiazide (SALT) double-blind crossover trial in patients with low-renin hypertension and elevated aldosterone-renin ratio. *Circulation.* 2007;116(3):268-75.
30. Kohvakka A, Eisalo A, Manninen V. Maintenance of potassium balance during diuretic therapy. *Acta Med Scand.* 1979;205(4):319-24.
31. Larochelle P, Logan AG. Hydrochlorothiazide-amiloride versus hydrochlorothiazide alone for essential hypertension: effects on blood pressure and serum potassium level. *Can Med Assoc J.* 1985;132(7):801-5.
32. Dean S, Spencer-Mills L. Hydrochlorothiazide in combination with potassium-sparing agents in the treatment of hypertension. *Curr Med Res Opin.* 1984;9(5):287-9.
33. Maxwell MH, Brachfeld J, Itskovitz H, et al. Blood pressure lowering and potassium conservation by triamterene-hydrochlorothiazide and amiloride-hydrochlorothiazide in hypertension. *Clin Pharmacol Ther.* 1985;37(1):61-5.
34. Hansson L, Lindholm LH, Ekblom T, Dahlöf B, Lanke J, Scherstén B, et al. Randomized trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 (STOP) study. *Lancet.* 1999 Nov 20;354(9192):1751-6.
35. Williams RL, Clark T, Blume CD. Clinical experience with a new combination formulation of triamterene and hydrochlorothiazide (Maxzide) in patients with mild to moderate hypertension. *Am J Med.* 1984 Nov 5;77(5A):62-6.
36. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, et al. Randomized trial of effects of calcium antagonists compared with diuretics and  $\beta$ -blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet.* 2000 Jul 29;356(9227):359-65.
37. Messerli FH, Grossman E, Goldbourt U. Are beta-blockers efficacious as first-line therapy for hypertension in the elderly? A systematic review. *JAMA.* 1998 Jun 17;279(23):1903-7.
38. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet.* 2005 Oct 29-Nov 4;366(9496):1545-53.
39. Wiyongse CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev.* 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Thiazide Diuretics  
AHFS Class 402820  
May 4, 2022**

**I. Overview**

The thiazide diuretics are approved for the treatment hypertension and edema due to renal dysfunction. They are also approved as adjunctive therapy for the management of edema associated with congestive heart failure, hepatic cirrhosis, as well as corticosteroid and estrogen therapy.<sup>1-3</sup> The thiazide diuretics inhibit the reabsorption of sodium and chloride in the cortical thick ascending limb of the loop of Henle and the early distal tubules. This action leads to an increase in the urinary excretion of sodium and chloride in approximately equivalent amounts.<sup>1-5</sup> Additionally, increased potassium and bicarbonate excretion, decreased calcium excretion, and uric acid retention may be observed. During initial thiazide therapy a reduction in cardiac output and extracellular volume occurs. However, with chronic therapy cardiac output normalizes and both peripheral vascular resistance and extracellular volume are reduced. In general, similar therapeutic and adverse effects are seen when equipotent doses are used. Thiazide diuretics are generally recommended when the glomerular filtration rate is above 30 mL/min.<sup>4-6</sup>

The thiazide diuretics that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the agents are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Thiazide Diuretics Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Chlorothiazide	injection*, suspension	Diuril®	chlorothiazide
Hydrochlorothiazide	capsule, tablet	N/A	hydrochlorothiazide

\*Generic is available in at least one dosage form or strength.

PDL=Preferred Drug List

N/A=Not available

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the thiazide diuretics are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Thiazide Diuretics**

Clinical Guideline	Recommendation(s)
American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022) <sup>7</sup>	<p><b>Treatment of Stage A heart failure (HF)</b></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><b>Treatment of Stage B heart failure</b></p> <ul style="list-style-type: none"> <li>In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</p> <ul style="list-style-type: none"> <li>• In patients with a recent MI and LVEF ≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF ≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><u>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</u></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three β-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is &gt;30 mL/min/1.73 m<sup>2</sup> and serum potassium is &lt;5.0 mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at &lt;5.5 mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> </ul>



Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</li> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><u>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</u></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><b>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</b></p> <ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><b>Treatment of Stage D (advanced/refractory) HF</b></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>8</sup></p>	<p><b>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</b></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq 35\%</math>, in sinus rhythm and a resting heart rate <math>\geq 70</math> bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor (patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</p> <ul style="list-style-type: none"> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HFrEF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HFmrEF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HFmrEF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HFmrEF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HFpEF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HFpEF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HFpEF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>disease or with CV disease in order to prevent HF hospitalizations.</p> <ul style="list-style-type: none"> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>9</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>initial drug should be increased or second drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</p> <ul style="list-style-type: none"> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: Global Hypertension Practice Guidelines (2020)<sup>10</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease</li> </ul> </li> </ul>

Clinical Guideline	Recommendation(s)
	<p>(CKD), diabetes mellitus (DM), or hypertension-mediated organ damage (HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</p> <ul style="list-style-type: none"> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> <li>● Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: <b>2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)</b><sup>11</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>● Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt;</math>60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> </ul> </li> </ul>



Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul style="list-style-type: none"> <li>• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li>• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li>• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li>• Possible reasons for poor response to therapy should be considered.</li> <li>• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li>• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li>• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li>• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li>• Short-acting nifedipine should not be used.</li> <li>• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><b>Guidelines for patients with hypertension who have had a recent myocardial infarction</b></p> <ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><b>Treatment of hypertension in association with heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction &lt;40%), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq</math>30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><b>Treatment of hypertension in association with stroke</b></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><b>Treatment of hypertension in association with LVH</b></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or</li> </ul>



Clinical Guideline	Recommendation(s)
	<p><b>minoxidil should not be used.</b></p> <p><b>Treatment of hypertension in association with nondiabetic chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><b>Treatment of hypertension in association with renovascular disease</b></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><b>Treatment of hypertension in association with diabetes mellitus</b></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><b>Resistant hypertension</b></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>significantly, with the greatest BP-lowering shown with spironolactone.</p> <ul style="list-style-type: none"> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90<sup>th</sup> percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>12</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a CCB or diuretic.</p> <ul style="list-style-type: none"> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (≥65 years) are treated with the same recommendations in non-older patient population. In very old patients (≥80 years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should be avoided.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Treatment strategies in hypertensive women</u></p> <ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• RAS blockers are more effective in reducing albuminuria than other antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</li> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq</math>220 mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq</math>140/90 mmHg.</li> <li>• In patients with HFrEF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HFpEF, BP treatment threshold and target values should be the same as for HFrEF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq</math>130 mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis</b></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq</math>160 mmHg) the same treatment as people with both raised systolic and diastolic blood</li> </ul>

Clinical Guideline	Recommendation(s)
<p><b>and management (2019)<sup>13</sup></b></p>	<p>pressure.</p> <ul style="list-style-type: none"> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>14</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease</b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD)</u></p>

Clinical Guideline	Recommendation(s)
(2021) <sup>15</sup>	<p><u>not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adults kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)<sup>16</sup></b></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) ≥130 mmHg or an average diastolic blood pressure (DBP) of ≥80 mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of ≥10% and an average SBP of ≥130 mmHg or an average ≥80 mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of ≥140 mmHg or a DBP of ≥90 mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of ≥10%, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide</li> </ul>



Clinical Guideline	Recommendation(s)
	<p>diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</p> <ul style="list-style-type: none"> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>• Adults with SIHD and hypertension (BP ≥130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (≥300 mg/d, or ≥300 mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</p> <ul style="list-style-type: none"> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq</math>220/120 mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq</math>140/90 mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (<math>\geq 65</math> years of age) with an average SBP of <math>\geq 130</math> mmHg.</li> <li>• For older adults (<math>\geq 65</math> years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP <math>\geq 180</math> mmHg or DBP <math>\geq 110</math> mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>• Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>17</sup></p>	<p><b>Hypertension/blood pressure control</b></p> <ul style="list-style-type: none"> <li>• Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq 140/90</math>) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq 180/110</math> and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>• All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>• For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>• For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of <math>&lt;130/80</math> mmHg may be appropriate, if it can be safely attained.</li> <li>• For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;15\%</math>), treat to blood pressure target of <math>&lt;140/90</math>.</li> <li>• In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>• For patients with blood pressure <math>&gt;120/80</math>, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>• Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>• Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>• Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>• An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>• For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>• Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Association for the Study of Liver Diseases: <b>Diagnosis, Evaluation, and Management of</b></p>	<p><b>Treatment of ascites</b></p> <ul style="list-style-type: none"> <li>• Moderate sodium restriction (2 g or 90 mmol/day) and diuretics (spironolactone with or without furosemide) are the first-line treatment in patients with cirrhosis and grade 2 ascites.</li> <li>• After ascites is adequately mobilized, attempts should be made to taper the diuretics to the lowest dose necessary to maintain minimal or no ascites to prevent the development of adverse effects.</li> </ul>

Clinical Guideline	Recommendation(s)
Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance (2021) <sup>18</sup>	<ul style="list-style-type: none"> <li>Fluid restriction is not necessary unless serum sodium is &lt;125 mmol/L.</li> <li>In patients receiving diuretics, body weight and serum creatinine and sodium should be regularly monitored to assess response and to detect the development of adverse effects.</li> <li>Human albumin solution (20 to 40 g/week) or baclofen administration (10 mg/day, with a weekly increase of 10 mg/day, up to 30 mg/day) can be considered in cases of severe muscle cramps.</li> <li>Large-volume paracentesis is the first-line treatment of grade 3 ascites. After paracentesis, sodium restriction and diuretics should be started.</li> <li>Referral for liver transplant evaluation should be considered in patients with grade 2 or 3 ascites.</li> <li>Nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers should be avoided in patients with cirrhosis and ascites.</li> <li>Aminoglycosides should be avoided whenever possible in the treatment of bacterial infections.</li> <li>For patients with cirrhosis and diuretic-responsive ascites, controversial data suggest potential benefits of long-term infusion of human albumin solution. At present, no recommendation can be made for its use in routine clinical practice.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the thiazide diuretics are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Thiazide Diuretics<sup>1-4</sup>**

Indication	Chlorothiazide*	HCTZ*
<b>Edema</b>		
Adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, and corticosteroid and estrogen therapy	✓	✓ (tablet)
<b>Hypertension</b>		
Treatment of hypertension	✓ † (oral)	✓ †

\*Has been found useful in edema due to various forms of renal dysfunction such as nephrotic syndrome, acute glomerulonephritis, and chronic renal failure.

†Alone or in combination with other antihypertensive agents.

HCTZ=hydrochlorothiazide

### IV. Pharmacokinetics

The pharmacokinetic parameters of the thiazide diuretics are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Thiazide Diuretics<sup>5</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life
Chlorothiazide	Poor	Not reported	Not metabolized	Renal (96)	45 to 120 minutes
HCTZ	60 to 80	40	Not metabolized	Renal (50 to 70)	10 to 12 hours

HCTZ=hydrochlorothiazide

## V. Drug Interactions

Major drug interactions with the thiazide diuretics are listed in Table 5.

**Table 5. Major Drug Interactions with the Thiazide Diuretics<sup>5</sup>**

Generic Name(s)	Interaction	Mechanism
Thiazide diuretics (chlorothiazide, HCTZ)	Amphetamines	Concurrent use of amphetamines and thiazide diuretics may result in increased exposure to amphetamine.
Thiazide diuretics (chlorothiazide, HCTZ)	Digitalis glycosides	Thiazide diuretics may induce electrolyte disturbances which may predispose patients to digitalis-induced arrhythmias.
Thiazide diuretics (chlorothiazide, HCTZ)	Dofetilide	Thiazide diuretics may induce hypokalemia which may increase the risk of torsades de pointes.
Thiazide diuretics (chlorothiazide, HCTZ)	Lithium	Thiazide diuretics may promote enhanced proximal tubular reabsorption of lithium leading to elevated serum concentrations. Thiazide diuretics may increase the therapeutic and toxic effects of lithium.
Thiazide diuretics (chlorothiazide, HCTZ)	Loop diuretics	Both groups have synergistic effects that may result in profound diuresis and serious electrolyte abnormalities
Thiazide diuretics (chlorothiazide, HCTZ)	NSAIDs	Concurrent use of NSAIDs and thiazide diuretics may result in reduced diuretic effectiveness and possible nephrotoxicity.
Thiazide diuretics (chlorothiazide)	Bepridil	Concurrent use of chlorothiazide and bepridil may result in hypokalemia and subsequent cardiotoxicity (torsades de pointes).
Chlorothiazide	Flecainide	Concurrent use of chlorothiazide and flecainide may result in increased risk of electrolyte imbalance and subsequent cardiotoxicity.
Hydrochlorothiazide	Methotrexate	Concurrent use of hydrochlorothiazide and methotrexate may result in increased methotrexate exposure and enhanced myelosuppression.

HCTZ=hydrochlorothiazide, NSAIDs=nonsteroidal anti-inflammatory drugs

## VI. Adverse Drug Events

The most common adverse drug events reported with the thiazide diuretics are listed in Table 6.

**Table 6. Adverse Drug Events (%) Reported with the Thiazide Diuretics<sup>1-5</sup>**

Adverse Events	Chlorothiazide	HCTZ
<b>Cardiovascular</b>		
Hypotension	✓	✓
Necrotizing angiitis	✓	✓
Orthostatic hypotension	✓	✓
<b>Central Nervous System</b>		
Dizziness	✓	✓
Fever	✓	✓
Headache	✓	✓
Restlessness	✓	✓
Vertigo	✓	✓
<b>Dermatological</b>		
Alopecia	✓	✓
Cutaneous vasculitis	-	✓
Erythema multiforme	✓	✓
Exfoliative dermatitis	✓	✓
Photosensitivity	✓	✓
Purpura	✓	✓

Adverse Events	Chlorothiazide	HCTZ
Rash	✓	✓
Stevens-Johnson syndrome	✓	✓
Toxic epidermal necrolysis	✓	✓
Urticaria	✓	✓
Vasculitis	-	✓
<b>Gastrointestinal</b>		
Abdominal cramping	✓	✓
Anorexia	✓	✓
Constipation	✓	✓
Diarrhea	✓	✓
Gastric irritation	✓	✓
Nausea	✓	✓
Pancreatitis	✓	✓
Sialadenitis	✓	✓
Vomiting	✓	✓
<b>Genitourinary</b>		
Impotence	✓	✓
<b>Hematologic</b>		
Agranulocytosis	✓	✓
Aplastic anemia	✓	✓
Hemolytic anemia	✓	✓
Leukopenia	✓	✓
Thrombocytopenia	✓	✓
<b>Hepatic</b>		
Jaundice	✓	✓
<b>Laboratory Test Abnormalities</b>		
Cholesterol increased	✓	-
Electrolyte imbalance	✓	✓
Hypercalcemia	-	✓
Hyperglycemia	✓	✓
Hyperuricemia	✓	✓
Hypochloremic alkalosis	✓	-
Hypokalemia	✓	-
Hypomagnesemia	✓	-
Hyponatremia	✓	-
Triglycerides increased	✓	-
<b>Musculoskeletal</b>		
Muscle spasm	✓	✓
Paresthesia	✓	✓
Weakness	✓	✓
<b>Ocular</b>		
Blurred vision	✓	✓
Xanthopsia	✓	✓
<b>Renal</b>		
Glycosuria	✓	✓
Interstitial nephritis	✓	✓
Renal dysfunction	✓	✓
Renal failure	✓	✓
<b>Respiratory</b>		
Pneumonitis	✓	✓
Pulmonary edema	✓	✓
Respiratory distress	✓	✓
<b>Other</b>		
Anaphylactic reactions	✓	✓



Adverse Events	Chlorothiazide	HCTZ
Systemic lupus erythematosus	✓	-

✓ Percent not specified  
-Event not reported  
HCTZ=hydrochlorothiazide

## VII. Dosing and Administration

The usual dosing regimens for the thiazide diuretics are listed in Table 7.

**Table 7. Usual Dosing Regimens for the Thiazide Diuretics<sup>1-5</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Chlorothiazide	<p><u>Edema:</u> Injection, suspension, tablet: 0.5 to 1 g once or twice daily, often administered on alternate days or on three to five days each week</p> <p><u>Hypertension:</u> Injection, suspension, tablet: initial, 0.5 or 1 g/day as a single dose or in divided dose(s); maintenance, adjust according to blood pressure response, some patients may require up to 2 g/day in divided doses</p>	<p><u>Diuresis and hypertension:</u> Suspension, tablet: the usual pediatric dosage is 5 to 10 mg per pound (10 to 20 mg/kg) per day in single or two divided doses, not to exceed 375 mg per day in infants up to 2 years of age or 1 g per day in children 2 to 12 years of age. In infants less than 6 months of age, doses up to 15 mg per pound (30 mg/kg) per day in two divided doses may be required</p>	<p>Injection: 500 mg</p> <p>Suspension 250 mg/5 mL</p> <p>Tablet: 250 mg 500 mg</p>
HCTZ	<p><u>Edema:</u> Capsule, tablet: maintenance, 25 to 100 mg/day in a single or divided dose(s)</p> <p><u>Hypertension:</u> Capsule, tablet: initial, 12.5 to 25 mg once daily; maintenance, 50 to 100 mg daily in a single or divided dose(s)</p>	<p><u>Diuresis and hypertension:</u> Tablet: the usual pediatric dosage is 0.5 to 1 mg per pound (1 to 2 mg/kg) per day in single or two divided doses, not to exceed 37.5 mg per day in infants up to 2 years of age or 100 mg per day in children 2 to 12 years of age. In infants less than 6 months of age, doses up to 1.5 mg per pound (3 mg/kg) per day in two divided doses may be required.</p>	<p>Capsule: 12.5 mg</p> <p>Tablet: 12.5 mg 25 mg 50 mg</p>

HCTZ=hydrochlorothiazide

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the thiazide diuretics are summarized in Table 8.

**Table 8. Comparative Clinical Trials with the Thiazide Diuretics**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Edema</b>				
Rengo et al. <sup>19</sup> (1979)  HCTZ 150 mg QD  vs  amiloride 15 mg QD  vs  amiloride and HCTZ 15-150 mg QD (fixed-dose combination product)	RCT  Patients 35 to 60 years of age with liver cirrhosis and ascites or CHF	N=30  15 days	Primary: Body weight, 24 hour diuresis, serum sodium, serum potassium, sodium and potassium urinary loss  Secondary: Not reported	Primary: All treatment groups had a significant reduction in body weight from baseline (P<0.001 for all). Amiloride and HCTZ-treated patients achieved a significantly greater reduction compared to amiloride-treated patients (P<0.001).  All treatment groups significantly differed from baseline in 24 hour diuresis (P<0.01). Amiloride and HCTZ- and HCTZ-treated patients achieved greater diuresis compared to amiloride-treated patients (P<0.001 for both).  Serum sodium was reduced from baseline in all treatment groups. HCTZ-treated patients had a significantly greater reduction than amiloride- (P<0.01) and amiloride and HCTZ-treated patients (P<0.001). Sodium urinary loss was seen with all treatments at day two, amiloride and HCTZ therapy had maintained the loss at day five (P<0.001 for both).  Serum potassium decreased in HCTZ-treated patients but increased in amiloride- and amiloride and HCTZ-treated patients. HCTZ-treated patients had a marked increase in potassium urinary loss (P<0.001).  Secondary: Not reported
Cheitlin et al. <sup>20</sup> (1991)  Amiloride 5 or 10 mg QD for 7 days, followed by placebo plus HCTZ 50 or 100	DB, PC, RCT, XO  Patients with a history CHF and ≥1 episode of pulmonary edema (NYHA class 2 to 3) who were not	N=11  21 days	Primary: Hemodynamic changes at rest and exercise  Secondary: Not reported	Primary: At rest, there were no significant differences between placebo- and amiloride-treated patients in right atrial pressure, pulmonary atrial pressure, heart rate, pulmonary artery wedge pressure, systemic arterial pressure, right ventricular stroke work index, left ventricular stroke work index, systemic vascular resistance, cardiac index or stroke volume index (P values not reported).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD for 14 days</p> <p>vs</p> <p>placebo for 14 days, followed by amiloride 5 or 10 mg plus HCTZ 50 or 100 mg QD for the next 7 days</p>	<p>previously treated</p>			<p>During exercise, there were significant differences between placebo- and amiloride-treated patients at the 50-watt stage in right atrial pressure (15.0 vs 10.5 mm Hg), pulmonary artery wedge pressure (28.6 vs 22.1 mm Hg), pulmonary artery diastolic pressure (32.2 vs 21.6 mm Hg), mean pulmonary artery pressure (44.4 vs 38.9 mm Hg), left ventricular stroke work index (69.5 vs 77.9 g-m/m<sup>2</sup>) and stroke volume index (44.9 vs 46.2 cc/beat/m<sup>2</sup>), respectively (P values not reported).</p> <p>There were no significant differences between placebo and amiloride therapy during exercise in right ventricular stroke work index, heart rate, aortic pressure, cardiac index and total systemic vascular resistance (P values not reported).</p> <p>Secondary: Not reported</p>
<p>Kohvakka<sup>21</sup> (1988)</p> <p>HCTZ 50 mg BID</p> <p>vs</p> <p>amiloride 5 mg BID plus HCTZ 50 mg BID</p> <p>vs</p> <p>triamterene 75 mg BID plus HCTZ 50 mg BID</p> <p>vs</p> <p>KCl 1,000 mg BID plus HCTZ 50 mg BID</p>	<p>RCT, XO</p> <p>Patients 41 to 69 years of age with CHF (NYHA class 2 to 3) who developed persistent hypokalemia on HCTZ alone</p>	<p>N=25</p> <p>5 months</p>	<p>Primary: Changes in weight, blood pressure, serum sodium, serum potassium and total body potassium</p> <p>Secondary: Percentage with hypokalemia, median days until hypokalemia detection, serum magnesium</p>	<p>Primary: Weight loss was significant in amiloride plus HCTZ- and triamterene plus HCTZ-treated patients (P=0.05 for both), but not in KCl plus HCTZ-treated patients (P value not reported), compared to HCTZ-treated patients.</p> <p>No significant changes in blood pressure were observed (P values not reported).</p> <p>No differences in serum sodium were observed in amiloride plus HCTZ- or triamterene plus HCTZ-treated patients (P values not reported). Serum sodium levels were slightly higher in KCl plus HCTZ-patients compared to HCTZ-treated patients (P=0.01).</p> <p>Serum potassium was found to be significantly higher in all combination treated-patients compared to HCTZ-treated patients (P=0.01 for all comparisons). Total body potassium was significantly higher in amiloride plus HCTZ- and triamterene plus HCTZ-treated patients (P=0.05 for both), but not in KCl plus HCTZ-treated patients (P value not reported), compared to HCTZ-treated patients.</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The percentages of patients that became hypokalemic were 39, 52 and 52% in amiloride plus HCTZ-, triamterene plus HCTZ- and KCl plus HCTZ-treated patients (P values not reported).</p> <p>The median days until hypokalemia detection were 114.0, 75.0 and 51.5 for amiloride plus HCTZ-, triamterene plus HCTZ- and KCl plus HCTZ-treated patients (P values not reported).</p> <p>Serum magnesium was maintained at a significantly higher rate in amiloride plus HCTZ- and triamterene plus HCTZ- patients compared to KCl plus HCTZ-treated patients (P values not reported).</p>
<p>Faris et al.<sup>22</sup> (2006)</p> <p>Thiazide diuretics (chlorothiazide), loop diuretics (furosemide, bumetanide), or potassium-sparing diuretics (amiloride, triamterene)</p> <p>vs</p> <p>placebo or active control (ACE inhibitors, digoxin)</p>	<p>MA (14 trials)</p> <p>Adult patients with chronic heart failure</p>	<p>N=525</p> <p>2 to 52 weeks</p>	<p>Primary: Mortality</p> <p>Secondary: Effect of diuretic withdrawal on worsening of heart failure and exercise capacity</p>	<p>Primary: Pooled data from three PC trials (n=202) reporting on mortality revealed that mortality was lower for diuretic-treated patients compared to placebo-treated patients (2.7 vs 10.9%, respectively; OR, 0.24; 95% CI, 0.07 to 0.83; P=0.02). The difference represents an absolute risk reduction of 8% in mortality in diuretic-treated patients (NNT, 12.5).</p> <p>Secondary: Pooled data from two PC trials (n=169) reporting on the effect of diuretics on worsening heart failure revealed lower admission rates for worsening heart failure in diuretic-treated patients compared to placebo-treated patients (OR, 0.07; 95% CI, 0.01 to 0.52; P=0.01).</p> <p>Pooled data from two parallel RCTs (n=43) reporting on the effect of diuretics on exercise capacity revealed that diuretic therapy improved exercise capacity compared to active control (WMD, 0.74; 95% CI, 0.37 to 1.11; P&lt;0.0001). Pooled data from two XO RCTs (n=48) revealed similar results (WMD, 0.67; 95% CI, 0.02 to 1.31; P=0.04). In total (n=91), diuretic therapy improved exercise capacity in patients with chronic heart failure (WMD, 0.72; 95% CI, 0.40 to 1.04; P&lt;0.0001).</p>
<b>Hypertension</b>				
<p>Hua et al.<sup>23</sup> (1976)</p> <p>Chlorothiazide up to 5 g BID</p>	<p>XO</p> <p>Patients with HTN</p>	<p>N=20</p> <p>Duration not specified</p>	<p>Primary: Blood pressure, serum potassium</p> <p>Secondary:</p>	<p>Primary: Blood pressures on metolazone tended to be lower than on chlorothiazide, but the difference was not statistically significant.</p> <p>Both agents significantly lowered serum potassium concentrations and</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs metolazone 5 mg QD			Not reported	total body potassium to a similar degree. However, the serum potassium did not fall below the normal range in any patient and no potassium supplements were required.  Secondary: Not reported
Carter et al. <sup>24</sup> (2004)  HCTZ 12.5 to 450 mg/day  vs  chlorthalidone 12.5 to 600 mg/day	MA  Included trials which evaluate the pharmacokinetic and blood pressure lowering effects of chlorthalidone and HCTZ	N=200  Duration varied per study	Primary: Blood pressure  Secondary: Serum potassium	Primary: In a dose equivalence study comparing HCTZ 100 mg QD to chlorthalidone 50 mg QD, blood pressure (SBP/DBP) reduced by 18/8 and 25/10 mm Hg compared to baseline, respectively.  In another study comparing HCTZ 25 mg and triamterene 50 mg QD, HCTZ 50 mg and triamterene 100 mg QD, and chlorthalidone 50 mg QD, the blood pressure reduction was 15/8, 18/12, and 25/16 mm Hg, respectively.  One other dose equivalence study comparing HCTZ 50 mg BID and chlorthalidone 50 mg QD, blood pressure reduction was 22/16 and 18/15 mm Hg, respectively.  All available studies were inspected and it was concluded that HCTZ 50 mg is approximately equivalent to chlorthalidone 25 to 37 mg. Furthermore, it was suggested that chlorthalidone doses should generally be approximately 50% to 75% of the typical HCTZ dose.  Secondary: In a study comparing HCTZ 100 mg QD and chlorthalidone 50 mg QD, potassium increased slightly with chlorthalidone (0.02 mEq/L) and decreased significantly with HCTZ (0.22 mEq/L; P=0.009).  However, in another study comparing HCTZ 50 mg BID and chlorthalidone 50 mg QD, serum potassium decreased by 0.38 mEq/L with HCTZ and by 0.03 mEq/L with chlorthalidone. The difference was not statistically significant (P<0.07).
Ernst et al. <sup>25</sup> (2006)	RCT, SB, XO  Men and women	N=30  8 weeks plus 4	Primary: Comparison of the change in 24-hour	Primary: At week eight, there was a greater reduction in 24-hr mean SBP with chlorthalidone 25 mg/day compared to HCTZ 50 mg/day compared to

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 25 mg in the morning</p> <p>vs</p> <p>chlorthalidone 12.5 mg in the morning</p> <p>At week 4, both HCTZ and chlorthalidone were titrated to 50 mg in the morning and 25 mg in the morning, respectively for the remainder of the trial.</p>	<p>aged 18 to 79 years with pre-HTN or a new or established diagnosis of HTN (stage 1 or 2), not receiving antihypertensive medications, and had an average office blood pressure value in the last 6 months between 140 and 179 mm Hg systolic or 90 and 109 mm Hg diastolic</p>	<p>week washout period</p>	<p>mean SBP and DBP from baseline to week 8</p> <p>Secondary: Comparison of changes in mean SBP and mean DBP for office blood pressure at each visit, change in ambulatory daytime and nighttime mean SBP and DBP from baseline to week 8, development of hypokalemia</p>	<p>baseline (-12.4±1.8 vs -7.4±1.7 mm Hg, respectively; P=0.054).</p> <p>Secondary: There was a trend in favor of greater reduction in SBP with chlorthalidone than with HCTZ at each office visit. However, the difference was only statistically significant at week 2 (-15.7±2.2 vs -4.5±2.1 mm Hg, respectively; P=0.001).</p> <p>Although mean reductions in DBP was also greater with chlorthalidone compared to HCTZ at each study visit, the differences were not statistically significant at any visit (P&gt;0.89 for all).</p> <p>The reduction in SBP during nighttime hours was -13.5±1.9 mm Hg for chlorthalidone and -6.4±1.7 mm Hg for HCTZ (P=0.009). The reduction in daytime mean SBP between both groups was not significantly different (-11.4±2.0 vs -8.1±1.9 mm Hg, respectively; P=0.230).</p> <p>Changes in serum potassium were similar between treatment groups (P=0.76). The incidence of hypokalemia was 50% in patients taking HCTZ and 46% in patients taking chlorthalidone (P=0.682).</p>
<p>Finnerty et al.<sup>26</sup> (1980)</p> <p>HCTZ 50 mg plus reserpine 0.125 mg</p> <p>vs</p> <p>chlorthalidone 50 mg plus reserpine 0.25 mg</p>	<p>DB</p> <p>Patients with essential HTN unresponsive to diet control and diuretic therapy</p>	<p>N=57</p> <p>6 weeks</p>	<p>Primary: The change in mean DBP from baseline</p> <p>Secondary: Incidence of frequent or severe side effects</p>	<p>Primary: The chlorthalidone plus reserpine group had a mean decrease in DBP of 17.0 mm Hg at study endpoint compared with a mean decrease of 18.6 mm Hg in the HCTZ plus reserpine group.</p> <p>At study completion both treatment groups achieved diastolic control of at least 5 mm Hg below the targeted diastolic goal of 90 mm Hg.</p> <p>Secondary: There were no reports of frequent or severe side effects in either treatment group.</p>
<p>Bakris et al.<sup>27</sup> (2012)</p> <p>Azilsartan medoxomil and chlorthalidone</p>	<p>DB, RCT</p> <p>Patients aged ≥18 years with stage 2 primary HTN</p>	<p>N=609</p> <p>10 weeks (after 2 week placebo run-in)</p>	<p>Primary: Change in trough, seated clinic systolic blood pressure at weeks 6 and 10</p>	<p>Primary: Change in SBP at week six demonstrated a mean difference of -5.6 mm Hg (95% CI, -8.3 to -2.9; P&lt;0.001) in favor of the chlorthalidone group. Fewer patients in the chlorthalidone group required titration to a higher dose of diuretic (P&lt;0.001). At the end of week 10, a greater mean SBP reduction was maintained in the chlorthalidone group compared to the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(single pill)</p> <p>vs</p> <p>azilsartan medoxomil and HCTZ (co-administered)</p> <p>Treatments were titrated to a target of &lt;140/90 mm Hg (or &lt;130/80 mm Hg if diabetes of chronic kidney disease)</p>			<p>Secondary: Change from baseline in clinic DBP and 24-hour mean systolic and diastolic blood pressures by ambulatory blood pressure monitoring</p>	<p>HCTZ group (-5.0 mm Hg; 95% CI, -7.5 to -2.5; P&lt;0.001).</p> <p>Secondary: The chlorthalidone group demonstrated a significantly greater reduction in 24-hour mean SBP at weeks six and 10. For both clinica and 24-hour mean DBP, greater blood pressure reduction was observed in the chlorthalidone group compared to the HCTZ group at both study points.</p>
<p>Valmin et al.<sup>28</sup> (1975)</p> <p>HCTZ 12.5 mg BID</p> <p>vs</p> <p>furosemide 12.5, 25, or 40 mg BID</p> <p>vs</p> <p>placebo</p>	<p>DB, RCT, XO, 5 experimental periods each of 4 weeks</p> <p>Men and women with essential HTN</p>	<p>N=34</p> <p>20 weeks</p>	<p>Primary: Blood pressure, urinary output, serum electrolytes, safety and tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: When compared to placebo, there was a significant reduction of blood pressure with HCTZ 12.5 mg BID and furosemide 12.5 mg BID (P&lt;0.05).</p> <p>Paired comparison showed that HCTZ 12.5 mg BID and furosemide 25 and 40 mg BID had a similar hypotensive effect, irrespective of the initial blood pressure (P&gt;0.10).</p> <p>When compared to placebo, the urinary output increased significantly with furosemide 12.5, 25, or 40 mg BID (P&lt;0.05, P&lt;0.01 and P&lt;0.001, respectively) but not with the HCTZ group (P&gt;0.10).</p> <p>Sodium level did not alter during the various treatment periods when compared with the placebo period, or between the individual treatment periods (P&gt;0.10).</p> <p>Potassium level fell significantly during the HCTZ period (P&lt;0.001) and furosemide 25 mg and 40 mg BID period (P&lt;0.01 and P&lt;0.001, respectively). Potassium level was not significantly affected with furosemide 12.5 mg BID (P&gt;0.10).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Araoye et al. <sup>29</sup> (1978)  HCTZ 50 mg BID  vs  furosemide 40 mg BID	DB, XO  Patients with HTN	N=not specified  3 months	Primary: Blood Pressure  Secondary: Not reported	Primary: Furosemide and HCTZ significantly reduced blood pressure. The decrease in blood pressure was consistently greater in the HCTZ group than with furosemide; however the difference was significant in regards to SBP only.  Secondary: Not reported
Madkour et al. <sup>30</sup> (1996)  HCTZ 50 mg QD  vs  indapamide 2.5 mg QD	RCT  Patients aged 32 to 70 years with impaired renal function for 1 to 15 years and moderate HTN for 2 to 27 years, initial creatinine clearance between 32 and 80 mL/min/1.73 m <sup>2</sup> BSA	N=28  24 months	Primary: Blood pressure, changes in creatinine clearance  Secondary: Not reported	Primary: Blood pressure normalized in all patients taking either indapamide or HCTZ. There were no significant differences in SBP or DBP between groups.  At 24 months, creatinine clearance progressively increased from 58±4.4 to 72±4.4 mL/min/1.73 m <sup>2</sup> BSA in patients treated with indapamide (P<0.01).  Creatinine clearance progressively decreased from 65±3.0 to 53±3.0 mL/min/1.73 m <sup>2</sup> BSA in patients treated with HCTZ (P<0.01). Creatinine clearance significantly increased by 28.5±4.4% with indapamide and decreased by 17.4±3.0% with thiazide therapy (P<0.01).  Secondary: Not reported
Ames <sup>31</sup> (1996)  HCTZ ≤25 mg (or its equivalent in other thiazides) up to 112.5 mg QD  vs	MA (13 trials)  Patients with HTN	N=1,547  1 to 25 months	Primary: Comparison of the effects of thiazides and indapamide on blood lipids and blood pressure  Secondary: Not reported	Primary: The mean change from baseline was 1.4% for TC, 5.5% for HDL-C, and -0.5% for TG with indapamide. None of the differences were statistically significant.  Low-dose thiazide therapy did not decrease TC at any data point. The mean percent increase in TC was 3.8%, in HDL-C was 3.1%, and in TG was 10.8% with low-dose HCTZ. The increases in TC and TG from baseline was statistically significant (P<0.01).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
indapamide 2.5 mg QD				<p>The mean change in TC was 6.3%, in HDL-C was -0.5%, and in TGs was 19.5% for higher doses of HCTZ. Increases from baseline in TC and TG were statistically significant.</p> <p>SBP decreased more with higher doses of HCTZ than with low-dose thiazide therapy (P&lt;0.05). The effects of indapamide on systolic arterial pressure were intermediate between, and not statistically different from, either thiazide dose. Decreases in DBP did not differ among groups.</p> <p>Secondary: Not reported</p>
<p>Larochelle et al.<sup>32</sup> (1985)</p> <p>HCTZ 50 mg</p> <p>vs</p> <p>amiloride 5 mg/day and HCTZ 50 mg/day</p>	<p>DB, RCT</p> <p>Ambulant patients 18 to 70 years of age with essential HTN who after not being treated for ≥2 weeks prior to the trial had a supine DBP of 95 to 109 mm Hg and a serum potassium level of &gt;3.5 mmol/L</p>	<p>N=266</p> <p>8 weeks</p>	<p>Primary: Blood pressure, serum potassium concentration</p> <p>Secondary: Not reported</p>	<p>Primary: At eight weeks, there were no differences between the two treatments in the mean blood pressure reductions (P value not reported).</p> <p>During the eight weeks of treatment, the HCTZ plus amiloride-treated patients experienced a decrease in mean supine blood pressure (159/99 to 138/88 mm Hg) and serum potassium levels (4.23 to 3.91 mmol/L) (P values not reported).</p> <p>During the eight weeks of treatment, HCTZ-treated patients experienced a reduction in mean supine blood pressure (157/99 to 138/87 mm Hg) and serum potassium levels (4.16 to 3.69 mmol/L) (P values not reported).</p> <p>Hypokalemia occurred less frequently in HCTZ plus amiloride-treated patients compared to HCTZ-treated patients (14 and 29%, respectively; P=0.0026). However, the proportions of patients with a potassium level exceeding 4.5 mmol/L were similar (4.5 vs 3.9%, respectively; P value not reported).</p> <p>Secondary: Not reported</p>
<p>Salmela et al.<sup>33</sup> (1986)</p> <p>HCTZ 25 mg daily</p>	<p>DB, MC, PG, RCT</p> <p>Adult patients with mild to moderate HTN</p>	<p>N=40</p> <p>12 weeks</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary:</p>	<p>Primary: At the end of the first treatment period (four weeks), mean supine SBP and DBP was 161 and 91 mm Hg in amiloride plus HCTZ-treated patients (P&lt;0.01 and P&lt;0.001, respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs</p> <p>amiloride 2.5 mg/day and HCTZ 25 mg/day</p>			<p>Not reported</p>	<p>At the end of the first treatment period (four weeks), mean supine SBP and DBP was 165 and 96 mm Hg in HCTZ-treated patients (P&lt;0.01 for both).</p> <p>At the end of the second treatment period (eight weeks), mean supine SBP and DBP was 154 and 86 mm Hg in amiloride plus HCTZ-treated patients (P&lt;0.01 and P&lt;0.001).</p> <p>At the end of the second treatment period (eight weeks), mean supine SBP and DBP was 155 and 90 mm Hg in HCTZ-treated patients (P&lt;0.001 and P&lt;0.001).</p> <p>There were no significant differences in blood pressure reduction between the two treatments (P value not reported).</p> <p>Secondary: Not reported</p>
<p>Multicenter Diuretic Cooperative Study Group<sup>34</sup> (1981)</p> <p>HCTZ 50 mg QD</p> <p>vs</p> <p>amiloride and HCTZ 5-50 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>amiloride 5 mg QD</p>	<p>DB, MC, RCT</p> <p>Patients 21 to 69 years of age with mild to moderate essential HTN (supine DBP 95 to 115 mm Hg)</p>	<p>N=179</p> <p>12 weeks</p>	<p>Primary: Change from baseline in average supine SBP and DBP</p> <p>Secondary: Heart rate, body weight, serum potassium</p>	<p>Primary: Baseline vs 12 week average supine blood pressure was 153/101 vs 139/93 for amiloride-, 160/100 vs 137/90 for amiloride and HCTZ- and 154/101 vs 134/89 mm Hg for HCTZ-treated patients. Reductions in supine blood pressure were significant with all treatments (P&lt;0.01). The SBP reduction was significantly greater with amiloride and HCTZ-treated patients compared to amiloride-treated patients at all weeks and HCTZ-treated patients at four and eight weeks (P&lt;0.05, both).</p> <p>Secondary: No significant changes from baseline in heart rate were observed in amiloride and HCTZ-treated patients (P values not reported). An increase in heart rate of 3.3 bpm was observed in these patients (P&lt;0.05).</p> <p>Changes in body weight from baseline were -1.17 kg in amiloride and HCTZ-, -0.72 kg in HCTZ- and 0.045 kg in amiloride-treated patients (P&lt;0.05, for amiloride plus HCTZ only).</p> <p>Changes in serum potassium from baseline were 0.23 in amiloride- (P&lt;0.01), -0.38 in amiloride and HCTZ- (P&lt;0.01) and -0.59 mEq/L in HCTZ-treated patients (P&lt;0.01). The change in HCTZ-treated patients</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>was statistically greater than the change in the amiloride and HCTZ-treated patients (P&lt;0.05). Twenty three, two and zero percent of HCTZ-, amiloride and HCTZ- and amiloride-treated patients experienced hypokalemia.</p>
<p>Wray et al.<sup>35</sup> (2010)</p> <p>HCTZ 12.5 to 50 mg QD</p> <p>vs</p> <p>spironolactone 25 to 100 mg QD</p> <p>Patients also received potassium 0 to 40 mEq to maintain blinding.</p>	<p>DB, RCT</p> <p>Patients ≥60 years of age with stage 1 HTN</p>	<p>N=36</p> <p>6 months</p>	<p>Primary: Blood pressure, sympathetic nervous system activity</p> <p>Secondary: Not reported</p>	<p>Primary: Arterial blood pressure decreased significantly with spironolactone (SBP: 160 to 134 mm Hg and DBP: 77 to 68 mm Hg) and with HCTZ (SBP: 161 to 145 mm Hg and 78 to 73 mm Hg). There was no significant difference between the groups.</p> <p>Sympathetic nervous system activity was significantly reduced after spironolactone (plasma norepinephrine: 378 to 335 pg/mL; P=0.04; [<sup>3</sup>H]-norepinephrine release rate: 2.74 to 1.97 μg/min/m<sup>2</sup>; P=0.04), but not with HCTZ (plasma norepinephrine: 368 to 349 pg/mL; P=0.47; [<sup>3</sup>H]-norepinephrine release rate: 2.63 to 2.11 μg/min/m<sup>2</sup>; P=0.21).</p> <p>There were no instances of hyperkalemia, and no other adverse effects were reported.</p> <p>Secondary: Not reported</p>
<p>Nash et al.<sup>36</sup> (1977)</p> <p>HCTZ 50 mg BID</p> <p>vs</p> <p>spironolactone 50 mg BID</p> <p>vs</p> <p>spironolactone 100 mg BID</p> <p>vs</p>	<p>DB, RCT</p> <p>Male outpatients between the ages of 21 to 65 years, with essential HTN, DBP between 90 to 114 mm Hg</p>	<p>N=79</p> <p>12 weeks</p>	<p>Primary: Change in SBP, DBP, blood urea nitrogen, serum potassium, gynecomastia</p> <p>Secondary: Not reported</p>	<p>Primary: At week 12, all study groups exhibited significant reductions in SBP and DBP from baseline (P&lt;0.05).</p> <p>At week 12, all three spironolactone monotherapy groups exhibited statistically significant increases in blood urea nitrogen from baseline (P&lt;0.05).</p> <p>At week 12, the HCTZ monotherapy group was associated with a statistically significant decrease in serum potassium levels (P&lt;0.001).</p> <p>At week 12, all three spironolactone monotherapy groups exhibited statistically significant increases in serum potassium levels from baseline (P&lt;0.05).</p> <p>At week 12, the spironolactone and HCTZ combination group was not</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>spironolactone 200 mg BID</p> <p>vs</p> <p>spironolactone and HCTZ 25-25 mg BID (fixed-dose combination product)</p>				<p>associated with statistically significant increases in serum potassium levels from baseline.</p> <p>A dose-related risk of gynecomastia was observed in the spironolactone-treated patients. Among patients treated with spironolactone 50, 100, or 200 mg BID; 5.5, 11.8, and 40% reported gynecomastia symptoms. Of the patients randomized to spironolactone and HCTZ combination product, 7.7% reported gynecomastia symptoms.</p> <p>Secondary: Not reported</p>
<p>Schrijver et al.<sup>37</sup> (1979)</p> <p>Spironolactone 50 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IA)</p> <p>vs</p> <p>spironolactone 50 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IB)</p> <p>vs</p>	<p>DB</p> <p>Patients, between 24 to 63 years of age, with DBP between 90 to 114 mm Hg</p>	<p>N=49</p> <p>20 weeks (4-week placebo run-in, 8-week single drug therapy, 4-week two-drug therapy, 4-week recovery)</p>	<p>Primary: Change in MABP, serum potassium, uric acid level, blood glucose, blood urea nitrogen, creatinine, plasma renin activity, aldosterone, side effects</p> <p>Secondary: Not reported</p>	<p>Primary: Following eight weeks of therapy with a single drug, all study groups exhibited a statistically significant reduction in MABP from baseline (P&lt;0.01). There were no significant differences in MABP reduction among the study groups.</p> <p>The addition of a second drug to the antihypertensive regimen was not associated with a significant improvement in MABP. At the end of the two-drug treatment period, there were no differences in MABP among any of the study groups.</p> <p>Spironolactone therapy was associated with a significant decrease in serum potassium concentration from baseline (P&lt;0.001).</p> <p>Spironolactone regimens were not associated with a significant change in potassium levels from baseline.</p> <p>Following eight weeks of therapy with a single drug, HCTZ-treated patients experienced a statistically significant increase in uric acid from baseline (P&lt;0.001). Groups IIA and IIB also experienced a significant but smaller increase in uric acid level from baseline (P&lt;0.05) with no change in groups I and IV.</p> <p>During the single-drug treatment phase, patients randomized to group I experienced a significant increase in blood glucose from baseline (P&lt;0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>spironolactone 100 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IIA)</p> <p>vs</p> <p>spironolactone 100 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IIB)</p> <p>vs</p> <p>spironolactone 200 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IIIA)</p> <p>vs</p> <p>spironolactone 200</p>				<p>During the single-drug treatment phase, all patients except those randomized to group I experienced a significant increase in blood urea nitrogen from baseline (P&lt;0.05).</p> <p>During the single-drug treatment phase, patients randomized to groups I and II experienced a significant increase in serum creatinine from baseline (P&lt;0.05).</p> <p>During the single-drug treatment phase, all treatment groups experienced a significant increase in plasma renin activity from baseline (P&lt;0.01). The addition of HCTZ in the two-drug study phase was associated with a rise in plasma renin activity in all study groups (P&lt;0.05).</p> <p>All treatment groups experienced a significant increase in plasma aldosterone from baseline (P&lt;0.05).</p> <p>Gynecomastia was reported only by patients randomized to the higher-dose spironolactone groups.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IIIB)</p> <p>vs</p> <p>HCTZ 50 mg BID for 8 weeks (single drug phase), with the addition of a placebo for subsequent 4 weeks (group IVA)</p> <p>vs</p> <p>HCTZ 50 mg BID for 8 weeks (single drug phase), subsequently HCTZ 50 mg BID was added to the regimen for an additional 4 weeks (group IVB)</p>				
<p>Johnson et al.<sup>38</sup> (2009)</p> <p>HCTZ 12.5 to 25 mg QD for 9</p>	<p>RCT</p> <p>Patients 17 to 65 years of age mild to moderate essential</p>	<p>N=368</p> <p>15 to 18 weeks</p>	<p>Primary: Blood pressure lowering effect of drug initiation order: the addition</p>	<p>Primary: When analyzed by order of initiation of the two drugs, the response to HCTZ and atenolol was greater overall than that seen for atenolol and HCTZ (P=0.0007 and P&lt;0.0001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>weeks, followed by HCTZ 12.5 to 25 mg QD and atenolol 50 to 100 mg QD for 9 weeks</p> <p>vs</p> <p>atenolol 50 to 100 mg QD for 9 weeks, followed by atenolol 50 to 100 mg QD and HCTZ 12.5 to 25 mg QD for 9 weeks</p>	<p>HTN</p>		<p>of a <math>\beta</math>-blocker to a thiazide versus the addition of a thiazide to a <math>\beta</math>-blocker</p> <p>Secondary: Not reported</p>	<p>This study suggests that initiation of HCTZ followed by atenolol results in greater blood pressure lowering as compared with initiation in the reverse order, with differences that are potentially clinically important.</p> <p>Secondary: Not reported</p>
<p>Dahlöf et al.<sup>39</sup> (1991) Hypertension (STOP)</p> <p>Atenolol 50 mg QD, HCTZ 25 mg QD plus amiloride 2.5 mg QD, metoprolol 100 mg QD, or pindolol 5 mg QD</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, RCT</p> <p>Swedish men and women 70 to 84 years old with treated or untreated essential HTN defined as SBP <math>\geq</math>180 mm Hg with a DBP of <math>\geq</math>90 mm Hg, or DBP <math>&gt;</math>105 mm Hg irrespective of the SBP measured on 3 separate occasions during a 1-month placebo run-in phase in previously untreated patients</p>	<p>N=1,627</p> <p>25 months</p>	<p>Primary: Frequency of stroke, MI, and other cardiovascular death</p> <p>Secondary: Not reported</p>	<p>Primary: The active treatments significantly reduced the number of all primary endpoints (94 vs 58; RR, 0.60; 95% CI, 0.43 to 0.85; P=0.0031), frequency of stroke (53 vs 29; RR, 0.53; 95% CI, 0.33 to 0.86; P=0.0081) and frequency of other cardiovascular deaths (13 vs 4; RR, 0.30; 95% CI, 0.07 to 0.97) compared to placebo.</p> <p>There was not a statistically significant decrease observed in the rate of MI between the active treatments and placebo (28 vs 25; RR, 0.87; 95% CI, 0.49 to 1.56).</p> <p>Secondary: Not reported</p>
<p>Frishman et al.<sup>40</sup></p>	<p>DB, MC, PC, RCT</p>	<p>N=512</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1994)</p> <p>HCTZ 6.25 or 25 mg QD</p> <p>vs</p> <p>bisoprolol 2, 5, 10, or 40 mg QD</p> <p>vs</p> <p>bisoprolol plus HCTZ, all possible combinations</p>	<p>Patients 21 years and older with mild to moderate essential HTN whose weight was 35% of the ideal for height and frame and mean sitting DBP was stable and between 95 to 115 mm Hg</p>	<p>12 weeks</p>	<p>Changes in DBP and SBP</p> <p>Secondary: Not reported</p>	<p>All treatment groups (all doses) of bisoprolol, HCTZ and the combination of bisoprolol and HCTZ significantly reduced sitting DBP from baseline (P&lt;0.01).</p> <p>The reduction in blood pressure was significantly greater as the doses of the bisoprolol, HCTZ and the combination of bisoprolol-HCTZ were increased (P&lt;0.05).</p> <p>The combination bisoprolol and HCTZ significantly reduced sitting DBP compared to the separate agents as monotherapy (P&lt;0.01).</p> <p>With higher doses of HCTZ, there was a significantly higher incidence of hypokalemia, defined as potassium &lt;3.5 mmol/L (P&lt;0.01). Incidence of hyperuricemia also significantly increased with the increase in HCTZ dose (P&lt;0.01). Adverse events associated with hypokalemia and hyperuricemia were not reported.</p> <p>As the dose of bisoprolol was increased, the frequency and severity of adverse events reported significantly increased (P&lt;0.05). Adverse events reported included asthenia, diarrhea, dyspepsia and somnolence, but severity of effects was not reported.</p> <p>Secondary: Not reported</p>
<p>Frishman et al.<sup>41</sup> (1995)</p> <p>HCTZ 25 mg QD</p> <p>vs</p> <p>bisoprolol 5 mg QD</p> <p>vs</p> <p>bisoprolol and</p>	<p>DB, MC, PC, PG, RCT</p> <p>Patients ≥21 years with mild to moderate (stage II or II) systemic HTN whose body weight was not &gt;10% below or 35% above the ideal weight for height and frame, and were off all</p>	<p>N=547</p> <p>10 weeks</p>	<p>Primary: Changes in blood pressure and adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: All active treatment groups significantly reduced sitting DBP and SBP from baseline compared to placebo (P&lt;0.01).</p> <p>Addition of HCTZ 6.25 mg contributed significantly to the blood pressure lowering effects of bisoprolol 5 mg.</p> <p>The combination bisoprolol and HCTZ 5-6.25 mg produced a significantly greater reduction in mean sitting DBP from baseline (-12.6±0.5 mm Hg) compared to bisoprolol 5 mg alone (-10.5±0.5 mm Hg; P=0.02) and HCTZ 25 mg alone (-8.5±0.5 mm Hg; P&lt;0.01). Bisoprolol 5 mg monotherapy was significantly better a reducing DBP compared to HCTZ 25 mg alone (P=0.03).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 5-6.25 mg QD (fixed-dose combination product)</p> <p>vs</p> <p>placebo</p>	<p>antihypertensive medications before study entry and sitting DBP was 95 to 115 mm Hg on 3 consecutive weekly visits</p>			<p>The combination bisoprolol and HCTZ 5-6.25 mg produced a significantly greater reduction in mean sitting SBP from baseline (-15.8 mm Hg) compared to bisoprolol 5 mg alone (-10 mm Hg; P&lt;0.01) and HCTZ 25 mg alone (-15.8 mm Hg; P&lt;0.01). There was not a significant difference in mean reduction between bisoprolol 5 mg alone and HCTZ 25 mg alone.</p> <p>Bisoprolol and HCTZ 5-6.25 mg in combination had a 73% response rate compared to 61% for the bisoprolol group and 47% for the HCTZ group.</p> <p>Bisoprolol and HCTZ 5-6.25 mg in combination was found to be significantly more effective compared to bisoprolol 5 mg or HCTZ 25 mg in all subgroups of patients regardless of age, race, gender, or smoking history (P&gt;0.05 for all comparisons).</p> <p>Bisoprolol and HCTZ 5-6.25 mg in combination did not have an increase in frequency or severity of adverse events. The adverse events were comparable to that in the placebo group and frequency among groups was not significant. The most common adverse events reported were headache, dizziness, fatigue, and cough.</p> <p>Significantly greater number patients in the HCTZ 25 mg group (6.5%) experienced hypokalemia (potassium &lt;3.4 mEq/L) compared to the bisoprolol 5 mg group (0.7%; P&lt;0.01), the bisoprolol and HCTZ combination group (0.7%; P&lt;0.01), and placebo (0%; P&lt;0.01).</p> <p>Hyperglycemia occurred in 7.4% of patients in the HCTZ 25 mg group, which was significantly higher than in the placebo group (5.2%; P=0.03). Also, the incidence of hyperuricemia (uric acid &gt;7.5 mg/dL) was significantly higher in the HCTZ 25 mg group (24.4%) compared to placebo (2.7%; P&lt;0.01).</p> <p>Secondary: Not reported</p>
<p>Dafgard et al.<sup>42</sup> (1981)</p>	<p>DB, MC, RCT</p> <p>Patients with</p>	<p>N=31</p> <p>32 weeks</p>	<p>Primary: Blood pressure, heart rate, adverse</p>	<p>Primary: After the eight week run-in period with HCTZ 25 mg alone, the mean supine blood pressure was significantly reduced from 183/110 to 172/103</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 50 mg QD in the morning</p> <p>vs</p> <p>HCTZ 25 mg QD in the morning</p> <p>vs</p> <p>metoprolol and HCTZ 200-25 mg QD in the morning (fixed-dose combination product)</p>	<p>essential HTN (WHO stages I or II) not adequately controlled (<math>\geq 160/95</math> mm Hg) on HCTZ 25 mg/day</p>		<p>events, laboratory values</p> <p>Secondary: Not reported</p>	<p>mm Hg (<math>P &lt; 0.01/P &lt; 0.01</math>). The increased dose of HCTZ 50 mg following the run-in period did not further significantly reduce the mean blood pressure (165/104 mm Hg).</p> <p>A small but statistically significant reduction in supine heart rate was seen when the HCTZ dose was increased from 25 to 50 mg (82 down to 78 bpm; <math>P &lt; 0.05</math>).</p> <p>After the 12 week double-blind period, the mean supine blood pressure was 153/98 mm Hg in the HCTZ 50 mg group. After the 12 week follow-up period, there was not any additional decrease in blood pressure (153/97 mm Hg).</p> <p>Fixed-dose combination product of metoprolol and HCTZ produced a significant reduction in supine blood pressure after 12 weeks of therapy from 172/105 mm Hg on HCTZ 25 mg alone to 154/97 mm Hg on the combination therapy (<math>P &lt; 0.001/P &lt; 0.01</math>). Similar results were found with the standing blood pressure reductions, from 165/108 to 147/97 mm Hg (<math>P &lt; 0.001/P &lt; 0.001</math>).</p> <p>After the eight week run-in period, the supine heart rate was 80 bpm which decreased to 64 bpm with the metoprolol and HCTZ fixed-dose combination (<math>P &lt; 0.001</math>). The values for standing heart rate demonstrated similar significant reductions (85 to 66 bpm; <math>P &lt; 0.001</math>).</p> <p>After the additional 12 week follow-up, the patients in the metoprolol and HCTZ fixed-dose combination group did not demonstrate a significant further reduction in heart rate or blood pressure in any position.</p> <p>Both agents were tolerated and the most common adverse events reported included insomnia, headache, tiredness, and shortness of breath. The majority of events were mild, few were moderate, and none were severe. The only significant changes in laboratory values occurred with the HCTZ 25 and 50 mg groups, where an increase in serum uric acid was observed from 0.30 to 0.34 and 0.35 mmol/L, respectively (<math>P &lt; 0.01</math> and <math>P &lt; 0.05</math>; respectively).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
<p>Smilde et al.<sup>43</sup> (1983)</p> <p>Metoprolol 400 mg QD in the morning for 5 weeks, followed by metoprolol and HCTZ 200-25 mg QD in the morning (fixed-dose combination product) (group 1)</p> <p>vs</p> <p>metoprolol and HCTZ 200-25 mg QAM for 5 weeks (fixed-dose combination product), followed by metoprolol 400 mg QD in the morning for 5 weeks (group 2)</p>	<p>DB, PG, RCT, XO</p> <p>Patients &lt;65 years with essential HTN (supine DBP <math>\geq</math>95 mm Hg) not controlled on metoprolol 200 mg alone</p>	<p>N=37</p> <p>15 weeks</p>	<p>Primary: Changes in DBP, SBP, and heart rate</p> <p>Secondary: Not reported</p>	<p>Primary: Both group 1 and 2 significantly reduced DBP (P&lt;0.01) from baseline and the two groups were not significantly different from each other.</p> <p>The combination products significantly reduced SBP from baseline (P&lt;0.05, P&lt;0.01 depending on comparison)</p> <p>Group 2 significantly reduced heart rate at the end of the study compared to baseline (P&lt;0.05).</p> <p>Clinically relevant changes in laboratory parameters or mean body weight were not observed between the groups.</p> <p>Secondary: Not reported</p>
<p>Stevens et al.<sup>44</sup> (1982)</p> <p><u>Dose-finding phase:</u> propranolol and HCTZ 80-50, 160-50, 240-50, 320-50 mg/day in 2</p>	<p>DB, PG, RCT</p> <p>Patients with mild to moderate essential HTN (DBP 100 to 125 mm Hg)</p>	<p>N=158</p> <p>25 weeks</p>	<p>Primary: Mean changes of SBP and DB, heart rate, lab values</p> <p>Secondary: Not reported</p>	<p>Primary: After the 12 week dose finding-phase, 94% of patients had a decrease <math>\geq</math>10 mm Hg in DBP. The mean SBP and DBP reduced from 158.0 (<math>\pm</math>17.3)/105.6 (<math>\pm</math>6.0) mm Hg to 131.5 (<math>\pm</math>14.4)/86.4 (<math>\pm</math> 6.7) mm Hg (P&lt;0.001).</p> <p>After the 10 week portion of the study, there were significantly greater increases (P&lt;0.05) in mean SBP or DBP with propranolol and HCTZ alone vs the combination product of propranolol and HCTZ from the end</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>divided doses (fixed-dose combination product)</p> <p>vs</p> <p>propranolol 80, 160, 240, or 320 mg/day in 2 divided doses</p> <p><u>Double-blind phase:</u> HCTZ</p> <p>vs</p> <p>propranolol</p> <p>vs</p> <p>propranolol and HCTZ (fixed-dose combination product)</p>				<p>of the dose-finding to the last four biweekly visits to the mean of those visits, and to the last visit. The mean increases of SBP and DBP at the endpoint were: propranolol, 10.2/6.3 mm Hg; HCTZ 13.1/9.3 mm Hg; propranolol-HCTZ combination product 3/1.5 mm Hg.</p> <p>There was a significant decrease in heart rate as the dose of propranolol was increased though the trial (<math>P&gt;0.30</math>).</p> <p>The only lab value that showed a statistically significant change was serum chloride. The percent of patients that fell outside of the normal range were as follows: propranolol 6/36 (17%), HCTZ 14/37 (38%), and combination 4/28 (14%); <math>P&lt;0.05</math>.</p> <p>Secondary: Not reported</p>
<p>Borhani et al.<sup>45</sup> (1996) MIDAS</p> <p>HCTZ 12.5 to 25 mg QD</p> <p>vs</p> <p>isradipine 2.5 to 5 mg BID</p>	<p>DB, MC, positive-control, RCT</p> <p>Patients, average of 58.5 years old, with HTN</p>	<p>N=883</p> <p>3 years</p>	<p>Primary: Rate of progression of intimal-medial thickness in carotid arteries</p> <p>Secondary: Rate of cardiovascular events (MI, stroke, CHF, angina,</p>	<p>Primary: There was no difference in the rate of progression of intimal-medial thickness between the treatment groups (<math>P=0.68</math>).</p> <p>Secondary: The rate of cardiovascular events was greater in the isradipine group than in the HCTZ group (5.65 vs 3.17%; <math>P=0.07</math>).</p> <p>The rate of non-major cardiovascular events was greater in the isradipine group than in the HCTZ group (9.05 vs 5.22%; <math>P=0.02</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			sudden death), rate of non-major cardiovascular events and procedures (TIAs, dysrhythmia, aortic valve replacement, femoral popliteal bypass graft), blood pressure	There was a significant decrease in SBP in the HCTZ group as compared to isradipine (-19.5 vs -16.0 mm Hg; P=0.002).  There was no difference in change in DBP (both groups, -13.0 mm Hg).
Manyemba et al. <sup>46</sup> (1997)  HCTZ 25 mg QD plus reserpine 0.25 mg QD  vs  HCTZ 25 mg QD plus nifedipine SR 20 mg BID	OL, RCT, XO  African American patients aged 21 to 65 years with HTN (blood pressure >140/95 mm Hg) after 4 weeks of daily HCTZ therapy	N=32  10 weeks	Primary: The change in blood pressure from baseline to the end of each 4-week treatment period  Secondary: Not reported	Primary: Reserpine reduced SBP by 15.9 mm Hg (95% CI, 8.4 to 23.4) and DBP by 11.1 mm Hg (95% CI, 7.5 to 14.6).  Nifedipine SR reduced SBP by 18.9 mm Hg (95% CI, 12.1 to 25.7) and DBP by 9.6 mm Hg (95% CI, 7.2 to 12.0).  There was no significant difference between the two groups.  Secondary: Not reported
Jamerson et al. <sup>47</sup> (2007) ACCOMPLISH  HCTZ 12.5 to 25 mg QD and benazepril 20 to 40 mg QD  vs  benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD	DB, MC, RCT  Patients >60 years of age with HTN and at high risk of cardiovascular events	N=10,704  Analysis performed at 6 months (complete trial duration 5 years)	Primary: Changes in mean SBP from baseline to 6 months, blood pressure control rates (SBP/DBP <140/90 mm Hg or <130/89 mm Hg for patients with diabetes and chronic kidney disease)  Secondary: Not reported	Primary: At baseline, 97% of subjects were treated with antihypertensive medications at entry, but only 37% of participants had blood pressure control.  Mean blood pressure fell from 145/80 to 132/74 mm Hg after six months of treatment with either combination regimen (P<0.001).  The six month blood pressure control rate was 73% in the overall trial (78% in the United States), 43% in diabetics, and 40% in patients with renal disease. Of the patients uncontrolled, 61% were not on maximal medications.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Jamerson et al.<sup>48</sup> (2008) ACCOMPLISH</p> <p>HCTZ 12.5 to 25 mg QD and benazepril 20 to 40 mg QD</p> <p>vs</p> <p>benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Patients &gt;60 years of age with HTN and at high risk of cardiovascular events</p>	<p>N=11,506</p> <p>36 months (mean)</p>	<p>Primary: The composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for angina, resuscitation after sudden cardiac arrest, and coronary revascularization.</p> <p>Secondary: Death from cardiovascular causes, nonfatal MI, and nonfatal stroke</p>	<p>Primary: There were 552 primary-outcome events in the benazepril plus amlodipine group (9.6%) and 679 events in the benazepril plus HCTZ group (11.8%). The absolute risk reduction with benazepril plus amlodipine therapy was 2.2% and the relative risk reduction was 19.6% compared to benazepril plus HCTZ (HR, 0.80; 95% CI, 0.72 to 0.90; P&lt;0.001).</p> <p>Secondary: For the secondary end point of death from cardiovascular causes, nonfatal MI, and nonfatal stroke, there were 288 (5%) events in the benazepril plus amlodipine group compared to 364 (6.3%) events in the benazepril plus HCTZ group. The absolute risk reduction with benazepril plus amlodipine therapy was 1.3% and the RR reduction was 21.2% compared to benazepril plus HCTZ (HR, 0.79; 95% CI, 0.67 to 0.92; P=0.002).</p>
<p>Bakris et al.<sup>49</sup> (2013) ACCOMPLISH</p> <p>HCTZ 12.5 to 25 mg QD and benazepril 20 to 40 mg QD (B+H)</p> <p>vs</p> <p>benazepril 20 to 40 mg QD and amlodipine 5 to 10 mg QD (B+A)</p>	<p>Post hoc analysis</p> <p>Patients included in the ACCOMPLISH trial (&gt;60 years of age with HTN and at high risk of cardiovascular events) stratified by presence of known CAD at baseline</p>	<p>N=11,506</p> <p>36 months (mean)</p>	<p>Primary: The composite of death from cardiovascular causes, nonfatal MI, nonfatal stroke, hospitalization for angina, resuscitation after sudden cardiac arrest, and coronary revascularization.</p>	<p>Primary: Among the patients with CAD, 13% in the B+A group and 16% in the B+H group reached the primary end point, representing an absolute risk reduction of 3% and a hazard reduction of 18%. The difference in event rates of the composite primary end point between the B+A and B+H groups was significant (HR, 0.80; 95% CI, 0.69 to 0.92; P=0.0016).</p> <p>Among the patients without CAD, fewer patients in the B+A treatment arm (204 of 3,096) reached the primary end point compared with those in the B+H arm (251 of 3,095). The difference in event rates between the B+A and B+H groups was significant (HR, 0.81; 95% CI, 0.67 to 0.98; P=0.026).</p> <p>A comparison of patients with and without CAD event rates for the primary end points demonstrated that the patients with CAD had a greater</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Secondary: Death from cardiovascular causes, nonfatal MI, and nonfatal stroke	CV event rate than those without CAD (15 vs 7%; P<0.0001).  Secondary: The composite secondary end point of CV mortality, MI, and stroke occurred in 5.74% in the B+A group and 8% in the B+H group, resulting in an absolute risk reduction of 1.95% and a hazard reduction of 25% (HR, 0.73; 95% CI, 0.59 to 0.9; P=0.033). The rate of all-cause mortality differed significantly between the treatment arms (HR, 0.77; 95% CI, 0.6 to 0.99; P=0.042). Among the patients without CAD, the rates of CV mortality, MI, and stroke did not differ between the two arms (HR, 0.86; 95% CI, 0.68 to 1.08). The secondary end point events were lower in the group of patients without CAD.
Wing et al. <sup>50</sup> (2003) ANBP2  HCTZ  vs  enalapril  The choice of the specific agent and dose was made by the family practitioner.	MC, OL, PRO, RCT  Patients 65 to 84 years of age with average SBP while sitting of $\geq 160$ mm Hg or an average DBP of $\geq 90$ mm Hg (if the SBP was $\geq 140$ mm Hg)	N=6,083  4.1 years (median)	Primary: All cardiovascular events or death from any cause (both initial and subsequent fatal and nonfatal cardiovascular events)  Secondary: Not reported	Primary: By the end of the study, blood pressure had decreased to a similar extent in both groups (a decrease of 26/12 mm Hg).  There were 695 cardiovascular events or deaths from any cause in the ACE inhibitor group (56.1 per 1,000 patient-years; HR, 0.89; 95% CI, 0.79 to 1.0; P=0.05) compared to 736 in the diuretic group (59.8 per 1,000 patient-years).  The beneficial effects of ACE inhibitor treatment were more evident in male subjects (HR, 0.83; 95% CI, 0.71 to 0.97; P=0.02).  The rates of nonfatal cardiovascular events and MI decreased with ACE inhibitor treatment, whereas a similar number of strokes occurred in each group (although there were more fatal strokes in the ACE inhibitor group).  Secondary: Not reported
Poldermans et al. <sup>51</sup> (2007)  HCTZ 12.5 mg QD and lisinopril 10 to 20 mg	AC, DB, MC, PG, RCT  Males and females, ages 18 years and older with HTN (mean DBP $\geq 110$ )	N=130  6 weeks	Primary: Safety/adverse events, vital signs, hematology, biochemistry variables	Primary: Both treatments were well tolerated, 26 (40.6%) of patients receiving amlodipine and valsartan and 21 (31.8%) of patients receiving lisinopril and HCTZ reported an adverse events and most were not considered drug related.  Peripheral edema was reported more often in the amlodipine and valsartan

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs amlodipine 5 to 10 mg QD and valsartan 160 mg QD	mm Hg and <120 mm Hg)		Secondary: Efficacy (mean DBP, response rate, proportion of patients with mean DBP <90 mm Hg or a $\geq$ 10 mm Hg reduction from baseline)	group than the lisinopril and HCTZ group (7.7 vs 1.5%) and cough was reported less often in the amlodipine and valsartan group than the receiving lisinopril and hydrochlorothiazide group (1.6 vs 3.0%).  No difference was found between the treatments in changes in laboratory values or biochemistry variables.  Secondary: Both treatments led to a reduction in mean SBP and DBP (P<0.0001 for both from baseline) but were not significantly different from each other. Mean blood pressure for each group at study end: amlodipine and valsartan 135.0/83.6 mm Hg and lisinopril and HCTZ 138.7/85.2 mm Hg.  The response rate was similar among the groups (100 vs 95.5%; P value not significant).
Fogari et al. <sup>52</sup> (2007) CANDIA  HCTZ 12.5 mg QD and candesartan 16 mg  vs amlodipine 10 mg QD	DB, MC, RCT  Patients, 20 to 80 years old, with mild to moderate uncomplicated HTN not controlled on monotherapy with an antihypertensive (SBP <180 mg Hg and DBP 90 to 110 mg Hg)	N=203  8 weeks	Primary: Decrease in DBP  Secondary: Sitting SBP, reduction of the orthostatic blood pressure at least two minutes after standing, change in heart rate, percentage of patients normalized (DBP <90 mm Hg and SBP <140 mm Hg), percentage of responders (reduction in DBP $\geq$ 5 mm Hg)	Primary: There was no significant difference in the mean decrease in DBP between treatment groups; the difference in final DBP was -0.02 mm Hg (95% CI, -1.48 to 1.52 mm Hg; P=0.979).  Secondary: There was no significant difference between the groups at week eight for the following: sitting SBP (P=0.835), heart rate (P<0.500), orthostatic SBP (P=0.883), orthostatic DBP (P=0.264), percentage of patients normalized (P=10), percentage of responders (P=0.900).  The number of patients reporting an adverse event was greater in the amlodipine group (P=0.001).  The number of patients reporting an adverse drug-related event was greater in the amlodipine group (P<0.001).  Changes in blood chemistry and other secondary measurements were not significantly different between the treatment groups.
Neutel et al. <sup>53</sup> (2008)	AC, DB, RCT	N=538	Primary: Change in SBP	Primary: At week eight, there was a reduction in SBP of 27.1 mm Hg with



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
HCTZ 25 mg QD and irbesartan 300 mg vs irbesartan 300 mg QD vs HCTZ 25 mg QD	Patients with moderate HTN (seated SBP 160 to 179 mm Hg when DBP <110 mm Hg; or DBP 100 to 109 mm Hg when SBP <180 mm Hg)	12 weeks	after week 8  Secondary: Change from baseline in DBP at weeks 8 and 12, SBP at week 12, proportion of responders (SBP <140 mm Hg and DBP <90 m Hg) at weeks 8 and 12	irbesartan and HCTZ compared to 22.1 mm Hg with irbesartan monotherapy (P=0.0016) and 15.7 mm Hg with HCTZ (P<0.0001).  Secondary: At week eight, there was a reduction in DBP of 14.6 mm Hg with irbesartan and HCTZ compared to 11.6 mm Hg with irbesartan monotherapy (P=0.0013) and 7.3 mm Hg with HCTZ (P<0.0001).  A significantly greater percentage of patients reached a treatment goal of SBP <140 mm Hg and DBP <90 mm Hg by week eight with irbesartan and HCTZ (53.4%) compared with irbesartan (40.6%; P=0.0254) and HCTZ (20.2%; P<0.0001) alone.  Treatment was well tolerated in all three treatment groups with a slight increase in adverse events in the combination therapy group.
Salerno et al. <sup>54</sup> (2004)  HCTZ 12.5 mg QD and losartan 50 mg  vs  losartan 50 to 100 mg QD  Doses were titrated as needed to reach blood pressure goal (<90 mm Hg).	DB, RCT  Patients with severe HTN	N=585  6 weeks	Primary: Proportion of patients achieving goal blood pressure  Secondary: Adverse events	Primary: Almost twice as many patients achieved goal blood pressure at four weeks on losartan 50 mg and HCTZ 12.5 mg vs losartan 50 to 100 mg monotherapy (P=0.002).  Almost three times as many patients achieved goal blood pressure at six weeks with losartan and HCTZ vs losartan monotherapy (P<0.001).  Adverse experiences on losartan and HCTZ (43%) were significantly less than with losartan monotherapy (53%).
Minami et al. <sup>55</sup> (2007)  HCTZ 12.5 mg/day and losartan 50 mg/day	OL  Japanese outpatients with essential HTN treated for ≥2 months with either	N=15  12 months	Primary: Changes in blood pressure  Secondary: Not reported	Primary: In patients who had previously received candesartan, 24-hour blood pressure decreased significantly from 137/89 mm Hg to 126/81 mm Hg after three months (P<0.05/P<0.001) and to 123/81 mm Hg after 12 months (P<0.01/P<0.001) of treatment with losartan and HCTZ.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>vs candesartan 8 mg QD or amlodipine 5 mg QD</p>	<p>candesartan or amlodipine and 24-hour ambulatory blood pressure <math>\geq 135/80</math> mm Hg</p>			<p>In patients who had previously received amlodipine, 24-hour blood pressure decreased significantly from 137/81 to 125/75 mm Hg after three months (<math>P &lt; 0.05/P &lt; 0.05</math>) and to 124/77 mm Hg after 12 months (<math>P &lt; 0.05/P</math> value not significant) of treatment with losartan and HCTZ.</p> <p>There were significant decreases in SBP during the daytime, nighttime and early morning after 12 months in both groups.</p> <p>No adverse changes in the indices of glucose or lipid metabolism were observed in either group.</p> <p>Secondary: Not reported</p>
<p>Chrysant et al.<sup>56</sup> (2004)  HCTZ 12.5 to 25 mg QD and olmesartan 10 to 40 mg QD  vs  olmesartan 10 to 40 mg QD  vs  HCTZ 12.5 to 25 mg QD  vs  placebo</p>	<p>DB, RCT, factorial design  Patients with a baseline mean seated DBP of 110 to 115 mm Hg</p>	<p>N=502  8 weeks</p>	<p>Primary: Change in DBP at week 8  Secondary: Change in SBP at week 8</p>	<p>Primary: Olmesartan and HCTZ produced greater reductions in seated DBP at week eight than did monotherapy with either component. All olmesartan and HCTZ combinations significantly reduced DBP compared with placebo in a dose-dependent manner.</p> <p>Reductions in mean trough DBP were 8.2, 16.4, and 21.9 mm Hg with placebo, olmesartan 20 mg plus HCTZ 12.5 mg, and olmesartan 40 mg plus HCTZ 25 mg, respectively.</p> <p>Secondary: Olmesartan and HCTZ produced greater reductions in seated SBP at week eight than did monotherapy with either component. All olmesartan and HCTZ combinations significantly reduced DBP compared with placebo in a dose-dependent manner.</p> <p>Reductions in mean trough SBP were 3.3, 20.1, and 26.8 mm Hg with placebo, olmesartan 20 mg plus HCTZ 12.5 mg, and olmesartan 40 mg plus HCTZ 25 mg, respectively.</p> <p>All treatments were well tolerated.</p>
<p>White et al.<sup>57</sup> (2008) Val-DICTATE</p>	<p>DB, MB, RCT  Patients with stage 1</p>	<p>4 weeks  Duration not</p>	<p>Primary: Percentage of patients whose</p>	<p>Primary: A significantly higher proportion of hypertensive patients met blood pressure control levels in the valsartan and HCTZ group (37%) compared</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>HCTZ 25 mg QD vs valsartan and HCTZ 160-12.5 mg QD (fixed-dose combination product)</p>	<p>to 2 HTN whose BP remained uncontrolled on HCTZ 12.5 mg</p>	<p>reported</p>	<p>clinic blood pressure values were &lt;140/90 mm Hg and blood pressure values  Secondary: Not reported</p>	<p>with the HCTZ group (16%; P&lt;0.001).  Changes in SBP and DBP were significantly greater with valsartan and HCTZ (-12.4/-7.5 mm Hg) compared to HCTZ (-5.6/-2.1 mm Hg; P&lt;0.001).  Secondary: Not reported</p>
<p>White et al.<sup>58</sup> (2008)  HCTZ 25 mg QD and valsartan 160 mg  vs  HCTZ 25 mg QD and telmisartan 80 mg  vs  placebo</p>	<p>DB, PC, RCT  Hypertensive patients</p>	<p>N=1,181  8 weeks</p>	<p>Primary: Changes in DBP and SBP at 8 weeks  Secondary: Safety</p>	<p>Primary: Changes from baseline in blood pressure following telmisartan and HCTZ (-24.6/-18.2 mm Hg) were significantly greater than both valsartan and HCTZ (-22.5/-17.0 mm Hg; P=0.017 for SBP and P=0.025 for DBP), and placebo (-4.1/-6.1 mm Hg; P&lt;0.0001).  Secondary: The total number of patients with at least one adverse event reported was similar among the three treatment groups and was 37% for valsartan and HCTZ, 36% for telmisartan and HCTZ, and 42% for placebo.</p>
<p>Waeber et al.<sup>59</sup> (2001)  Valsartan 80 mg QD, which was switched to valsartan 80 mg and HCTZ 12.5 mg QD or valsartan 80 mg and benazepril 10</p>	<p>OL, RCT  Patients with mild-to-moderate uncontrolled HTN (DBP ≥90) while on valsartan monotherapy</p>	<p>N=327  4 weeks</p>	<p>Primary: Efficacy and safety  Secondary: Not reported</p>	<p>Primary: The two combinations produced an additional blood pressure reduction compared to monotherapy (P&lt;0.001 for both), with similar DBP reductions reported for the two combination groups (-4.5 mm Hg with valsartan plus HCTZ and -3.3 mm Hg with valsartan plus benazepril).  SBP reductions of -6.7 and -3.2 mm Hg with valsartan plus HCTZ and valsartan plus benazepril, respectively, were reported (P=0.1).  At the end of the trial, the blood pressure of the responders to valsartan monotherapy was lower than that of patients requiring combination</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg QD				<p>therapy.</p> <p>Valsartan given alone or in association with HCTZ or benazepril was well tolerated.</p> <p>Secondary: Not reported</p>
<p>Izzo Jr et al.<sup>60</sup> (2011) ValVET</p> <p>Valsartan and HCTZ 160-12.5 mg QD (fixed-does combination product)</p> <p>vs</p> <p>valsartan 160 mg QD</p> <p>vs</p> <p>HCTZ 12.5 mg QD</p> <p>All patients were allowed to up titrate study medication if blood pressure did not improve.</p>	<p>DB, RCT</p> <p>Patients <math>\geq 70</math> years of age with systolic HTN</p>	<p>N=384</p> <p>16 weeks</p>	<p>Primary: Change in baseline SBP at week 4</p> <p>Secondary: Time to blood pressure control</p>	<p>Primary: At week four, reductions in baseline SBP were significantly greater with combination therapy (-17.3 mm Hg) compared to valsartan (-8.6 mm Hg; <math>P &lt; 0.001</math>). At this time, reductions with combination therapy and HCTZ were similar (-17.3 vs -13.6 mm Hg; <math>P = 0.096</math>).</p> <p>Secondary: Median time to blood pressure control was significantly shorter with combination therapy compared to HCTZ (four vs eight weeks; <math>P &lt; 0.05</math>) and valsartan (four vs 12 weeks; <math>P &lt; 0.0001</math>).</p>
<p>Duprez et al.<sup>61</sup> (abstract) (2011) ValVET</p>	<p>Subgroup analysis</p> <p>Patients <math>\geq 70</math> years of age with systolic</p>	<p>N=108</p> <p>Duration not specified</p>	<p>Primary: Change in ambulatory SBP</p>	<p>Primary: Initiation of treatment with combination valsartan and HCTZ reduced ambulatory blood pressure more effectively compared to monotherapy with either valsartan or HCTZ throughout daytime, night-time, and 24 hr</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Valsartan and HCTZ 160-12.5 mg QD (fixed-does combination product)</p> <p>vs</p> <p>valsartan 160 mg QD</p> <p>vs</p> <p>HCTZ 12.5 mg QD</p> <p>All patients were allowed to up titrate study medication if blood pressure did not improve.</p>	<p>HTN</p>		<p>Secondary: Safety</p>	<p>monitoring periods, as well as during the last four to six hour dosing periods.</p> <p>Twenty-four hour ambulatory blood pressure was reduced from 141.1/76.5 to 125.8/69.2 mm Hg by week four with combination valsartan and HCTZ compared to reductions from 142.2/78.7 to 139.1/77.5 mm Hg with HCTZ and 142.2/78.3 to 136.4/75.1 mm Hg with valsartan (P&lt;0.01 for all).</p> <p>Secondary: In the overall study, tolerability was similar among the three treatment groups.</p>
<p>Schmieder et al.<sup>62</sup> (2009)</p> <p>HCTZ 12.5 to 25 mg QD (with optional addition of amlodipine 5 to 10 mg QD)</p> <p>vs</p> <p>aliskiren 150 to 300 mg QD (with optional addition</p>	<p>AC, DB, RCT</p> <p>Adults with essential HTN</p>	<p>N=1,124</p> <p>12 months</p>	<p>Primary: Safety and change in mean sitting DBP</p> <p>Secondary: Change in mean sitting SBP</p>	<p>Primary: The proportion of patients who experienced adverse events during the six week placebo-controlled period was similar in the aliskiren monotherapy, HCTZ monotherapy, and placebo groups (26.4, 24.5, and 28.5%, respectively).</p> <p>During the 52 week double-blind treatment period, adverse events were reported by a similar proportion of patients receiving the aliskiren and hydrochlorothiazide regimens. Most adverse events were mild or moderate in intensity.</p> <p>At week 26, the aliskiren regimen provided significantly greater reductions from baseline in DBP compared to HCTZ (-14.2 and -13.0 mm Hg, respectively; P&lt;0.05). The greater reduction in DBP with the aliskiren</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>of amlodipine 5 to 10 mg QD)</p> <p>vs</p> <p>placebo for 6 weeks, then randomized to either aliskiren 300 mg QD or HCTZ 25 mg QD</p>				<p>regimen compared with the HCTZ regimen was maintained at week 52 (-16.0 and -15.0 mm Hg, respectively; <math>P&lt;0.05</math>).</p> <p>Secondary: At week 26, the aliskiren regimen provided significantly greater reductions from baseline in SBP compared to HCTZ (-20.3 and -18.6 mm Hg, respectively; <math>P&lt;0.05</math>). Reductions in SBP at week 52 were not inferior to those of HCTZ (-22.1 and -21.2 mm Hg, respectively; <math>P&lt;0.0001</math> for non-inferiority).</p>
<p>Schmieder et al.<sup>63</sup> (2009)</p> <p>HCTZ 12.5 mg QD, followed by 25 mg QD after 3 weeks</p> <p>vs</p> <p>aliskiren 150 mg QD, followed by 300 mg QD after 3 weeks</p> <p>vs</p> <p>placebo, followed by aliskiren 300 mg QD or HCTZ 25 mg QD after 6 weeks</p>	<p>Subgroup analysis of obese patients in Schmieder et al.</p> <p>Patients 18 years of age and older with essential HTN, a mean sitting DBP <math>\geq 90</math> and <math>&lt;110</math> mm Hg; at randomization, patients had to have a mean sitting DBP <math>\geq 95</math> and <math>&lt;110</math> mm Hg and show a difference of <math>\leq 10</math> mm Hg since the previous visit</p>	<p>N=1,124</p> <p>52 weeks</p>	<p>Primary: Mean sitting DBP</p> <p>Secondary: Mean sitting SBP at week 26, mean sitting DBP and SBP at week 52, proportion of patients with response to treatment, blood pressure control at weeks 26 and 52, and safety</p>	<p>Primary: The least squares mean DBP and SBP reductions at week 12 were significantly greater with aliskiren compared to HCTZ (<math>P&lt;0.0001</math> and <math>P=0.001</math> respectively).</p> <p>Secondary: At week 52, aliskiren resulted in significantly greater mean sitting DBP reductions compared to HCTZ (<math>P&lt;0.001</math>).</p> <p>Blood pressure response rates were significantly greater with aliskiren compared to HCTZ at both week 12 and week 52 (<math>P&lt;0.05</math>).</p> <p>Significantly more obese patients achieved blood pressure control with aliskiren compared to HCTZ at week 12 (<math>P=0.0013</math>). Blood pressure control rates were similar between groups at week 52 (<math>P</math> value not reported).</p>
<p>Villamil et al.<sup>64</sup> (2007)</p> <p>HCTZ 6.25 to 25</p>	<p>DB, MC, PC, RCT</p> <p>Men and women <math>\geq 18</math> years with</p>	<p>N=2,776</p> <p>8 weeks</p>	<p>Primary: Change in mean sitting DBP</p>	<p>Primary: Aliskiren monotherapy significantly reduced mean sitting DBP (<math>P=0.0002</math>). Only the aliskiren 150 and 300 mg doses were more effective than placebo (<math>P=0.09</math> for aliskiren 75 mg). HCTZ monotherapy</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg QD</p> <p>vs</p> <p>aliskiren 75 to 300 mg QD</p> <p>vs</p> <p>aliskiren and HCTZ (every dose combination except aliskiren 300 mg and HCTZ 6.25 mg) QD</p> <p>vs</p> <p>placebo</p>	<p>mild-to-moderate essential HTN</p>		<p>Secondary: Change in mean sitting SBP, dose-response efficacy for all treatment groups, proportion achieving a successful response (DBP &lt;90 mm Hg or ≥10 mm Hg), proportion achieving blood pressure control (&lt;140/90 mm Hg), plasma renin activity, renin concentrations, safety</p>	<p>significantly reduced DBP from baseline (P&lt;0.01 for all vs placebo).</p> <p>All combinations were more effective than placebo (P&lt;0.0001) with reductions in DBP ranging from 10.4 to 14.3 mm Hg. Most combination regimens were more effective than monotherapy with the individual components (exceptions were aliskiren 150 mg and HCTZ 6.25 mg vs monotherapy, and aliskiren 75 mg and HCTZ 12.5 mg vs HCTZ monotherapy).</p> <p>Secondary: After eight weeks of therapy, aliskiren 150 and 300 mg regimens (both P&lt;0.0001) were more effective than placebo in lowering mean sitting SBP, but the 75 mg dose was not (P=0.151).</p> <p>Combination therapy was consistently more effective in reducing SBP than monotherapy with the individual components, with the exception of aliskiren 75 mg plus HCTZ 12.5 vs HCTZ monotherapy. Reductions in SBP with combination therapy ranged from 14.3 to 21.2 mm Hg.</p> <p>Blood pressure reductions were related to the doses of both aliskiren and HCTZ.</p> <p>Responder rates were significantly higher with aliskiren 300 mg (63.9%; P=0.0005), HCTZ 12.5 and 25 mg (60.6 and 59.0%, respectively; both P&lt;0.02) and all combination doses (58.4 to 80.6%; all P&lt;0.05) than placebo (45.8%). Responder rates for all combinations of aliskiren and HCTZ 25 mg, and aliskiren 300 mg and HCTZ 12.5 mg were higher than both monotherapies (P&lt;0.05), while aliskiren 75 mg and HCTZ 12.5 mg and aliskiren 150 mg and HCTZ 12.5 mg were more effective than their respective aliskiren monotherapies (P&lt;0.05).</p> <p>In the aliskiren and HCTZ monotherapy groups, only aliskiren 300 mg led to statistically significantly greater control rates than placebo (46.7 vs 28.1%; P=0.0001). Control rates for all combinations, with the exception of aliskiren 75 mg and HCTZ 6.25 mg, were higher than placebo (all P&lt;0.02). There was a trend towards improved control rates with combination therapy (37.4 to 59.5%) compared to aliskiren monotherapy</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>(29.0 to 46.7%) or HCTZ monotherapy (32.5 to 37.8%). Combinations utilizing the higher doses of one or both drugs (aliskiren 75 to 300 mg with HCTZ 25 mg or aliskiren 150 to 300 mg with HCTZ 12.5 mg) yielded control rates that were significantly higher than monotherapy with either component.</p> <p>While all doses of aliskiren decreased plasma renin activity and all doses of HCTZ increased plasma renin activity, combination therapy resulted in decreased plasma renin activity of 46.1 to 63.5%. Renin concentrations increased in all monotherapy and combination regimens with the exception of HCTZ 6.25 and 12.5 mg.</p> <p>All active treatments were well tolerated with 37.3 to 39.2% of patients experiencing adverse events with aliskiren monotherapy, 38.7 to 42.0% with HCTZ monotherapy, 34.6 to 45.3% with aliskiren and HCTZ, and 44% with placebo. Hypokalemia (serum potassium &lt;3.5 mmol/L) occurred with the highest frequency with HCTZ 12.5 and 25 mg (3.9 and 5.2%, respectively). When administered in combination with aliskiren, the frequency of hypokalemia was 0.7 to 2.0% with HCTZ 12.5 mg and 2.2% to 3.4% with HCTZ 25 mg.</p>
<p>Blumenstein et al.<sup>65</sup> (2009)</p> <p>HCTZ 25 mg (existing therapy)</p> <p>vs</p> <p>aliskiren and HCTZ 300-25 mg QD (fixed-dose combination product)</p> <p>vs</p>	<p>DB, MC, RCT</p> <p>Patients with HTN and an inadequate response to HCTZ (mean sitting DBP &gt;90 and ≤110 mm Hg following 4 weeks of HCTZ 25 mg)</p>	<p>N=722</p> <p>8 weeks</p>	<p>Primary:</p> <p>Changes in mean sitting SBP/DBP, proportion of patients achieving blood pressure control (mean sitting blood pressure &lt;140/90 mm Hg), and blood pressure response rates (msDBP &lt;90 mm Hg or a ≥10 mm Hg decrease from baseline)</p>	<p>Primary:</p> <p>The mean reductions in mean sitting SBP/DBP from baseline with aliskiren and HCTZ 300-25 and 150-25 mg were significantly greater compared to those achieved with HCTZ monotherapy (P&lt;0.001 for all).</p> <p>Rates of blood pressure control were significantly higher with aliskiren and HCTZ 300-25 and 150-25 mg compared to HCTZ monotherapy (P&lt;0.001 for both).</p> <p>Aliskiren and HCTZ 300-25 mg provided significantly greater reductions in mean sitting SBP/DBP and rates of blood pressure control compared to aliskiren and HCTZ 150-25 mg dose (P&lt;0.05 for all).</p> <p>Blood pressure response rates were significantly higher with aliskiren and HCTZ 300-25mg (78.5%) and aliskiren and HCTZ 150-25 mg (67.4%) compared to HCTZ monotherapy (47.1%; P&lt;0.001 for both comparisons).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
aliskiren and HCTZ 150-25 mg QD (fixed-dose combination product)			Secondary: Not reported	<p>All treatments were generally well-tolerated and the proportion of patients experiencing adverse events was similar across treatment groups. The majority of adverse events were mild and transient. Adverse events reported in &gt;2% of patients were nasopharyngitis, dizziness, back pain, and vertigo.</p> <p>The proportion of patients with serum potassium &lt;3.5 mmol/L was lower with aliskiren and HCTZ (1.3 to 2.2%) compared to HCTZ monotherapy (3.4%). Hyperkalemia (serum potassium &gt;5.5 mmol/L) was observed in only one patient receiving aliskiren and HCTZ and two patients in the HCTZ monotherapy group. No patient had increases in serum creatinine above the pre-specified clinically significant threshold.</p> <p>Secondary: Not reported</p>
<p>Jordan et al.<sup>66</sup> (2007)</p> <p>HCTZ 25 mg QD (existing therapy)</p> <p>vs</p> <p>aliskiren 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)</p> <p>vs</p> <p>amlodipine 5 to 10 mg QD, added to existing HCTZ therapy (single entity products)</p>	<p>DB, DD, MC, PG, RCT</p> <p>Obese men and women (BMI ≥30 kg/m<sup>2</sup>) ≥18 years with essential HTN (mean sitting DBP 95 to 109 mm Hg and SBP &lt;180 mm Hg) who had not responded to 4 weeks of treatment with HCTZ 25 mg</p>	<p>N=489</p> <p>12 weeks</p>	<p>Primary: Change in mean sitting DBP with aliskiren 300 mg plus HCTZ vs HCTZ alone at 8 weeks</p> <p>Secondary: Comparisons of mean sitting DBP and SBP with aliskiren plus HCTZ vs the other treatment groups, percentage of responders (mean sitting DBP &lt;90 mm Hg or ≥10 mm Hg reduction from baseline), proportion of</p>	<p>Primary: Aliskiren 300 mg added to HCTZ 25 mg significantly reduced mean sitting DBP compared with HCTZ alone at week eight (mean difference, -4.0; P&lt;0.0001).</p> <p>Secondary: Aliskiren 300 mg added to HCTZ caused numerically larger reductions in mean sitting DBP and SBP compared with amlodipine 10 mg plus HCTZ and irbesartan 300 mg plus HCTZ at week eight, but there were no statistically significant differences between treatment groups (P&gt;0.05).</p> <p>Responder rates were significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0193) and week 12 (P=0.004) but comparable to responder rates observed with amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>The proportion of patients achieving blood pressure control was significantly higher with aliskiren plus HCTZ than HCTZ alone at week eight (P=0.0005) and week 12 (P=0.0001) but not statistically different than amlodipine plus HCTZ (P&gt;0.05) and irbesartan plus HCTZ (P&gt;0.05).</p> <p>Plasma renin activity significantly increased (P&lt;0.05) during four weeks</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  irbesartan 150 to 300 mg QD, added to existing HCTZ therapy (single entity products)			patients achieving blood pressure control (mean sitting blood pressure <140/90 mm Hg), plasma renin activity, safety and tolerability	of HCTZ monotherapy. Combination with aliskiren neutralized this increase and led to an overall significant reduction in plasma renin activity compared with pretreatment baseline (P<0.05) whereas amlodipine and irbesartan led to further significant increases (P<0.05).  All of the study treatments were generally well tolerated. Amlodipine plus HCTZ (45.2%) was associated with a higher incidence of adverse events than the other treatment groups (36.1 to 39.3%), largely due to a higher rate of peripheral edema (11.1 vs 0.8 to 1.6%).
Geiger et al. <sup>67</sup> (2009)  HCTZ 25 mg QD  vs  aliskiren 150 to 300 mg QD, added to existing HCTZ therapy  vs  valsartan 160 to 320 mg QD, added to existing HCTZ therapy  vs  aliskiren 150 to 300 mg and valsartan 160 to 320 mg QD, added to existing HCTZ therapy	AC, DB, RCT  Patients ≥18 years of age with mild to moderate essential HTN who were taking HCTZ for 4 weeks with a DBP ≥95 mm Hg	N=641  8 weeks	Primary: Change in DBP at week 8  Secondary: Change SBP at week 8, change in DBP and SBP at week 4, proportion of patients achieving blood pressure control (SBP/DBP <140/90 mm Hg), change in plasma renin activity, plasma renin concentration	Primary: After eight weeks of therapy, the triple therapy showed significantly greater reductions in SBP and DBP compared with the other groups. The additional SBP and DBP reductions were 7 and 5 mm Hg, respectively compared to aliskiren and HCTZ (P<0.0001), 3 and 2 mm Hg compared to valsartan and HCTZ (P<0.01), and 15 and 10 mm Hg compared to HCTZ monotherapy (P<0.001).  Aliskiren and HCTZ and valsartan and HCTZ combination therapies were more effective compared to HCTZ monotherapy. Valsartan and HCTZ was more effective than aliskiren and HCTZ. SBP and DBP were reduced by 15 and 11 mm Hg, respectively in the aliskiren and HCTZ group. SBP and DBP were reduced by 18 and 14 mm Hg, respectively, in the valsartan and HCTZ group.  Secondary: Blood pressure control rate was significantly higher with triple therapy compared to aliskiren and HCTZ (40.9%, P<0.001), valsartan and HCTZ (48.7%, P<0.001), and HCTZ monotherapy (20.5%, P<0.001).  At week four, a significantly greater blood pressure control rate was observed for the triple therapy group at lower doses (150-160-25 mg) compared to the respective doses of the other groups: aliskiren and valsartan and HCTZ (300-320-25 mg) group (56%) compared to aliskiren and HCTZ (36.6%, P<0.05), valsartan and HCTZ (42.2%, P<0.05), and HCTZ monotherapy (19.9%, P<0.01).  At week eight, plasma renin concentration was unchanged in the HCTZ

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>group, but was significantly increased in other groups. A significant decrease in plasma renin activity from baseline was observed in the aliskiren and HCTZ group (<math>P&lt;0.001</math>) and a significant increase was observed in the valsartan and HCTZ (<math>P&lt;0.001</math>). In the HCTZ and triple therapy groups, there was no change in plasma renin activity (both <math>P&gt;0.75</math>).</p>
<p>O'Brien et al.<sup>68</sup> (2007)</p> <p>Aliskiren 150 mg QD for 3 weeks, then HCTZ 25 mg QD was added for an additional 3 weeks (if ABPM remained <math>\geq 135/85</math> mm Hg)</p> <p>vs</p> <p>irbesartan 150 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg QD added for 3 weeks</p> <p>vs</p> <p>ramipril 5 mg QD for 3 weeks, then aliskiren 75 mg QD added for 3 weeks, then aliskiren 150 mg</p>	<p>3 OL studies</p> <p>Men and women 18 to 80 years with ambulatory SBP <math>\geq 140</math> and <math>\leq 180</math> mm Hg without treatment</p>	<p>N=67</p> <p>6 to 9 weeks</p>	<p>Primary: Change in daytime systolic ABPM with combination therapy compared with monotherapy</p> <p>Secondary: Change in daytime diastolic ABPM, nighttime systolic and diastolic ABPM, daytime and nighttime heart rates, plasma renin activity</p>	<p>Primary: Aliskiren coadministered with HCTZ (<math>P=0.0007</math>) or ramipril (<math>P=0.03</math>) led to significantly greater reductions in daytime systolic ABPM compared to monotherapy. There was a trend for a reduction in daytime systolic ABPM with the addition of aliskiren to irbesartan; however, this trend was not statistically significant.</p> <p>Secondary: Aliskiren plus HCTZ significantly lowered daytime diastolic ABPM compared to aliskiren monotherapy (<math>P=0.0006</math>). Changes in nighttime systolic and diastolic ABPM followed similar trends but did not achieve statistical significance (<math>P=0.06</math> and <math>P=0.09</math>, respectively). No changes in heart rate were observed with either aliskiren regimen.</p> <p>Aliskiren added to irbesartan did not significantly change diastolic ABPM compared to irbesartan monotherapy; however, nighttime systolic and diastolic ABPM were significantly reduced (<math>P&lt;0.05</math> for all). No changes in heart rate were observed with either irbesartan regimen.</p> <p>Mean diastolic ABPM was significantly decreased with the addition of aliskiren 150 mg (<math>P&lt;0.05</math>) but not aliskiren 75 mg to ramipril monotherapy. Both aliskiren doses significantly decreased nighttime systolic and diastolic ABPM (<math>P&lt;0.05</math> for all). No changes in heart rate were observed with either ramipril regimen.</p> <p>Aliskiren alone significantly inhibited plasma renin activity by 65% (<math>P&lt;0.0001</math>), while ramipril and irbesartan monotherapy increased renin activity by 90 and 175%, respectively. When aliskiren was coadministered with HCTZ, ramipril or irbesartan, plasma renin activity remained similar to baseline levels or decreased.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>QD added for 3 weeks</p>				
<p>Pepine et al.<sup>69</sup> (2003) INVEST</p> <p>Verapamil SR 240 mg/day (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist strategy)</p> <p>vs</p> <p>atenolol 50 mg/day (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)</p> <p>Trandolapril was recommended for all patients with heart failure, diabetes, or renal insufficiency.</p>	<p>MC, OL, RCT</p> <p>Patients with essential HTN</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: First occurrence of death (all cause), nonfatal MI or stroke</p> <p>Secondary: Cardiovascular death, angina, cardiovascular hospitalization, angina, blood pressure control (SBP/DBP &lt;140/90 mm Hg or &lt;130/85 mm Hg if diabetic or renal impairment), safety</p>	<p>Primary: At 24 months, in the calcium antagonist strategy subgroup, 81.5% of patients were taking verapamil SR, 62.9% trandolapril, and 43.7% HCTZ. In the non-calcium antagonist strategy, 77.5% of patients were taking atenolol, 60.3% HCTZ, and 52.4% trandolapril.</p> <p>After a follow-up of 61,835 patient-years (mean, 2.7 years per patient), 2,269 patients had a primary outcome event with no statistically significant difference between treatment strategies (9.93% in calcium antagonist strategy vs 10.17% in non-calcium antagonist strategy; RR, 0.98; 95% CI, 0.90 to 16; P=0.57).</p> <p>Secondary: There was no significant difference in the rate of cardiovascular death (P=0.94) or cardiovascular hospitalization (P=0.59) between the two treatment groups.</p> <p>At 24 months, angina episodes decreased in both groups, but the mean frequency was lower in the calcium antagonist strategy group (0.77 episodes/week) compared to the non-calcium antagonist strategy group (0.88 episodes/week; P=0.02).</p> <p>Two-year blood pressure control was similar between groups. The blood pressure goals were achieved by 65.0% (systolic) and 88.5% (diastolic) of calcium antagonist strategy patients and 64.0% (systolic) and 88.1% (diastolic) of non-calcium antagonist strategy patients. A total of 71.7% of calcium antagonist strategy patients and 70.7% of non-calcium antagonist strategy patients achieved an SBP &lt;140 mm Hg and DBP &lt;90 mm Hg.</p> <p>Both regimens were generally well tolerated. Patients in the calcium antagonist strategy group reported constipation and cough more frequently than patients in the non-calcium antagonist strategy group, while non-calcium antagonist strategy patients experienced more dyspnea, lightheadedness, symptomatic bradycardia and wheezing (all were statistically significant with P≤0.05).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Hansson et al.<sup>70</sup> (1999) STOP-Hypertension</p> <p>Atenolol 50 mg or metoprolol 100 mg or pindolol 5 mg QD and/or HCTZ 25 mg with amiloride 2 to 5 mg QD</p> <p>vs</p> <p>enalapril 10 mg or lisinopril 10 mg QD</p> <p>vs</p> <p>felodipine 2.5 mg or isradipine 2.5 mg QD</p>	<p>MC, OL, PRO, RCT</p> <p>Men and women, age 70 to 84 years with HTN (SBP <math>\geq</math>180 mm Hg or DBP <math>\geq</math>105 mm Hg or both)</p>	<p>N=6,614</p> <p>4 years</p>	<p>Primary: Fatal stroke, fatal MI, other fatal cardiovascular events</p> <p>Secondary: Blood pressure</p>	<p>Primary: The rate of prevention of cardiovascular deaths was similar in all groups (RR, 0.97 to 1.4; 95% CI, 0.86 to 1.26).</p> <p>Fatal cardiovascular events, including fatal stroke and fatal myocardial infarction MI, occurred in 19.8 per 1,000 patient-years in the <math>\beta</math>-blocker and/or HCTZ group, in the felodipine or isradipine group and in the enalapril or lisinopril group (RR, 0.99; 95% CI, 0.84 to 1.16).</p> <p>The RR of cardiovascular death in patients in the enalapril or lisinopril group as compared to the felodipine or isradipine group was 1.4 (95% CI, 0.86 to 1.26; P=0.67.)</p> <p>Secondary: Decreases in blood pressure were similar among the groups.</p>
<p>Pepine et al.<sup>71</sup> (2006) INVEST</p> <p>Verapamil SR (step 1), then add trandolapril if needed (step 2), then increase doses of both (step 3), then add HCTZ (step 4) (calcium antagonist</p>	<p>Post hoc analysis of INVEST</p> <p>Patients with essential HTN</p>	<p>N=22,576</p> <p>24 months</p>	<p>Primary: Risk for adverse outcome associated with baseline factors, follow-up blood pressure and drug treatments</p> <p>Secondary: Not reported</p>	<p>Primary: Previous heart failure (adjusted HR, 1.96), as well as diabetes (HR, 1.77), increased age (HR, 1.63), United States residency (HR, 1.61), renal impairment (HR, 1.50), stroke/TIA (HR, 1.43), smoking (HR, 1.41), MI (HR, 1.34), PVD (HR, 1.27), and revascularization (HR, 1.15) predicted increased risk.</p> <p>Follow-up SBP &lt;140 mm Hg (HR, 0.82) or DBP &lt;90 mm Hg (HR, 0.70) and trandolapril with verapamil SR (HR, 0.78 and 0.79) were associated with reduced risk.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
strategy)  vs  atenolol (step 1), then add HCTZ if needed (step 2), then increase doses of both (step 3), then add trandolapril (step 4) (non-calcium antagonist strategy)				
Conlin et al. <sup>72</sup> (2000) PREVAIL  Low-dose HCTZ plus ARB  vs  candesartan 8 to 16 mg QD, irbesartan 150 to 300 mg QD, losartan 50 to 100 mg QD, and valsartan 80 to 160 mg QD  vs  another ARB	MA  Patients with HTN	N=11,281 (43 trials)  Duration varied	Primary: Weighted average for SBP and DBP reduction with ARB monotherapy, dose titration, and with the addition of low-dose HCTZ were calculated; responder rates  Secondary: Not reported	Primary: The absolute weighted-average reductions in DBP (8.2 to 8.9 mm Hg) and SBP (10.4 to 11.8 mm Hg) for ARB monotherapy were comparable for all ARBs. Responder rates for ARB monotherapy were 48 to 55%.  Dose titration resulted in slightly greater blood pressure reductions and an increase in responder rates of 53 to 63%.  ARB and HCTZ combinations produced substantially greater reductions in SBP (16.1 to 20.6 mm Hg) and DBP (9.9 to 13.6 mm Hg) than ARB monotherapy. Responder rates for ARB and HCTZ combinations were 56 to 70%.  The authors concluded that candesartan, irbesartan, losartan, and valsartan produced comparable antihypertensive efficacy when administered at their recommended doses, a near flat dose response when titrating from starting to maximum recommended dose, and substantial potentiation of the antihypertensive effect with addition of HCTZ.  Secondary: Not reported
Stanton et al. <sup>73</sup> (2010)	MA	N=4,877 (8 trials)	Primary: Paradoxical blood	Primary: There were no significant differences among the pooled aliskiren,

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Irbesartan, losartan, valsartan, ramipril, HCTZ, placebo</p> <p>vs</p> <p>aliskiren 300 mg QD</p>	<p>Adults with mild to moderate essential HTN</p>	<p>4 to 12 weeks</p>	<p>pressure rises, as well as the percentage of patients with SBP increases (&gt;10 or &gt;20 mm Hg) or DBP increases (&gt;5 or &gt;10 mm Hg) from baseline</p> <p>Secondary: Not reported</p>	<p>irbesartan, losartan, valsartan, ramipril, and HCTZ groups in the incidence of SBP increases &gt;10 mm Hg (P=0.30) and &gt;20 mm Hg (P=0.28) or DBP increases &gt;5 mm Hg (P=0.65) and &gt;10 mm Hg (P=0.5).</p> <p>Increases in SBP and DBP occurred significantly more frequently in the pooled placebo group than the aliskiren group (P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Hansson et al.<sup>74</sup> (2000) NORDIL</p> <p>Conventional therapy (diuretic, β-blocker or both)</p> <p>vs</p> <p>diltiazem 180 to 360 mg QD</p>	<p>BE, MC, OL, PRO, RCT</p> <p>Patients 50 to 74 years of age with DBP ≥100 mm Hg and previously untreated</p>	<p>N=10,881</p> <p>4.5 years</p>	<p>Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death</p> <p>Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI</p>	<p>Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/β-blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).</p> <p>Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/β-blocker patients (P=0.04).</p> <p>Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/β-blocker patients (P=0.17).</p> <p>Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57 and congestive heart failure (P=0.42).</p>
<p>Messerli et al.<sup>75</sup> (1998)</p> <p>Diuretics (amiloride, chlorthalidone, HCTZ, HCTZ and triamterene [fixed-dose combination product], or</p>	<p>MA</p> <p>10 RCTs lasting ≥1 year, which used as first line agents diuretics and/or β-blockers and reported morbidity and mortality</p>	<p>N=16,164</p> <p>1 year</p>	<p>Primary: Cardiovascular morbidity and mortality, all-cause morbidity</p> <p>Secondary: Not reported</p>	<p>Primary: Diuretic treatment significantly reduced the odds for cardiovascular mortality by 25% (OR, 0.75; 95% CI, 0.64 to 0.87), while β-blockers did not reduce cardiovascular mortality (OR, 0.98; 95% CI, 0.78 to 1.23; P values not reported).</p> <p>Diuretic treatment significantly reduced the odds for all-cause mortality by 14% (OR, 0.86; 95% CI, 0.77 to 0.96), while β-blockers did not reduce all-cause mortality (OR, 1.05; 95% CI, 0.88 to 1.25; P values not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
thiazide)  vs  β-blockers (atenolol, metoprolol or pindolol)	outcomes in patients ≥60 years of age with HTN			Secondary: Not reported
Baguet et al. <sup>76</sup> (2007)  Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)  Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.	MA  Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)	N=10,818  8 to 12 weeks	Primary: Weighted average reductions in SBP and DBP  Secondary: Not reported	Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).  The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the β-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).  The weighted average reduction of SBP and DBP for each drug class were as follows: Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively. β-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively. Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively. ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively. ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively. Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials relating to these agents satisfied all inclusion criteria.</p>				<p>Secondary: Not reported</p>
<p>Lindholm et al.<sup>77</sup> (2005)</p> <p>Other antihypertensive therapies (amiloride, amlodipine, bendroflumethiazide*, captopril, diltiazem, enalapril, felodipine, HCTZ, isradipine, lacidipine, lisinopril, losartan, or verapamil)</p> <p>or</p> <p>placebo</p> <p>vs</p> <p>β-blocker therapy (atenolol,</p>	<p>MA</p> <p>13 RCTs evaluating the treatment of primary HTN with a β-blocker as first-line treatment (in ≥50% of all patients in one treatment group) and outcome data for all-cause mortality, cardiovascular morbidity or both</p>	<p>N=105,951</p> <p>2.1 to 10.0 years</p>	<p>Primary: Stroke, MI, all-cause mortality</p> <p>Secondary: Not reported</p>	<p>Primary: The RR of stroke was 16% higher with β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 1.04 to 1.30; P=0.009). The RR of stroke was the highest with atenolol (26% higher) compared to other non β-blockers (RR, 1.26%; 95% CI, 15 to 38; P&lt;0.0001).</p> <p>The relative risk of MI was 2% higher for β-blocker therapy than for the comparator therapies (RR, 1.02; 95% CI, 0.93 to 1.12), which was not significant (P value not reported).</p> <p>The RR of all-cause mortality was 3% higher for β-blocker therapy than for the comparator therapies (RR, 1.16; 95% CI, 0.99 to 1.08; P=0.14).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
metoprolol, oxprenolol*, pindolol, or propranolol)				
Hilleman et al. <sup>78</sup> (1999)  Monotherapy (atenolol, HCTZ, captopril, enalapril, lisinopril, amlodipine, diltiazem, nifedipine, verapamil)  vs  amlodipine-benazepril (fixed-dose combination)	MA (82 trials)  Patients with mild-to-moderate essential HTN	N=not reported  ≥4 weeks	Primary: Absolute change in supine DBP from baseline  Secondary: Percent of patients who achieved blood pressure control, safety	Primary: The mean absolute decrease in supine DBP ranged from 9.7 to 13.3 mm Hg with verapamil showing the greatest effect and captopril the least. When studies were weighted by sample size, amlodipine and benazepril, atenolol, lisinopril, and verapamil showed the greatest blood pressure effect.  Secondary: The average percentage of patients defined as controlled after treatment varied from 53.5 to 79.0%, with amlodipine and benazepril (74.3%) and lisinopril (79.0%) showing the highest percentage control (P=0.096).  The incidence of adverse events ranged from 12.1 to 41.8%, with lisinopril and verapamil showing the lowest incidences (12.1% and 14.1%, respectively) and nifedipine the highest incidence. Lisinopril demonstrated significantly less overall side effects compared to nifedipine (P=0.030).  Nifedipine demonstrated a higher withdrawal rate due to side effects compared to atenolol, HCTZ, enalapril, amlodipine, and diltiazem (P=0.002). Although amlodipine and benazepril had the lowest rate of withdrawals due to adverse events, lack of significant change was due to the low number of cohorts available for analysis.
Wysong et al. <sup>79</sup> (2007)  Other antihypertensive therapies (i.e., placebo, diuretics, calcium channel blockers, or renin-angiotensin system inhibitors)	MA  13 RCTs evaluating patients ≥18 years of age with HTN	N=91,561  Duration varied	Primary: All-cause mortality  Secondary: Stroke, CHD, cardiovascular death, total cardiovascular disease, adverse reactions	Primary: There was not a significant difference observed in all-cause mortality between β-blocker therapy and placebo (RR, 0.99; 95% CI, 0.88 to 1.11; P value not reported), diuretics (RR, 1.04; 95% CI, 0.91 to 1.19; P value not reported) or renin-angiotensin system inhibitors (RR, 1.10; 95% CI, 0.98 to 1.24; P value not reported). There was a significantly higher rate in all-cause mortality with β-blocker therapy compared to calcium channel blockers (RR, 1.07; 95% CI, 1.00 to 1.14; P=0.04).  Secondary: There was a significant decrease in stroke observed with β-blocker therapy

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  β-blockers (atenolol, metoprolol, oxprenolol*, or propranolol)				<p>compared to placebo (RR, 0.80; 95% CI, 0.66 to 0.96). Also there was a significant increase in stroke with β-blocker therapy compared to calcium channel blockers (RR, 1.24; 95% CI, 1.11 to 1.40) and renin-angiotensin system inhibitors (RR, 1.30; 95% CI, 1.11 to 1.53), but there was no difference observed compared to diuretics (RR, 1.17; 95% CI, 0.65 to 2.09).</p> <p>CHD risk was not significantly different between β-blocker therapy and placebo (RR, 0.93; 95% CI, 0.81 to 1.07), diuretics (RR, 1.12; 95% CI, 0.82 to 1.54), calcium channel blockers (RR, 1.05; 95% CI, 0.96 to 1.15) or renin-angiotensin system inhibitors (RR, 0.90; 95% CI, 0.76 to 1.06).</p> <p>The risk of total cardiovascular disease was lower with β-blocker therapy compared to placebo (RR, 0.88; 95% CI, 0.79 to 0.97). The effect of β-blocker therapy on cardiovascular disease was significantly worse than that of calcium channel blockers (RR, 1.18; 95% CI, 1.08 to 1.29), but was not significantly different from that of diuretics (RR, 1.13; 95% CI, 0.99 to 1.28) or renin-angiotensin system inhibitors (RR, 1.00; 95% CI, 0.72 to 1.3).</p> <p>There was a significantly higher rate of discontinuation due to side effects with β-blocker therapy compared to diuretics (RR, 1.86; 95% CI, 1.39 to 2.50) and renin-angiotensin system inhibitors (RR, 1.41; 95% CI, 1.29 to 1.54), but there was no significant difference compared to calcium channel blockers (RR, 1.20; 95% CI, 0.71 to 2.04). Actual side effects were not reported.</p>

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, QD=once daily, SR=sustained-release

Study Design abbreviations: AC=active comparator, BE=blinded endpoint, DB=double blind, DD=double dummy, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, SB=single blind, XO=cross over

Miscellaneous abbreviations: ACE inhibitors=angiotensin converting enzyme inhibitors, ABPM=ambulatory blood pressure monitoring, BSA=body surface area, CHD=coronary heart disease, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, HCTZ=hydrochlorothiazide, HDL-C=high-density lipoprotein cholesterol, HR=hazard ratio, HTN=hypertension, KCl=potassium chloride, MI=myocardial infarction, NNT=number needed to treat, NYHA=New York Heart Association, OR=odds ratio, PVD=peripheral arterial disease, RR=relative risk, SBP=systolic blood pressure, TC=total cholesterol, TG=triglyceride, TIA=transient ischemic attack, WHO=World Health Organization, WMD=weighted mean difference

### Additional Evidence

#### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

#### Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

## IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 9. Relative Cost of the Thiazide Diuretics**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Chlorothiazide	injection*, suspension, tablet*	Diuril®	\$\$	\$\$
HCTZ	capsule, tablet	N/A	N/A	\$

\*Generic is available in at least one dosage form or strength.

HCTZ=hydrochlorothiazide, N/A=Not available

## X. Conclusions

The thiazide diuretics are approved for the treatment of hypertension and edema due to renal dysfunction. They are also approved as adjunctive therapy for the management of edema associated with congestive heart failure, hepatic cirrhosis, as well as corticosteroid and estrogen therapy.<sup>1-3</sup> All of the agents are available in a generic formulation.

Guidelines recommend the use of diuretics and sodium restriction for the management of ascites due to cirrhosis. Spironolactone is recommended as first-line therapy, either as monotherapy or in combination with furosemide. Amiloride or eplerenone are alternative treatment options in patients experiencing gynecomastia with spironolactone.<sup>18</sup>

For the treatment of chronic heart failure, guidelines recommend the use of diuretics in all patients who have evidence of volume overload. Loop diuretics are generally recommended as initial therapy in patients with left

ventricular dysfunction. For those with persistent fluid retention despite treatment with a loop diuretic, a thiazide diuretic or metolazone may be added to the regimen. In patients with normal left ventricular function, either a thiazide diuretic or loop diuretic may be used as initial therapy to manage fluid overload.<sup>7,8</sup>

There are several published guidelines on the treatment of hypertension. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>9-15</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>9</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>9-17</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>9-15</sup>

In clinical trials, the thiazide diuretics have been shown to effectively lower blood pressure.<sup>23-79</sup> There were no studies found in the medical literature that directly compared the efficacy and safety of the thiazide diuretics for the treatment of hypertension.

There is insufficient evidence to support that one brand thiazide diuretic is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand thiazide diuretics within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand thiazide diuretic is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Diuril® Oral Suspension [package insert]. Morrisville (NC): Salix Pharm, Inc; 2020 Jun.
2. Microzide® [package insert]. Morristown (NJ): Watson Pharma, Inc; 2011 Feb.
3. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
4. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
5. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
6. Mann JFE and Hilgers KF. Use of thiazide diuretics in patients with primary (essential) hypertension. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA, 2021.
7. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
8. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
9. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
10. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6..
11. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, Butalia S, Leung AA, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
12. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
13. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
14. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
15. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
16. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
17. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl).
18. Biggins SW, Angeli P, Garcia-Tsao G, Ginès P, Ling SC, Nadim MK, Wong F, Kim WR. Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology*. 2021 Aug;74(2):1014-1048. doi: 10.1002/hep.31884.
19. Rengo F, Trimarco B, Bonaduce D, Petretta M, Ferrara N, D'Ascia C, et al. Potassium-sparing effect of amiloride in patients receiving diuretics: a quantitative review. *Acta Cardiol*. 1979;34(4)259-67.

20. Cheitlin MD, Byrd R, Benowitz N, Liu E, Modin G. Amiloride improves hemodynamics in patients with chronic congestive heart failure treated with chronic digoxin and diuretics. *Cardiovasc Drugs Ther.* 1991 Aug;5(4):719-25.
21. Kohvakka A. Maintenance of potassium balance during long-term diuretic therapy in chronic heart failure patients with thiazide-induced hypokalemia: comparison of potassium supplementation with potassium chloride and potassium-sparing agents, amiloride and triamterene. *Int J Clin Pharmacol Ther Toxicol.* 1988 May;26(5):273-7.
22. Faris R, Flather MD, Purcell H, Poole-Wilson PA, Coats AJ. Diuretics for heart failure. *Cochrane Database Syst Rev.* 2006 Jan 25;(1):CD003838. DOI: 10.1002/14651858.CD003838.pub2.
23. Hua AS, Kincaid-Smith P. A comparison of the effects of chlorothiazide and of metolazone in the treatment of hypertension. *Clin Sci Mol Med Suppl.* 1976 Dec;3:627s-9s.
24. Carter BL, Ernst ME, Cohen JD. Hydrochlorothiazide versus chlorthalidone: evidence supporting their interchangeability. *Hypertension.* 2004 Jan;43(1):4-9.
25. Ernst ME, Carter BL, Goerdt CJ, Steffensmeier JJ, Phillips BB, Zimmerman MB, Bergus GR. Comparative antihypertensive effects of hydrochlorothiazide and chlorthalidone on ambulatory and office blood pressure. *Hypertension.* 2006 Mar;47(3):352-8.
26. Finnerty FA Jr. Chlorthalidone plus reserpine versus hydrochlorothiazide plus reserpine in a stepped-care approach to the treatment of essential hypertension. *J Clin Pharmacol.* 1980;20(5-6 Pt 1):357-63.
27. Bakris GL, Sica D, White WB, et al. Antihypertensive efficacy of hydrochlorothiazide vs chlorthalidone combined with azilsartan medoxomil. *Am J Med.* 2012 Dec;125(12):1229.e1-1229.e10.
28. Valmin K, Hansen T. Treatment of benign essential hypertension: comparison of furosemide and hydrochlorothiazide. *Eur J Clin Pharmacol.* 1975;8(6):393-401.
29. Araoye MA, Chang MY, Khatri IM, et al. Furosemide compared with hydrochlorothiazide. Long-term treatment of hypertension. *JAMA* 1978;240(17):1863.
30. Madkour H, Gadallah M, Riveline B, Plante GE, Massry SG. Indapamide is superior to thiazide in the preservation of renal function in patients with renal insufficiency and systemic hypertension. *Am J Cardiol.* 1996 Feb 22;77(6):23B-5B.
31. Ames RP. A comparison of blood lipid and blood pressure responses during the treatment of systemic hypertension with indapamide and with thiazides. *Am J Cardiol.* 1996 Feb 22;77(6):12b-6b.
32. Larochelle P, Logan AG. Hydrochlorothiazide-amiloride versus hydrochlorothiazide alone for essential hypertension: effects on blood pressure and serum potassium level. *Can Med Assoc J.* 1985;132(7):801-5.
33. Salmela PI, Juustila H, Kinnunen O, et al. Comparison of low doses of hydrochlorothiazide plus amiloride and hydrochlorothiazide alone in hypertension in elderly patients. *Ann Clin Res.* 1986;18(2):88-92.
34. Multiclinic comparison of amiloride, hydrochlorothiazide, and hydrochlorothiazide plus amiloride in essential hypertension. Multicenter Diuretic Cooperative Study Group. *Arch Intern Med.* 1981 Mar;141(4):482-6.
35. Wray DW, Supiano MA. Impact of aldosterone receptor blockade compared with thiazide therapy on sympathetic nervous system function in geriatric hypertension. *Hypertension* 2010;55:1217-23.
36. Nash DT. Antihypertensive effect and serum potassium homeostasis: comparison of hydrochlorothiazide and spironolactone alone and in combination. *J Med.* 1977;8(5):367-77.
37. Schrijver G, Weinberger MH. Hydrochlorothiazide and spironolactone in hypertension. *Clin Pharmacol Ther.* 1979;25(1):33-42.
38. Johnson JA, Gong Y, Bailey KR, et al. Hydrochlorothiazide and atenolol combination antihypertensive therapy: Effects of drug initiation order. *Clin Pharmacol Ther* 2009;86:533-9.
39. Dahlöf B, Lindholm LH, Hansson L, Scherstén B, Ekblom T, Wester PO. Morbidity and mortality in the Swedish Trial in Old Patients with Hypertension (STOP-Hypertension). *Lancet.* 1991 Nov 23;338(8778):1281-5.
40. Frishman WH, Bryzinski BS, Coulson LR, et al. A multifactorial trial design to assess combination therapy in hypertension. Treatment with bisoprolol and hydrochlorothiazide. *Arch Intern Med.* 1994 Jul 11;154(13):1461-8.
41. Frishman WH, Burris JF, Mroczek WJ, et al. First-line therapy option with low-dose bisoprolol fumarate and low-dose hydrochlorothiazide in patients with stage I and stage II systemic hypertension. *J Clin Pharmacol.* 1995 Feb;35(2):182-8.
42. Dafgard T, Forsen B, Lindahl T. Comparative study of hydrochlorothiazide and a fixed combination of metoprolol and hydrochlorothiazide essential hypertension. *Ann Clin Res.* 1981;13 Suppl 30:37-44.
43. Smilde JG. Comparison of the antihypertensive effect of a double dose of metoprolol versus the addition of hydrochlorothiazide to metoprolol. *Eur J Clin Pharmacol.* 1983;25(5):581-3.

44. Stevens JD, Mullane JF. Propranolol-hydrochlorothiazide combination in essential hypertension. *Clin Ther.* 1982;4(6):497-509.
45. Borhani NO, Mercuri M, Borhani PA, et al. Final outcome results of the Multicenter Isradipine Diuretic Atherosclerosis Study (MIDAS): a randomized controlled trial. *JAMA.* 1996; 276:785-791.
46. Manyemba J. A randomized crossover comparison of reserpine and sustained-release nifedipine in hypertension. *Cent Afr J Med.* 1997 Dec;43(12):344-9.
47. Jamerson K, Bakris GL, Dahlof B, et al; for the ACCOMPLISH Investigators. Exceptional early blood pressure control rates: the ACCOMPLISH trial. *Blood Press.* 2007;16(2):80-6.
48. Jamerson K, Weber MA, Bakris GL, et al. Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high risk patients. *N Engl J Med* 2008;359:2417-28.
49. Bakris G, Briasoulis A, Dahlof B, et al. Comparison of benazepril plus amlodipine or hydrochlorothiazide in high-risk patients with hypertension and coronary artery disease. *Am J Cardiol.* 2013 Jul 15;112(2):255-259.
50. Wing LMH, Reid CM, Ryan P, et al; for the Second Australian National Blood Pressure Study Group. A comparison of outcomes with angiotensin-converting enzyme inhibitors and diuretics for hypertension in the elderly. *N Engl J Med.* 2003 Feb 13;348(7):583-92.
51. Poldermans D, Glazer R, Karagiannis S, et al. Tolerability and blood pressure-lowering efficacy of the combination of amlodipine plus valsartan compared with lisinopril plus hydrochlorothiazide in adult patients with stage 2 hypertension. *Clin Ther.* 2007 Feb;29(2):279-89.
52. Fogari R, Mugellini A, Derosa G; CANDIA Study Group. Efficacy and tolerability of candesartan cilexetil/hydrochlorothiazide and amlodipine in patients with poorly controlled mild-to-moderate essential hypertension. *J Renin Angiotensin Aldosterone Syst.* 2007 Sep;8(3):139-44.
53. Neutel JM, Franklin SS, Lapuerta P, Bhaumik A, Ptaszynska A. A comparison of the efficacy and safety of irbesartan/HCTZ combination therapy with irbesartan and HCTZ monotherapy in the treatment of moderate hypertension. *J Hum Hypertens* 2008;22:266-74.
54. Salerno CM, Demopoulos L, Mukherjee R, Gradman AH. Combination angiotensin receptor blocker/hydrochlorothiazide as initial therapy in the treatment of patients with severe hypertension. *J Clin Hypertens (Greenwich).* 2004 Nov;6(11):614-20.
55. Minami J, Abe C, Akashiba A, Takahashi T, Kameda T, Ishimitsu T, Matsuoka H. Long-term efficacy of combination therapy with losartan and low-dose hydrochlorothiazide in patients with uncontrolled hypertension. *Int Heart J.* 2007 Mar;48(2):177-86.
56. Chrysant SG, Weber MA, Wang AC, Hinman DJ. Evaluation of antihypertensive therapy with the combination of olmesartan medoxomil and hydrochlorothiazide. *Am J Hypertens.* 2004 Mar;17(3):252-9.
57. White WB, Calhoun DA, Samuel R, et al. Improving blood pressure control: increase the dose of diuretic or switch to a fixed-dose angiotensin receptor blocker/diuretic? The valsartan hydrochlorothiazide diuretic for initial control and titration to achieve optimal therapeutic effect (Val-DICTATE) trial. *J Clin Hypertens (Greenwich)* 2008;10:450-8.
58. White WB, Murwin D, Chrysant SG, et al. Effects of the angiotensin II receptor blockers telmisartan versus valsartan in combination with hydrochlorothiazide: a large, confirmatory trial. *Blood Press Monit* 2008;13:21-7.
59. Waeber B, Aschwanden R, Sadecky L, Ferber P. Combination of hydrochlorothiazide or benazepril with valsartan in hypertensive patients unresponsive to valsartan alone. *J Hypertens.* 2001 Nov;19(11):2097-104.
60. Izzo JL Jr, Weintraub HS, Duprez DA, Purkayastha D, Zappe D, Smauel R, et al. Treating systolic hypertension in very elderly with valsartan-hydrochlorothiazide vs either monotherapy: ValVET primary results. 2011;13:722-730.
61. Duprenz DA, Weintraub HS, Cushman WC, Purkayastha D, Zappe D, Samuel R, et al. Effect of valsartan, hydrochlorothiazide, and their combination on 24-h ambulatory blood pressure response in elderly patients with systolic hypertension: a ValVET substudy. *Blood Press Monitor.* 2011 Aug;16(4):186-96.
62. Schmieder RE, Philipp T, Guerediaga J, et al. Long-term antihypertensive efficacy and safety of the oral direct renin inhibitor aliskiren: a 12-month randomized, double-blind comparator trial with hydrochlorothiazide. *Circulation* 2009;119:417-25.
63. Schmieder R, Philipp T, Guerediaga J, Gorostidi M, Bush C, Keefe D. Aliskiren-based therapy lowers blood pressure more effectively than hydrochlorothiazide-based therapy in obese patients with hypertension: sub-analysis of a 52-week, randomized, double-blind trial. *J Hypertens.* 2009;27:1493-1501.
64. Villamil A, Chrysant SG, Calhoun D, et al. Renin inhibition with aliskiren provides additive antihypertensive efficacy when used in combination with hydrochlorothiazide. *J Hypertens.* 2007 Jan;25(1):217-26.



65. Blumenstein M, Romaszko J, Calderón A, et al. Antihypertensive efficacy and tolerability of aliskiren/hydrochlorothiazide (HCT) single-pill combinations in patients who are non-responsive to HCT 25 mg alone. *Curr Med Res Opin* 2009;25:903-910.
66. Jordan J, Engeli S, Boye S, et al. Direct renin inhibition with aliskiren in obese patients with arterial hypertension. *Hypertension*. 2007 May;49:1047-55.
67. Geiger H, Barranco E, Gorostidi M, et al. Combination therapy with various combinations of aliskiren, valsartan, and hydrochlorothiazide in hypertensive patients not adequately responsive to hydrochlorothiazide alone. *J Clin Hypertens (Greenwich)* 2009;11:324-32.
68. O'Brien E, Barton J, Nussberger J, et al. Aliskiren reduces blood pressure and suppresses plasma renin activity in combination with a thiazide diuretic, an angiotensin-converting enzyme inhibitor, or an angiotensin receptor blocker. *Hypertension*. 2007 Feb;49(2):276-84.
69. Pepine CJ, Handberg EM, Cooper-DeHoff RM, et al. A calcium antagonist vs a non-calcium antagonist hypertension treatment strategy for patients with coronary artery disease: the international verapamil-trandolapril study (INVEST): a randomized controlled trial. *JAMA*. 2003 Dec 3;290(21):2805-16.
70. Hansson L, Lindholm LH, Ekblom T, et al. Randomized trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet*. 1999 Nov 20;354(9192):1751-6.
71. Pepine CJ, Kowey PR, Kupfer S, et al. Predictors of adverse outcome among patients with hypertension and coronary artery disease. *J Am Coll Cardiol*. 2006;47(3):547-51.
72. Conlin PR, Spence JD, Williams B, Ribeiro AB, Saito I, Benedict C, et al. Angiotensin II antagonists for hypertension: are there differences in efficacy? *Am J Hypertens*. 2000 Apr;13(4 Pt 1):418-26.
73. Stanton AV, Gradman AH, Schmieder RE, et al. Aliskiren monotherapy does not cause paradoxical blood pressure rises. Meta-analysis of data from 8 clinical trials. *Hypertension* 2010;55:54-60.
74. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, et al. Randomized trial of effects of calcium antagonists compared with diuretics and  $\beta$ -blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet*. 2000 Jul 29;356(9227):359-65.
75. Messerli FH, Grossman E, Goldbourt U. Are beta-blockers efficacious as first-line therapy for hypertension in the elderly? A systematic review. *JAMA*. 1998 Jun 17;279(23):1903-7.
76. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig*. 2007;27(11):735-53.
77. Lindholm LH, Carlberg B, Samuelsson O. Should beta blockers remain first choice in the treatment of primary hypertension? A meta-analysis. *Lancet*. 2005 Oct 29-Nov 4;366(9496):1545-53.
78. Hilleman DE, Ryschon KL, Mohiuddin SM, Wurdeman RL. Fixed-dose combination vs monotherapy in hypertension: a meta-analysis evaluation. *J Hum Hypertens*. 1999;13:477-83.
79. Wiysonge CS, Bradley H, Mayosi BM, Maroney R, Mbewu A, Opie LH, et al. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD002003. doi: 10.1002/14651858.CD002003.pub2.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Thiazide-Like Diuretics  
AHFS Class 402824  
May 4, 2022**

**I. Overview**

The thiazide-like diuretics are approved for the treatment of edema and hypertension.<sup>1-2</sup> They inhibit sodium reabsorption in the distal convoluted tubule of the nephron. This results in an initial modest reduction in plasma volume and cardiac output. However, long-term maintenance of decreased blood pressure has been shown to be associated with partial reversal of the hemodynamic changes as plasma volume and cardiac output return to baseline. Although thiazide-like diuretics are pharmacologically similar to thiazide diuretics, there are chemical differences in the molecular structure that differentiate these agents. Indapamide may produce an independent vascular action, which results in a reduction in total peripheral resistance. Metolazone may produce diuresis in patients with glomerular filtration rates below 20 mL/minute.<sup>1-4</sup>

The thiazide-like diuretics that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All of the agents are available in a generic formulation. This class was last reviewed in February 2020.

**Table 1. Thiazide-Like Diuretics Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Chlorthalidone	tablet*	Thalitone®	chlorthalidone
Indapamide	tablet*	N/A	indapamide
Metolazone	tablet*	N/A	metolazone

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List  
N/A=Not available

**II. Evidence-Based Medicine and Current Treatment Guidelines**

Current treatment guidelines that incorporate the use of the thiazide-like diuretics are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Thiazide-Like Diuretics**

Clinical Guideline	Recommendation(s)
American Heart Association/American College of Cardiology/Heart Failure Society of America: 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure (2022) <sup>5</sup>	<p><b>Treatment of Stage A heart failure (HF)</b></p> <ul style="list-style-type: none"> <li>Hypertension should be controlled in accordance with guideline-directed medication therapy for hypertension to prevent symptomatic HF. (LoE: A)</li> <li>In patients with type 2 diabetes and either established CVD or at high cardiovascular risk, SGLT-2 inhibitors should be used to prevent hospitalizations for HF. (LoE: A)</li> <li>In the general population, healthy lifestyle habits such as regular physical activity, maintaining normal weight, healthy dietary patterns, and avoiding smoking are helpful to reduce future risk of HF. (LoE: B)</li> <li>Patients at risk of developing HF, natriuretic peptide biomarker-based screening followed by team-based care, including a cardiovascular specialist optimizing therapy, can be useful to prevent the development of LV dysfunction (systolic or diastolic) or new-onset HF. (LoE: B)</li> </ul> <p><b>Treatment of Stage B heart failure</b></p> <ul style="list-style-type: none"> <li>In patients with LVEF≤40%, ACE inhibitors should be used to prevent HF and reduce mortality. (LoE: A)</li> <li>In patients with a recent or remote history of MI or ACS, statins should be used to prevent symptomatic HF and adverse cardiovascular events. (LoE: A)</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In patients with a recent MI and LVEF ≤40% who are intolerant to ACE inhibitors, ARB should be used to prevent symptomatic HF and reduce mortality. (LoE: B)</li> <li>• In patients with a recent or remote history of MI or ACS and LVEF ≤40%, evidence-based beta blockers should be used to reduce mortality. (LoE: B)</li> <li>• In patients who are at least 40 days post-MI with LVEF ≤30% and NYHA class I symptoms while receiving guideline-directed medication therapy and have reasonable expectation of meaningful survival for greater than one year, an implantable cardioverter-defibrillator is recommended for primary prevention of sudden cardiac death to reduce total mortality. (LoE: B)</li> <li>• In patients with LVEF ≤40%, beta blockers should be used to prevent symptomatic HF. (LoE: C)</li> <li>• In patients with LVEF ≤50%, thiazolidinediones should not be used because they increase the risk of HF, including hospitalizations. (LoE: B)</li> <li>• In patients with LVEF ≤50%, nondihydropyridine calcium channel blockers with negative inotropic effects may be harmful. (LoE: C)</li> </ul> <p><u>Treatment for Stage C Heart Failure with Reduced Ejection Fraction (HFrEF)</u></p> <ul style="list-style-type: none"> <li>• For patients with stage C HF, avoiding excessive sodium intake is reasonable to reduce congestive symptoms. (LoE: C)</li> <li>• In patients with HF who have fluid retention, diuretics are recommended to relieve congestion, improve symptoms, and prevent worsening HF. (LoE: B)</li> <li>• For patients with HF and congestive symptoms, addition of a thiazide (e.g., metolazone) to treatment with a loop diuretic should be reserved for patients who do not respond to moderate or high dose loop diuretics to minimize electrolyte abnormalities. (LoE: B)</li> <li>• In patients with HFrEF and NYHA class II to III symptoms, the use of an ARNI is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, the use of an ACE inhibitor is beneficial to reduce morbidity and mortality when the use of an ARNI is not feasible. (LoE: A)</li> <li>• In patients with previous or current symptoms of chronic HFrEF, who are intolerant to an ACE inhibitor because of cough or angioedema, and when the use of an ARNI is not feasible, the use of an ARB is recommended to reduce morbidity and mortality. (LoE: A)</li> <li>• In patients with chronic symptomatic HFrEF NYHA class II or III who tolerate an ACE inhibitor or ARB, replacement by an ARNI is recommended to further reduce morbidity and mortality. (LoE: B)</li> <li>• ARNIs should not be administered concomitantly with ACE inhibitors or within 36 hours of the last dose of an ACE inhibitor. (LoE: B)</li> <li>• ARNI or ACE inhibitors should not be administered in patients with a history of angioedema. (LoE: C)</li> <li>• In patients with HFrEF, with current or previous symptoms, use of one of the three β-blockers proven to reduce mortality (e.g., bisoprolol, carvedilol, sustained-release metoprolol succinate) is recommended to reduce mortality and hospitalizations. (LoE: A)</li> <li>• In patients with HFrEF and NYHA class II to IV symptoms, a mineralocorticoid receptor antagonist (spironolactone or eplerenone) is recommended to reduce morbidity and mortality, if eGFR is &gt;30 mL/min/1.73 m<sup>2</sup> and serum potassium is &lt;5.0 mEq/L. Careful monitoring of potassium, renal function, and diuretic dosing should be performed at initiation and closely followed thereafter to minimize risk of hyperkalemia and renal insufficiency. (LoE: A)</li> <li>• In patients taking a mineralocorticoid receptor antagonist whose serum potassium cannot be maintained at &lt;5.5 mEq/L, mineralocorticoid receptor antagonist should be discontinued to avoid life threatening hyperkalemia. (LoE: B)</li> <li>• In patients with symptomatic chronic HFrEF, SGLT-2 an inhibitor is</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>recommended to reduce hospitalization for HF and cardiovascular mortality, irrespective of the presence of type 2 diabetes. (LoE: A)</p> <ul style="list-style-type: none"> <li>• The combination of hydralazine and isosorbide dinitrate is recommended to improve symptoms and reduce morbidity and mortality for patients self-identified as African Americans with NYHA class III to IV HFrEF receiving optimal therapy. (LoE: A)</li> <li>• In patients with current or previous symptomatic HFrEF who cannot be given first-line agents, such as an ARNI, ACE inhibitor or ARB because of drug intolerance or renal insufficiency, a combination of hydralazine and isosorbide dinitrate might be considered to reduce morbidity and mortality. (LoE: C)</li> <li>• In patients with HF class II to IV symptoms, omega-3 polyunsaturated fatty acid supplementation may be reasonable to use as adjunctive therapy to reduce mortality and cardiovascular hospitalizations. (LoE: B)</li> <li>• In patients with chronic HFrEF without specific indication (e.g., venous thromboembolism, atrial fibrillation, a previous thromboembolic event, or a cardioembolic source, anticoagulation is not recommended. (LoE: B)</li> <li>• Dihydropyridine and non-dihydropyridine calcium channel blockers are not recommended for patients with HFrEF. (LoE: A)</li> <li>• In patients with HFrEF, vitamins, nutritional supplements, and hormonal therapy are not recommended other than to correct specific deficiencies. (LoE: B)</li> <li>• In patients with HFrEF, class IC antiarrhythmic medication and dronedarone may increase risk of mortality. (LoE: A)</li> <li>• In patients with HFrEF, thiazolidinediones increase the risk of worsening HF symptoms and hospitalizations. (LoE: A)</li> <li>• In patients with type 2 diabetes and high cardiovascular risk, the DPP-4 inhibitors, saxagliptin and alogliptin, increase the risk of HF hospitalization and should be avoided in patients with HF. (LoE: B)</li> <li>• In patients with HFrEF, nonsteroidal anti-inflammatory drugs worsen HF symptoms and should be avoided or withdrawn whenever possible. (LoE: B)</li> <li>• For patients with symptomatic (NYHA class II to III) stable chronic HFrEF (LVEF <math>\leq</math>35%) who are receiving guideline-directed medical therapy, including a maximally tolerated dose of a beta blocker, and who are in sinus rhythm with a heart rate of <math>\geq</math>70 beats per minute at rest, ivabradine can be beneficial to reduce HF hospitalizations and cardiovascular death. (LoE: B)</li> <li>• In patients with symptomatic HFrEF despite guideline-directed medical therapy or who are unable to tolerate guideline-directed medical therapy, digoxin might be considered to decrease hospitalizations for HF. (LoE: B)</li> <li>• In select high-risk patients with HFrEF and recent worsening of HF already on guideline-directed medical therapy, an oral soluble guanylate cyclase stimulator (vericiguat) may be considered to reduce HF hospitalization and cardiovascular death. (LoE: B)</li> </ul> <p><b>Pharmacological treatment for Stage C HF with mildly reduced EF and improved EF</b></p> <ul style="list-style-type: none"> <li>• In patients with HF with mildly reduced EF, SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• Among patients with current or previous symptomatic HF with mildly reduced EF (LVEF, 41 to 49%), use of evidence-based beta blockers for HFrEF, ARNI, ACE inhibitor or ARB, and mineralocorticoid receptor antagonists may be considered to reduce the risk of HF hospitalization and cardiovascular mortality, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• In patients with HF with improved EF after treatment, guideline-directed medical therapy should be continued to prevent relapse of HF and LV dysfunction, even in patients who may become asymptomatic. (LoE: B)</li> </ul> <p><b>Pharmacological treatment for Stage C HF with preserved EF (HFpEF)</b></p>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Patients with hypertension and HFpEF should have medication titrated to attain blood pressure targets in accordance with published clinical practice guidelines to prevent morbidity. (LoE: C)</li> <li>• SGLT-2 inhibitors can be beneficial in decreasing HF hospitalizations and cardiovascular mortality. (LoE: B)</li> <li>• In patients with HFpEF, the use of a mineralocorticoid receptor antagonist, ARB, or ARNI may be considered to decrease hospitalizations, particularly among patients with LVEF on the lower end of this spectrum. (LoE: B)</li> <li>• Routine use of nitrates or phosphodiesterase-5 inhibitors to increase activity or quality of life in patients with HFpEF is ineffective. (LoE: B)</li> </ul> <p><u>Treatment of Stage D (advanced/refractory) HF</u></p> <ul style="list-style-type: none"> <li>• For patients with advanced HF and hyponatremia, the benefit of fluid restriction to reduce congestive symptoms is uncertain. (LoE: C)</li> <li>• Continuous intravenous inotropic support is reasonable as “bridge therapy” in patients with HF (Stage D) refractory to guideline-directed medical therapy and device therapy who are eligible for and awaiting mechanical circulatory support or cardiac transplantation. (LoE: B)</li> <li>• In select patients with HF Stage D, despite optimal guideline-directed medical therapy and device therapy who are ineligible for either mechanical circulatory support or cardiac transplantation, continuous intravenous inotropic support may be considered as palliative therapy for symptom control and improvement in functional status. (LoE: B)</li> <li>• Long-term use of either continuous or intermittent, intravenous inotropic agents, for reasons other than palliative care or as a bridge to advanced therapies, is potentially harmful. (LoE: B)</li> </ul>
<p>European Society of Cardiology: <b>Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure (2021)</b><sup>6</sup></p>	<p><u>Pharmacological treatments indicated in patients with (NYHA Class II-IV) heart failure with reduced ejection fraction</u></p> <ul style="list-style-type: none"> <li>• An ACE inhibitor is recommended, in addition to a <math>\beta</math>-blocker, for symptomatic patients with HFrEF to reduce the risk of HF hospitalization and death.</li> <li>• A mineralocorticoid receptor antagonist (MRA) is recommended for patients with HFrEF, who remain symptomatic despite treatment with an ACE inhibitor and a <math>\beta</math>-blocker, to reduce the risk of HF hospitalization and death.</li> <li>• Dapagliflozin or empagliflozin are recommended for patients with HFrEF to reduce the risk of HF hospitalization and death. Dapagliflozin or empagliflozin are recommended, in addition to optimal medical therapy with an ACE-I/ARNI, a <math>\beta</math>-blocker and an MRA, for patients with HFrEF regardless of diabetes status.</li> <li>• Sacubitril-valsartan is recommended as a replacement for an ACE inhibitor to further reduce the risk of HF hospitalization and death in ambulatory patients with HFrEF who remain symptomatic despite optimal treatment with an ACE inhibitor, a <math>\beta</math>-blocker, and a mineralocorticoid receptor antagonist.</li> <li>• Diuretics are recommended in order to improve symptoms and exercise capacity in patients with signs and/or symptoms of congestion.</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization or cardiovascular death in symptomatic patients with LVEF <math>\leq</math>35%, in sinus rhythm and a resting heart rate <math>\geq</math>70 bpm despite treatment with an evidence-based dose of <math>\beta</math>-blocker (or maximum tolerated dose below that), ACE inhibitor (or ARB), and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• Ivabradine should be considered to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients with LVEF <math>\leq</math>35%, in sinus rhythm and a resting heart rate <math>\geq</math>70 bpm who are unable to tolerate or have contraindications for a <math>\beta</math>-blocker. Patients should also receive an ACE inhibitor (or ARB) and a mineralocorticoid receptor antagonist (or ARB).</li> <li>• An ARB is recommended to reduce the risk of HF hospitalization and cardiovascular death in symptomatic patients unable to tolerate an ACE inhibitor</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>(patients should also receive a <math>\beta</math>-blocker and mineralocorticoid receptor antagonist).</p> <ul style="list-style-type: none"> <li>• An ARB may be considered to reduce the risk of HF hospitalization and death in patients who are symptomatic despite treatment with a <math>\beta</math>-blocker who are unable to tolerate a mineralocorticoid receptor antagonist.</li> <li>• Vericiguat may be considered in patients in NYHA class II-IV who have had worsening HF despite treatment with an ACE-I (or ARNI), a <math>\beta</math>-blocker and an MRA to reduce the risk of CV mortality or HF hospitalization.</li> <li>• Hydralazine and isosorbide dinitrate should be considered in self-identified black patients with LVEF <math>\leq 35\%</math> or with an LVEF <math>&lt; 45\%</math> combined with a dilated LV in NYHA Class III-IV despite treatment with an ACE-I a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist to reduce the risk of HF hospitalization and death.</li> <li>• Hydralazine and isosorbide dinitrate may be considered in symptomatic patients with HF<sub>r</sub>EF who can tolerate neither an ACE inhibitor nor an ARB (or they are contraindicated) to reduce the risk of death.</li> <li>• Digoxin is a treatment with less-certain benefits and may be considered in symptomatic patients in sinus rhythm despite treatment with an ACE inhibitor (or ARB), a <math>\beta</math>-blocker and a mineralocorticoid receptor antagonist, to reduce the risk of hospitalization (both all-cause and HF-hospitalizations).</li> </ul> <p><u>Recommendations for treatment of patients with (NYHA class II-IV) heart failure with mildly reduced ejection fraction (HF<sub>mr</sub>EF)</u></p> <ul style="list-style-type: none"> <li>• Diuretics are recommended in patients with congestion and HF<sub>mr</sub>EF in order to alleviate symptoms and signs.</li> <li>• An ACE inhibitor may be considered for patients with HF<sub>mr</sub>EF to reduce the risk of HF hospitalization and death.</li> <li>• An ARB may be considered for patients with HF<sub>mr</sub>EF to reduce the risk of HF hospitalization and death.</li> <li>• A <math>\beta</math>-blocker may be considered for patients with HF<sub>mr</sub>EF to reduce the risk of HF hospitalization and death.</li> <li>• An MRA may be considered for patients with HF<sub>mr</sub>EF to reduce the risk of HF hospitalization and death.</li> <li>• Sacubitril/valsartan may be considered for patients with HF<sub>mr</sub>EF to reduce the risk of HF hospitalization and death.</li> </ul> <p><u>Recommendations for treatment of patients with heart failure with preserved ejection fraction (HF<sub>p</sub>EF)</u></p> <ul style="list-style-type: none"> <li>• It is recommended to screen patients with HF<sub>p</sub>EF for both cardiovascular and noncardiovascular comorbidities, which, if present, should be treated provided safe and effective interventions exist to improve symptoms, well-being and/or prognosis.</li> <li>• Diuretics are recommended in congested patients with HF<sub>p</sub>EF in order to alleviate symptoms and signs.</li> </ul> <p><u>Recommendations for the primary prevention of heart failure in patients with risk factors for its development</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension is recommended to prevent or delay the onset of HF and prolong life.</li> <li>• Treatment with statins is recommended in patients at high risk of CV disease or with CV disease in order to prevent or delay the onset of HF, and to prevent HF hospitalizations.</li> <li>• SGLT2 inhibitors (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) are recommended in patients with diabetes at high risk of CV disease or with CV disease in order to prevent HF hospitalizations.</li> </ul>



Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Counselling against sedentary habit, obesity, cigarette smoking, and alcohol abuse is recommended to prevent or delay the onset of HF.</li> </ul> <p><u>Recommendations for the initial management of patients with acute heart failure – pharmacotherapy</u></p> <ul style="list-style-type: none"> <li>• Intravenous loop diuretics are recommended for all patients with acute HF admitted with signs/symptoms of fluid overload to improve symptoms. It is recommended to regularly monitor symptoms, urine output, renal function and electrolytes during use of intravenous diuretics.</li> <li>• Combination of a loop diuretic with thiazide type diuretic should be considered in patients with resistant oedema who do not respond to an increase in loop diuretic doses.</li> <li>• In patients with acute HF and SBP &gt;110 mmHg, intravenous vasodilators may be considered as initial therapy to improve symptoms and reduce congestion.</li> <li>• Inotropic agents may be considered in patients with SBP &lt;90 mmHg and evidence of hypoperfusion who do not respond to standard treatment, including fluid challenge, to improve peripheral perfusion and maintain end-organ function.</li> <li>• Inotropic agents are not recommended routinely, due to safety concerns, unless the patient has symptomatic hypotension and evidence of hypoperfusion.</li> <li>• A vasopressor, preferably norepinephrine, may be considered in patients with cardiogenic shock to increase blood pressure and vital organ perfusion.</li> <li>• Thromboembolism prophylaxis (e.g. with LMWH) is recommended in patients not already anticoagulated and with no contraindication to anticoagulation, to reduce the risk of deep venous thrombosis and pulmonary embolism.</li> <li>• Routine use of opiates is not recommended, unless in selected patients with severe/intractable pain or anxiety.</li> </ul>
<p>Eighth Joint National Committee (JNC 8): <b>2014 Evidence-based Guideline for the Management of High Blood Pressure in Adults (2014)</b><sup>7</sup></p>	<ul style="list-style-type: none"> <li>• Pharmacologic treatment should be initiated in patients ≥60 years of age to lower blood pressure at systolic blood pressure ≥150 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;150 mm Hg and goal diastolic blood pressure &lt;90 mm Hg. Adjustment of treatment is not necessary if treatment results in lower blood pressure and treatment is well tolerated and without adverse effects on health or quality of life.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at diastolic blood pressure ≥90 mm Hg to a goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• In patients &lt;60 years of age, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥150 mm Hg to a goal diastolic blood pressure &lt;140 mm Hg.</li> <li>• For patients ≥18 years of age with chronic kidney disease or diabetes, pharmacologic treatment should be initiated to lower blood pressure at systolic blood pressure ≥140 mm Hg or diastolic blood pressure ≥90 mm Hg and to a goal systolic blood pressure &lt;140 mm Hg and goal diastolic blood pressure &lt;90 mm Hg.</li> <li>• Initial antihypertensive treatment for the general nonblack population, including those with diabetes, should include thiazide-type diuretic, calcium channel blocker (CCB), ACE inhibitor, or ARB.</li> <li>• Initial antihypertensive treatment for the general black population, including those with diabetes, should include thiazide-type diuretic or CCB.</li> <li>• For patients ≥18 years of age with chronic kidney disease regardless of race or diabetes status, initial (or add-on) treatment should include an ACE inhibitor or ARB to improve kidney outcomes.</li> <li>• The main goal of antihypertensive treatment is to attain and maintain goal blood pressure.</li> <li>• If goal blood pressure is not attained within a month of treatment, the dose of the initial drug should be increased or second drug from the thiazide-type diuretic,</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>CCB, ACE inhibitor, or ARB classes should be added.</p> <ul style="list-style-type: none"> <li>• If goal is not achieved with two drugs, a third drug from the thiazide-type diuretic, CCB, ACE inhibitor, or ARB classes should be added.</li> <li>• An ACE inhibitor and ARB should not be used together.</li> <li>• Antihypertensive classes can be used if the patient is unable to achieve goal blood pressure with three agents or had a contraindication to a preferred class.</li> <li>• If blood pressure is not able to be achieved or in complicated patients, referral to a hypertension specialist may be indicated.</li> </ul>
<p>International Society of Hypertension: <b>Global Hypertension Practice Guidelines (2020)</b><sup>8</sup></p>	<p><b>Lifestyle modifications</b></p> <ul style="list-style-type: none"> <li>• Lifestyle modification is the first line of antihypertensive treatment.</li> <li>• Salt reduction: Reduce salt added when preparing foods, and at the table. Avoid or limit consumption of high salt foods such as soy sauce, fast foods and processed food including breads and cereals high in salt.</li> <li>• Healthy diet: Eating a diet that is rich in whole grains, fruits, vegetables, polyunsaturated fats and dairy products and reducing food high in sugar, saturated fat and trans fats, such as the DASH diet.</li> <li>• Healthy drinks: Moderate consumption of coffee, green and black tea. Other beverages that can be beneficial include karkadé (hibiscus) tea, pomegranate juice, beetroot juice and cocoa.</li> <li>• Moderation of alcohol consumption: The recommended daily limit for alcohol consumptions is 2 standard drinks for men and 1.5 for women (10 g alcohol/standard drink). Avoid binge drinking.</li> <li>• Weight reduction: Body weight control is indicated to avoid obesity. Particularly abdominal obesity should be managed. Ethnic-specific cut-offs for BMI and waist circumference should be used. Alternatively, a waist-to-height ratio &lt;0.5 is recommended for all populations.</li> <li>• Smoking cessation: Smoking is a major risk factor for cardiovascular disease (CVD), COPD and cancer. Smoking cessation and referral to smoking cessation programs are advised.</li> <li>• Regular physical activity: Studies suggest that regular aerobic and resistance exercise may be beneficial for both the prevention and treatment of hypertension. Moderate intensity aerobic exercise (walking, jogging, cycling, yoga, or swimming) for 30 minutes on five to seven days per week Strength training also can help reduce blood pressure. Performance of resistance/strength exercises on two to three days per week.</li> <li>• Reduce stress and induce mindfulness: Stress should be reduced and mindfulness or meditation introduced into the daily routine.</li> <li>• Reduce exposure to air pollution and cold temperature: Evidence from studies support a negative effect of air pollution on blood pressure in the long-term.</li> </ul> <p><b>Pharmacological Treatment</b></p> <ul style="list-style-type: none"> <li>• Ideal characteristics of drug treatment <ul style="list-style-type: none"> <li>○ Treatments should be evidence-based in relation to morbidity/mortality prevention.</li> <li>○ Use a once-daily regimen which provides 24-hour blood pressure control.</li> <li>○ Treatment should be affordable and/or cost-effective relative to other agents.</li> <li>○ Treatments should be well-tolerated.</li> <li>○ Evidence of benefits of use of the medication in populations to which it is to be applied.</li> </ul> </li> <li>• General scheme for drug treatment <ul style="list-style-type: none"> <li>○ Lifestyle modification is the first line of antihypertensive treatment.</li> <li>○ Grade 1 hypertension (BP 140 to 159/ 90 to 99): Immediate drug treatment in high-risk patients or those with CVD, chronic kidney disease (CKD), diabetes mellitus (DM), or hypertension-mediated organ damage</li> </ul> </li> </ul>



Clinical Guideline	Recommendation(s)
	<p>(HMOD). Drug treatment after three to six months of lifestyle intervention if BP still not controlled in low to moderate risk patients.</p> <ul style="list-style-type: none"> <li>○ Grade 2 hypertension (BP <math>\geq</math>160/100): Immediate drug treatment in all patients.</li> <li>● Core drug-treatment strategy <ul style="list-style-type: none"> <li>○ Step 1: Dual low-dose combination (ACE inhibitor or ARB plus dihydropyridine-calcium channel blockers (DHP-CCB)); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance; consider ACE/ARB plus DHP-CCB or DHP-CCB plus thiazide-like diuretics in black patients.</li> <li>○ Step 2: Dual full-dose combination (ACE inhibitor or ARB plus DHP-CCB); consider monotherapy in low-risk grade 1 hypertension or in very old (<math>\geq</math>80 years) or frail patients; consider ACE/ARB plus thiazide-like diuretic in post-stroke, very elderly, incipient HF, or CCB intolerance.</li> <li>○ Step 3: Triple combination (ACE/ARB plus DHP-CCB plus thiazide-like diuretic).</li> <li>○ Step 4, resistant hypertension: triple combination plus spironolactone or other drug, alternatives include amiloride, doxazosin, eplerenone, clonidine, or <math>\beta</math>-blocker. Caution with spironolactone or other potassium sparing diuretics when estimated GFR <math>&lt;</math>45 mL/min/1.73m<sup>2</sup> or K<sup>+</sup> <math>&gt;</math>4.5 mmol/L.</li> <li>○ Consider <math>\beta</math>-blockers at any treatment step when there is a specific indications for their use, e.g., heart failure, angina, post-MI, atrial fibrillation, or younger women planning pregnancy or pregnant.</li> </ul> </li> </ul>
<p>Hypertension Canada: <b>2020 Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children (2020)</b><sup>9</sup></p>	<p><u>Indications for drug therapy for adults with hypertension without compelling indications for specific agents</u></p> <ul style="list-style-type: none"> <li>● Antihypertensive therapy should be prescribed for average diastolic blood pressure (DBP) measurements of <math>\geq</math>100 mmHg or average systolic blood pressure (SBP) measurements of <math>\geq</math>160 mmHg in patients without macrovascular target organ damage or other cardiovascular risk factors.</li> <li>● Antihypertensive therapy should be strongly considered for average DPB readings <math>\geq</math>90 mmHg or for average SBP readings <math>\geq</math>140 mmHg in the presence of macrovascular target organ damage or other independent cardiovascular risk factors.</li> <li>● For high-risk patients, aged 50 years or older, with SBP levels <math>\geq</math>130 mmHg, intensive management to target a SBP <math>&lt;</math>120 mmHg should be considered. Patient selection for intensive management is recommended and caution should be taken in certain high-risk groups.</li> </ul> <p><u>Indications for drug therapy for adults with diastolic and with or without systolic hypertension</u></p> <ul style="list-style-type: none"> <li>● Initial therapy should be with either monotherapy or single pill combination (SPC). <ul style="list-style-type: none"> <li>○ Recommended monotherapy choices are: <ul style="list-style-type: none"> <li>▪ A thiazide/thiazide-like diuretic, with longer-acting diuretics preferred;</li> <li>▪ A <math>\beta</math>-blocker (in patients <math>&lt;</math>60 years of age);</li> <li>▪ An angiotensin-converting enzyme (ACE) inhibitor (in nonblack patients);</li> <li>▪ An angiotensin receptor blocker (ARB); or</li> <li>▪ A long-acting calcium channel blocker (CCB).</li> </ul> </li> <li>○ Recommended SPC choices are those in which an ACE inhibitor is combined with a CCB, ARB with a CCB, or ACE inhibitor or ARB with a diuretic.</li> <li>○ Hypokalemia should be avoided in patients treated with thiazide/thiazide-like</li> </ul> </li> </ul>

Clinical Guideline	Recommendation(s)
	<p data-bbox="581 205 821 233">diuretic monotherapy.</p> <ul data-bbox="488 237 1406 699" style="list-style-type: none"> <li data-bbox="488 237 1406 415">• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line choices. Useful choices include a thiazide/thiazide-like diuretic or CCB with either: ACE inhibitor, ARB, or <math>\beta</math>-blocker. Caution should be exercised in combining a nondihydropyridine CCB and a <math>\beta</math>-blocker. The combination of an ACE inhibitor and an ARB is not recommended.</li> <li data-bbox="488 420 1406 478">• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other antihypertensive drugs may be added.</li> <li data-bbox="488 483 1406 510">• Possible reasons for poor response to therapy should be considered.</li> <li data-bbox="488 514 1406 699">• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated hypertension; <math>\beta</math>-blockers are not recommended as first-line therapy for uncomplicated hypertension in patients <math>\geq 60</math> years of age; and ACE inhibitors are not recommended as first-line therapy for uncomplicated hypertension in black patients. However, these agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p data-bbox="488 730 1271 758"><u>Indications for drug therapy for adults with isolated systolic hypertension</u></p> <ul data-bbox="488 762 1422 1287" style="list-style-type: none"> <li data-bbox="488 762 1422 884">• Initial therapy should be single-agent therapy with a thiazide/thiazide-like diuretic, a long-acting dihydropyridine CCB, or an ARB. If there are adverse effects, another drug from this group should be substituted. Hypokalemia should be avoided in patients treated with thiazide/thiazide-like diuretic monotherapy.</li> <li data-bbox="488 888 1422 974">• Additional antihypertensive drugs should be used if target BP levels are not achieved with standard-dose monotherapy. Add-on drugs should be chosen from first-line options.</li> <li data-bbox="488 978 1422 1100">• If BP is still not controlled with a combination of two or more first-line agents, or there are adverse effects, other classes of drugs (such as <math>\alpha</math>-blockers, ACE inhibitors, centrally acting agents, or nondihydropyridine CCBs) may be combined or substituted.</li> <li data-bbox="488 1104 1422 1131">• Possible reasons for poor response to therapy should be considered.</li> <li data-bbox="488 1136 1422 1287">• <math>\alpha</math>-Blockers are not recommended as first-line agents for uncomplicated isolated systolic hypertension; and <math>\beta</math>-blockers are not recommended as first-line therapy for isolated systolic hypertension in patients <math>\geq 60</math> years of age. However, both agents may be used in patients with certain comorbid conditions or in combination therapy.</li> </ul> <p data-bbox="488 1318 1263 1346"><u>Guidelines for hypertensive patients with coronary artery disease (CAD)</u></p> <ul data-bbox="488 1350 1422 1854" style="list-style-type: none"> <li data-bbox="488 1350 1422 1409">• For most hypertensive patients with CAD, an ACE inhibitor or ARB is recommended.</li> <li data-bbox="488 1413 1422 1472">• For hypertensive patients with CAD, but without coexisting systolic heart failure, the combination of an ACE inhibitor and ARB is not recommended.</li> <li data-bbox="488 1476 1422 1598">• For high-risk hypertensive patients, when combination therapy is being used, choices should be individualized. The combination of an ACE inhibitor and a dihydropyridine CCB is preferable to an ACE inhibitor and a thiazide/thiazide-like diuretic in selected patients.</li> <li data-bbox="488 1602 1422 1688">• For patients with stable angina pectoris but without previous heart failure, myocardial infarction, or coronary artery bypass surgery, either a <math>\beta</math>-blocker or CCB can be used as initial therapy.</li> <li data-bbox="488 1692 1422 1719">• Short-acting nifedipine should not be used.</li> <li data-bbox="488 1724 1422 1854">• When decreasing SBP to target levels in patients with established CAD (especially if isolated systolic hypertension is present), be cautious when the DBP is <math>\leq 60</math> mmHg because of concerns that myocardial ischemia might be exacerbated, especially in patients with left ventricular hypertrophy (LVH).</li> </ul> <p data-bbox="488 1885 1422 1913"><u>Guidelines for patients with hypertension who have had a recent myocardial infarction</u></p>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Initial therapy should include a <math>\beta</math>-blocker as well as an ACE inhibitor.</li> <li>• An ARB can be used if the patient is intolerant of an ACE inhibitor.</li> <li>• CCBs may be used in patients after myocardial infarction when <math>\beta</math>-blockers are contraindicated or not effective. Nondihydropyridine CCBs should not be used when there is heart failure, evidenced by pulmonary congestion on examination or radiography.</li> </ul> <p><b>Treatment of hypertension in association with heart failure</b></p> <ul style="list-style-type: none"> <li>• In patients with systolic dysfunction (ejection fraction &lt;40%), ACE inhibitors and <math>\beta</math>-blockers are recommended for initial therapy. Aldosterone antagonists (mineralocorticoid receptor antagonists) may be combined in treatment for patients with a recent cardiovascular hospitalization, acute myocardial infarction, elevated B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level, or New York Heart Association (NYHA) Class II-IV symptoms. Careful monitoring for hyperkalemia is recommended when combining an aldosterone antagonist with ACE inhibitor or ARB treatment. Other diuretics are recommended as additional therapy if needed. Beyond considerations of BP control, doses of ACE inhibitors or ARBs should be titrated to those reported to be effective in trials unless adverse effects become manifest.</li> <li>• An ARB is recommended if ACE inhibitors are not tolerated.</li> <li>• A combination of hydralazine and isosorbide dinitrate is recommended if ACE inhibitors and ARBs are contraindicated or not tolerated.</li> <li>• For hypertensive patients whose BP is not controlled, an ARB may be combined with an ACE inhibitor and other antihypertensive drug treatment. Careful monitoring should be used if combining an ACE inhibitor and an ARB because of potential adverse effects such as hypotension, hyperkalemia, and worsening renal function. Additional therapies may also include dihydropyridine CCBs.</li> <li>• An angiotensin receptor-neprilysin inhibitor (ARNI) should be used in place of an ACE inhibitor or ARB for patients with HFrEF (&lt;40%) who remain symptomatic despite treatment with appropriate dose of guideline directed HF therapy. Eligible patients must have a serum potassium &lt;5.2 mmol/L, an eGFR <math>\leq</math>30 mL/min/1.73m<sup>2</sup> and close surveillance of serum potassium and creatinine.</li> </ul> <p><b>Treatment of hypertension in association with stroke</b></p> <ul style="list-style-type: none"> <li>• BP management in acute ischemic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> <li>• BP management after acute ischemic stroke <ul style="list-style-type: none"> <li>○ Strong consideration should be given to the initiation of antihypertensive therapy after the acute phase of a stroke or transient ischemic attack.</li> <li>○ After the acute phase of a stroke, BP-lowering treatment is recommended to a target of consistently &lt;140/90 mmHg.</li> <li>○ Treatment with an ACE inhibitor and thiazide/thiazide-like diuretic combination is preferred.</li> <li>○ For patients with stroke, the combination of an ACE inhibitor and ARB is not recommended.</li> </ul> </li> <li>• BP management in hemorrhagic stroke (onset to 72 hours) <ul style="list-style-type: none"> <li>○ Please refer to Stroke Best Practice recommendations.</li> </ul> </li> </ul> <p><b>Treatment of hypertension in association with LVH</b></p> <ul style="list-style-type: none"> <li>• Hypertensive patients with LVH should be treated with antihypertensive therapy to decrease the rate of subsequent cardiovascular events.</li> <li>• The choice of initial therapy can be influenced by the presence of LVH. Initial therapy can be drug treatment using ACE inhibitors, ARBs, long-acting CCBs, or thiazide/thiazide-like diuretics. Direct arterial vasodilators such as hydralazine or minoxidil should not be used.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Treatment of hypertension in association with nondiabetic chronic kidney disease</u></p> <ul style="list-style-type: none"> <li>• Individualize BP targets in patients with chronic kidney disease. Consider intensive targets (SBP &lt;120 mmHg) in appropriate patients.</li> <li>• For patients with hypertension and proteinuric chronic kidney disease (urinary protein &gt;150 mg per 24 hours or albumin to creatinine ratio &gt;30 mg/mmol), initial therapy should be an ACE inhibitor or an ARB if there is intolerance to ACE inhibitors.</li> <li>• In most cases, combination therapy with other antihypertensive agents might be needed to reach target BP levels.</li> <li>• The combination of an ACE inhibitor and ARB is not recommended for patients with nonproteinuric chronic kidney disease.</li> </ul> <p><u>Treatment of hypertension in association with renovascular disease</u></p> <ul style="list-style-type: none"> <li>• Patients with hypertension attributable to atherosclerotic renal artery stenosis should be primarily medically managed because renal angioplasty and stenting offers no benefit over optimal medical therapy alone.</li> <li>• Renal artery angioplasty and stenting for atherosclerotic hemodynamically significant renal artery stenosis could be considered for patients with uncontrolled hypertension resistant to maximally tolerated pharmacotherapy, progressive renal function loss, and acute pulmonary edema.</li> <li>• Patients with confirmed renal fibromuscular dysplasia (FMD) should be referred to a hypertension specialist.</li> <li>• Renal artery angioplasty without stenting is recommended for treatment of FMD-related renal artery stenosis. Stenting is not recommended unless needed because of a periprocedural dissection. Surgical revascularization should be considered in cases of complex lesions less amendable to angioplasty, stenosis associated with complex aneurysm, and restenosis despite two unsuccessful attempts of angioplasty.</li> </ul> <p><u>Treatment of hypertension in association with diabetes mellitus</u></p> <ul style="list-style-type: none"> <li>• Persons with diabetes mellitus should be treated to attain SBP of &lt;130 mmHg and DBP of &lt;80 mmHg.</li> <li>• For persons with cardiovascular or kidney disease, including microalbuminuria, or with cardiovascular risk factors in addition to diabetes and hypertension, an ACE inhibitor or an ARB is recommended as initial therapy.</li> <li>• For persons with diabetes and hypertension not included in other guidelines in this section, appropriate choices include (in alphabetical order): ACE inhibitors, ARBs, dihydropyridine CCBs, and thiazide/thiazide-like diuretics.</li> <li>• If target BP levels are not achieved with standard-dose monotherapy, additional antihypertensive therapy should be used. For persons in whom combination therapy with an ACE inhibitor is being considered, a dihydropyridine CCB is preferable to a thiazide/thiazide-like diuretic.</li> </ul> <p><u>Resistant hypertension</u></p> <ul style="list-style-type: none"> <li>• Resistant hypertension is defined as BP above target despite three or more BP-lowering drugs at optimal doses preferably including a diuretic (and usually a renin-angiotensin-aldosterone system blocker and a CCB).</li> <li>• Accurate office and out-of-office BP measurement is essential.</li> <li>• Other reasons for apparent resistant hypertension should be eliminated before diagnosing true resistant hypertension, including nonadherence, white coat effect, and secondary hypertension.</li> <li>• Pharmacotherapy with the additional use of spironolactone, bisoprolol, doxazosin, amiloride, eplerenone, or clonidine with the baseline regimen decreases BP significantly, with the greatest BP-lowering shown with spironolactone.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Patients with resistant hypertension should be referred to providers with expertise in diagnosis and management of hypertension.</li> </ul> <p><u>Hypertension and pediatrics</u></p> <ul style="list-style-type: none"> <li>• BP should be measured regularly in children three years of age or older; the auscultatory method is the gold-standard at present.</li> <li>• Initial therapy should be monotherapy, with an ACE inhibitor or ARB (not first-line in black children), or a long-acting dihydropyridine CCB.</li> <li>• The treatment goal is systolic and diastolic office BP and/or ABPM &lt; 95th percentile or &lt;90th percentile in children with risk factors or target organ damage.</li> <li>• Complex cases should be referred to an expert in pediatric hypertension.</li> </ul> <p><u>Hypertension and pregnancy</u></p> <ul style="list-style-type: none"> <li>• Up to 7% of pregnancies are complicated by a hypertensive disorder of pregnancy, and approximately 5% of women will have chronic hypertension when they become pregnant.</li> <li>• The prevalence of hypertension in pregnancy is expected to increase with women becoming pregnant later in their reproductive years and the increasing prevalence of cardiovascular comorbidities such as increased preconception body mass index and maternal diabetes.</li> <li>• The possibility of pregnancy should be considered when managing women with hypertension who are of reproductive age.</li> <li>• Preconception counselling should be offered to all women with hypertension who are considering pregnancy.</li> <li>• ACE inhibitor and ARB therapy should be avoided before conception and during pregnancy unless there is a compelling indication for their use (i.e., proteinuric kidney disease).</li> <li>• Hypertension during pregnancy can increase the risk of adverse maternal and fetal outcomes, including an increased risk of preeclampsia, placental abruption, prematurity, small for gestational age infants, stillbirth, and maternal renal and retinal injury, thus generally requires involvement of an interdisciplinary team including obstetrical care providers.</li> <li>• Antihypertensive therapy is recommended for average SBP measurements of <math>\geq 140</math> mmHg or DBP measurements of <math>\geq 90</math> mmHg in pregnant women with chronic hypertension, gestational hypertension, or preeclampsia. Initial antihypertensive therapy should be monotherapy from the following first-line drugs: oral labetalol, oral methyldopa, long-acting oral nifedipine, or other oral <math>\beta</math>-blockers (acebutolol, metoprolol, pindolol, and propranolol). Other antihypertensive drugs can be considered as second-line drugs including: clonidine, hydralazine, and thiazide diuretics.</li> <li>• Women with severe hypertension with SBP <math>\geq 160</math> mmHg or DBP <math>\geq 110</math> mmHg in pregnancy or postpartum require urgent antihypertensive therapy because it is considered an obstetrical emergency.</li> <li>• Antihypertensive drugs used in breastfeeding women include: labetalol, methyldopa, long-acting nifedipine, enalapril, or captopril.</li> </ul>
<p>European Society of Hypertension/ European Society of Cardiology: <b>2018 Guidelines for the management of arterial hypertension (2018)</b><sup>10</sup></p>	<p><u>Treatment strategies and choice of antihypertensive drugs</u></p> <ul style="list-style-type: none"> <li>• Diuretics (including thiazides, chlorthalidone, and indapamide), <math>\beta</math>-blockers, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) are all suitable for the initiation and maintenance of antihypertensive treatment, either as monotherapy or in some combinations.</li> <li>• Some agents should be considered as the preferential choice in specific conditions because used in trials in those conditions or because of greater effectiveness in specific types of organ damage. Preferred combinations should comprise a renin-angiotensin system (RAS) blocker (either an ACE inhibitor or an ARB) with a</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>CCB or diuretic.</p> <ul style="list-style-type: none"> <li>• It is recommended that beta-blockers are combined with any of the other major drug classes when there are specific clinical situations, e.g. angina, post-myocardial infarction, heart failure, or heart rate control.</li> <li>• Initiation of antihypertensive therapy with a two-drug combination may be considered in patients with markedly high baseline blood pressure (BP) or at high cardiovascular (CV) risk.</li> <li>• The combination of two antagonists of the RAS is not recommended and should be discouraged.</li> <li>• Other drug combinations should be considered and probably are beneficial in proportion to the extent of BP reduction. However, combinations that have been successfully used in trials may be preferable.</li> <li>• Combinations of two antihypertensive drugs at fixed doses in a single tablet may be recommended and favored, because reducing the number of daily pills improves adherence, which is low in patients with hypertension. Exceptions are frail older patients and those at low risk and with grade one hypertension (particularly if systolic BP is &lt;150 mmHg).</li> <li>• It is recommended that if BP is not controlled with a three-drug combination, treatment should be increased by the addition of spironolactone or, if not tolerated, other diuretics such as amiloride or higher doses of other diuretics, a beta-blocker, or an alpha-blocker.</li> </ul> <p><u>Treatment strategies in white-coat and masked hypertension</u></p> <ul style="list-style-type: none"> <li>• In white-coat hypertensives without additional risk factors, therapeutic intervention may be limited to lifestyle changes only, but this decision should be accompanied by close follow-up.</li> <li>• In white-coat hypertensives with a higher CV risk or hypertension-mediated organ damage, drug treatment may be considered in addition to lifestyle changes.</li> <li>• In masked hypertension, both lifestyle measures and antihypertensive drug treatment should be considered, because this type of hypertension has been consistently found to have a CV risk very close to that of in- and out-of-office hypertension.</li> <li>• Antihypertensive drug up-titration should be considered in treated patients whose out-of-office BP is not controlled (i.e., masked uncontrolled hypertension), because of the high CV risk of these patients.</li> </ul> <p><u>Antihypertensive treatment strategies in the elderly</u></p> <ul style="list-style-type: none"> <li>• It is recommended that older patients (≥65 years) are treated with the same recommendations in non-older patient population. In very old patients (≥80 years), it may be appropriate to initiate treatment with monotherapy.</li> <li>• In all older patients, when combination therapy is used, it is recommended that this is initiated at the lowest available doses. In all older patients, and especially very old or frail patients, the possible occurrence of postural BP should be closely monitored and symptoms of possible hypotensive episodes checked by ambulatory blood pressure monitoring.</li> <li>• Unless required for concomitant diseases, loop diuretics and alpha-blockers should be avoided because of their association with injurious falls.</li> <li>• Renal function should be frequently assessed to detect possible increases in serum creatinine and reductions in estimated glomerular filtration rate (eGFR) as a result of BP-related reductions in renal perfusion.</li> <li>• When treated, BP should be lowered to a systolic value of 130 to 139 mmHg and a diastolic value of &lt;80 mmHg if tolerated. Treated values of &lt;130 mmHg should be avoided.</li> </ul> <p><u>Treatment strategies in hypertensive women</u></p>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• Hormone therapy and selective estrogen receptor modulators are not recommended and should not be used for primary or secondary prevention of CVD.</li> <li>• In women with gestational hypertension, pre-existing hypertension superimposed by gestational hypertension, or with hypertension and subclinical organ damage or symptoms, initiation of drug treatments is recommended when SBP is <math>\geq 140</math> mmHg or DBP <math>\geq 90</math> mmHg. In all other cases, initiation of drug treatment is recommended when SBP is <math>\geq 150</math> mmHg or DBP is <math>\geq 95</math> mmHg.</li> <li>• Methyldopa, labetalol, and CCBs are recommended as the drugs of choice for the treatment of hypertension in pregnancy.</li> <li>• ACE inhibitors, ARBs, or direct renin inhibitors are not recommended during pregnancy.</li> <li>• SBP <math>\geq 170</math> mmHg or DBP <math>\geq 110</math> mmHg in a pregnant woman is an emergency, and admission to hospital is recommended.</li> <li>• In severe hypertension, drug treatment with intravenous (IV) labetalol, oral methyldopa, or nifedipine is recommended.</li> <li>• The recommended treatment for hypertensive crisis is IV labetalol or nicardipine and magnesium.</li> <li>• In pre-eclampsia associated with pulmonary edema, nitroglycerin given as an IV infusion is recommended.</li> <li>• In women with gestational hypertension or mild-pre-eclampsia, delivery is recommended at 37 weeks.</li> <li>• It is recommended to expedite delivery in pre-eclampsia with adverse conditions, such as visual disturbances or hemostatic disorders</li> <li>• In women at high risk of pre-eclampsia, provided they are at low risk of gastrointestinal hemorrhage, treatment with low dose aspirin from 12 weeks until delivery may be considered.</li> <li>• Methyldopa, labetalol, and nifedipine should be considered preferential antihypertensive drugs in pregnancy. Intravenous labetalol or infusion of nitroprusside should be considered in case of emergency (pre-eclampsia).</li> </ul> <p><u>Treatment strategies in patients with diabetes</u></p> <ul style="list-style-type: none"> <li>• Antihypertensive drug treatment is recommended for people with diabetes when office BP is <math>\geq 140/90</math> mmHg.</li> <li>• A SBP goal <math>&lt; 130</math> mmHg is recommended in patients with diabetes and <math>&lt; 130</math> mmHg if tolerated, but not <math>&lt; 120</math> mmHg.</li> <li>• In older people, the target SBP range is 130 to 139 mmHg.</li> <li>• The DBP target in patients with diabetes is recommended to be <math>&lt; 80</math> mmHg, but not <math>&lt; 70</math> mmHg.</li> <li>• It is recommended to initiate treatment with a combination of a RAS blocker with a CCB or thiazide/thiazide-like diuretic.</li> <li>• It is recommended that individual drug choice takes comorbidities into account.</li> <li>• Simultaneous administration of two blockers of the RAS is not recommended and should be avoided in patients with diabetes.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with nephropathy</u></p> <ul style="list-style-type: none"> <li>• In patients with diabetic and non-diabetic CKD, it is recommended that an office BP of <math>\geq 140/90</math> mmHg be treated with lifestyle advice and BP-lowering medication.</li> <li>• In patients with diabetic or non-diabetic CKD, it is recommended to lower SBP to a range of 130 to 139 mmHg. And individualized treatments should be considered according to its tolerability and impact on renal function and electrolytes.</li> <li>• A combination of a RAS blocker with a CCB or a diuretic is recommended as initial therapy.</li> <li>• RAS blockers are more effective in reducing albuminuria than other</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>antihypertensive agents, and are indicated in hypertensive patients in the presence of microalbuminuria or overt proteinuria.</p> <ul style="list-style-type: none"> <li>• Combination of two RAS blockers, though potentially more effective in reducing proteinuria, is not recommended.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with cerebrovascular disease</u></p> <ul style="list-style-type: none"> <li>• In patients with acute intracerebral hemorrhage, immediate BP lowering is not recommended for patients with SBP &lt;220 mmHg. In patients with SBP <math>\geq</math>220 mmHg, care acute BP lowering with IV therapy to &lt;180 mmHg should be considered.</li> <li>• In acute ischemic stroke, routine BP lowering with antihypertensive therapy is not recommended with the exceptions with the exceptions: in patients with acute ischemic stroke who are eligible for IV thrombolysis, BP should be carefully lowered and maintained at &lt;180/105 mmHg for at least the first 24 hours after thrombolysis; In patients with markedly elevated BP who do not receive fibrinolysis, drug therapy may be considered, based on clinical judgement, to reduce BP by 15% during the first 24 hours after the stroke onset.</li> <li>• In hypertensive patients with an acute cerebrovascular event, antihypertensive treatment is recommended immediately for transient ischemic attack (TIA) and after several days in ischemic stroke.</li> <li>• In all hypertensive patients with ischemic stroke or TIA, an SBP target range of 120 to 130 mmHg should be considered.</li> <li>• The recommended antihypertensive drug treatment strategy for stroke prevention is a RAS blocker plus a CCB or a thiazide-like diuretic.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with heart failure (HF) or left ventricular hypertrophy (LVH)</u></p> <ul style="list-style-type: none"> <li>• In hypertensive patients with heart failure (with reduced or preserved ejection fraction), BP-lowering treatment should be considered if <math>\geq</math>140/90 mmHg.</li> <li>• In patients with HF<sub>r</sub>EF, it is recommended that BP-lowering treatment comprises an ACE inhibitor or ARB, and a beta-blocker and a diuretic and/or mineralocorticoid receptor antagonist (MRA) if required.</li> <li>• Dihydropyridine CCBs may be added if BP control is not achieved.</li> <li>• In patients with HF<sub>p</sub>EF, BP treatment threshold and target values should be the same as for HF<sub>r</sub>EF).</li> <li>• Because no specific drug has proven its superiority, all major agents can be used.</li> <li>• In all patients with LVH, it is recommended to treat with a RAS blocker or in combination with a CCB or diuretic and SBP should be lowered to a range of 120 to 130 mmHg.</li> </ul> <p><u>Therapeutic strategies in hypertensive patients with coronary artery disease (CAD)</u></p> <ul style="list-style-type: none"> <li>• In patients with CAD receiving BP-lowering drugs, it is recommended to target SBP to <math>\leq</math>130 mmHg if tolerated, but not &lt;120 mmHg. In older patients, the target is an SBP range of 130 to 140 mmHg. The target DBP is &lt;80 mmHg, but not &lt;70 mmHg.</li> <li>• In hypertensive patients with a history of myocardial infarction, beta-blockers and RAS blockers are recommended as part of treatment.</li> <li>• In patients with asymptomatic angina, beta-blockers and/or CCBs are recommended.</li> </ul>
<p>National Institute for Health and Clinical Excellence: <b>Hypertension in adults: diagnosis</b></p>	<p><u>Choosing antihypertensive drug treatment (for people with or without type II diabetes)</u></p> <ul style="list-style-type: none"> <li>• Where possible, recommend treatment with drugs taken only once a day.</li> <li>• Prescribe non-proprietary drugs where these are appropriate and minimize cost.</li> <li>• Offer people with isolated systolic hypertension (systolic blood pressure <math>\geq</math>160 mmHg) the same treatment as people with both raised systolic and diastolic blood</li> </ul>



Clinical Guideline	Recommendation(s)
<p><b>and management (2019)<sup>11</sup></b></p>	<p>pressure.</p> <ul style="list-style-type: none"> <li>• Offer antihypertensive drug treatment to women of child-bearing potential with diagnosed hypertension in line with recommendations in this guideline. For women considering pregnancy or who are pregnant or breastfeeding, manage hypertension in line with the recommendations on Management of pregnancy with chronic hypertension and Breastfeeding in 'Hypertension in pregnancy'.</li> <li>• When choosing antihypertensive drug treatment for adults of black African or African-Caribbean family origin, consider an angiotensin II receptor blocker, in preference to an angiotensin-converting enzyme inhibitor.</li> </ul> <p><u>Step one treatment</u></p> <ul style="list-style-type: none"> <li>• Patients &lt;55 years of age should be offered a step one antihypertensive with an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).</li> <li>• Offer an ACE inhibitor or an ARB to adults starting step 1 antihypertensive treatment who have type II diabetes and are of any age or family origin or those aged &lt;55 years but not of black African or African-Caribbean family origin.</li> <li>• If an ACE inhibitor is not tolerated, offer an ARB.</li> <li>• Do not combine an ACE inhibitor with an ARB for the treatment of hypertension.</li> <li>• Offer a calcium channel blocker (CCB) to adults starting step 1 antihypertensive treatment who are &gt;55 years of age and do not have diabetes and are of black African or African-Caribbean family origin and do not have type II diabetes and of any age.</li> <li>• If a CCB is not suitable, for example because of edema or intolerance, or if there is evidence of heart failure or a high risk of heart failure, offer a thiazide-like diuretic.</li> <li>• If diuretic treatment is to be initiated or changed, offer a thiazide-like diuretic, such as indapamide in preference to a conventional thiazide diuretic such as bendroflumethiazide or hydrochlorothiazide.</li> <li>• For adults with hypertension who are already receiving treatment with bendroflumethiazide or hydrochlorothiazide, who have stable, well-controlled blood pressure, continue with their treatment.</li> </ul> <p><u>Step two treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on "Medicines adherence: involving patients decisions about prescribed medicines and supporting adherence".</li> <li>• If hypertension is not controlled with a step one treatment of an ACE inhibitor or ARB, offer choice of one of the following drugs in addition to the step one treatment: a CCB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults taking step one treatment of a CCB, offer the choice of one of the following drugs in addition to the step one treatment: an ACE inhibitor or an ARB or a thiazide-like diuretic.</li> <li>• If hypertension is not controlled in adults of black African or African-Caribbean family origin who do not have type 2 diabetes taking step one treatment, consider an ARB, in preference to an ACE inhibitor, in addition to step one treatment.</li> </ul> <p><u>Step three treatment</u></p> <ul style="list-style-type: none"> <li>• Before considering step three treatment, review the person's medications to ensure they are being taken at the optimal doses and discuss adherence (see recommendation under step two).</li> <li>• If hypertension is not controlled in adults taking step two treatment, offer a combination of an ACE inhibitor or ARB and a CCB and a thiazide-like diuretic.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><u>Step four treatment</u></p> <ul style="list-style-type: none"> <li>• If hypertension is not controlled in adults taking the optimal tolerated doses of an ACE inhibitor or an ARB plus a CCB and a thiazide-like diuretic, regard them as having resistant hypertension.</li> <li>• Before considering further treatment for a person with resistant hypertension, confirm elevated clinic blood pressure measurements using ambulatory or home blood pressure recordings, assess for postural hypotension, and discuss adherence.</li> <li>• For people with confirmed resistant hypertension, consider adding a fourth antihypertensive drug as step four treatment or seeking specialist advice.</li> <li>• Consider further diuretic therapy with low-dose spironolactone for adults with resistant hypertension starting step four treatment who have a blood potassium level of 4.5 mmol/l or less. Use particular caution in people with a reduced estimated glomerular filtration rate because they have an increased risk of hyperkalemia.</li> <li>• When using further diuretic therapy for step four treatment of resistant hypertension, monitor blood sodium and potassium and renal function within one month of starting treatment and repeat as needed thereafter.</li> <li>• Consider an alpha-blocker or beta-blocker for adults with resistant hypertension starting step four treatment who have a blood potassium level of more than 4.5 mmol/l.</li> <li>• If blood pressure remains uncontrolled in people with resistant hypertension taking the optimal tolerated doses of four drugs, seek specialist advice.</li> </ul>
<p>International Society on Hypertension in Blacks: <b>Management of High Blood Pressure in Blacks (2010)</b><sup>12</sup></p>	<ul style="list-style-type: none"> <li>• To attain and maintain blood pressure (BP) below target levels, multiple antihypertensive drugs will be required in most hypertensive blacks.</li> <li>• Use of two-drug combination therapy when SBP is &gt;15 mm Hg and/or DBP is &gt;10 mm Hg above goal levels is increasingly recommended as first-line therapy.</li> <li>• Two-drug regimens have generally contained a thiazide-type diuretic; however, the combination of a calcium channel blocker (CCB) with either an ACE inhibitor or an ARB has been shown equally efficacious in BP lowering but with demonstrated superiority (CCB+ACE) for hard clinical outcomes compared with the same ACE inhibitor plus a thiazide-type diuretic.</li> <li>• In secondary prevention patients, the combination therapy should include a drug(s) with the appropriate compelling indications.</li> <li>• Certain classes of antihypertensive medications, specifically diuretics and CCBs, lower BP on average more than <math>\beta</math>-blockers and renin-angiotensin system (RAS) blockers in black patients when used as monotherapies.</li> <li>• In the absence of compelling indications, when BP is near goal levels, monotherapy with a diuretic or a CCB is preferred.</li> <li>• Lifestyle modifications should be initiated in all patients with hypertension, whether or not pharmacotherapy is planned.</li> <li>• ACE inhibitors or ARBs are recommended as alternative monotherapy options in the treatment of hypertension in blacks. The rationale for their lower tier monotherapy recommendation is because they have consistently achieved lesser average reductions in BP relative to that observed with monotherapy using either a diuretic or CCB.</li> </ul>
<p>Kidney Disease Improving Clinical Outcomes Group; <b>KDIGO Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease</b></p>	<p><u>Blood pressure measurement</u></p> <ul style="list-style-type: none"> <li>• The Work Group recommends standardized office blood pressure (BP) measurement in preference to routine office BP measurements for the management of high BP in adults.</li> <li>• The Work Group suggests that out-of-office BP measurements with ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM) be used to complement standardized office BP readings for the management of high BP.</li> </ul> <p><u>Lifestyle interventions for lowering BP in patients with chronic kidney disease (CKD)</u></p>

Clinical Guideline	Recommendation(s)
(2021) <sup>13</sup>	<p><u>not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests targeting a sodium intake &lt;2 grams of sodium per day (or &lt;90 mmol of sodium per day, or &lt;5 grams of sodium chloride per day) in patients with high BP and CKD.</li> <li>The Work Group suggests that patients with high BP and CKD be advised to undertake moderate-intensity physical activity for a cumulative duration of at least 150 minutes per week, or to a level compatible with their cardiovascular and physical tolerance.</li> </ul> <p><u>BP management in patients with CKD, with or without diabetes, not receiving dialysis</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that adults with high BP and CKD be treated with a target systolic BP (SBP) of &lt;120 mmHg, when tolerated, using standardized office BP measurement.</li> <li>The Work Group recommends starting renin-angiotensin-system inhibitors (RASi) (angiotensin-converting enzyme inhibitor [ACEi] or angiotensin II receptor blocker [ARB]) for people with high BP, CKD, and severely increased albuminuria without diabetes.</li> <li>The Work Group suggests starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately increased albuminuria without diabetes.</li> <li>The Work Group recommends starting RASi (ACEi or ARB) for people with high BP, CKD, and moderately-to-severely increased albuminuria with diabetes.</li> <li>The Work Group recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor (DRI) therapy in patients with CKD, with or without diabetes.</li> </ul> <p><u>BP management in kidney transplant recipients</u></p> <ul style="list-style-type: none"> <li>Treat adult kidney transplant recipients with high BP to a target BP of &lt;130 mmHg systolic and &lt;80 mmHg diastolic using standardized office BP measurement.</li> <li>The Work Group recommends that a dihydropyridine calcium channel blocker (CCB) or an ARB be used as the first-line antihypertensive agent in adult kidney transplant recipients.</li> </ul> <p><u>BP management in children with CKD</u></p> <ul style="list-style-type: none"> <li>The Work Group suggests that in children with CKD, 24-hour mean arterial pressure by ABPM should be lowered to &lt;50<sup>th</sup> percentile for age, sex, and height.</li> </ul>
<p>American College of Cardiology/ American Heart Association Task Force: <b>Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)<sup>14</sup></b></p>	<p><u>Initiation of Blood Pressure (BP) Treatment for Overall Cardiovascular Disease (CVD) Risk</u></p> <ul style="list-style-type: none"> <li>Use of BP-lowering medications is recommended for secondary prevention of recurrent CVD events in patients with clinical CVD and an average systolic blood pressure (SBP) ≥130 mmHg or an average diastolic blood pressure (DBP) of ≥80 mmHg and for primary prevention in adults with an estimated 10-year atherosclerotic cardiovascular disease (ASCVD) risk of ≥10% and an average SBP of ≥130 mmHg or an average ≥80 mmHg.</li> <li>Use of BP-lowering medication is recommended for primary prevention of CVD in adults with no history of CVD and with an estimated 10-year ASCVD risk &lt;10% and an SBP of ≥140 mmHg or a DBP of ≥90 mmHg.</li> <li>Simultaneous use of an angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), and/or renin inhibitor is potentially harmful and is not recommended to treat adults with hypertension.</li> <li>For adults with confirmed hypertension and known CVD or 10-year ASCVD risk of ≥10%, a BP target &lt;130/80 mmHg is recommended. For adults with confirmed hypertension without additional markers of increased CVD risk, a BP target &lt;130/80 mmHg may be reasonable.</li> <li>For initiation of antihypertensive drug therapy, first-line agents include thiazide</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>diuretics, calcium channel blockers (CCBs), and ACE inhibitors or ARBs.</p> <ul style="list-style-type: none"> <li>• Initiation of antihypertensive drug therapy with two first-line agents of different classes, either as separate agents or in a fixed-dose combination, is recommended in adults with stage 2 hypertension and an average BP &gt;20/10 mmHg above their BP target.</li> <li>• Initiation of antihypertensive drug therapy with a single antihypertensive drug is reasonable in adults with stage 1 hypertension and BP goal &lt;130/80 mmHg with dosage titration and sequential addition of other agents to achieve the BP target.</li> </ul> <p><u>Stable Ischemic Heart Disease (SIHD)</u></p> <ul style="list-style-type: none"> <li>• In adults with SIHD and hypertension, a BP target &lt;130/80 is recommended.</li> <li>• Adults with SIHD and hypertension (BP ≥130/80 mmHg) should be treated with medications [e.g., guideline-directed medical therapy (GDMT) beta-blockers, ACE inhibitors, or ARBs] for compelling indications [e.g., previous myocardial infarction (MI), stable angina] as first-line therapy, with the addition of other drugs (e.g., dihydropyridine CCBs, thiazide diuretics, and/or mineralocorticoid receptor antagonists) as needed to further control hypertension.</li> <li>• In adults with SIHD with angina and persistent uncontrolled hypertension, the addition of dihydropyridine CCBs to GDMT beta-blockers is recommended.</li> <li>• In adults who have had a MI or acute coronary syndrome, it is reasonable to continue GDMT beta-blockers beyond three years as long-term therapy for hypertension.</li> <li>• Beta-blockers and/or CCBs might be considered to control hypertension in patients with coronary artery disease (CAD) had an MI more than three years ago and have angina.</li> </ul> <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> <li>• In adults with increased risk of HF, the optimal BP in those with hypertension should be &lt;130 mmHg.</li> <li>• Adults with HFrEF and hypertension should be prescribed GDMT titrated to attain a BP &lt;130/80 mmHg.</li> <li>• Non-dihydropyridine CCBs are not recommended in the treatment of hypertension in adults with HFrEF.</li> <li>• In adults with HFpEF who present with symptoms of volume overload, diuretics should be prescribed to control hypertension.</li> <li>• Adults with HFpEF and persistent hypertension after management of volume overload should be prescribed ACE inhibitors or ARBs and beta-blockers titrated to attain SBP &lt;130 mmHg.</li> </ul> <p><u>CKD</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and CKD should be treated to a BP goal &lt;130/80 mmHg.</li> <li>• In adults with hypertension and CKD [stage 3 or higher or stage 1 or 2 with albuminuria (≥300 mg/d, or ≥300 mg/g albumin-to-creatinine ratio or the equivalent in the first morning void)], treatment with an ACE inhibitor is reasonable to slow kidney disease progression. Treatment with an ARB may be reasonable if an ACE inhibitor is not tolerated.</li> <li>• After kidney transplantation, it is reasonable to treat patients with hypertension to a BP goal &lt;130/80 mmHg and with a CCB on the basis of improved glomerular filtration rate (GFR) and kidney survival.</li> </ul> <p><u>Cerebrovascular Disease</u></p> <ul style="list-style-type: none"> <li>• In adults with intracerebral hemorrhage (ICH) who present with SBP &gt;220 mmHg, it is reasonable to use continuous intravenous (IV) drug infusion and close BP monitoring to lower levels. Immediate lowering of SBP to &lt;140 mmHg in adults with spontaneous ICH who present within six hours of the acute event and have an</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>SBP between 150 mmHg and 220 mmHg is not of benefit to reduce death or severe disability and can be potentially harmful.</p> <ul style="list-style-type: none"> <li>• Adults with acute ischemic stroke and elevated BP who are eligible for treatment with IV tissue plasminogen activator (tPA) should have their BP slowly lowered to &lt;185/110 mmHg before thrombolytic therapy is initiated.</li> <li>• In adults with an acute ischemic stroke, BP should be &lt;185/110 mmHg before administration of IV tPA and should be maintained below 180/105 mmHg for at least the first 24 hours after initiation drug therapy.</li> <li>• Starting or restarting antihypertensive therapy during hospitalization in patients with BP &gt;140/90 mmHg who are neurologically stable is safe and reasonable to improve long-term BP control, unless contraindicated.</li> <li>• In patient with BP <math>\geq</math>220/120 mmHg who did not receive IV alteplase or endovascular treatment and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke. In patients with BP &lt;220/120 mmHg with the same conditions, initiating or reinitiating treatment of hypertension within the first 48 to 72 hours after an acute ischemic stroke is not effective to prevent death or dependency.</li> <li>• Adults with previously treated stroke or transient ischemic attack should be restarted on antihypertensive treatment after the first few days of the index event to reduce the risk of recurrent stroke and other vascular events. Treatment with a thiazide diuretic, ACE inhibitor, or ARB, or combination treatment consisting of a thiazide diuretic plus ACE inhibitor, is useful.</li> <li>• Adults not previously treated for hypertension who experienced a stroke or transient ischemic attack and have an established BP <math>\geq</math>140/90 mmHg should be prescribed antihypertensive treatment a few days after the index event to reduce the risk of recurrent stroke and other vascular event.</li> <li>• For adults who experience a stroke or transient ischemic attack, selection of specific drugs should be individualized on the basis of patient comorbidities and agent pharmacological class.</li> <li>• For adults who experience a stroke or transient ischemic attack, a BP goal &lt;130/80 mmHg may be reasonable.</li> <li>• For adults with a lacunar stroke, a target SBP goal &lt;130 mmHg may be reasonable.</li> <li>• In adults previously untreated for hypertension who experience an ischemic stroke or transient ischemic attack and have an SBP &lt;140 mmHg and a DBP &lt;90 mmHg, the usefulness of initiating antihypertensive treatment is not well established.</li> </ul> <p><u>Peripheral Artery Disease (PAD)</u></p> <ul style="list-style-type: none"> <li>• Adults with hypertension and PAD should be treated similarly to patients with hypertension without PAD.</li> </ul> <p><u>Diabetes Mellitus (DM)</u></p> <ul style="list-style-type: none"> <li>• In adults with DM and hypertension, antihypertensive drug treatment should be initiated at a BP of <math>\geq</math>130/80 mmHg with a treatment goal &lt;130/80 mmHg.</li> <li>• In adults with DM and hypertension, all first-line classes of antihypertensive agents (i.e., diuretics, ACE inhibitors, ARBs, and CCBs) are useful and effective.</li> <li>• In adults with DM and hypertension, ACE inhibitors or ARBs may be considered in the presence of albuminuria.</li> </ul> <p><u>Atrial Fibrillation, Valvular Heart Disease, and Aortic disease</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension can be useful for prevention of recurrence of AF.</li> <li>• In adults with asymptomatic aortic stenosis, hypertension should be treated with pharmacotherapy, starting at a low dose and gradually titrating upward as needed.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>• In patients with chronic aortic insufficiency, treatment of systolic hypertension with agents that do not slow the heart rate (i.e., avoid beta-blockers) is reasonable.</li> <li>• Beta-blockers are recommended as the preferred antihypertensive agents in patients with hypertension and thoracic aortic disease.</li> </ul> <p><u>Racial and Ethnic Differences in Treatment</u></p> <ul style="list-style-type: none"> <li>• In black adults with hypertension but without HF or CKD, including those with DM, initial antihypertensive treatment should include a thiazide-type diuretic or CCB. Two or more antihypertensive medications are recommended to achieve a BP target &lt;130/80 mmHg in most adults with hypertension, especially in black adults with hypertension.</li> </ul> <p><u>Pregnancy</u></p> <ul style="list-style-type: none"> <li>• Women with hypertension who become pregnant, or are planning to become pregnant, should be transitioned to methyldopa, nifedipine, and/or labetalol during pregnancy.</li> <li>• Women with hypertension who become pregnant should not be treated with ACE inhibitors, ARBs, or direct renin inhibitors.</li> </ul> <p><u>Older Persons</u></p> <ul style="list-style-type: none"> <li>• Treatment of hypertension with an SBP treatment goal &lt;130 mmHg is recommended for noninstitutionalized ambulatory community-dwelling adults (≥65 years of age) with an average SBP of ≥130 mmHg.</li> <li>• For older adults (≥65 years of age) with hypertension and a higher burden of comorbidity and limited life expectancy, clinical judgment, patient preference, and a team-based approach to assess risk/benefit is reasonable for decisions regarding intensity of BP lowering and choice of antihypertensive drugs.</li> </ul> <p><u>Hypertensive Crises</u></p> <ul style="list-style-type: none"> <li>• In adults with a hypertensive emergency, admission to an intensive care unit is recommended for continuous monitoring of BP and target organ damage and for parenteral administration of an appropriate agent.</li> <li>• For adults with a compelling condition (i.e., aortic dissection, severe pre-eclampsia or eclampsia, or pheochromocytoma crisis), SBP should be reduced to &lt;140 mmHg during the first hour and to &lt;120 mmHg in aortic dissection.</li> <li>• For adults without a compelling condition, SBP should be reduced by no more than 25% within the first hours; then, if stable, to 160/100 mmHg within the next two to six hours; and then cautiously to normal during the following 24 to 48 hours.</li> </ul> <p><u>Cognitive Decline and Dementia</u></p> <ul style="list-style-type: none"> <li>• In adults with hypertension, BP lowering is reasonable to prevent cognitive decline and dementia.</li> </ul> <p><u>Patients Undergoing Surgical Procedures</u></p> <ul style="list-style-type: none"> <li>• In patients with hypertension undergoing major surgery who have been on beta-blockers chronically, beta-blockers should be continued.</li> <li>• In patients with hypertension undergoing planned elective major surgery, it is reasonable to continue medical therapy for hypertension until surgery.</li> <li>• In patients with hypertension undergoing major surgery, discontinuation of ACE inhibitors or ARBs perioperatively may be considered.</li> <li>• In patients with planned elective major surgery and SBP ≥180 mmHg or DBP ≥110 mmHg, deferring surgery may be considered.</li> <li>• For patients undergoing surgery, abrupt pre-operative discontinuation of beta-blockers or clonidine is potentially harmful.</li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>Beta-blockers should not be started on the day of surgery in beta-blocker-naïve patients.</li> <li>Patients with intraoperative hypertension should be managed with IV medications until such time as oral medications can be resumed.</li> </ul>
<p>American Diabetes Association: <b>Standards of Medical Care in Diabetes (2022)</b><sup>15</sup></p>	<p><b>Hypertension/blood pressure control</b></p> <ul style="list-style-type: none"> <li>Blood pressure should be measured at every routine visit. When possible, patients found to have elevated blood pressure (<math>\geq 140/90</math>) should have blood pressure confirmed using multiple readings, including measurements on a separate day, to diagnose hypertension. Patients with blood pressure <math>\geq 180/110</math> and cardiovascular disease could be diagnosed with hypertension at a single visit.</li> <li>All hypertensive patients with diabetes should monitor their blood pressure at home.</li> <li>For patients with diabetes and hypertension, with diabetes and hypertension, blood pressure targets should be individualized through a shared decision-making process that addresses cardiovascular risk, potential adverse effects of antihypertensive medications, and patient preferences.</li> <li>For individuals with diabetes and hypertension at higher cardiovascular risk (existing atherosclerotic cardiovascular disease [ASCVD] or 10-year ASCVD risk <math>\geq 15\%</math>), a blood pressure target of <math>&lt;130/80</math> mmHg may be appropriate, if it can be safely attained.</li> <li>For individuals with diabetes and hypertension at lower risk for cardiovascular disease (10-year atherosclerotic cardiovascular disease risk <math>&lt;15\%</math>), treat to blood pressure target of <math>&lt;140/90</math>.</li> <li>In pregnant patients with diabetes and preexisting hypertension, a blood pressure target of 110 to 135/85 is suggested in the interest of reducing the risk for accelerated maternal hypertension and minimizing impaired fetal growth.</li> <li>For patients with blood pressure <math>&gt;120/80</math>, lifestyle intervention consists of weight loss when indicated, a Dietary Approaches to Stop Hypertension (DASH)-style eating pattern including reducing sodium and increasing potassium intake, moderation of alcohol intake, and increased physical activity.</li> <li>Patients with confirmed office-based blood pressure <math>\geq 140/90</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of pharmacologic therapy to achieve blood pressure goals.</li> <li>Patients with confirmed office-based blood pressure <math>\geq 160/100</math> mmHg should, in addition to lifestyle therapy, have prompt initiation and timely titration of two drugs or a single pill combination of drugs demonstrated to reduce cardiovascular events in patients with diabetes.</li> <li>Treatment for hypertension should include drug classes demonstrated to reduce cardiovascular events in patients with diabetes (ACE inhibitors, angiotensin receptor blockers, thiazide-like diuretics, or dihydropyridine calcium channel blockers). ACE inhibitors or angiotensin receptor blockers are recommended first-line therapy for hypertension in people with diabetes and coronary artery disease.</li> <li>Multiple-drug therapy is generally required to achieve blood pressure targets (but not a combination of ACE inhibitors and angiotensin receptor blockers).</li> <li>An ACE inhibitor or angiotensin receptor blocker, at the maximum tolerated dose indicated for blood pressure treatment, is the recommended first-line treatment for hypertension in patients with diabetes and urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine or 30 to 299 mg/g creatinine. If one class is not tolerated, the other should be substituted.</li> <li>For patients treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually.</li> <li>Patients with hypertension who are not meeting blood pressure targets on three classes of antihypertensive medications (including a diuretic) should be considered for mineralocorticoid receptor antagonist therapy.</li> </ul>

Clinical Guideline	Recommendation(s)
	<p><b>Chronic kidney disease</b></p> <ul style="list-style-type: none"> <li>• At least once a year, assess urinary albumin (e.g., spot urinary albumin-to-creatinine ratio) and estimated glomerular filtration rate in patients with type 1 diabetes with duration of five or more years and in all patients with type 2 diabetes.</li> <li>• Patients with diabetes and urinary albumin <math>\geq 300</math> mg/g creatinine and/ or an estimated glomerular filtration rate 30 to 60 mL/min/1.73 m<sup>2</sup> should be monitored twice annually to guide therapy.</li> <li>• Optimize glucose control to reduce the risk or slow the progression of chronic kidney disease (CKD).</li> <li>• For patients with type 2 diabetes and diabetic kidney disease, use of a sodium-glucose cotransporter 2 inhibitor in patients with an estimated glomerular filtration rate <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> and urinary albumin <math>\geq 300</math> mg/g creatinine is recommended to reduce CKD progression and cardiovascular events.</li> <li>• In patients with type 2 diabetes and CKD, consider use of sodium-glucose cotransporter 2 inhibitors additionally for cardiovascular risk reduction when estimated glomerular filtration rate and urinary albumin creatinine are <math>\geq 25</math> mL/min/1.73 m<sup>2</sup> or <math>\geq 300</math> mg/g, respectively.</li> <li>• In patients with CKD who are at increased risk for cardiovascular events or CKD progression and are unable to use a sodium-glucose cotransporter 2 inhibitor, a nonsteroidal mineralocorticoid receptor antagonist (finerenone) is recommended to reduce CKD progression and cardiovascular events.</li> <li>• Optimize blood pressure control and reduce blood pressure variability to reduce the risk or slow the progression of CKD.</li> <li>• Do not discontinue renin-angiotensin system blockade for minor increases in serum creatinine (<math>\leq 30\%</math>) in the absence of volume depletion.</li> <li>• For people with nondialysis-dependent stage 3 or higher CKD, dietary protein intake should be a maximum of 0.8 g/kg body weight per day (the recommended daily allowance). For patients on dialysis, higher levels of dietary protein intake should be considered, since malnutrition is a major problem in some dialysis patients.</li> <li>• In nonpregnant patients with diabetes and hypertension, either an ACE inhibitor or an angiotensin receptor blocker is recommended for those with modestly elevated urinary albumin-to-creatinine ratio (30 to 299 mg/g creatinine) and is strongly recommended for those with urinary albumin-to-creatinine ratio <math>\geq 300</math> mg/g creatinine and/or estimated glomerular filtration rate <math>&lt; 60</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Periodically monitor serum creatinine and potassium levels for the development of increased creatinine or changes in potassium when ACE inhibitors, angiotensin receptor blockers, or diuretics are used.</li> <li>• An ACE inhibitor or an angiotensin receptor blocker is not recommended for the primary prevention of CKD in patients with diabetes who have normal blood pressure, normal urinary albumin-to-creatinine ratio (<math>&lt; 30</math> mg/g creatinine), and normal estimated glomerular filtration rate.</li> <li>• Patients should be referred for evaluation by a nephrologist if they have an estimated glomerular filtration rate <math>&lt; 30</math> mL/min/1.73 m<sup>2</sup>.</li> <li>• Promptly refer to a nephrologist for uncertainty about the etiology of kidney disease, difficult management issues, and rapidly progressing kidney disease.</li> </ul>
<p>American Association for the Study of Liver Diseases: <b>Diagnosis, Evaluation, and Management of</b></p>	<p><b>Treatment of ascites</b></p> <ul style="list-style-type: none"> <li>• Moderate sodium restriction (2 g or 90 mmol/day) and diuretics (spironolactone with or without furosemide) are the first-line treatment in patients with cirrhosis and grade 2 ascites.</li> <li>• After ascites is adequately mobilized, attempts should be made to taper the diuretics to the lowest dose necessary to maintain minimal or no ascites to prevent the development of adverse effects.</li> </ul>



Clinical Guideline	Recommendation(s)
Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance (2021) <sup>16</sup>	<ul style="list-style-type: none"> <li>Fluid restriction is not necessary unless serum sodium is &lt;125 mmol/L.</li> <li>In patients receiving diuretics, body weight and serum creatinine and sodium should be regularly monitored to assess response and to detect the development of adverse effects.</li> <li>Human albumin solution (20 to 40 g/week) or baclofen administration (10 mg/day, with a weekly increase of 10 mg/day, up to 30 mg/day) can be considered in cases of severe muscle cramps.</li> <li>Large-volume paracentesis is the first-line treatment of grade 3 ascites. After paracentesis, sodium restriction and diuretics should be started.</li> <li>Referral for liver transplant evaluation should be considered in patients with grade 2 or 3 ascites.</li> <li>Nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers should be avoided in patients with cirrhosis and ascites.</li> <li>Aminoglycosides should be avoided whenever possible in the treatment of bacterial infections.</li> <li>For patients with cirrhosis and diuretic-responsive ascites, controversial data suggest potential benefits of long-term infusion of human albumin solution. At present, no recommendation can be made for its use in routine clinical practice.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the thiazide-like diuretics are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Thiazide-Like Diuretics<sup>1-3</sup>**

Indication	Chlorthalidone*	Indapamide	Metolazone
<b>Edema</b>			
Adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, and corticosteroid and estrogen therapy	✓		
Treatment of salt and fluid retention associated with congestive heart failure		✓	
Treatment of salt and water retention, including edema accompanying congestive heart failure			✓
Treatment of salt and water retention, including edema accompanying renal disease, including the nephrotic syndrome and states of diminished renal function			✓
<b>Hypertension</b>			
Treatment of hypertension	✓ †	✓ †	✓ †

\*Chlorthalidone is also useful in the treatment of edema due to various forms of renal dysfunction such as nephrotic syndrome, acute glomerulonephritis, and chronic renal failure.

†Alone or in combination with other antihypertensive agents.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the thiazide-like diuretics are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Thiazide-Like Diuretics<sup>4</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
Chlorthalidone	65	75	Not reported	Renal (50 to 74)	40 to 60
Indapamide	100	71 to 79	Liver, extensive (% not reported)	Bile (23) Feces (16 to 20) Renal (60 to 70)	14 to 15
Metolazone	40 to 65	Not reported	Not reported	Renal (56)	8 to 14

## V. Drug Interactions

Major drug interactions with the thiazide-like diuretics are listed in Table 5.

**Table 5. Major Drug Interactions with the Thiazide-Like Diuretics<sup>4</sup>**

Generic Name(s)	Interaction	Mechanism
Thiazide-like diuretics (chlorthalidone, indapamide, metolazone)	Dofetilide	Increased potassium excretion caused by thiazide diuretic administration Hypokalemia may occur.
Thiazide-like diuretics (chlorthalidone, indapamide, metolazone)	Lithium	Thiazide-like diuretics may decrease the renal excretion of lithium and produce elevated serum lithium concentrations with toxicity.
Thiazide-like diuretics (chlorthalidone, indapamide, metolazone)	Digitalis glycosides	Excretion of potassium and magnesium is increased by thiazide-like diuretics. Potassium and magnesium depletion can sensitize the myocardium to the toxic effects of digitalis glycosides.
Thiazide-like diuretics (chlorthalidone, indapamide, metolazone)	Loop diuretics (Bumetanide, ethacrynic acid, furosemide, torsemide)	Both groups have synergistic effects that may result in profound diuresis and serious electrolyte abnormalities
Thiazide-like diuretics (chlorthalidone, indapamide, metolazone)	NSAIDs	Concurrent use of NSAIDs and thiazide-like diuretics may result in reduced diuretic effectiveness and possible nephrotoxicity.
Thiazide-like diuretics (chlorthalidone, metolazone)	Bepridil	Concurrent use of thiazide-like diuretics and bepridil may result in hypokalemia and subsequent cardiotoxicity (torsades de pointes).
Thiazide-like diuretics (chlorthalidone, metolazone)	Flecainide	Concurrent use of thiazide-like diuretics and flecainide may result in increased risk of electrolyte imbalance and subsequent cardiotoxicity.

NSAIDs=nonsteroidal anti-inflammatory drugs

## VI. Adverse Drug Events

The most common adverse drug events reported with the thiazide-like diuretics are listed in Table 6. The boxed warning for metolazone is listed in Table 7.

**Table 6. Adverse Drug Events (%) Reported with the Thiazide-Like Diuretics<sup>1-4</sup>**

Adverse Event(s)	Chlorthalidone	Indapamide	Metolazone
<b>Cardiovascular</b>			
Chest pain	-	<5	✓
Irregular heartbeat	-	<5	-
Orthostatic hypotension	✓	<5	✓
Palpitations	-	<5	✓
Peripheral edema	-	<5	-
Premature ventricular contractions	-	<5	-

Adverse Event(s)	Chlorthalidone	Indapamide	Metolazone
Venous thrombosis	-	-	✓
Volume depletion	-	-	✓
<b>Central Nervous System</b>			
Anxiety	-	≥5	-
Blurred vision	-	<5	✓
Depression	-	<5	-
Dizziness	✓	≥5	✓
Drowsiness	-	<5	✓
Fatigue	-	≥5	✓
Headache	✓	≥5	✓
Insomnia	-	<5	-
Lethargy	-	≥5	-
Lightheadedness	-	<5	✓
Nervousness	-	<5	-
Neuropathy	-	-	✓
Paresthesia	✓	<5	✓
Restlessness	✓	-	✓
Syncope	-	-	✓
Tension	-	≥5	-
Vertigo	✓	<5	✓
Weakness	✓	≥5	✓
Xanthopsia	✓	-	-
<b>Dermatological</b>			
Dermatitis	-	-	✓
Petechiae	-	-	✓
Photosensitivity	✓	-	✓
Pruritus	-	<5	✓
Purpura	✓	-	✓
Rash	✓	<5	✓
Skin necrosis	-	-	✓
Stevens-Johnson syndrome	-	-	✓
Toxic epidermal necrolysis	✓	✓	✓
Urticaria	✓	<5	✓
<b>Gastrointestinal</b>			
Abdominal pain	-	<5	✓
Anorexia	✓	<5	✓
Constipation	✓	<5	✓
Cramping	✓	-	-
Diarrhea	✓	<5	✓
Dry mouth	-	<5	✓
Dyspepsia	-	<5	-
Epigastric distress	-	-	✓
Gastric irritation	✓	<5	-
Nausea	✓	<5	✓
Pancreatitis	✓	-	✓
Vomiting	✓	<5	✓
<b>Genitourinary</b>			
Impotence	✓	<5	✓
Nocturia	-	<5	✓
Polyuria	-	<5	-
<b>Hematologic</b>			
Agranulocytosis	✓	-	✓
Aplastic anemia	✓	-	✓
Leukopenia	✓	-	✓

Adverse Event(s)	Chlorthalidone	Indapamide	Metolazone
Thrombocytopenia	✓	-	✓
<b>Laboratory Test Abnormalities</b>			
Blood urea nitrogen increased	-	<5	✓
Hypercalcemia	✓	✓	✓
Hyperglycemia	✓	<5	✓
Hyperlipidemia	✓	-	-
Hyperuricemia	✓	<5	✓
Hypochloremia	-	<5	✓
Hypokalemia	✓	3 to 7	✓
Hypomagnesemia	✓	✓	✓
Hyponatremia	✓	<5	-
Hypophosphatemia	-	-	✓
Serum creatinine increased	-	-	✓
<b>Musculoskeletal</b>			
Asthenia	-	<5	-
Back pain	-	≥5	-
Joint pain	-	-	✓
Hypertonia	-	<5	-
Muscle spasm	✓	≥5	✓
<b>Renal</b>			
Glycosuria	✓	<5	✓
<b>Respiratory</b>			
Cough	-	<5	-
Pharyngitis	-	<5	-
Rhinitis	-	≥5	-
Sinusitis	-	<5	-
<b>Other</b>			
Chills	-	-	✓
Conjunctivitis	-	<5	-
Gout	-	-	✓
Hemoconcentration	-	-	✓
Hepatitis	-	-	✓
Infection	-	≥5	-
Jaundice	✓	-	✓
Necrotizing angiitis/vasculitis	-	-	✓
Vasculitis	✓	<5	-
Weight loss	-	<5	-

✓ Percent not specified  
- Event not reported

**Table 7. Boxed Warning for Metolazone<sup>1</sup>**

<b>WARNING</b>
Do not interchange Zaroxolyn <sup>®</sup> tablets and other formulations of metolazone that share its slow and incomplete bioavailability and are not therapeutically equivalent at the same doses to Mykrox <sup>®*</sup> tablets, a more rapidly available and completely bioavailable metolazone product. Formulations bioequivalent to Zaroxolyn <sup>®</sup> and formulations bioequivalent to Mykrox <sup>®*</sup> should not be interchanged for one another.

\* Mykrox<sup>®</sup> is no longer available in the United States.

## VII. Dosing and Administration

The usual dosing regimens for the thiazide-like diuretics are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Thiazide-Like Diuretics<sup>1-4</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
Chlorthalidone	<u>Edema:</u> Tablet: initial, 50 to 100 mg/day or 100 mg on alternate days; maintenance, 90 to 120 mg on alternate days or 120 mg/day  <u>Hypertension:</u> Tablet: initial, 12.5 to 25 mg/day; maintenance, 25 to 100 mg/day	Safety and efficacy in children have not been established.	Tablet: 15 mg 25 mg 50 mg
Indapamide	<u>Edema:</u> Tablet: initial, 2.5 mg/day; maintenance, 2.5 to 5 mg/day  <u>Hypertension:</u> Tablet: initial, 1.25 mg/day; maintenance, 2.5 to 5 mg/day	Safety and efficacy in children have not been established.	Tablet: 1.25 mg 2.5 mg
Metolazone	<u>Edema:</u> Tablet: 5 to 20 mg/day  <u>Hypertension:</u> Tablet: initial, 2.5 to 5 mg/day, maintenance, 5 to 20 mg/day	Safety and efficacy in children have not been established.	Tablet: 2.5 mg 5 mg 10 mg

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the thiazide-like diuretics are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Thiazide-Like Diuretics**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>SHEP Cooperative Research Group<sup>17</sup> and Kostis et al.<sup>18</sup> (1991 and 1995) SHEP</p> <p>Chlorthalidone 12.5 mg QD</p> <p>vs</p> <p>placebo</p> <p>Dosage was doubled for patients failing to achieve SBP goals. If SBP goal was not reached with chlorthalidone 25 mg QD, atenolol 25 mg QD or matching placebo was added to the drug regimen. Reserpine 0.05 mg QD or matching placebo was substituted in patients with contraindications</p>	<p>DB, MC, PC, RCT</p> <p>Patients aged ≥60 years with SBP between 160 and 219 mm Hg and DBP &lt;90 mm Hg</p>	<p>N=4,736</p> <p>Mean 4.5 years</p>	<p>Primary: Total stroke</p> <p>Secondary: Sudden or rapid cardiac death (defined as death within 1 hour or within 1 to 24 hours of the onset of severe cardiac symptoms), nonfatal or fatal MI, other cardiovascular death, TIA</p>	<p>Primary: With a mean follow-up of 4.5 years, the stroke occurred in 103 patients in the active treatment group compared to 159 patients in the placebo group (RR, 0.64; 95% CI, 0.5 to 0.82; P=0.0003).</p> <p>Stroke incidence was lower in patients taking active treatment compared to placebo in all baseline age groups: 60 to 69 years (34 vs 47 events, respectively), 70 to 79 years (48 vs 74 events, respectively), 80+ years (21 vs 38 events, respectively).</p> <p>The results were stratified according to whether patients had had previous antihypertensive therapy or not. In both stratified groups, there was a decrease in the risk of stroke with active treatment compared to placebo. For patients who were not receiving antihypertensive medication at initial contact, the RR, of stroke was 0.69 (95% CI, 0.51 to 0.95; P=0.02).</p> <p>For patients who had been receiving antihypertensive medication at initial contact, the RR, of stroke was 0.57 (95% CI, 0.38 to 0.85; P=0.01).</p> <p>Secondary: There were 23 sudden and 21 rapid deaths in the active treatment group compared to 23 sudden and 24 rapid deaths in the placebo group (RR, 1.0; 95% CI, 0.56 to 1.78 vs RR, 0.87; 95% CI, 0.48 to 1.56, respectively).</p> <p>There were 50 nonfatal and 15 fatal MIs in the active treatment group compared to 74 nonfatal and 26 fatal MIs in the placebo group (RR, 0.67; 95% CI, 0.47 to 0.96 vs RR, 0.57; 95% CI, 0.30 to 1.18, respectively).</p> <p>There were 21 other cardiovascular deaths in the active treatment group compared to 25 in the placebo group (RR, 0.87; 95% CI, 0.49 to 1.55).</p> <p>There were 62 TIAs in the active treatment group compared to 82 in the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
to atenolol.				<p>placebo group (RR, 0.75; 95% CI, 0.54 to 1.4).</p> <p>In the combined endpoints, the RR, of nonfatal MI or coronary heart disease death was 0.73 (95% CI, 0.57 to 0.94), CHD was 0.75 (95% CI, 0.60 to 0.94), cardiovascular disease was 0.68 (95% CI, 0.58 to 0.79).</p> <p>The RR, for atenolol were 0.84 (95% CI, 0.54 to 1.30) for death, 1.34 (95% CI, 0.80 to 2.28) for stroke, and 1.17 (95% CI, 0.71 to 1.61) for cardiovascular disease.</p> <p>The RR, for reserpine were 0.65 (95% CI, 0.26 to 1.59) for death, 0.27 (95% CI, 0.04 to 2.26) for stroke, and 0.55 (95% CI, 0.20 to 1.49) for cardiovascular disease.</p>
<p>ALLHAT Collaborative Research Group<sup>19</sup> (2000) ALLHAT</p> <p>Chlorthalidone 12.5 to 25 mg QD</p> <p>vs</p> <p>doxazosin 2 to 8 mg QD</p>	<p>AC, DB, MC, RCT</p> <p>Patients aged 55 years or older who had stage 1 or stage 2 HTN with ≥1 additional risk factor for CHD events (including previous MI or stroke &gt;6 months ago, left ventricular hypertrophy or echocardiography, history of type 2 diabetes, current cigarette smoking, high density lipoprotein cholesterol &lt;35 mg/dL, or documentation of other atherosclerotic cardiovascular</p>	<p>N=24,335</p> <p>Median 3.3 years</p>	<p>Primary: Fatal CHD or nonfatal MI combined</p> <p>Secondary: All cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease, cancer, end-stage renal disease</p>	<p>Primary: There was no significant difference in the primary outcome between doxazosin and chlorthalidone treatments (risk ratio, 1.13; 95% CI, 0.90 to 1.17; P=0.71).</p> <p>Secondary: Total mortality did not differ between the doxazosin and chlorthalidone treatments (four year rates, 9.62 and 9.08%, respectively; RR, 1.13; 95% CI, 0.90 to 1.15; P=0.56).</p> <p>The doxazosin group, compared with the chlorthalidone group, had a higher risk of stroke (RR 1.19; 95% CI, 1.01 to 1.40; P=0.04) and combined cardiovascular disease (four year rates 25.45 vs 21.76%; RR, 1.25; 95% CI, 1.17 to 1.33; P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	disease)			
Black et al. <sup>20</sup> (2008) ALLHAT  Chlorthalidone 12.5 to 25 mg QD  vs  amlodipine 2.5 to 10 mg QD  vs  lisinopril 10 to 40 mg QD	MC, RCT  Men and women, age 55 years old and older, with HTN and metabolic syndrome	N=17,515  4.9 years (mean)	Primary: Fatal coronary heart disease and nonfatal MI  Secondary: All cause mortality, fatal and nonfatal stroke, combined coronary heart disease, combined cardiovascular disease	Primary: For patients with metabolic syndrome, there was no significant difference in rates of coronary heart disease and nonfatal MI with amlodipine vs chlorthalidone (RR, 0.96; 95% CI, 0.79 to 1.16), or lisinopril vs chlorthalidone (RR, 1.15; 95% CI, 0.88 to 1.27).  Secondary: For patients with metabolic syndrome, there were no significant differences found between amlodipine vs chlorthalidone in all secondary endpoints (P value not significant).  For patients without metabolic syndrome, amlodipine treatment was associated with significantly more heart failure, but in patients with metabolic syndrome, there was no difference (P=0.03).  Patients with metabolic syndrome who received lisinopril experienced more heart failure and cardiovascular disease than those who received chlorthalidone (RR, 1.31; 95% CI, 1.04 to 1.64 and RR, 1.19; 95% CI, 0.97 to 1.32).
ALLHAT Collaborative Research Group <sup>21</sup> (2002) ALLHAT  Chlorthalidone 12.5 to 25 mg/day  vs  amlodipine 2.5 to 10 mg/day  vs  lisinopril 10 to 40	DB, MC, RCT  Patients ≥55 years with HTN and ≥1 additional CHD risk factor	N=33,357  4.9 years (mean)	Primary: Combined fatal CHD or nonfatal MI  Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization,	Primary: There were no significant differences in the primary outcome between lisinopril (11.4%), amlodipine (11.3%), and chlorthalidone (11.5%).  Secondary: All-cause mortality did not differ between groups.  Five year SBPs were significantly higher in the lisinopril (2 mm Hg; P<0.001) and amlodipine groups (0.8 mm Hg; P=0.03) compared to chlorthalidone, and five year DBPs were significantly lower with amlodipine (0.8 mm Hg; P<0.001).  Amlodipine had a higher six year rate of heart failure compared to chlorthalidone (10.2 vs 7.7%; RR, 1.38; 95% CI, 1.25 to 1.52).  Lisinopril had a higher six year rate of combined cardiovascular disease (33.3 vs 30.9%; RR, 1.10; 95% CI, 1.05 to 1.16); stroke (6.3 vs 5.6%; RR,



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day			heart failure, and PAD)	1.15; 95% CI, 1.02 to 1.30) and heart failure (8.7 vs 7.7%; RR, 1.19; 95% CI, 1.07 to 1.31).
Rahman et al. <sup>22</sup> (2012) ALLHAT  Chlorthalidone 12.5 to 25 mg/day  vs  amlodipine 2.5 to 10 mg/day  vs  lisinopril 10 to 40 mg/day	Long-term, post-trial, follow-up  Patients in ALLHAT stratified based on eGFR	N=31,350  4 to 8 years	Primary: Cardiovascular mortality  Secondary: Total mortality, CHD, cardiovascular disease, stroke, heart failure, ESRD	Primary: After an average of 8.8 years of follow-up, total mortality was significantly higher in patients with moderate/severe eGFR reduction (eGFR <60 mL/min/1.73 m <sup>2</sup> ) compared to patients with normal/increased (eGFR ≥90 mL/min/1.73 m <sup>2</sup> ) and mildly reduced eGFR (eGFR 60 to 89 mL/min/1.73 m <sup>2</sup> ) (P<0.001).  In patients with moderate/severe eGFR reduction, there was no significant difference in cardiovascular mortality between chlorthalidone and amlodipine (P=0.64), or chlorthalidone and lisinopril (P=0.56).  Secondary: No significant differences were observed for any of the secondary endpoints among eGFR reduction groups.
Muntner et al. <sup>23</sup> (2014) ALLHAT  Chlorthalidone 12.5 to 25 mg/day  vs  amlodipine 2.5 to 10 mg/day  vs  lisinopril 10 to 40 mg/day	Post-hoc analysis of ALLHAT  Patients in ALLHAT with 5, 6, or 7 visits in 6 to 28 months of follow-up	N=24,004  6 to 28 months	Primary: Visit-to-visit variability (VVV) of blood pressure  Secondary: Not reported	Primary: Each measure of VVV of SBP was lower among participants randomized to chlorthalidone and amlodipine compared with those randomized to lisinopril. All four VVV of SBP metrics were lower among participants randomized to amlodipine vs chlorthalidone after full multivariable adjustment.  After multivariable adjustment including mean SBP across visits and compared with participants randomized to chlorthalidone, participants randomized to amlodipine had a 0.36 (standard error [SE]: 0.07) lower standard deviation (SD) of SBP and participants randomized to lisinopril had a 0.77 (SE=0.08) higher SD of SBP. Results were consistent using other VVV of SBP metrics. These data suggest chlorthalidone and amlodipine are associated with lower VVV of SBP than lisinopril.  Secondary: Not reported

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Bangalore et al.<sup>24</sup> (2017) ALLHAT</p> <p>Chlorthalidone 12.5 to 25 mg/day</p> <p>vs</p> <p>amlodipine 2.5 to 10 mg/day</p> <p>vs</p> <p>lisinopril 10 to 40 mg/day</p>	<p>Post-hoc analysis of ALLHAT</p> <p>Patients in ALLHAT with average blood pressure <math>\geq 140</math> mmHg systolic or <math>\geq 90</math> mm Hg diastolic on <math>\geq 3</math> antihypertensive medications, or blood pressure <math>&lt; 140/90</math> mmHg on <math>\geq 4</math> antihypertensive medications (i.e., identified as having apparent treatment-resistant hypertension) at 2-year follow up</p>	<p>N=14,684</p> <p>4.9 years (mean)</p>	<p>Primary: Combined fatal CHD or nonfatal MI</p> <p>Secondary: All-cause mortality, fatal and nonfatal stroke, combined CHD, combined cardiovascular disease (combined CHD, stroke, treated angina without hospitalization, heart failure, and PAD)</p>	<p>Primary: Of participants assigned to chlorthalidone, amlodipine, or lisinopril, 9.6%, 11.4%, and 19.7%, respectively, had treatment-resistant hypertension. During mean follow-up of 2.9 years, primary outcome incidence was similar for those assigned to chlorthalidone compared with amlodipine or lisinopril (amlodipine- vs chlorthalidone-adjusted HR, 0.86; 95% CI, 0.53 to 1.39; P=0.53; lisinopril- vs chlorthalidone-adjusted HR, 1.06; 95% CI, 0.70 to 1.60; P=0.78).</p> <p>Secondary: Secondary outcome risks were similar for most comparisons except coronary revascularization, which was higher with amlodipine than with chlorthalidone (HR, 1.86; 95% CI, 1.11 to 3.11; P=0.02). An as-treated analysis based on diuretic use produced similar results.</p>
<p>Pupita et al.<sup>25</sup> (1983)</p> <p>Chlorthalidone 50 mg QD</p> <p>vs</p> <p>furosemide 25 mg QD</p>	<p>RCT, XO</p> <p>Men and women with a mean age of <math>53.9 \pm 9.2</math> years with mild to moderate HTN</p>	<p>N=36</p> <p>12 months</p>	<p>Primary: Blood pressure</p> <p>Secondary: Plasma electrolytes, adverse events</p>	<p>Primary: Patients taking chlorthalidone had significantly lower SBP at each monthly measurement compared to baseline (P&lt;0.01). However, only DBP values at month five were significant compared to baseline (P&lt;0.05).</p> <p>Patients taking furosemide had significantly lower SBP at months three, four, and five compared to baseline (P&lt;0.05 for month three, and P&lt;0.01 for months four and five). DBP values were significantly lower at all monthly measurements compared to baseline in patients taking furosemide (P&lt;0.01).</p> <p>At month one, SBP decreased by 19.4 mm Hg with chlorthalidone and by 21.2 mm Hg with furosemide (P&lt;0.001). DBP decreased by 11 mm Hg with chlorthalidone and by 12.6 mm Hg with furosemide at month one (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Secondary: There were no significant changes in serum sodium levels with either chlorthalidone or furosemide. Patients taking chlorthalidone had significantly lower serum chloride levels compared to baseline at all points (P&lt;0.01), whereas patients taking furosemide had significantly lower levels only at month six (P&lt;0.05). Both chlorthalidone and furosemide significantly reduced serum potassium levels at all points compared to baseline (P&lt;0.01).</p> <p>Patient taking chlorthalidone reported adverse effects including dizziness, transient abdominal disorder, and slight weakness. Patients taking furosemide reported transient early weakness and irritability. The rate of adverse events was not statistically significant in either treatment group.</p>
<p>Bakris et al.<sup>26</sup> (2012)</p> <p>Azilsartan medoxomil and chlorthalidone (single pill)</p> <p>vs</p> <p>azilsartan medoxomil and HCTZ (co-administered)</p> <p>Treatments were titrated to a target of &lt;140/90 mm Hg (or &lt;130/80 mm Hg if diabetes of chronic kidney disease)</p>	<p>DB, RCT</p> <p>Patients aged ≥18 years with stage 2 primary HTN</p>	<p>N=609</p> <p>10 weeks (after 2 week placebo run-in)</p>	<p>Primary: Change in trough, seated clinic systolic blood pressure at weeks 6 and 10</p> <p>Secondary: Change from baseline in clinic DBP and 24-hour mean systolic and diastolic blood pressures by ambulatory blood pressure monitoring</p>	<p>Primary: Change in SBP at week six demonstrated a mean difference of -5.6 mm Hg (95% CI, -8.3 to -2.9; P&lt;0.001) in favor of the chlorthalidone group. Fewer patients in the chlorthalidone group required titration to a higher dose of diuretic (P&lt;0.001). At the end of week 10, a greater mean SBP reduction was maintained in the chlorthalidone group compared to the HCTZ group (-5.0 mm Hg; 95% CI, -7.5 to -2.5; P&lt;0.001).</p> <p>Secondary: The chlorthalidone group demonstrated a significantly greater reduction in 24-hour mean SBP at weeks six and 10. For both clinica and 24-hour mean DBP, greater blood pressure reduction was observed in the chlorthalidone group compared to the HCTZ group at both study points.</p>
<p>Ernst et al.<sup>27</sup> (2006)</p>	<p>RCT, SB, XO</p>	<p>N=30</p>	<p>Primary: Comparison of the</p>	<p>Primary: At week eight, there was a greater reduction in 24-hour mean SBP with</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Chlorthalidone 12.5 mg in the morning</p> <p>vs</p> <p>HCTZ 25 mg in the morning</p> <p>At week 4, both HCTZ and chlorthalidone were titrated to 50 mg in the morning and 25 mg in the morning, respectively for the remainder of the trial.</p>	<p>Men and women aged 18 to 79 years with pre-HTN or a new or established diagnosis of HTN (stage 1 or 2), not receiving antihypertensive medications, and had an average office blood pressure value in the last 6 months between 140 and 179 mm Hg systolic or 90 and 109 mm Hg diastolic</p>	<p>8 weeks plus 4 week washout period</p>	<p>change in 24-hour mean SBP and DBP from baseline to week 8</p> <p>Secondary: Comparison of changes in mean SBP and mean DBP for office blood pressure at each visit, change in ambulatory daytime and nighttime mean SBP and DBP from baseline to week 8, development of hypokalemia</p>	<p>chlorthalidone 25 mg/day compared to HCTZ 50 mg/day compared to baseline (-12.4±1.8 vs -7.4±1.7 mm Hg, respectively; P=0.054).</p> <p>Secondary: There was a trend in favor of greater reduction in SBP with chlorthalidone than with HCTZ at each office visit. However, the difference was only statistically significant at week 2 (-15.7±2.2 vs -4.5±2.1 mm Hg, respectively; P=0.001).</p> <p>Although mean reductions in DBP was also greater with chlorthalidone compared to HCTZ at each study visit, the differences were not statistically significant at any visit (P&gt;0.89 for all).</p> <p>The reduction in SBP during nighttime hours was -13.5±1.9 mm Hg for chlorthalidone and -6.4±1.7 mm Hg for HCTZ (P=0.009). The reduction in daytime mean SBP between both groups was not significantly different (-11.4±2.0 vs -8.1±1.9 mm Hg, respectively; P=0.230).</p> <p>Changes in serum potassium were similar between treatment groups (P=0.76). The incidence of hypokalemia was 50% in patients taking HCTZ and 46% in patients taking chlorthalidone (P=0.682).</p>
<p>Carter et al.<sup>28</sup> (2004)</p> <p>Chlorthalidone 12.5 to 600 mg/day</p> <p>vs</p> <p>HCTZ 12.5 to 450 mg/day</p>	<p>MA</p> <p>Included trials which evaluate the pharmacokinetic and blood pressure lowering effects of chlorthalidone and HCTZ</p>	<p>N=200</p> <p>Duration varied per study</p>	<p>Primary: Blood pressure</p> <p>Secondary: Serum potassium</p>	<p>Primary: In a dose equivalence study comparing HCTZ 100 mg QD to chlorthalidone 50 mg QD, blood pressure (SBP/DBP) reduced by 18/8 and 25/10 mm Hg compared to baseline, respectively.</p> <p>In another study comparing HCTZ 25 mg and triamterene 50 mg QD, HCTZ 50 mg and triamterene 100 mg QD, and chlorthalidone 50 mg QD, the blood pressure reduction was 15/8, 18/12, and 25/16 mm Hg, respectively.</p> <p>One other dose equivalence study comparing HCTZ 50 mg BID and chlorthalidone 50 mg QD, blood pressure reduction was 22/16 and 18/15 mm Hg, respectively.</p> <p>All available studies were inspected and it was concluded that HCTZ 50 mg is approximately equivalent to chlorthalidone 25 to 37 mg.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Furthermore, it was suggested that chlorthalidone doses should generally be approximately 50% to 75% of the typical HCTZ dose.</p> <p>Secondary: In a study comparing HCTZ 100 mg QD and chlorthalidone 50 mg QD, potassium increased slightly with chlorthalidone (0.02 mEq/L) and decreased significantly with HCTZ (0.22 mEq/L; P=0.009).</p> <p>However, in another study comparing HCTZ 50 mg BID and chlorthalidone 50 mg QD, serum potassium decreased by 0.38 mEq/L with HCTZ and by 0.03 mEq/L with chlorthalidone. The difference was not statistically significant (P&lt;0.07).</p>
<p>Karotsis et al.<sup>29</sup> (2006)</p> <p>Chlorthalidone 12.5 mg QD</p> <p>vs</p> <p>felodipine 5 mg QD</p> <p>vs</p> <p>lisinopril 10 mg QD</p> <p>vs</p> <p>valsartan 80 mg QD</p> <p>All patients also received diltiazem 240 mg QD.</p>	<p>RCT</p> <p>Patients 25 to 79 years of age with uncontrolled HTN (average office blood pressure &gt;140/90 mm Hg for all or &gt;153/85 mm Hg for diabetics or patients &lt;65 years of age, confirmed on 2 office visits ≥1 week apart) after ≥4 weeks of OL monotherapy with diltiazem at 240 mg QD</p>	<p>N=211</p> <p>8 weeks</p>	<p>Primary: Blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: There was a significant decline in both office and home SBP and DBP during the trial with all treatments. The antihypertensive effect was more pronounced and reached significance when home blood pressure monitoring was used in comparison to office blood pressure without the white-coat effect (P&lt;0.001 for all blood pressure changes). With or without the white-coat effect, blood pressure still declined and the differences were significant (P&lt;0.0001 for all blood pressure changes).</p> <p>Secondary: Not reported</p>
<p>Nissinen et al.<sup>30</sup></p>	<p>DB, RCT</p>	<p>N=23</p>	<p>Primary:</p>	<p>Primary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1980)</p> <p>Atenolol 100 mg QD plus chlorthalidone 25 mg in the morning</p> <p>vs</p> <p>atenolol and chlorthalidone 100-25 mg in the morning (fixed-dose combination product)</p> <p>vs</p> <p>placebo</p>	<p>Patients with newly diagnosed mild to moderate HTN (supine DBP 100 mm Hg on <math>\geq 3</math> occasions)</p>	<p>16 weeks</p>	<p>Changes in blood pressure and heart rate</p> <p>Secondary: Not reported</p>	<p>Each of the active drug combinations lowered standing, supine, and post-exercise blood pressure significantly compared to placebo at two and four weeks (<math>P &lt; 0.001</math>, <math>P &lt; 0.01</math> and <math>P &lt; 0.05</math>). There was not a statistical difference between the active treatment regimens (<math>P</math> value not significant).</p> <p>Each of the active drug combinations lowered standing, supine, and post-exercise heart rate significantly compared to placebo at two and four weeks (<math>P &lt; 0.001</math>, <math>P &lt; 0.01</math> and <math>P &lt; 0.05</math>). There was not a statistical difference between the active treatment regimens (<math>P</math> value not significant).</p> <p>Side effects did not differ between treatment groups and placebo in terms of frequency or severity. Reported side effects included dizziness, headache and tiredness.</p> <p>Secondary: Not reported</p>
<p>Fogari et al.<sup>31</sup> (1984)</p> <p><u>Weeks 1 to 4:</u> chlorthalidone 12.5 mg QD</p> <p>vs</p> <p>atenolol 50 mg QD</p> <p><u>Weeks 5 to study end:</u> atenolol and chlorthalidone 50-12.5 mg QD (fixed-dose combination</p>	<p>RCT, SB</p> <p>Patients 61 to 80 years inadequately controlled (SBP <math>&gt; 170</math> mm Hg and/or DBP <math>&gt; 100</math> mm Hg) on antihypertensive medications</p>	<p>N=38</p> <p>6 months</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary: After the first four weeks, atenolol (from 177.5 to 161.1 mm Hg) significantly reduced blood pressure compared to baseline, but chlorthalidone did not (from 176.6 to 179.1 mm Hg).</p> <p>The combination atenolol-chlorthalidone therapy significantly reduced mean standing SBP and DBP, supine SBP and DBP, supine and standing heart rate, compared to previous therapies (<math>P &lt; 0.001</math> for all comparisons).</p> <p>The combination atenolol-chlorthalidone therapy significantly reduced mean standing SBP and DBP, supine SBP and DBP, supine and standing heart rate, compared to atenolol and chlorthalidone monotherapy (<math>P &lt; 0.001</math> or <math>P &lt; 0.01</math> for all comparisons).</p> <p>Mean blood pressure reduction obtained by the atenolol and chlorthalidone combination product was 30/15 mm Hg in the standing position (<math>P &lt; 0.001</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
product)				<p>Serum potassium increased with atenolol-chlorthalidone (4.45 mEq/L) compared to chlorthalidone alone (4.01 mEq/L; P&lt;0.001).</p> <p>Secondary: Not reported</p>
<p>Leonetti et al.<sup>32</sup> (1986)</p> <p>Atenolol 50 mg QD</p> <p>vs</p> <p>atenolol 100 mg QD</p> <p>vs</p> <p>chlorthalidone 12.5 mg QD</p> <p>vs</p> <p>atenolol and chlorthalidone 50-12.5 mg QD (fixed-dose combination product)</p>	<p>DB, RCT</p> <p>Patients 24 to 68 years with mild to moderate HTN (WHO stage I or II), with supine DBP ≥95 mm Hg at the end of the 4-week washout period</p>	<p>N=28</p> <p>16 weeks</p>	<p>Primary: Changes in blood pressure</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>Mean supine blood pressure was significantly reduced in all treatment groups compared to placebo: 153±18/93±9 mm Hg for atenolol 50 mg patients, 155±22/91±8 mm Hg for atenolol 100 mg patients, 148±17/93±11 mm Hg for chlorthalidone 12.5 mg patients, and 144±16/89±6 mm Hg for the atenolol-chlorthalidone combination patients. All of the changes in blood pressure were significant (P&lt;0.01) versus placebo.</p> <p>Supine SBP was lower with atenolol-chlorthalidone than with the atenolol 100 mg alone (P&lt;0.05).</p> <p>Upright SBP was lower with atenolol-chlorthalidone than with atenolol 50 mg alone (P&lt;0.05) and atenolol 100 mg alone (P&lt;0.05).</p> <p>Mean supine heart rate was 77±7 bpm after placebo which decreased to 69±10 bpm (P&lt;0.01) after atenolol 50 mg, to 67±6 bpm (P&lt;0.01) after atenolol 100 mg, to 77±10 bpm (P=not significant, was not reported) after chlorthalidone alone.</p> <p>Chlorthalidone alone demonstrated a significant reduction in serum potassium levels compared to placebo (3.88 vs 4.09 mEq/L; P&lt;0.05) and no change when the atenolol-chlorthalidone combination was compared to placebo (3.98 vs 4.09; P=not significant, value was not reported).</p> <p>Chlorthalidone alone and atenolol-chlorthalidone demonstrated a significant increase in serum uric acid levels compared to placebo (4.90±1.52 mg/dL, 5.07±1.33 mg/dL, respectively, vs 4.24±1.12 for placebo; P&lt;0.05 for both).</p> <p>All treatments were well tolerated. Some adverse events reported included dyspnea, precordial discomfort and cold extremities. Incidence, severity</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				and P values were not reported.
Finnerty et al. <sup>33</sup> (1980)  Chlorthalidone 50 mg plus reserpine 0.25 mg  vs  HCTZ 50 mg plus reserpine 0.125 mg	DB  Patients with essential HTN unresponsive to diet control and diuretic therapy	N=57  6 weeks	Primary: The change in mean DBP from baseline  Secondary: Incidence of frequent or severe side effects	Primary: The chlorthalidone plus reserpine group had a mean decrease in DBP of 17.0 mm Hg at study endpoint compared with a mean decrease of 18.6 mm Hg in the HCTZ plus reserpine group.  At study completion both treatment groups achieved diastolic control of at least 5 mm Hg below the targeted diastolic goal of 90 mm Hg.  Secondary: There were no reports of frequent or severe side effects in either treatment group.
Akram et al. <sup>34</sup> (2007) NATIVE  Indapamide SR 1.5 mg QD added to background antihypertensive therapy	OL  Patients remaining hypertensive (145 to 180/95 to 105 mm Hg) while receiving an ACE inhibitor, $\beta$ -blocker, calcium-channel blocker, ARBs, $\alpha$ -blocker, or other therapy	N=1,941  3 months	Primary: Blood pressure  Secondary: Glucose and cholesterol levels	Primary: At three months, SBP and DBP both decreased significantly compared to baseline. SBP had a change from 166 $\pm$ 16 mm Hg at baseline to 132 $\pm$ 12 mm Hg at three months. DBP had a change from 102 $\pm$ 8 mm Hg at baseline to 83 $\pm$ 6 mm Hg at three months (P<0.0001 for both).  At study end, 84% of patients achieved target SBP of $\leq$ 140 mm Hg and 61% achieved blood pressure normalization (SBP/DBP <140/90 mm Hg).  Secondary: Glucose and cholesterol levels were unaffected by indapamide SR.
Beckett et al. <sup>35</sup> (2008) HYVET  Indapamide 1.5 mg/day  vs  placebo  Perindopril 2 to 4 mg/day or matching placebo	DB, MC, PC, RCT  Patients $\geq$ 80 years (mean age 84 years) with sustained SBP $\geq$ 160 mm Hg	N=3,845  1.8 years (mean)	Primary: Fatal or nonfatal stroke  Secondary: Death from any cause, death from cardiovascular causes, death from stroke	Primary: At two years, 73.4% of patients in the active-treatment groups were receiving indapamide plus perindopril. Mean blood pressure while sitting was 15.0/6.1 mm Hg lower with active-treatment than placebo.  Active treatment was associated with a 30% reduction in the rate of fatal or nonfatal stroke (95% CI, -1 to 51; P=0.06).  Secondary: Active treatment was associated with a 21% reduction in the rate of death from any cause (95% CI, 4 to 35; P=0.02), a 23% reduction in the rate of death from cardiovascular causes (95% CI, -1 to 40; P=0.06) and a 39% reduction in the rate of death from stroke (95% CI, 1 to 62; P=0.05).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
was added if necessary to achieve the target blood pressure of 150/80 mm Hg.				<p>Active treatment was associated with a 64% reduction in the rate of heart failure (95% CI, 42 to 78; P&lt;0.001).</p> <p>Fewer serious adverse events were reported in the active-treatment group (358 vs 448; P=0.001).</p>
<p>Milia et al.<sup>36</sup> (2006)</p> <p>Indapamide 2.5 mg QD</p> <p>vs</p> <p>bendroflumethiazide* 2.5 mg QD</p>	<p>DB, PG, PRO, RCT</p> <p>Ambulant patients with a first-ever minor hemispheric ischemic stroke or TIA</p>	<p>N=26</p> <p>28 days</p>	<p>Primary: Blood pressure, cerebral blood flow</p> <p>Secondary: Not reported</p>	<p>Primary: Both indapamide and bendroflumethiazide significantly reduced blood pressure from baseline (-14.7±12.5 mm Hg and -7.7±9.16 mm Hg, respectively; P&lt;0.001 and P=0.02, respectively).</p> <p>A nonsignificant trend toward greater blood pressure reduction was seen in patients taking indapamide. There were no statistically significant differences in blood pressure reduction between both treatment groups.</p> <p>There was a nonsignificant trend toward increases in blood flow in both treatment groups. However, there was no statistically significant differences in carotid blood flow between both treatment groups (P=0.04 for between-group comparison).</p> <p>Secondary: Not reported</p>
<p>Madkour et al.<sup>37</sup> (1996)</p> <p>Indapamide 2.5 mg QD</p> <p>vs</p> <p>HCTZ 50 mg QD</p>	<p>RCT</p> <p>Patients aged 32 to 70 years with impaired renal function for 1 to 15 years and moderate HTN for 2 to 27 years, initial creatinine clearance between 32 and 80 mL/min/1.73 m<sup>2</sup> BSA</p>	<p>N=28</p> <p>24 months</p>	<p>Primary: Blood pressure, changes in creatinine clearance</p> <p>Secondary: Not reported</p>	<p>Primary: Blood pressure normalized in all patients taking either indapamide or HCTZ. There were no significant differences in SBP or DBP between groups.</p> <p>At 24 months, creatinine clearance progressively increased from 58±4.4 to 72±4.4 mL/min/1.73 m<sup>2</sup> BSA in patients treated with indapamide (P&lt;0.01).</p> <p>Creatinine clearance progressively decreased from 65±3.0 to 53±3.0 mL/min/1.73 m<sup>2</sup> BSA in patients treated with HCTZ (P&lt;0.01). Creatinine clearance significantly increased by 28.5±4.4% with indapamide and decreased by 17.4±3.0% with thiazide therapy (P&lt;0.01).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>PROGRESS<sup>38</sup> (2001)</p> <p>Perindopril 4 mg/day</p> <p>vs</p> <p>perindopril 4 mg/day and indapamide 2 to 2.5 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients with a history of prior stroke or TIA within the previous 5 years</p>	<p>N=6,105</p> <p>4 years</p>	<p>Primary: Fatal or nonfatal stroke</p> <p>Secondary: Fatal or disabling stroke, total major vascular events comprising the composite of nonfatal stroke, nonfatal MI, or death due to any vascular cause (including unexplained sudden death); total and cause specific deaths; hospital admissions</p>	<p>Primary: Patients receiving active treatment experienced a 28% reduction in nonfatal or fatal stroke (95% CI, 17 to 38; P&lt;0.0001).</p> <p>There were similar reductions in the risk of stroke in hypertensive and non-hypertensive subgroups (32 vs 27%; P&lt;0.01)</p> <p>A trend towards a greater effect of active treatment among patients treated with combination therapy (43% risk reduction) than in those treated with single drug therapy (5% risk reduction) was reported.</p> <p>Secondary: There was a 33% reduction in fatal or disabling strokes in the active treatment group.</p> <p>Active treatment reduced the risk of total major vascular events by 26% (P=0.02).</p> <p>There were no significant differences between active treatment and placebo in total deaths from vascular or nonvascular causes.</p> <p>Among those assigned active treatment, there was a 9% RR reduction in hospitalization, with a median reduction of 2.5 days in the time spent in the hospital during follow-up.</p> <p>Combination therapy with perindopril plus indapamide reduced blood pressure by 12/5 mm Hg and stroke risk by 43%. Single drug therapy reduced blood pressure by 5/3 mm Hg and produced no discernible reduction in the risk of stroke.</p>
<p>Hua et al.<sup>39</sup> (1976)</p> <p>Metolazone 5 mg QD</p> <p>vs</p>	<p>XO</p> <p>Patients with HTN</p>	<p>N=20</p> <p>Duration not specified</p>	<p>Primary: Blood pressure, serum potassium</p> <p>Secondary: Not reported</p>	<p>Primary: Blood pressures on metolazone tended to be lower than on chlorothiazide, but the difference was not statistically significant.</p> <p>Both agents significantly lowered serum potassium concentrations and total body potassium to a similar degree. However, the serum potassium did not fall below the normal range in any patient and no potassium supplements were required.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
chlorothiazide up to 5 g BID				Secondary: Not reported
ADVANCE Collaborative Group <sup>40</sup> (2007)  Perindopril (2 to 4 mg) and indapamide (0.625 to 1.25 mg) QD  vs  placebo	DB, MC, PC, RCT  Adults 55 years of age or older who were diagnosed with type 2 diabetes at age 30 or older, and a history of cardiovascular disease or ≥1 other risk factor for cardiovascular disease	N=11,140  Mean 4.3 years	Primary: Composites of major macrovascular and microvascular events (death from cardiovascular disease, nonfatal stroke, nonfatal MI, or new renal or diabetic eye disease)  Secondary: Macrovascular and microvascular endpoints analyzed separately	Primary: The relative risk of a major macrovascular or microvascular event was reduced by 9% (861 [15.5%] active vs 938 [16.8%] placebo; HR, 0.91, 95% CI 0.83 to 1.0, P=0.04).  Secondary: The RR of death from cardiovascular disease was reduced by 18% (211 [3.8%] active vs 257 [4.6%] placebo; 0.82, 0.68-0.98, p=0.03) and death from any cause was reduced by 14% (408 [7.3%] active vs 471 [8.5%] placebo; 0.86, 0.75-0.98, P=0.03).
Hansson et al. <sup>41</sup> (2000) NORDIL  Conventional therapy (diuretic, β-blocker or both)  vs  diltiazem 180 to 360 mg QD	BE, MC, OL, PRO, RCT  Patients 50 to 74 years of age with DBP ≥100 mm Hg and previously untreated	N=10,881  4.5 years	Primary: Combined fatal and nonfatal stroke, fatal and nonfatal MI, other cardiovascular death  Secondary: Fatal plus nonfatal stroke and fatal plus nonfatal MI	Primary: The primary endpoint occurred in 403 of the diltiazem patients and 400 of the diuretic/β-blocker patients (RR, 1.00; 95% CI, 0.87 to 1.15; P=0.97).  Secondary: Rates of secondary endpoints were similar between the groups. Fatal plus nonfatal stroke occurred in 159 of the diltiazem patients and 196 of the diuretic/β-blocker patients (P=0.04).  Fatal plus nonfatal MI occurred in 183 of the diltiazem patients and 157 of the diuretic/β-blocker patients (P=0.17).  Other endpoints were not statistically different between the groups including cardiovascular death (P=0.41), all cardiac events (P=0.57) and congestive heart failure (P=0.42).
Ames <sup>42</sup>	MA (13 trials)	N=1,547	Primary:	Primary:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(1996)</p> <p>Indapamide 2.5 mg QD</p> <p>vs</p> <p>HCTZ <math>\leq</math>25 mg or its equivalent in other thiazides, up to 112.5 mg QD</p>	<p>Patients with HTN</p>	<p>1 to 25 months</p>	<p>Comparison of the effects of thiazides and indapamide on blood lipids and blood pressure</p> <p>Secondary: Not reported</p>	<p>The mean change from baseline was 1.4% for TC, 5.5% for HDL-C, and -0.5% for TG with indapamide. None of the differences were statistically significant.</p> <p>Low-dose thiazide therapy did not decrease TC at any data point. The mean percent increase in TC was 3.8%, in HDL-C was 3.1%, and in TG was 10.8% with low-dose HCTZ. The increases in TC and TG from baseline was statistically significant (P&lt;0.01).</p> <p>The mean change in TC was 6.3%, in HDL-C was -0.5%, and in TGs was 19.5% for higher doses of HCTZ. Increases from baseline in TC and TG were statistically significant.</p> <p>SBP decreased more with higher doses of HCTZ than with low-dose thiazide therapy (P&lt;0.05). The effects of indapamide on systolic arterial pressure were intermediate between, and not statistically different from, either thiazide dose. Decreases in DBP did not differ among groups.</p> <p>Secondary: Not reported</p>
<p>Messerli et al.<sup>43</sup> (1998)</p> <p>Diuretics (amiloride, chlorthalidone, HCTZ, HCTZ and triamterene [fixed-dose combination product], or thiazide)</p> <p>vs</p> <p><math>\beta</math>-blockers (atenolol, metoprolol or</p>	<p>MA</p> <p>10 RCTs lasting <math>\geq</math>1 year, which used as first line agents diuretics and/or <math>\beta</math>-blockers and reported morbidity and mortality outcomes in patients <math>\geq</math>60 years of age with HTN</p>	<p>N=16,164</p> <p>1 year</p>	<p>Primary: Cardiovascular morbidity and mortality, all-cause morbidity</p> <p>Secondary: Not reported</p>	<p>Primary: Diuretic treatment significantly reduced the odds for cardiovascular mortality by 25% (OR, 0.75; 95% CI, 0.64 to 0.87), while <math>\beta</math>-blockers did not reduce cardiovascular mortality (OR, 0.98; 95% CI, 0.78 to 1.23; P values not reported).</p> <p>Diuretic treatment significantly reduced the odds for all-cause mortality by 14% (OR, 0.86; 95% CI, 0.77 to 0.96), while <math>\beta</math>-blockers did not reduce all-cause mortality (OR, 1.05; 95% CI, 0.88 to 1.25; P values not reported).</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>pindolol)</p> <p>Baguet et al.<sup>44</sup> (2007)</p> <p>Antihypertensive drugs (enalapril, ramipril, trandolapril, candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, HCTZ, indapamide SR*, atenolol, amlodipine, lercanidipine*, manidipine*, enalapril, ramipril, trandolapril, and aliskiren)</p> <p>Drugs were used as monotherapy, either at a fixed daily dosage or in increasing dosages.</p> <p>Although cicletanine*, furosemide and spironolactone were considered for inclusion, none of the trials</p>	<p>MA</p> <p>Patients greater than 18 years of age with mild or moderate essential HTN (SBP 140 to 179 mm Hg and/or DBP 90 to 109 mm Hg)</p>	<p>N=10,818</p> <p>8 to 12 weeks</p>	<p>Primary: Weighted average reductions in SBP and DBP</p> <p>Secondary: Not reported</p>	<p>Primary: Data did not reflect outcomes from direct, head-to-head comparative trials or formal comparisons between drugs. Diuretics (-19.2 mm Hg; 95% CI, -20.3 to -18.0), calcium channel blockers (-16.4 mm Hg; 95% CI, -17.0 to -15.8) and ACE inhibitors (-15.6 mm Hg; 95% CI, -17.6 to -13.6) produced the greatest reductions in SBP from baseline (P values not reported).</p> <p>The magnitude of DBP reductions were generally similar among all drug classes; however, the greatest reductions in DBP from baseline were observed with the <math>\beta</math>-blocker, atenolol (-11.4 mm Hg; 95% CI, -12.0 to -10.9), calcium channel blockers (-11.4 mm Hg; 95% CI, -11.8 to -11.1) and diuretics (-11.1 mm Hg; 95% CI, -11.7 to -10.5) (P values were not reported).</p> <p>The weighted average reduction of SBP and DBP for each drug class were as follows:  Diuretics: -19.2 (95% CI, -20.3 to -18.0) and -11.1 mm Hg (95% CI, -11.7 to -10.5), respectively.  <math>\beta</math>-blockers: -14.8 (95% CI, -15.9 to -13.7) and -11.4 mm Hg (95% CI, -12.0 to -10.9), respectively.  Calcium channel blockers: -16.4 (95% CI, -17.0 to -15.8) and -11.4 mm Hg (95% CI, -11.8 to -11.1), respectively.  ACE inhibitors: -15.6 (95% CI, -17.6 to -13.6) and -10.8 mm Hg (95% CI, -11.9 to -9.7), respectively.  ARBs: -13.2 (95% CI, -13.6 to -12.9) and -10.3 mm Hg (95% CI, -10.5 to -10.1), respectively.  Renin inhibitor: -13.5 (95% CI, -14.2 to -12.9) and -11.3 mm Hg (95% CI, -11.7 to -10.9), respectively.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
relating to these agents satisfied all inclusion criteria.				

\*Agent not available in the United States.

Drug regimen abbreviations: BID=twice daily, QD=once daily, SR=systematic review

Study design abbreviations: BE=blinded endpoint, DB=double blind, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, PG=parallel group, PRO=prospective, RCT=randomized controlled trial, SB=single blind, XO=cross over

Miscellaneous abbreviations=ACE inhibitors=angiotensin converting enzyme inhibitors, ARB=angiotensin II receptor blocker, BSA=body surface area, CHD=coronary heart disease, CI=confidence interval, DBP=diastolic blood pressure, HCTZ=hydrochlorothiazide, HR=hazard ratio, HTN=hypertension, MI=myocardial infarction, PAD=peripheral artery disease, RR=relative risk, SBP=systolic blood pressure, TC=total cholesterol, TG=triglycerides, TIA=transient ischemic attack, WHO=World Health Organization

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Thiazide-Like Diuretics**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Chlorthalidone	tablet*	Thalitone®	\$\$\$\$	\$
Indapamide	tablet*	N/A	N/A	\$
Metolazone	tablet*	N/A	N/A	\$\$

\*Generic is available in at least one dosage form or strength.  
N/A=Not available

**X. Conclusions**

The thiazide-like diuretics are approved for the treatment of hypertension and edema associated with congestive heart failure. Chlorthalidone and metolazone are also indicated for the treatment of edema due to renal dysfunction. Additionally, chlorthalidone is approved for the adjunctive treatment of edema associated with hepatic cirrhosis, as well as corticosteroid and estrogen therapy.<sup>1-4</sup> All of the agents are available in a generic formulation.

Guidelines recommend the use of diuretics and sodium restriction for the management of ascites due to cirrhosis. Spironolactone is recommended as first-line therapy, either as monotherapy or in combination with furosemide. Amiloride or eplerenone are alternative treatment options in patients experiencing gynecomastia with spironolactone.<sup>16</sup>

For the treatment of chronic heart failure, guidelines recommend the use of diuretics in all patients who have evidence of volume overload. Loop diuretics are generally recommended as initial therapy in patients with left ventricular dysfunction. For those with persistent fluid retention despite treatment with a loop diuretic, a thiazide diuretic or metolazone may be added to the regimen. In patients with normal left ventricular function, either a thiazide diuretic or loop diuretic may be used as initial therapy to manage fluid overload.<sup>7-9</sup>

There are several national and international organizations that have published guidelines on the treatment of hypertension. Thiazide-type diuretics are frequently recommended as initial therapy in patients with uncomplicated hypertension.<sup>7-13</sup> According to the National Heart, Lung, and Blood Institute's Eighth Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8), thiazide-type diuretics should be utilized first-line for most patients with hypertension, either alone or in combination with another hypertensive from a different medication class (e.g., ACE inhibitors, ARBs,  $\beta$ -blockers, calcium channel blockers).<sup>7</sup> Several guidelines consistently recommend that the selection of an antihypertensive agent be based on compelling indications for use.<sup>7-15</sup> Most patients will require more than one antihypertensive medication to achieve blood pressure goals.<sup>7-13</sup>

In clinical trials, the thiazide-like diuretics have been shown to effectively lower blood pressure.<sup>17-44</sup> There were no studies found in the medical literature that directly compared the efficacy and safety of the thiazide-like diuretics for the treatment of hypertension.

There is insufficient evidence to support that one brand thiazide-like diuretic is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand thiazide-like diuretics within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand thiazide-like diuretic is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.



## XII. References

1. Daily Med [database on the internet]. Bethesda (MD): National Library of Medicine; 2022 [cited 2022 Jan]. Available at: <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.
2. Mann JFE. Choice of drug therapy in primary (essential) hypertension. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA, 2021.
3. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
4. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
5. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Apr 1;45. DOI: 10.1161/CIR0000000000001063.
6. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al, ESC Scientific Document Group, 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC, *European Heart Journal*, Volume 42, Issue 36, 21 September 2021, Pages 3599–3726, <https://doi.org/10.1093/eurheartj/ehab368>.
7. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014 Feb 5;311(5):507-20.
8. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, Ramirez A, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020 Jun;75(6):1334-1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. Epub 2020 May 6.
9. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, Butalia S, Leung AA, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596-624. doi: 10.1016/j.cjca.2020.02.086.
10. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2018 ESH/ESC Guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39(33):3021–3104.
11. National Institute for Health and Clinical Excellence (NICE). Hypertension in adults: diagnosis and management [guideline on the Internet]. London (UK): NICE; 2019 Aug [cited 2019 Sep]. Available from: <https://www.nice.org.uk/guidance/ng136>.
12. Flack JM, Sica DA, Bakris G, et al. Management of High Blood Pressure in Blacks: An Update of the International Society on Hypertension in Blacks Consensus Statement. *Hypertension*. 2010; 56:780-800.
13. Kidney Disease: Improving Global Outcomes (KDIGO) Blood Pressure Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int*. 2021 Mar;99(3S):S1-S87. doi: 10.1016/j.kint.2020.11.003.
14. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun; 71(6): 1269-1324.
15. American Diabetes Association Professional Practice Committee. Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2022;45(Suppl.1).
16. Biggins SW, Angeli P, Garcia-Tsao G, Ginès P, Ling SC, Nadim MK, Wong F, Kim WR. Diagnosis, Evaluation, and Management of Ascites, Spontaneous Bacterial Peritonitis and Hepatorenal Syndrome: 2021 Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology*. 2021 Aug;74(2):1014-1048. doi: 10.1002/hep.31884.
17. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. Final results of the Systolic Hypertension in the Elderly Program (SHEP). SHEP Cooperative Research Group. *JAMA*. 1991 Jun 26;265(24):3255-64.
18. Kostis JB, Berge KG, Davis BR, Hawkins CM, Probstfield J. Effect of atenolol and reserpine on selected events in the systolic hypertension in the elderly program (SHEP). *Am J Hypertens*. 1995 Dec;8(12 Pt 1):1147-53.

19. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major cardiovascular events in hypertensive patients randomized to doxazosin versus chlorthalidone. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2000;283:1967-75.
20. Black HR, Davis B, Barzilay J, et al; Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Metabolic and clinical outcomes in nondiabetic individuals with the metabolic syndrome assigned to chlorthalidone, amlodipine, or lisinopril as initial treatment for hypertension: a report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Diabetes Care*. 2008 Feb;31(2):353-60.
21. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium-channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288:2981-97.
22. Rahman M, Ford CE, Cutler JA, Davis BR, Piller LB, Whelton PK, et al. Long-term renal and cardiovascular outcomes in antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT) participants by baseline estimated GFR. *Clin J Am Soc Nephrol*. 2012;7:989-1002.
23. Muntner P, Levitan EB, Lynch AI, et al. Effect of chlorthalidone, amlodipine, and lisinopril on visit-to-visit variability of blood pressure: results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. *J Clin Hypertens (Greenwich)*. 2014 May;16(5):323-330.
24. Bangalore S, Davis BR, Cushman WC, Pressel SL, Muntner PM, Calhoun DA, et al. Treatment-Resistant Hypertension and Outcomes Based on Randomized Treatment Group in ALLHAT. *Am J Med*. 2017 Apr;130(4):439-448.e9.
25. Pupita F, Belogi M, Ansuini R, Campolucci G. Long-acting and short-acting diuretics in the treatment of hypertension. *Pharmatherapeutica*. 1983;3(7):475-81.
26. Bakris GL, Sica D, White WB, et al. Antihypertensive efficacy of hydrochlorothiazide vs chlorthalidone combined with azilsartan medoxomil. *Am J Med*. 2012 Dec;125(12):1229.e1-1229.e10.
27. Ernst ME, Carter BL, Goerdts CJ, Steffensmeier JJ, Phillips BB, Zimmerman MB, Bergus GR. Comparative antihypertensive effects of hydrochlorothiazide and chlorthalidone on ambulatory and office blood pressure. *Hypertension*. 2006 Mar;47(3):352-8.
28. Carter BL, Ernst ME, Cohen JD. Hydrochlorothiazide versus chlorthalidone: evidence supporting their interchangeability. *Hypertension*. 2004 Jan;43(1):4-9.
29. Karotsis AK, Symeonidis A, Mastorantonakis SE, Stergiou GS; Home-Di-Plus Study Group. Additional antihypertensive effect of drugs in hypertensive subjects uncontrolled on diltiazem monotherapy: a randomized controlled trial using office and home blood pressure monitoring. *Clin Exp Hypertens*. 2006 Oct;28(7):655-62.
30. Nissinen A, Tuomilehto J. Evaluation of the antihypertensive effect of atenolol in fixed or free combination with chlorthalidone. *Pharmatherapeutica*. 1980;2(7):462-8.
31. Fogari R, Zoppi A. Half-strength atenolol-chlorthalidone combination (Tenoretic mite) in the treatment of elderly hypertensive patients. *Int J Clin Pharmacol Ther Toxicol*. 1984 Jul;22(7):386-93.
32. Leonetti G, Pasotti C, Capra A. Low-dose atenolol-chlorthalidone combination for treatment of mild hypertension. *Int J Clin Pharmacol Ther Toxicol*. 1986 Jan;24(1):43-7.
33. Finnerty FA Jr. Chlorthalidone plus reserpine versus hydrochlorothiazide plus reserpine in a stepped-care approach to the treatment of essential hypertension. *J Clin Pharmacol*. 1980;20(5-6 Pt 1):357-63.
34. Akram J, Sheikh UE, Mahmood M, Donnelly R. Antihypertensive efficacy of indapamide SR in hypertensive patients uncontrolled with a background therapy: the NATIVE study. *Curr Med Res Opin*. 2007 Dec;23(12):2929-36.
35. Beckett NS, Peters R, Fletcher AE, et al; for the HYVET Study Group. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med*. 2008 May 1;358(18):1887-98.
36. Milia P, Muir S, Alem M, Lees K, Walters M. Comparison of the effect of diuretics on carotid blood flow in stroke patients. *J Cardiovasc Pharmacol*. 2006 Mar;47(3):446-9.
37. Madkour H, Gadallah M, Riveline B, Plante GE, Massry SG. Indapamide is superior to thiazide in the preservation of renal function in patients with renal insufficiency and systemic hypertension. *Am J Cardiol*. 1996 Feb 22;77(6):23B-25B.
38. PROGRESS Collaborative group. Randomized trial of a perindopril-based blood-pressure-lowering regimen among 6105 individuals with previous stroke or transient ischemic attack. *Lancet*. 2001 Sep 29;358:1033-41.
39. Hua AS, Kincaid-Smith P. A comparison of the effects of chlorothiazide and of metolazone in the treatment of hypertension. *Clin Sci Mol Med Suppl*. 1976 Dec;3:627s-9s.

40. ADVANCE Collaborative Group. Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): A randomized controlled trial. *Lancet* 2007;370:829-40.
41. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, et al. Randomized trial of effects of calcium antagonists compared with diuretics and  $\beta$ -blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet*. 2000 Jul 29;356(9227):359-65.
42. Ames RP. A comparison of blood lipid and blood pressure responses during the treatment of systemic hypertension with indapamide and with thiazides. *Am J Cardiol*. 1996 Feb 22;77(6):12b-6b.
43. Messerli FH, Grossman E, Goldbourt U. Are beta-blockers efficacious as first-line therapy for hypertension in the elderly? A systematic review. *JAMA*. 1998 Jun 17;279(23):1903-7.
44. Baguet JP, Legallicier B, Auquier P, Robitail S. Updated meta-analytical approach to the efficacy of antihypertensive drugs in reducing blood pressure. *Clin Drug Investig*. 2007;27(11):735-53.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Vasopressin Antagonists  
AHFS Class 402828  
May 4, 2022**

**I. Overview**

Conivaptan is an injectable product that is Food and Drug Administration (FDA)-approved for the treatment of euvolemic and hypervolemic hyponatremia in hospitalized patients. Conivaptan is not indicated for the treatment of congestive heart failure as the effectiveness of this agent has not been established in such patients.<sup>1</sup> Tolvaptan is an oral vasopressin antagonist that is FDA-approved for the treatment of clinically significant euvolemic and hypervolemic hyponatremia (serum sodium <125 mEq/L or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction), including patients with heart failure and Syndrome of Inappropriate Antidiuretic Hormone (SIADH).<sup>2</sup> The major disorders associated with euvolemic hyponatremia include SIADH, nephrogenic syndrome of inappropriate antidiuresis (NSIAD), glucocorticoid deficiency, hypothyroidism, exercise-associated hyponatremia (EAH), low solute intake, and primary polydipsia. Hypervolemic hyponatremia is most often caused by heart failure, cirrhosis, nephrotic syndrome, as well as acute and chronic renal failure.<sup>3</sup> Tolvaptan is now also available under the brand name Jynarque®, which is indicated to slow kidney function decline in adults at risk of rapidly progressing autosomal dominant polycystic kidney disease (ADPKD).<sup>4</sup> ADPKD is a hereditary disease characterized by renal cysts that are visible by ultrasonographic imaging studies. Patients with ADPKD can present with hypertension, hematuria, proteinuria, or kidney function impairment, detected by routine laboratory examinations. Flank pain is the most common symptom reported by patients.<sup>5</sup> ADPKD slowly progresses to chronic kidney disease and ultimately end-stage renal disease.<sup>5</sup>

Hyponatremia is frequently associated with elevated plasma levels of arginine vasopressin (AVP). AVP is normally secreted in response to increased plasma osmolality, decreased blood volume, or decreased blood pressure. Suppression of AVP secretion occurs when osmolality falls below a certain threshold, which results in renal excretion of free water. Failure to suppress AVP secretion may result in water retention and hyponatremia.<sup>3</sup> The use of traditional diuretics leads to both water and electrolyte excretion (diuresis); whereas, the use of tolvaptan leads to an increase in water excretion only (aquaresis), a decrease in urine osmolality, and an increase in serum sodium concentration. Urinary excretion of sodium and potassium, as well as plasma potassium concentrations, are not significantly affected by tolvaptan.<sup>2</sup>

The management of hyponatremia depends on the clinical presentation and duration of the disease (acute versus chronic hyponatremia). Therapeutic options include treating the underlying disease (if possible), fluid restriction, sodium chloride administration, and diuresis. Patients with chronic mild hyponatremia are often asymptomatic and treatment consists of fluid restriction or isotonic saline administration.<sup>6</sup> Acute severe hyponatremia requires more aggressive initial therapy as it may increase morbidity and mortality. Treatment of hyponatremia must be approached carefully as overly rapid correction may cause osmotic demyelination. Symptoms of osmotic demyelination are often irreversible and include quadriparesis, paraparesis, dysphagia, dysarthria, diplopia, seizures, coma, and death.<sup>3,6</sup>

The vasopressin antagonists that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. **Tolvaptan is available in a generic formulation.** This class was last reviewed in February 2020.

**Table 1. Vasopressin Antagonists Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Conivaptan	injection <sup>^</sup>	Vaprisol®	none
Tolvaptan	tablet	Jynarque®, Samsca®*	<b>tolvaptan</b>

\*Generic is available in at least one dosage form or strength.

<sup>^</sup>Product is primarily administered in an institution.

PDL=Preferred Drug List

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the vasopressin antagonists are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Vasopressin Antagonists**

Clinical Guideline	Recommendation(s)
<p>American Journal of Medicine: <b>Diagnosis, Evaluation, and Treatment of Hyponatremia: Expert Panel Recommendations (2013)</b><sup>3</sup></p>	<p><u>General information</u></p> <ul style="list-style-type: none"> <li>• There are no data to suggest that the etiology of the hyponatremia, nor the methodology used to correct hyponatremia, alters the susceptibility for producing osmotic demyelination with overly rapid correction.</li> <li>• The rate of correction of hyponatremia must be taken into account before deciding on the most appropriate therapy for any patient with hyponatremia.</li> <li>• Patients with acute (&lt;48 hours) hyponatremia may present with alarming neurologic findings, and they sometimes die of brain herniation. When hyponatremia develops over several days, brain swelling is minimized so that patients with chronic (&lt;48 hours) hyponatremia have more modest symptoms and almost never die of brain herniation.</li> </ul> <p><u>Rate of correction of hyponatremia</u></p> <ul style="list-style-type: none"> <li>• To reverse serious manifestations of acute hyponatremia, increasing serum sodium by 4 to 6 mmol/L is sufficient to prevent brain herniation and neurological damage from cerebral ischemia.</li> <li>• The rate of correction does not need to be restricted in patients with true acute hyponatremia, nor is re-lowering of excessive corrections indicated; however, if there is any uncertainty as to whether the hyponatremia is chronic versus acute, then the limits for correction of chronic hyponatremia should be followed.</li> <li>• In patients with chronic hyponatremia, neurologic sequelae are associated with more rapid rates of correction. The osmotic demyelination syndrome (ODS) can usually be avoided by limiting correction of chronic hyponatremia to 4 to 8 mmol/L in 24 hours for those at low-risk of ODS and to 4 to 6 mmol/L/day for those at high-risk.</li> <li>• Limits not to exceed: 8 mmol/L in any 24-hour period for high-risk patients and 10 to 12 mmol/L in any 24-hour period or 18 mmol/L in any 48-hour period in patients at normal risk.</li> <li>• Factors that place patients at high risk of developing ODS include serum sodium concentration <math>\leq 105</math> mmol/L, hypokalemia, alcoholism, malnutrition, and advanced liver disease.</li> </ul> <p><u>Conventional therapy of euvolemic hyponatremia</u></p> <ul style="list-style-type: none"> <li>• Treatment of patients with euvolemic hyponatremia will vary greatly depending on their presentation. The single most important factor guiding initial therapy is the presence of neurologic symptoms.</li> <li>• Cases of acute hyponatremia (<math>\leq 48</math> hours in duration) are usually symptomatic if the hyponatremia is severe (<math>\leq 120</math> mmol/L). These patients are at greatest risk from neurologic complications from the hyponatremia itself and should be corrected to higher serum sodium levels promptly.</li> <li>• Patients with more chronic hyponatremia (<math>&gt;48</math> hours in duration) who have minimal neurologic symptomatology are at little risk from complications of hyponatremia itself, but can develop osmotic demyelination following rapid correction. There is no indication to correct these patients rapidly, and they should be treated using slower-acting therapies.</li> <li>• Syndrome of inappropriate antidiuretic hormone secretion:             <ul style="list-style-type: none"> <li>○ Correction of acute symptomatic hyponatremia is best accomplished with hypertonic (3%) saline given via bolus or continuous infusion. Intravenous furosemide 20 to 40 mg should be used to treat volume overload. Acute treatment should be discontinued when the patient's symptoms are abolished, a safe serum sodium level (<math>\geq 120</math> mmol/L) is achieved, or a total</li> </ul> </li> </ul>

Clinical Guideline	Recommendation(s)
	<p>correction of 18 mmol/L is achieved.</p> <ul style="list-style-type: none"> <li>○ For the treatment of mild-to-moderate chronic hyponatremia, fluid restriction represents the least toxic therapy, and has generally been the treatment of choice. Several days of restriction are usually necessary before a significant increase in plasma osmolality occurs.</li> <li>○ Pharmacologic interventions are reserved for refractory cases where the degree of fluid restriction required to avoid hypo-osmolality is so severe that the patient is unable, or unwilling, to maintain it. The preferred drug is demeclocycline, which causes a nephrogenic form of diabetes insipidus. Treatment must be continued for several days to achieve maximal diuretic effects. Other agents, such as lithium, have similar renal effects but are less desirable because of inconsistent results and significant side effects and toxicities. Urea is as an alternative treatment for syndrome of inappropriate antidiuretic hormone secretion.</li> </ul> <ul style="list-style-type: none"> <li>● Glucocorticoid deficiency: <ul style="list-style-type: none"> <li>○ Glucocorticoid replacement should be started immediately after completion of a rapid adrenocorticotropic hormone stimulation test. Several days of glucocorticoids are sometimes required for normalization of the plasma osmolality. Primary treatment of hyponatremia may be indicated if significant neurologic symptoms are present.</li> </ul> </li> <li>● Hypothyroidism: <ul style="list-style-type: none"> <li>○ The primary therapy of hypothyroidism is thyroid hormone replacement.</li> <li>○ Hyponatremia with hypothyroidism is infrequent and generally of mild severity; therefore, modest fluid restriction is generally the only treatment necessary.</li> <li>○ Symptomatic hyponatremia may be seen in patients with more severe hypothyroidism and altered mental status, primary treatment of hyponatremia may be indicated to ascertain whether the hyponatremia is contributing to the patient's neurologic symptoms.</li> </ul> </li> <li>● Exercise-associated hyponatremia (EAH): <ul style="list-style-type: none"> <li>○ EAH can be severe and life threatening as a result of cerebral edema and noncardiogenic pulmonary edema.</li> <li>○ Hyponatremia occurring in the setting of endurance exercise is acute, and treatment of symptomatic hyponatremia should be rapid.</li> <li>○ With significant central nervous system impairment, hypertonic saline should begin immediately and continued until the serum sodium reaches 125 mmol/L or symptoms resolve.</li> </ul> </li> <li>● Low solute intake: <ul style="list-style-type: none"> <li>○ Hyponatremia from low solute intake is corrected by instituting proper nutrition, with increased content of solute both as electrolytes and protein.</li> </ul> </li> <li>● Primary polydipsia: <ul style="list-style-type: none"> <li>○ Therapy should be directed at reducing fluid intake into the normal range.</li> <li>○ Fluid ingestion in patients with psychogenic causes of polydipsia responds variably to behavior modification and pharmacologic therapy (e.g., clozapine).</li> </ul> </li> </ul> <p><u>Conventional therapy of hypervolemic hyponatremia</u></p> <ul style="list-style-type: none"> <li>● For all diseases associated with edema formation, dietary sodium restriction and diuretic therapy are the mainstays of therapy.</li> <li>● Congestive heart failure (CHF): <ul style="list-style-type: none"> <li>○ For severely symptomatic patients with very low or rapidly falling serum sodium, treatment should consist of hypertonic (3%) NaCl combined with loop diuretics to prevent fluid overload; for patients with mild to moderate symptoms, begin with fluid restriction (1 L/d total) and, if signs of volume overload are present, administer loop diuretics.</li> <li>○ If the serum sodium does not correct to the desired level, lift the fluid</li> </ul> </li> </ul>

Clinical Guideline	Recommendation(s)
	<p>restriction and start either conivaptan (if intravenous route is preferred or required) or tolvaptan (if oral therapy is preferred).</p> <ul style="list-style-type: none"> <li>○ Hyponatremia in HF is almost always chronic, so current limits for rate of correction of chronic hyponatremias should be observed.</li> <li>○ If tolvaptan is used, it may be up-titrated from 15 to 30 to 60 mg/d as necessary to achieve the desired level of correction of serum sodium.</li> <li>○ Continue treatment until the serum sodium has either normalized, symptoms have improved, or the level of serum sodium is no longer compromising administration of needed diuretic therapy.</li> </ul> <ul style="list-style-type: none"> <li>● The stimuli for AVP secretion may be more dynamic than in other disease states; if prescribed after discharge, assessing the need for chronic therapy of hyponatremia by providing a window of observation off therapy two to four weeks after treatment initiation is a reasonable approach.</li> <li>● Cirrhosis: <ul style="list-style-type: none"> <li>○ There are no guidelines specifically regarding treatment of hyponatremia in cirrhosis.</li> <li>○ Demeclocycline is relatively contraindicated because of a high incidence of nephrotoxicity, and urea has not been used often. Fluid restriction is the usual approach, but without outcome studies to assess its effectiveness.</li> </ul> </li> <li>● Nephrotic syndrome, acute and chronic renal failure: <ul style="list-style-type: none"> <li>○ In patients with hyponatremia with advanced acute and chronic renal failure and glomerular filtration rate &lt;20 mL/min, fluid restriction to amounts less than insensible losses plus urine output is generally necessary to cause a negative solute-free water balance and correction of hyponatremia.</li> <li>○ Vaptans can be employed in selected cases where fluid restriction is not successful or not well tolerated.</li> </ul> </li> </ul> <p><u>Use of vasopressin receptor antagonists in hyponatremia</u></p> <ul style="list-style-type: none"> <li>● Exclude hypovolemic hyponatremia.</li> <li>● Do not use in conjunction with other treatments for hyponatremia.</li> <li>● Do not use immediately after cessation of other treatments for hyponatremia, particularly 3% NaCl.</li> <li>● Monitor serum sodium closely (every 6 to 8 hours) for the first 24 to 48 hours after initiating treatment.</li> <li>● Maintain ad libitum fluid intake during the first 24 to 48 hours of treatment; hyponatremia can correct too quickly with coincidental fluid restriction; in patients with a defective or impaired thirst mechanism (e.g., intubated or unconscious patients), provide sufficient fluid to prevent overly rapid correction due to unopposed aquaresis.</li> <li>● Increase the frequency of serum sodium monitoring and consider stopping the vaptan if there is a change or deterioration in the patient's condition (e.g., nothing-by-mouth status, intubation) that limits the ability to request, access, or ingest fluid.</li> <li>● Severe, symptomatic hyponatremia should be treated with 3% NaCl, as this provides a quicker and more certain correction of serum sodium than vaptans.</li> <li>● Currently, there are insufficient data for use of vaptans in severe asymptomatic hyponatremia (serum sodium &lt;120 mmol/L)—use vaptans with caution and with more frequent monitoring in these patients.</li> <li>● If overcorrection occurs, consider re-lowering the serum sodium to safe limits.</li> <li>● For the treatment of acute severe hyponatremia, there is insufficient data from clinical trials to know if sufficiently rapid correction can be achieved with vasopressin receptor antagonists without the use of hypertonic saline.</li> <li>● Most studies to date in patients with hyponatremia have only been of relatively short duration. The most appropriate way to use these agents, their long-term response rates, how important the role of water restriction will remain during chronic use, and whether correction of chronic hyponatremia will result in improved cognitive</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>function as suggested by 30-day studies of tolvaptan, and quality of life, or functional status, as suggested by initial studies of gait stability and falls, are unknown at the present time and will require additional study.</p> <ul style="list-style-type: none"> <li>• Safety issues must be considered carefully with any new class of drugs. The possibility of overcorrection has been of significant concern in all of the vasopressin receptor antagonist clinical trials, but to date osmotic demyelination has not been reported with any agent. The potential for serious drug interactions via interference with cytochrome P450 3A4-mediated metabolism of other drugs must also be recognized. Whether there will be any adverse effect of V<sub>2</sub> receptor inhibition in vascular endothelium is unknown.</li> <li>• Further studies will be needed to assess the appropriate use of vasopressin receptor antagonists, such as for correction of symptomatic hyponatremia either alone or in conjunction with hypertonic saline infusions; to assess the benefits of correction of hyponatremia in hospitalized patients in terms of disease outcomes and decreased lengths of intensive care unit and hospital stay; and for long-term treatment of minimally symptomatic hyponatremia in order to decrease the risks of neurocognitive dysfunction and gait instability.</li> </ul>
<p>Canadian Expert Consensus: <b>Updated Canadian Expert Consensus on Assessing Risk of Disease Progression and Pharmacological Management of Autosomal Dominant Polycystic Kidney Disease (2018)<sup>9</sup></b></p>	<ul style="list-style-type: none"> <li>• All patients with a diagnosis of Autosomal Dominant Polycystic Kidney Disease (ADPKD) or suspected ADPKD should be referred to a nephrologist for initial assessment. Initial assessment should include kidney imaging and, in some cases, genetic testing to determine the patient's risk of rapid progression and to determine what treatment should be initiated.</li> <li>• Patients with ADPKD who are &lt;50 years old with eGFR &gt;60 mL/min/1.73 m<sup>2</sup> and without significant cardiovascular comorbidities should have a target blood pressure of ≤110/75 mm Hg, realizing that in some patients an individual target may be needed.</li> <li>• Consider treatment with tolvaptan for patients who fulfill the enrollment criteria of the TEMPO 3:4 study: 18 to 50 years of age with total kidney volume (TKV) &gt;750 mL and eGFR &gt;45 mL/min/1.73 m<sup>2</sup>.</li> <li>• Treatment with tolvaptan is recommended for patients who fulfill the enrollment criteria of the REPRISE study: <ul style="list-style-type: none"> <li>○ 18 to 55 years of age with eGFR of 25 to 65 mL/min/1.73 m<sup>2</sup> OR</li> <li>○ 56 to 65 years of age with eGFR of 25 to 44 mL/min/1.73 m<sup>2</sup> with historical evidence of a decline in eGFR &gt;2.0 mL/min/1.73 m<sup>2</sup>/year.</li> <li>○ Although there were no inclusion criteria for kidney size, based on the abundance of evidence that increased size of kidneys is relevant, these REPRISE criteria relate to those patients with ADPKD who have enlarged kidneys. In those patients with advanced or rapidly progressive chronic kidney disease (CKD) without enlarged kidneys, an alternate diagnosis for CKD should be investigated.</li> </ul> </li> <li>• Treatment with tolvaptan is suggested for patients who, according to the Mayo Clinic Classification, are classified as 1D or 1E with eGFR in CKD stages 1 to 4 (eGFR &gt;25 mL/min). Treatment with tolvaptan may be considered for patients who are classified as 1C and are &lt;50 years old or have other risk factors for rapid progression, such as an annual decrease in eGFR of &gt;2.5 mL/min/1.73 m<sup>2</sup> and/or increase in TKV of &gt;5% per year.</li> <li>• Treatment with tolvaptan should be stopped when the patient develops ESRD. In the predialysis setting with eGFR &lt;25 mL/min/1.73 m<sup>2</sup>, there are no data to guide when treatment with tolvaptan should be stopped.</li> <li>• Additional considerations when giving tolvaptan: <ul style="list-style-type: none"> <li>○ Patients with ADPKD on treatment with tolvaptan should follow a sodium-restricted diet of ≤2.4 g/day (≤100 mmol/day).</li> <li>○ Titrating tolvaptan to the maximal tolerated dose or to achieve a uOSM &lt;250 mOsm/kg water is suggested. Consideration should be given to consultation with a dietician to minimize sodium and osmolal intake to help manage severe aquaretic adverse events.</li> </ul> </li> </ul>



### III. Indications

The Food and Drug Administration (FDA)-approved indications for the vasopressin antagonists are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Vasopressin Antagonists<sup>2</sup>**

Indication	Tolvaptan* <sup>†</sup>
To slow kidney function decline in adults at risk of rapidly progressing autosomal dominant polycystic kidney disease	✓ (Jynarque <sup>®</sup> )
Treatment of clinically significant hypervolemic and euvolemic hyponatremia (serum sodium <125 mEq/L or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction), including patients with heart failure and Syndrome of Inappropriate Antidiuretic Hormone	✓

\*Patients requiring intervention to raise serum sodium urgently to prevent or to treat serious neurological symptoms should not be treated with tolvaptan.

<sup>†</sup>It has not been established that tolvaptan provides a symptomatic benefit to patients.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the vasopressin antagonists are listed in Table 4.

**Table 4. Pharmacokinetic Parameters of the Vasopressin Antagonists<sup>7</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life (hours)
Tolvaptan	≥40	99	Liver, extensive (% not reported)	Non-renal routes	12

### V. Drug Interactions

Major drug interactions with the vasopressin antagonists are listed in Table 5. Tolvaptan is metabolized by cytochrome P450 (CYP) 3A, and use with strong CYP3A inhibitors causes a marked (5-fold) increase in exposure.<sup>1</sup> Tolvaptan is contraindicated in combination with strong cytochrome CYP3A inhibitors, such as clarithromycin, ketoconazole, itraconazole, ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, and telithromycin.<sup>1</sup> The use of tolvaptan in combination with CYP3A inducers and moderate CYP3A inhibitors should also be avoided.<sup>1</sup>

**Table 5. Major Drug Interactions with the Vasopressin Antagonists<sup>7</sup>**

Generic Name	Interaction	Mechanism
Vasopressin antagonists (tolvaptan)	HIV protease inhibitors	Inhibition of CYP3A4 by HIV protease inhibitors may decrease the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be increased by HIV protease inhibitors.
Vasopressin antagonists (tolvaptan)	Imidazoles	Inhibition of CYP3A4 by imidazoles may decrease the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be increased by imidazoles.
Vasopressin antagonists (tolvaptan)	Macrolides and ketolides	Inhibition of CYP3A4 and P-glycoprotein by macrolides and ketolides may decrease the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be increased by macrolides and ketolides.
Vasopressin antagonists (tolvaptan)	Nefazodone	Inhibition of CYP3A4 by nefazodone may decrease the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be increased by nefazodone.

Generic Name	Interaction	Mechanism
Vasopressin antagonists (tolvaptan)	Moderate CYP3A4 Inhibitors	Inhibition of CYP3A isoenzymes by moderate CYP3A4 inhibitors may decrease the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be increased by moderate CYP3A4 inhibitors.
Vasopressin antagonists (tolvaptan)	Rifamycins	Induction of CYP3A isoenzymes by rifamycins may increase the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be decreased by rifamycins compromising therapeutic effectiveness.
Vasopressin antagonists (tolvaptan)	St. John's wort	Induction of CYP3A isoenzymes by St. John's wort may increase the metabolic elimination of tolvaptan. Plasma concentrations and pharmacologic effects of tolvaptan may be decreased by St. John's wort compromising therapeutic effectiveness.

CYP=cytochrome P450 isoenzymes, HIV=human immunodeficiency virus

## VI. Adverse Drug Events

The most common adverse drug events reported with the vasopressin antagonists are listed in Table 6. The boxed warning for tolvaptan is listed in Table 7.

**Table 6. Adverse Drug Events (%) Reported with the Vasopressin Antagonists<sup>8</sup>**

Adverse Events	Tolvaptan
<b>Cardiovascular</b>	
Palpitations	4
Ventricular fibrillation	<2
<b>Central Nervous System</b>	
Cerebrovascular accident	<2
Dizziness	11
Fatigue	14
Pyrexia	4
<b>Endocrine and Metabolic</b>	
Diabetic ketoacidosis	<2
Hyperglycemia	6
Hypernatremia	<4
Hyperuricemia	4
<b>Gastrointestinal</b>	
Abdominal distention	5
Anorexia	4
Constipation	7
Diarrhea	13
Dyspepsia	8
Ischemic colitis	<2
Nausea	21
Xerostomia	7 to 16
<b>Genitourinary</b>	
Pollakiuria	4 to 11
Polyuria	4 to 70
Urethral bleeding	<2
Vaginal hemorrhage	<2
<b>Laboratory Abnormalities</b>	
Bilirubin increased	<1
Increased serum alanine aminotransferase	5
Prothrombin time prolonged	<2
<b>Musculoskeletal</b>	
Rhabdomyolysis	<2

Adverse Events	Tolvaptan
Weakness	9
<b>Respiratory</b>	
Pulmonary embolism	<2
Respiratory failure	<2
<b>Other</b>	
Deep vein thrombosis	<2
Dehydration	2 to 3
Disseminated intravascular coagulation	<2
Hepatotoxicity	≤4
Hypersensitivity reaction	<1
Skin rash	4
Thirst	12 to 64
Xeroderma	5

✓ Percent not specified  
- Event not reported

**Table 7. Boxed Warning for Tolvaptan<sup>2,4</sup>**

<b>WARNING</b>	
<b>Samsca®</b>	
Initiate and re-initiate in a hospital and monitor serum sodium Samsca® should be initiated and re-initiated in patients only in a hospital where serum sodium can be monitored closely.	
<ul style="list-style-type: none"> <li>• Too rapid correction of hyponatremia (e.g., &gt;12 mEq/L/24 hours) can cause osmotic demyelination resulting in dysarthria, mutism, dysphagia, lethargy, affective changes, spastic quadriparesis, seizures, coma and death. In susceptible patients, including those with severe malnutrition, alcoholism or advanced liver disease, slower rates of correction may be advisable.</li> </ul>	
Not for use for autosomal dominant polycystic kidney disease (ADPKD)	
<ul style="list-style-type: none"> <li>• Because of the risk of hepatotoxicity, tolvaptan should not be used for ADPKD outside of the FDA-approved REMS.</li> </ul>	
<b>Jynarque®</b>	
<ul style="list-style-type: none"> <li>• Jynarque® can cause serious and potentially fatal liver injury. Acute liver failure requiring liver transplantation has been reported.</li> <li>• Measure ALT, AST and bilirubin before initiating treatment, at two weeks and four weeks after initiation, then monthly for the first 18 months and every three months thereafter. Prompt action in response to laboratory abnormalities, signs, or symptoms indicative of hepatic injury can mitigate, but not eliminate, the risk of serious hepatotoxicity.</li> <li>• Because of the risks of serious liver injury, Jynarque® is available only through a restricted distribution program under a Risk Evaluation and Mitigation Strategy (REMS) called the Jynarque REMS Program.</li> </ul>	

## VII. Dosing and Administration

The usual dosing regimens for the vasopressin antagonists are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Vasopressin Antagonists<sup>2,4</sup>**

Generic Name	Usual Adult Dose	Usual Pediatric Dose	Availability
Tolvaptan	<p><u>Hypervolemic and euvoletic hyponatremia:</u> Tablet: initial, 15 mg once daily; maintenance, increase to 30 mg once daily after ≥24 hours as needed to achieve the desired level of serum sodium; maximum, 60 mg once daily</p> <p><u>Rapidly Progressing Autosomal Dominant Polycystic Kidney Disease</u></p>	Safety and effectiveness have not been established in pediatric patients.	Tablet: 15 mg 30 mg 45 mg 60 mg 90 mg

Generic Name	Usual Adult Dose	Usual Pediatric Dose	Availability
	Tablet: initial, 60 mg orally per day as 45 mg taken on waking and 15 mg taken 8 hours later; Titrate to 60 mg plus 30 mg then to 90 mg plus 30 mg per day if tolerated with at least weekly intervals between titrations		

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the vasopressin antagonists are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Vasopressin Antagonists**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Gheorghide et al.<sup>10</sup> (2006)</p> <p>Tolvaptan 10 mg/day, with titration to larger doses (15, 30, 45, and 60 mg/day) as needed to achieve serum sodium concentrations within normal limits</p> <p>vs</p> <p>fluid restriction (initially 1,200 mL/24 hrs) plus placebo</p>	<p>AC, MC, OL, RCT</p> <p>Patients ≥18 years, serum sodium &lt;135 mmol/L for ≥2 consecutive days, and normovolemia or signs of fluid overload</p>	<p>N=28</p> <p>Inpatient treatment: 14 days</p> <p>Outpatient treatment: 14 days</p> <p>Follow-up: 65 days</p>	<p>Primary: Normalization of serum sodium concentration (defined as ≥135 mmol/L or an increase of &gt;10% from baseline to the last inpatient assessment)</p> <p>Secondary: Changes in serum sodium from baseline to the last outpatient visit (day 65), urine osmolality, urine volume, urine sodium concentration, body weight, total fluid intake, thirst score from baseline to the last inpatient assessment</p>	<p>Primary: A higher proportion of subjects in the tolvaptan group had achieved the normalization of serum sodium compared to those in the fluid restriction group by the last inpatient visit (P=0.049). The normalization of serum sodium was achieved more rapidly in the tolvaptan group than in the fluid restriction group, occurring in 50% of tolvaptan-treated subjects by day four, compared to day eight in the fluid restriction group (P&lt;0.03).</p> <p>Patients in the tolvaptan group had a significantly greater increase in serum sodium concentration 4 hours after the first dose (1.6 mmol/L; P=0.016), at day 5 (5.2 mmol/L; P=0.019) and at the last inpatient visit (5.7 mmol/L; P=0.0065) compared to patients receiving fluid restriction (-0.8, 0.7, and 1.0, respectively).</p> <p>Secondary: At day 65, the mean change in serum sodium was 4.7 mmol/L in the tolvaptan group compared to -0.3 mmol/L in the placebo group (P=0.039).</p> <p>Urine sodium was significantly lower (P=0.021) and urine output was significantly greater (P=0.014) in the tolvaptan group compared to the placebo group.</p> <p>No significant differences in urine osmolality (P=0.058), serum potassium (P=0.45), blood pressure, heart rate, body weight (P value not significant), thirst score (P=0.8) or adverse events requiring drug discontinuation were observed between the treatment groups.</p>
<p>Schrier et al.<sup>11</sup> (2006)</p> <p>SALT-1 and SALT-2</p> <p>Tolvaptan 15</p>	<p>DB, MC, PC, RCT</p> <p>Patients ≥18 years of age with euvolemic or hypervolemic</p>	<p>N=102 (SALT-1)</p> <p>N=123 (SALT-2)</p>	<p>Primary: Change in the average daily AUC for the serum sodium from baseline to day 4</p>	<p>Primary: By day four, the increase in the average daily AUC for the serum sodium concentration was 3.62 and 4.33 for tolvaptan (SALT-1 and SALT-2, respectively) compared to 0.25 and 0.42 for placebo (P&lt;0.001 for all comparisons).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>mg/day for 30 days (dose could be titrated to 60 mg/day)</p> <p>vs</p> <p>placebo</p>	<p>hyponatremia (serum sodium &lt;135 mmol/L). Patients also had chronic heart failure, cirrhosis, or the syndrome of inappropriate antidiuretic hormone secretion (SIADH) in association with the hyponatremia.</p>	<p>37 days</p>	<p>and from baseline to day 30</p> <p>Secondary: Change in the AUC for the serum sodium in patients with marked hyponatremia, serum sodium concentration at each visit, time to normalization of the serum sodium, percent of patients with serum sodium concentrations that normalized at day 4 and day 30, serum sodium concentration on day 4 and day 30 for patients with mild or marked hyponatremia at baseline, change from baseline in scores on the Physical Component Summary and Mental component summary of the medical outcomes Study 12-item Short-Form General Health</p>	<p>By day 30, the increase in the average daily AUC for the serum sodium concentration was 6.22 and 6.20 for tolvaptan (SALT-1 and SALT-2, respectively) compared to 1.66 and 1.84 for placebo (P&lt;0.001 for all comparisons).</p> <p>Secondary: By day 30, the increase in the average daily AUC for the serum sodium concentration in patients with marked hyponatremia was 8.24 and 7.60 for tolvaptan (SALT-1 and SALT-2, respectively) compared to 2.54 and 2.72 for placebo (P&lt;0.001 for all comparisons).</p> <p>By day four, serum sodium concentrations were 133.9 and 135.3 mmol/L for tolvaptan (SALT-1 and SALT-2, respectively) compared to 129.7 and 129.6 mmol/L for placebo (P&lt;0.001 for all comparisons). By day 30, serum sodium concentrations were 135.7 and 135.9 mmol/L for tolvaptan (SALT-1 and SALT-2, respectively) compared to 131 and 131.5 mmol/L for placebo (P&lt;0.001 for all comparisons).</p> <p>By day four, 40 and 55% of patients receiving tolvaptan (SALT-1 and SALT-2, respectively) had normal serum sodium concentrations compared to 13 and 11% for placebo (P&lt;0.001 for all comparisons). By day 30, 53 and 58% of patients receiving tolvaptan (SALT-1 and SALT-2, respectively) had normal serum sodium concentrations compared to 25 and 25% for placebo (P&lt;0.001 for all comparisons).</p> <p>Scores on the Physical Component Summary did not differ significantly between groups. Scores for the Mental Component Summary improved in the tolvaptan group when the data from SALT-1 and SALT-2 were combined (P=0.02), as well as in SALT-1 (P=0.04). Scores improved significantly in the combined subgroup of patients with marked hyponatremia (P=0.04). There was no significant difference between the groups found in SALT-2 (P=0.14).</p> <p>Adverse event profiles in the two study groups were similar for all comparisons. The most common adverse events occurring during the study in the tolvaptan groups were thirst and dry mouth. Overall, there were 26 serious adverse events potentially related to the study treatment in SALT-1</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Survey	<p>and SALT-2. The number of deaths in the two study groups was similar (14 deaths among 223 patients in the tolvaptan groups and 13 deaths among 220 patients in the placebo groups), and they occurred within the defined observation period.</p> <p>In four of the patients in the tolvaptan group, the desirable rates of sodium correction were exceeded during the first 24 hours of the study (&gt;0.5 mmol/L per hour). In four patients (1.8%), the predefined serum sodium concentration (&gt;146 mmol per liter) was exceeded.</p>
<p>Berl et al.<sup>12</sup> (2010) SALTWATER</p> <p>Tolvaptan QD (dose varied based on response)</p>	<p>OL, ES (Extension of SALT-1 and SALT-2)</p> <p>Patients ≥18 years of age with euvolemic or hypervolemic hyponatremia (serum sodium &lt;135 mmol/L). Patients also had chronic heart failure, cirrhosis, or the SIADH in association with the hyponatremia</p>	<p>N=111</p> <p>4 years (mean 1.9 years)</p>	<p>Primary: Safety, efficacy</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>During the follow-up period, 105 of 111 patients experienced an adverse event. The most common adverse events that were potentially related to tolvaptan use were pollakiuria, thirst, fatigue, dry mouth, polydipsia, polyuria, hypotension, hypernatremia, dizziness, headache, peripheral edema, and acute renal failure.</p> <p>A total of 19 patients died during the follow-up period (9 deaths per 100 patient-years of exposure). The death rate during SALTWATER was lower than that observed for SALT (86.9 deaths per 100 patient-years of exposure).</p> <p>In five patients, serum sodium correction exceeded the rate of 1 mmol/L per h at the eight hour time point. There were 18 patients who had serum sodium levels &gt;145 mmol/L at individual time points.</p> <p>Correction of serum sodium levels during the first eight hours of therapy occurred at similar rates in SALTWATER compared to SALT-1 and SALT-2. After the initial titration period, mean serum sodium levels remained within the normal range throughout the four year treatment period.</p> <p>In all patient subgroups, serum sodium levels declined by seven days of withholding tolvaptan. On drug discontinuation, the proportion of patients who declined by ≥3 mEq/L was 68%, and an equal proportion fell from ≥135mEq/L to below this threshold of normal.</p> <p>The mean time to first fluid restriction was 122.3 and 162.5 days in the</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				mild and marked hyponatremia subgroups, respectively; 13.2% of patients in the mild hyponatremia group and 5.4% in the marked hyponatremia group required fluid restriction.  Secondary: Not reported
Cardenas et al. (abstract) <sup>13</sup> (2012) SALT-1 and SALT-2  Tolvaptan 15 mg/day for 30 days (dose could be titrated to 60 mg/day)  vs  placebo	Subgroup analysis  Patients with cirrhosis and hyponatremia	N=120  30 days	Primary: Change in the average daily AUC for the serum sodium from baseline to day 4 and from baseline to day 30  Secondary: Mental component summary of the medical outcomes Study 12-item Short-Form General Health Survey, safety	Primary: Treatment with tolvaptan effectively raised serum sodium. Average daily AUC for serum sodium was significantly greater with tolvaptan from baseline to day 4 (P<0.0001) and day 30 (P<0.0001) compared to placebo. Superiority of tolvaptan was maintained after stratification by baseline hyponatremia (mild and marked), eGFR ( $\leq 60$ and $>60$ mL/min), or serum creatinine levels ( $<1.5$ and $\geq 1.5$ mg/dL).  Hyponatremia recurred seven days after discontinuation of tolvaptan.  Secondary: Mean mental component summary scores of the Short-Form General Health Survey improved from baseline to day 30 with tolvaptan but not with placebo (4.68 vs 0.08; P=0.02).  Major adverse events with tolvaptan were dry mouth and thirst. Gastrointestinal bleeding occurred in 10 and 2% of patients receiving tolvaptan and placebo, respectively (P=0.11). Rates of adverse events, withdrawals, and deaths were similar with both treatments.
Udelson et al. <sup>14</sup> (2008)  Tolvaptan 15, 30, or 60 mg administered as a single dose  vs  placebo	DB, MC, PC, RCT  Patients $\geq 18$ years of age with symptomatic heart failure (NYHA class III or IV) of $\geq 3$ months' duration caused by LVEF $<40\%$ . Patients were also required to be on standard	N=181  12 hours	Primary: PCWP peak change from baseline within 3 to 8 h after treatment administration  Secondary: AUC for the change from baseline PCWP	Primary: The pairwise comparisons of 15, 30, and 60 mg tolvaptan versus placebo each showed a statistically significant decrease in peak change in PCWP from three to eight hours post-dose (P=0.003, P=0.044, and P=0.033, respectively).  Secondary: For the AUC <sub>0-8h</sub> , the 15 mg tolvaptan group was the only tolvaptan dose group that was statistically significantly different from placebo.  All tolvaptan doses produced statistically significantly greater changes than placebo in peak change in pulmonary artery pressure (P<0.01 for 15



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	background therapy for heart failure for $\geq 1$ month.		and other hemodynamic parameters over an 8 hour evaluation period and renal and electrolyte parameters	<p>mg; <math>P &lt; 0.05</math> for 30 and 60 mg).</p> <p>Tolvaptan 15 and 30 mg doses resulted in statistically significant reductions in peak change in right atrial pressure as compared to placebo (<math>P &lt; 0.01</math> and <math>P &lt; 0.05</math>, respectively).</p> <p>No significant changes in cardiac index, pulmonary vascular resistance, and systemic vascular resistance were observed after tolvaptan administration compared to placebo.</p> <p>The single dose of tolvaptan produced a dose-dependent increase in urine output (<math>P &lt; 0.0001</math> for all tolvaptan groups vs placebo). Urine osmolality was significantly reduced by all doses of tolvaptan relative to placebo (<math>P &lt; 0.0001</math> for all tolvaptan groups vs placebo). Free water clearance was significantly greater for all tolvaptan doses relative to placebo at all time points. Plasma osmolality increased in all of the tolvaptan-treated groups compared to placebo. Serum sodium levels showed a dose-related increase compared to placebo (1.2, 3.3, 4.6, and -0.7 mEq/L for the tolvaptan 15, 30, 60 mg, and placebo groups, respectively). Potassium levels were not different from placebo in any of the tolvaptan dosing groups. No significant changes in serum creatinine, blood urea nitrogen, serum potassium, and vital signs were observed after study drug administration.</p> <p>Tolvaptan was well tolerated relative to placebo. Patient-reported adverse events in this short-term study occurred in 45.5, 44.2, 54.3, and 33.3% of the 15, 30, and 60 mg tolvaptan and placebo groups, respectively.</p>
<p>Udelson et al.<sup>15</sup> (2007)</p> <p>Tolvaptan 30 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients <math>\geq 18</math> years of age with CHF (NYHA class II to III) with a LVEF <math>&lt; 30\%</math>. Patients were also required to be on standard background therapy for heart failure for</p>	<p>N=240</p> <p>55 weeks</p>	<p>Primary: Change from baseline in LVEDV index</p> <p>Secondary: Change from baseline in LVESV index, comparison of the change from baseline in</p>	<p>Primary: In the placebo group, there was no change in LVEDV index over the year of follow-up. After one year of tolvaptan therapy, there was a small reduction in LVEDV index; however, this was not significantly different from placebo (-1.8 mL/m<sup>2</sup>; <math>P = 0.21</math> vs placebo). There was also no difference in the change of volumes from baseline at the week 55 study.</p> <p>Secondary: In the placebo group, LVEDV index decreased 0.4 mL/m<sup>2</sup> compared to a decrease of 3.3 mL/m<sup>2</sup> in the tolvaptan group (<math>P = 0.09</math>). There was no difference in the change of LVESV index from baseline at week 55.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	<p>≥3 months before enrollment.</p>		<p>LVEDV index after drug withdrawal (week 55), assessment of symptoms (using subject-assessed symptom scales and the Minnesota Living With Heart Failure Questionnaire)</p>	<p>Ejection fraction changes were small and similar in both treatment groups.</p> <p>Only minor changes in blood pressure and heart rate were observed over the course of the trial; there were no significant differences in the tolvaptan versus placebo groups. There were no significant between-group differences in serum sodium or potassium across the course of the trial. There were also no differences in renal function parameters (BUN and serum creatinine) across the year of therapy.</p> <p>No statistically significant differences were observed between the tolvaptan group and the placebo group for the change from baseline in Minnesota Living With Heart Failure Questionnaire score or for the Visual Analog Scale assessment of global status or respiratory status. More subjects in the tolvaptan group reported a score of “better” in the subject-assessed overall treatment effect at each visit than did subjects in the placebo group; however, no statistically significant differences were observed between treatment groups.</p> <p>There were six deaths (5%) and 21 hospitalizations of patients with heart failure (18%) in the tolvaptan-treated group, compared to 11 deaths (9%) and 34 heart failure hospitalizations (28%) in the placebo-treated group (P&lt;0.03 for the composite of death and heart failure hospitalizations).</p> <p>Adverse events including urinary frequency, thirst, and dry mouth occurred more frequently with tolvaptan than with placebo therapy. There was no difference in the number of patients withdrawn from the trial as the result of bothersome side effects between the two randomization groups.</p>
<p>Gheorghide et al.<sup>16</sup> (2007) EVEREST  Tolvaptan 30 mg/day within 48 hours of admission  vs</p>	<p>DB, MC, PC, RCT  Patients ≥18 years of age with a history of chronic heart failure who had been hospitalized for worsening CHF and who had a LVEF ≤40%.</p>	<p>N=2,048 (Trial A)  N=2,085 (Trial B)  7 days</p>	<p>Primary: Composite score of changes from baseline in patient-assessed global clinical status and body weight at day 7 or discharge</p>	<p>Primary: The composite score of changes from baseline in patient-assessed global clinical status and body weight at day seven or discharge was greater with tolvaptan compared to placebo (Trial A, mean 16 vs 0.99; P&lt;0.001; Trial B, mean 17 vs 0.97; P&lt;0.001).</p> <p>Improvement in patient-assessed global clinical status (assessed alone), measured by a 100-point visual analog scale at day seven or discharge, was similar between the tolvaptan and placebo groups (Trial A, mean 18.25 vs 17.73; P=0.51; Trial B, mean 18.72 vs 18.28; P=0.52).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>placebo</p>	<p>Patients also received conventional heart failure therapy.</p>		<p>Secondary: Patient-assessed changes in dyspnea at day 1, global clinical status at day 7 or discharge, body weight at days 1 and 7 or discharge, and peripheral edema at day 7 or discharge</p>	<p>Mean body weight reductions at day seven or discharge in the tolvaptan and placebo groups were 3.35 vs 2.73 kg, respectively, in Trial A (P&lt;0.001) and 3.77 vs 2.79 kg, respectively, in Trial B (P&lt;0.001).</p> <p>Secondary: More patients in the tolvaptan groups (76.74% in Trial A and 72.06% in Trial B) reported an improvement dyspnea at day one (for those patients with dyspnea at baseline) compared to placebo (70.61% in Trial A and 65.32% in Trial B; P&lt;0.001 in both Trials).</p> <p>There was no significant difference in global clinical status at day seven or discharge between the tolvaptan or placebo treatment groups (Trial A, P=0.51; Trial B, P=0.52).</p> <p>Changes in mean body weight were significantly greater with tolvaptan at day one (Trial A, -1.71 kg; Trial B -1.82 kg) than with placebo (Trial A, -0.99 kg; Trial B, 0.95 kg; P&lt;0.001 in both trials).</p> <p>There was no difference in peripheral edema at inpatient day seven or discharge with tolvaptan vs placebo in Trial A. In Trial B, 73.67% of patients experienced at least a 2-grade improvement in pedal edema with tolvaptan compared to placebo (P=0.02).</p> <p>An overall in-hospital mortality rate of 2.4 and 2.9% was observed in the tolvaptan and placebo groups, respectively. Through day seven or discharge, adverse events were reported in 49.1 and 40.0% of patients in Trial A, and in 55.9 and 47.9% of patients in Trial B in the tolvaptan and placebo groups, respectively.</p>
<p>Konstam et al.<sup>17</sup> (2007) EVEREST  Tolvaptan 30 mg/day within 48 hours of admission</p>	<p>DB, MC, PC, RCT  Patients ≥18 years of age with a history of chronic heart failure who had been hospitalized for worsening CHF</p>	<p>N=4,133  ≥60 days</p>	<p>Primary: All-cause mortality, composite of cardiovascular death or hospitalization for heart failure</p>	<p>Primary: The median duration of follow-up was 9.9 months. A total of 537 patients in the tolvaptan group (25.9%) and 543 patients in the placebo group (26.3%) died (HR, 0.98; 95% CI, 0.87 to 1.11; P=0.68). A total of 871 patients in the tolvaptan group (42.0%) and 829 patients in the placebo group (40.2%) died from cardiovascular causes or had a first hospitalization for heart failure (HR, 1.14; 95% CI, 0.95 to 1.14; P=0.55).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo	and who had a LVEF ≤40%. Patients also received conventional heart failure therapy.		Secondary: Composite of cardiovascular mortality or cardiovascular hospitalization, incidence of cardiovascular mortality, incidence of clinical worsening of heart failure (death, hospitalization for heart failure, or unscheduled visit for heart failure), changes from baseline in body weight at day 1, serum sodium level at day 7 or discharge, edema score at day 7 or discharge, patient-assessed dyspnea at day 1, and Kansas City Cardiomyopathy Questionnaire overall summary score at outpatient week 1	<p>Secondary: The composite of cardiovascular death or cardiovascular hospitalization, the incidence of cardiovascular mortality, and the incidence of clinical worsening of heart failure did not differ between the two treatment groups (P=0.52, P=0.67 and P=0.62, respectively).</p> <p>In patients with dyspnea at baseline, patient-assessed dyspnea scores significantly improved at day one in patients receiving tolvaptan compared to placebo (P&lt;0.001), with 74.3% of the tolvaptan group and 68.0% of the placebo group demonstrating an improvement in dyspnea score.</p> <p>Mean body weight at day one was reduced by 1.76 kg in the tolvaptan group and by 0.97 kg in the placebo group (P&lt;0.001).</p> <p>Among patients with baseline serum sodium levels less than 134 mEq/L, mean serum sodium concentrations increased by 5.49 mEq/L at day 7 or discharge with tolvaptan compared to 1.85 mEq/L in the placebo group (P&lt;0.001). This effect was observed as early as day one and was maintained through 40 weeks of treatment.</p> <p>In patients with baseline pedal edema, edema scores significantly improved at day seven or discharge in patients receiving tolvaptan compared to placebo (P=0.003), with 73.8% of tolvaptan patients and 70.5% of placebo patients manifesting improvement in edema by at least two grades.</p> <p>A significant improvement in physician-assessed pedal edema was observed as early as day one and continued through post-discharge week four.</p> <p>No significant changes were observed at outpatient week one in the Kansas City Cardiomyopathy Questionnaire overall summary score. Statistically significant changes favoring tolvaptan were observed at the time of the last scheduled on-treatment assessment at study end for the quality-of life domain (P=0.003), the social limitation domain (P=0.05), and the overall summary score (P=0.02). The other domains (clinical summary, physical limitation, total symptom, symptom frequency,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>symptom burden, symptom stability, and self-efficacy) did not reach significance at the time of the last on-treatment assessment.</p> <p>Adverse events occurred in 89.0% of tolvaptan patients and 86.1% of placebo patients.</p>
<p>Pang et al.<sup>18</sup> (2009) EVEREST</p> <p>Tolvaptan 30 mg/day within 48 hours of admission</p> <p>vs</p> <p>placebo</p>	<p>Post-hoc analysis of EVEREST</p> <p>Patients <math>\geq 18</math> years of age with a history of chronic heart failure who had been hospitalized for worsening CHF with LVEF <math>\leq 40\%</math>. Patients also received conventional heart failure therapy.</p>	<p>N=3,664</p> <p>1 to 3 days</p>	<p>Primary: Patient-assessed dyspnea using a seven-point Likert scale administered on day 1 after randomization</p> <p>Secondary: Not reported</p>	<p>Primary: Tolvaptan was associated with improved patient-assessed dyspnea on inpatient day one compared to placebo (74.3 vs 68.0%; <math>P &lt; 0.0001</math>) as reported in the primary EVEREST analysis. The greatest treatment differences were seen in subjects with continuous dyspnea at baseline.</p> <p>Patients were divided post hoc into five groups, based on time (in hours) of dyspnea assessment after the first dose of tolvaptan. The percentage improvement with placebo stayed relatively constant, whereas improvement with tolvaptan was greatest when measured early (<math>P &lt; 0.05</math>). The majority of patients had an improvement in dyspnea at all time points relative to hospital admission; however, there was a significantly higher rate of improvement with tolvaptan compared to placebo (<math>P &lt; 0.05</math>).</p> <p>There was also a linear association between reductions in body weight and improvements in patient-assessed dyspnea.</p> <p>Secondary: Not reported</p>
<p>Hauptman et al.<sup>19</sup> (2013) EVEREST</p> <p>Tolvaptan 30 mg/day within 48 hours of admission</p> <p>vs</p> <p>placebo</p>	<p>Post-hoc analysis of EVEREST</p> <p>Patients <math>\geq 18</math> years of age with a history of chronic heart failure who had been hospitalized for worsening CHF with LVEF <math>\leq 40\%</math>. Patients also received conventional heart</p>	<p>N=475</p> <p><math>\geq 60</math> days</p>	<p>Primary: Body weight at day 1, serum sodium at day 7 or discharge in patients with a baseline serum sodium <math>&lt; 135</math> mEq/L, edema score at day 7 or discharge for those with peripheral edema at baseline, dyspnea at day 1</p>	<p>Primary: Mean change from baseline in serum sodium was 4.72 mEq/L vs 1.18 mEq/L at day one and 4.90 mEq/L vs 1.93 mEq/L at day seven in the tolvaptan and placebo groups, respectively (<math>P &lt; 0.0001</math> at each time point). Tolvaptan was more likely to lead to normalization of serum sodium defined by a value of <math>\geq 135</math> mEq/L at both day one and at discharge compared with placebo (58 vs 20% and 64 vs 29%, respectively; <math>P &lt; 0.001</math> for both comparisons).</p> <p>In patients with dyspnea and hyponatremia at baseline (<math>n=409</math>), the changes in dyspnea were more favorable in the tolvaptan group versus placebo (van Elteren analysis: 0.56, 95% CI 0.51 to 0.62; <math>P=0.028</math>).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	failure therapy. This analysis included the hyponatremic cohort.		<p>for those with dyspnea at baseline, long-term clinical course</p> <p>Secondary: Not reported</p>	<p>Mean body weight at day one was reduced by 1.69 kg in the tolvaptan group and by 0.96 kg in the placebo group (P&lt;0.0001). Changes observed in physician-assessed edema at day seven (or discharge if earlier) were not significantly different between the groups (P=0.79).</p> <p>Serum sodium increases observed in the short term among the patients with hyponatremia continued during the outpatient portion of the study, with results significantly favoring tolvaptan.</p> <p>There was a favorable effect of tolvaptan treatment on first occurrence of cardiovascular mortality or morbidity in those with more severe reduction in serum sodium at baseline (HR, 0.60; 95% CI, 0.37 to 0.98; P=0.04). There was no effect of tolvaptan on CV mortality or morbidity in the mild hyponatremia group, with baseline serum sodium 130 to 134 mEq/L (HR, 0.96; 95% CI, 0.74 to 1.25; P=0.77).</p> <p>Secondary: Not reported</p>
<p>Gheorghiade et al.<sup>20</sup> (2004) ACTIV IN CHF</p> <p>Tolvaptan 30, 60, or 90 mg/day</p> <p>vs placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients ≥18 years of age admitted for worsening CHF with LVEF &lt;40% within 1 year of admission and systemic congestion (JVD, rales, or peripheral edema after initial in-hospital therapy for heart failure). Patients also received conventional heart failure therapy.</p>	<p>N=319</p> <p>Inpatient: 10 days</p> <p>Outpatient: 7 weeks</p>	<p>Primary: Change in body weight at 24 hrs after the administration of the first dose of study drug; worsening heart failure at 60 days</p> <p>Secondary: Changes in dyspnea, JVD, rales, edema, body weight, urine output, serum electrolyte levels, length of hospital stay after</p>	<p><u>Inpatient Phase</u></p> <p>Primary: A greater median reduction in body weight was found in patients treated with tolvaptan compared to placebo 24 hrs after the administration of the first dose of study drug (-1.80, -2.10, -2.05, and -0.60 kg for tolvaptan 30, 60, and 90 mg, and placebo, respectively; P=0.002, P=0.002, and P=0.009 for the 3 tolvaptan groups compared to the placebo group).</p> <p>Secondary: The median body weight reductions from baseline to discharge were greater in the tolvaptan groups compared to the placebo group (-3.30, -2.80, -3.20, and -1.90 kg in the groups receiving tolvaptan 30, 60, and 90 mg, and placebo, respectively; P=0.006, P=0.002, and P=0.06 for the three tolvaptan groups compared to placebo).</p> <p>The mean urine output at 24 hrs was 4,056.2, 4,175.2, 4,127.3, and 2,296.5 mL for the tolvaptan 30, 60, and 90 mg, and placebo groups, respectively (P =0.02, P&lt;0.001, and P&lt;0.001 for the three tolvaptan groups compared to the placebo group).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			randomization, use of diuretics, and patient/physician-assessed symptom scales	<p>Signs and symptoms of heart failure improved in all patients during the period of hospitalization. There were no significant differences in JVD, and peripheral edema between the treatment groups (dyspnea P=0.04).</p> <p>Global assessment scales did not show a significant difference among the treatment groups.</p> <p>The median length of time between randomization and discharge was 4 days in both treatment groups.</p> <p><u>Outpatient Phase</u> Primary: There was no significant difference in worsening heart failure between the tolvaptan groups and the placebo group.</p> <p>Secondary: Diuretic use decreased in all patients after discharge. There was no significant difference in mean dose reduction between the treatment groups.</p>
<p>Gheorghide et al.<sup>21</sup> (2003)</p> <p>Tolvaptan 30, 45, or 60 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients ≥18 years of age with a diagnosis of CHF irrespective of LVEF. Patients also received conventional heart failure therapy.</p>	<p>N=254</p> <p>25 days</p>	<p>Primary: Changes in body weight</p> <p>Secondary: Ankle edema measurements, urine sodium excretion, urine volume, urine osmolality, safety</p>	<p>Primary: Mean decreases from baseline in body weight were observed on the first day of tolvaptan treatment at all doses and maintained throughout the study (P&lt;0.001 vs placebo). The decrease in body weight was similar in all tolvaptan-treated patients irrespective of the LVEF. Patients receiving placebo experienced an increase in body weight from baseline.</p> <p>Secondary: Improvements in ankle edema scores were significantly better with tolvaptan 45 mg compared to placebo (P&lt;0.05). None of the other doses studies differed significantly from placebo.</p> <p>Tolvaptan-treated patients had significantly greater mean total urinary sodium excretions (339.9, 373.0, and 355 mEq for the 30, 45, and 60 mg tolvaptan groups, respectively) than placebo-treated patients (193.7 mEq; P&lt;0.05).</p> <p>Urine volumes were greater in tolvaptan-treated patients (3,909, 4,232,</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>and 4,597 mL for the 30, 45, and 60 mg tolvaptan groups, respectively) than in placebo-treated patients (2,328 mL; P&lt;0.05).</p> <p>At day one, urine osmolality decreased by 15.5, 52.4, and 118.8 mOsm/kg in the 30, 45, and 60 mg tolvaptan groups, respectively compared to an increase of 135.8 mOsm/kg in the placebo group (P&lt;0.05 for all comparisons).</p> <p>No significant differences were found between the tolvaptan groups and the placebo group in the QOL assessment. No changes in heart rate or systolic or diastolic blood pressure, supine or standing, were observed in the tolvaptan groups during the study.</p> <p>Dry mouth, thirst, and polyuria, including urinary frequency, were higher in the tolvaptan-treated patients.</p>
<p>Salahudeen et al.<sup>22</sup> (2014)</p> <p>Tolvaptan vs placebo</p> <p>Both groups received the standard of care for hyponatremia, except that patients were allowed to drink to thirst</p>	<p>DB, RCT</p> <p>Adult patients with cancer who were admitted to MD Anderson and met the eligibility criteria for nonhypovolemic hyponatremia (125 to 130 mmol/L serum sodium)</p>	<p>N=30</p> <p>14 days</p>	<p>Primary: To compare the rate of tolvaptan-treated correction of hyponatremia with that of placebo on day 14</p> <p>Secondary: To compare the length of hospital stay and the change in mental test scores between the tolvaptan-treated and placebo groups</p>	<p>Primary: Sixteen of 17 patients in the tolvaptan group and one of 13 patients in the placebo group achieved the primary endpoint of serum sodium correction on day 14 (94 vs 8%, respectively; P&lt;0.001). The study met the predefined stopping rule of superiority for tolvaptan over placebo and further patient recruitment was halted.</p> <p>Secondary: The secondary endpoints between the tolvaptan and placebo groups (mean ± standard deviation) for length of stay (21 ± 15 vs 26 ± 15 days, respectively) and changes in the MMSE score (-0.35 ± 1.66 vs 0.31 ± 2.42, respectively) were not significantly different.</p>
<p>Dahl et al.<sup>23</sup> (2012)</p> <p>Vaptans (tolvaptan,</p>	<p>MA (12 RCTs)</p> <p>Patients with cirrhosis and hyponatremia or</p>	<p>N=2,266</p> <p>Duration not specified</p>	<p>Primary: Mortality</p> <p>Secondary: Complications to</p>	<p>Primary: No clear difference between vaptans and control was found regarding mortality (22 vs 20%; RR, 1.06; 95% CI, 0.90 to 1.26).</p> <p>Secondary:</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
satavaptan*, lixivaptan*)  vs  control (no intervention, placebo, other diuretics)	ascites		cirrhosis (variceal bleeding, hepatic encephalopathy, spontaneous bacterial peritonitis, and hepatorenal syndrome), renal failure, serum sodium levels, mobilization of ascites, safety	No clear differences between vaptans and control were found regarding complications to cirrhosis and renal failure.  Treatment with vaptans increased serum sodium levels (WMD, 1.8 mmol/L; 95% CI, 0.79 to 2.96).  Treatment with vaptans reduced weight (WMD, -1.82 kg; 95% CI, -2.86 to 0.79), time to first paracentesis (RR, 0.76; 95% CI, 0.63 to 0.90), and the clinical severity of ascites (RR, 0.71; 95% CI, 0.60 to 0.83).  Adverse events were more likely with vaptan therapy compared to control (RR, 3.97; 95% CI, 1.78 to 8.83), including an excessive urine volume (RR, 9.96; 95% CI, 1.38 to 71.68). Treatment with vaptans had no effect on SBP and DBP. Treatment with vaptans increased vasopressin and renin levels; however, there is no clear difference between treatments in aldosterone levels.
Torres et al. <sup>24</sup> (2011) TEMPO 3:4  Tolvaptan 45, 60, or 90 mg QAM and 15 or 30 mg QPM  vs  placebo	DB, MC, RCT  Adults 18 to 50 years of age with ADPKD, total kidney volume (TKV) ≥750 mL, CrCl ≥ 60 mL/min	N=1,445  36 months	Primary: Annual rate of percentage change in TKV  Secondary: ADPKD progression as measured by worsening kidney function, significant kidney pain, worsening hypertension, and worsening albuminuria	Primary: Over the three-year period, total kidney volume increased by 2.8% per year (95% CI, 2.5 to 3.1) with tolvaptan versus 5.5% per year (95% CI, 5.1 to 6.0) with placebo. Tolvaptan changed the rate of growth by -2.7 percentage points per year (95% CI, -3.3 to -2.1); the ratio of the geometric means of growth rate was 0.97 (95% CI, 0.97 to 0.98; P<0.001). Tolvaptan changed the rate of growth by -2.7 percentage points per year and a mixed-model repeated-measures analysis confirmed the analysis noting percent change of 9.65% with tolvaptan vs 18.85% with placebo; a difference of -9.2 percentage points (95% CI, -11.1 to -7.3; P<0.001).  Secondary: The relative rate of ADPKD-related events was decreased by 13.5% in tolvaptan treated patients as compared with placebo (44 vs 50 events per 100 person-years (HR, 0.87; 95% CI, 0.78 to 0.97; P<0.01). Effects on worsening kidney function and kidney pain events drove the result of secondary endpoint. Tolvaptan exhibited no effect on hypertension or albuminuria events.
Raina et al. <sup>25</sup> (2021) TEMPO 3:4	Post-hoc analysis of TEMPO 3:4	N=1,445  36 months	Primary: Annual rate of change in TKV	Primary: Over the three year period, TKV in the adolescents and young adults group treated with tolvaptan increased by 3.8% per year (95% CI, 3.8 to

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Tolvaptan 45, 60, or 90 mg QAM and 15 or 30 mg QPM vs placebo	Adults 18 to 50 years of age with ADPKD, total kidney volume (TKV) $\geq$ 750 mL, CrCl $\geq$ 60 mL/min		after three years in patients aged 18 to 24 years  Secondary: Long-term safety of tolvaptan in patients 18 to 24 years of age by assessing the number of patients who had liver toxicity as defined per Hy's law	5.2) compared to 6.5% per year (95% CI, 4.2 to 8.6) with placebo. Tolvaptan decreased the rate of growth by 2.7 percentage points per year (P=0.0491) compared to placebo and represents a treatment effect of 41.2%.  Secondary: In both the AYA and adult groups, none of the patients met the criteria for Hy's law of hepatotoxicity.
Torres et al. <sup>26</sup> REPRISE  Tolvaptan 45, 60, or 90 mg QAM and 15 or 30 mg QPM vs placebo	DB, MC, RCT  Adults 18 to 55 years of age with ADPKD with eGFR 25 to 65 mL/min or 56 to 65 years of age with eGFR 25 to 44 mL/min	N=1,370  12 months	Primary: Annualized mean change in eGFR from pre-treatment baseline to post-treatment follow-up  Secondary: Mean change in annualized eGFR slope	Primary: The change from baseline in the eGFR was -2.34 ml/min (95% CI, -2.81 to -1.87) in the tolvaptan group, as compared with -3.61 ml/min (95% CI, -4.08 to -3.14) in the placebo group (difference, 1.27 ml/min; 95% CI, 0.86 to 1.68; P<0.001).  Secondary: The difference in mean change in annualized eGFR slope between tolvaptan and placebo was 1.01 mL/min (95% CI, 0.62 to 1.40; P<0.001).

\*Drug not available in the United States.

Drug regimen abbreviations: QD

Study design abbreviations: AC=active comparator, DB=double blind, ES=extended study, MA=meta analysis, MC=multicenter, OL=open label, PC=placebo controlled, RCT=randomized controlled trial  
Miscellaneous abbreviations: ADPKD=autosomal dominant polycystic kidney disease, AUC=area under the curve, CHF=congestive heart failure, CI=confidence interval, DBP=diastolic blood pressure, eGFR=estimated glomerular filtration rate, HR=hazard ratio, JVD=jugular venous distention, LVEDV=left ventricular end diastolic volume, LVEF=left ventricular ejection fraction, LVESV=left ventricular end systolic volume, PCWP=pulmonary capillary wedge pressure, QOL=quality of life, RR=relative risk, SBP=systolic blood pressure, SIADH=syndrome of inappropriate antidiuretic hormone secretion, WMD=weighted mean difference

**Additional Evidence**

Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

Stable Therapy

A search of Medline and PubMed did not reveal data pertinent to this topic.

Impact on Physician Visits

A search of Medline and PubMed did not reveal data pertinent to this topic.

**IX. Cost**

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For branded products with little or no recent utilization data, the average cost per prescription is calculated by the average wholesale price (AWP) and the standard daily dosing per product labeling. For generic products with little or no recent utilization data, the average cost per prescription is calculated by the Alabama Medicaid maximum allowable cost (MAC) and the standard daily dosage per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Vasopressin Antagonists**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
Tolvaptan	tablet	Jynarque <sup>®</sup> , Samsca <sup>®*</sup>	\$\$\$\$\$	\$\$\$\$\$

N/A=not available

**X. Conclusions**

Tolvaptan (Samsca<sup>®</sup>) is FDA-approved for the treatment of clinically significant euvolemic and hypervolemic hyponatremia (serum sodium <125 mEq/L or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction), including patients with heart failure and SIADH.<sup>2</sup> The management of hyponatremia depends on the clinical presentation and duration of the disease. Treatment must be approached carefully as overly rapid correction of hyponatremia (>10 to 12 mEq/L per 24 hours) may cause osmotic demyelination.<sup>2,3,6</sup>

There are limited guidelines available that discuss the management of hyponatremia. An expert panel provided treatment recommendations in 2013, which includes fluid restriction, sodium chloride administration, and diuresis. The panel concluded that the current role for vasopressin antagonists in SIADH is in treating mild to moderate hyponatremia and asymptomatic severe hyponatremia. Because there is a paucity of data for patients with severely symptomatic hyponatremia, hypertonic saline remains the treatment of choice in this group until more evidence-based data are available. In patients with heart failure, a vasopressin antagonist is recommended if

serum sodium does not correct to the desired level with hypertonic saline or fluid restriction. The fluid restriction should be lifted before starting these agents.<sup>3</sup>

Three short-term trials evaluating the safety and efficacy of tolvaptan in a relatively small number of patients with euvolemic or hypervolemic hyponatremia demonstrated significant improvements in serum sodium concentrations compared to fluid restriction or placebo.<sup>10,11</sup> An open-label, long-term extension study (mean follow-up of 701 days) assessed the drug-related adverse effects of tolvaptan and maintenance of efficacy, and concluded that prolonged administration of tolvaptan maintained an increased serum sodium level with an acceptable margin of safety.<sup>12</sup> Evidence suggests that hyponatremia recurs after discontinuation of tolvaptan.<sup>12,13</sup> Several other studies have evaluated the use of tolvaptan in patients with congestive heart failure as an add-on to conventional treatments.<sup>14-17,19,21,22</sup> Significant changes in body weight have been observed; however, the long-term use of tolvaptan (median duration 9.9 months) failed to demonstrate any improvements in mortality or hospitalizations for worsening heart failure.<sup>17</sup> A meta-analysis also failed to demonstrate a benefit in mortality with vaptan therapy compared to control in patients with cirrhosis and hyponatremia or ascites.<sup>23</sup>

Data supporting the use of tolvaptan are limited. It has not been established that raising serum sodium with tolvaptan provides a symptomatic benefit to patients. Patients requiring intervention to raise serum sodium urgently should not be treated with tolvaptan. Hospitalization is required for initiation and reinitiation of tolvaptan therapy so that serum sodium can be monitored closely.<sup>2</sup>

Tolvaptan is now also available under the brand name Jynarque<sup>®</sup>, which is indicated to slow kidney function decline in adults at risk of rapidly progressing autosomal dominant polycystic kidney disease (ADPKD).<sup>4</sup> Jynarque<sup>®</sup> is the only FDA-approved treatment for ADPKD. Safety and efficacy were established in two phase III trials, treating almost 3,000 adult patients with early and late stage ADPKD. Tolvaptan slowed the increase in total kidney volume and decreased ADPKD-related events, including deterioration of kidney function and pain, as compared to placebo.<sup>24,25</sup> Guidelines for the use of tolvaptan in ADPKD are limited, but the Updated Canadian Expert Consensus on Assessing Risk of Disease Progression and Pharmacological Management of Autosomal Dominant Polycystic Kidney Disease does suggest using tolvaptan in line with the FDA-approved indication.<sup>9</sup>

There is insufficient evidence to conclude that tolvaptan offers a significant clinical advantage over other alternatives in general use. Since tolvaptan is not indicated as first-line therapy for the management of hyponatremia, it should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand vasopressin antagonists within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand vasopressin antagonist is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

## XII. References

1. Vaprisol® [package insert]. Nashville (TN): Cumberland Pharmaceuticals, Inc.; 2017 Sep.
2. Samsca® [package insert]. Tokyo: Otsuka Pharmaceutical Co., Ltd.; 2021 Apr.
3. Verbalis J, Goldsmith S, Greenberg A, et al. Diagnosis, Evaluation, and Treatment of Hyponatremia: Expert Panel Recommendations. *Am J Med* 2013;126(10;Suppl 1):S1-S42.
4. Jynarque® [package insert]. Tokyo: Otsuka Pharmaceutical Co., Ltd.; 2020 Oct.
5. Torres VE and Bennett WM. Autosomal dominant polycystic kidney disease (ADPKD) in adults: Epidemiology, clinical presentation, and diagnosis. In: UpToDate, Post TW (Ed), Waltham, MA, 2021.
6. Sterns RH. Overview of the treatment of hyponatremia in adults. UpToDate, Post TW (Ed), Waltham, MA, 2021.
7. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Micromedex; 2022 [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
8. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view.
9. Soroka S, Alam A, Bevilacqua M, et al. Updated Canadian Expert Consensus on Assessing Risk of Disease Progression and Pharmacological Management of Autosomal Dominant Polycystic Kidney Disease. *Can J Kidney Health Dis*. 2018;5:2054358118801589. Published 2018 Oct 12. doi:10.1177/2054358118801589.
10. Gheorghide M, Gottlieb S, Udelson J, et al. Vasopressin v(2) receptor blockade with tolvaptan versus fluid restriction in the treatment of hyponatremia. *Am J Cardiol* 2006;97:1064-7.
11. Schrier R, Gross P, Gheorghide M, et al. Tolvaptan, a selective oral vasopressin V2-receptor antagonist, for hyponatremia. *N Engl J Med* 2006;355:2099-112.
12. Berl T, Quittnat-Pelletier F, Verbalis JG, et al. Oral tolvaptan is safe and effective in chronic hyponatremia. *J Am Soc Nephrol* 2010;21:705-12.
13. Cardenas A, Gines P, Marotta P, Czerwiec F, Ouyang J, Guevara M, et al. Tolvaptan, an oral vasopressin antagonist, in the treatment of hyponatremia in cirrhosis (abstract). *J Hepatol*. 2012 Mar;56(3):571-8.
14. Udelson J, Orlandi C, Ouyang J, et al. Acute hemodynamic effects of tolvaptan, a vasopressin V2 receptor blocker, in patients with symptomatic heart failure and systolic dysfunction: an international, multicenter, randomized, placebo-controlled trial. *J Am Coll Cardiol* 2008;52:1540-5.
15. Udelson J, McGrew F, Flores E, et al. Multicenter, randomized, double-blind, placebo-controlled study on the effect of oral tolvaptan on left ventricular dilation and function in patients with heart failure and systolic dysfunction. *J Am Coll Cardiol* 2007;49:2151-9.
16. Gheorghide M, Konstam M, Burnett J, et al. Short-term clinical effects of tolvaptan, an oral vasopressin antagonist, in patients hospitalized for heart failure: the EVEREST Clinical Status Trials. *JAMA* 2007;297:1332-43.
17. Konstam M, Gheorghide M, Burnett J, et al. Effects of oral tolvaptan in patients hospitalized for worsening heart failure: the EVEREST Outcome Trial. *JAMA* 2007;297:1319-31.
18. Pang PS, Konstam MA, Krasa HB, et al. Effects of tolvaptan on dyspnoea relief from the EVEREST trials. *Eur Heart J* 2009;30:2233-40.
19. Hauptman PJ, Burnett J, Gheorghide M, et al. Clinical course of patients with hyponatremia and decompensated systolic heart failure and the effect of vasopressin receptor antagonism with tolvaptan. *J Card Fail*. 2013 Jun;19(6):390-397.
20. Gheorghide M, Gattis W, O'Connor C, et al. Effects of tolvaptan, a vasopressin antagonist, in patients hospitalized with worsening heart failure: a randomized controlled trial. *JAMA* 2004;291:1963-71.
21. Gheorghide M, Niazi I, Ouyang J, et al. Vasopressin V2-receptor blockade with tolvaptan in patients with chronic heart failure: results from a double-blind, randomized trial. *Circulation* 2003;107:2690-6.
22. Salahudeen AK, Ali N, George M, Lahoti A, Palla S. Tolvaptan in hospitalized cancer patients with hyponatremia: a double-blind, randomized, placebo-controlled clinical trial on efficacy and safety. *Cancer*. 2014 Mar 1;120(5):744-751.
23. Dahl E, Gluud LL, Kimer N, Krag A. Meta-analysis: the safety and efficacy of vaptans (tolvaptan, satavaptan and lixivaptan) in cirrhosis with ascites or hyponatremia. *Aliment Pharmacol Ther*. 2012;36:619-626.
24. Torres VE, Chapman AB, Devuyst O, Gansevoort RT, Grantham JJ, Higashihara E, et al. Tolvaptan in patients with autosomal dominant polycystic kidney disease. *N Engl J Med*. 2012 Dec 20;367(25):2407-18.
25. Raina R, Chakraborty R, DeCoy ME, Kline T. Autosomal-dominant polycystic kidney disease: tolvaptan use in adolescents and young adults with rapid progression. *Pediatr Res*. 2021 Mar;89(4):894-899. Torres VE, Chapman AB, Devuyst O, et al; for the REPRISE Trial Investigators. Tolvaptan in later-stage autosomal dominant polycystic kidney disease. *N Engl J Med*. 2017a Nov 16;377(20):1930-1942

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Diuretics, Miscellaneous  
AHFS Class 402892  
May 4, 2022**

**I. Overview**

In July 2010, conivaptan and tolvaptan were moved from the miscellaneous diuretics class (AHFS Class 402892) to the vasopressin antagonists class (AHFS Class 402828). Currently, there are no drugs classified by AHFS as miscellaneous diuretics.

**II. Conclusions**

There are no drugs available in the miscellaneous diuretics class (AHFS Class 402892).

**III. Recommendations**

No brand miscellaneous diuretic is recommended for preferred status. Alabama Medicaid should continue to include AHFS Class 402892 in the PDL screening process. If new outpatient miscellaneous diuretics are added, it is recommended that this class be re-reviewed at that time.

**Alabama Medicaid Agency  
Pharmacy and Therapeutics Committee Meeting  
Pharmacotherapy Review of Alzheimer's Agents  
Parasympathomimetic (Cholinergic) Agents, AHFS Class 120400  
Central Nervous System Agents, Miscellaneous, AHFS Class 289200  
May 4, 2022**

## I. Overview

Alzheimer's disease is a progressive neurodegenerative disorder in older adults that affects cognition, behavior, and activities of daily living.<sup>1,2</sup> It is the most common form of dementia and the average life expectancy from the onset of symptoms to death is approximately 10 years.<sup>1-3</sup> Diagnostic features include memory impairment and one or more of the following: aphasia, apraxia, agnosia, and/or disturbance in executive functioning.<sup>1</sup>

The pathophysiologic mechanisms are not entirely understood; however, the disease is characterized by the accumulation of intracellular neurofibrillary tangles and extracellular amyloid plaques in various regions of the brain. Inflammation and free radical processes lead to neuron dysfunction and death. It is thought that memory loss is partially the result of a deficiency of cholinergic neurotransmission.<sup>2-3</sup> Glutamate, an excitatory neurotransmitter, may also play a role in the pathophysiology of Alzheimer's disease. Glutamate activates N-methyl-D-aspartate (NMDA) receptors and is involved in learning and memory. However, excessive amounts of glutamate in the brain may lead to excitotoxicity and cell death.<sup>3</sup>

Agents approved for the treatment of dementia of Alzheimer's disease include cholinesterase inhibitors (donepezil, galantamine, and rivastigmine), an NMDA receptor antagonist (memantine), and a combination product (memantine-donepezil).<sup>4-11</sup> Although none of the agents delay the progression of neurodegeneration, they do delay the progression of symptoms. The cholinesterase inhibitors enhance cholinergic function by increasing the concentration of acetylcholine through reversible inhibition of its hydrolysis by acetylcholinesterase. Memantine blocks NMDA receptors and inhibits their overstimulation by glutamate. The combination product containing memantine and donepezil (Namzaric<sup>®</sup>) was launched in May 2015 with the indication for the treatment of moderate to severe dementia of the Alzheimer's type in patients stabilized on 10 mg of donepezil hydrochloride once daily.<sup>11</sup>

Aducanumab-avwa is the first disease modifying therapy approved for the treatment of Alzheimer's disease. Treatment with aducanumab-avwa should be initiated in patients with mild cognitive impairment or mild dementia stage of disease, the population in which treatment was initiated in clinical trials. There are no safety or effectiveness data on initiating treatment at earlier or later stages of the disease than were studied. The efficacy of aducanumab-avwa was evaluated in two double-blind, randomized, placebo-controlled, parallel group studies. Of note, aducanumab received FDA approval based on a surrogate endpoint (reduction in amyloid- $\beta$  plaques) and has not yet been shown to provide a clinical benefit. Aducanumab is a human immunoglobulin (IgG1) monoclonal antibody directed against aggregated soluble and insoluble forms of amyloid- $\beta$ , a defining pathophysiological feature of Alzheimer's disease.<sup>12</sup> None of the other available pharmacologic agents slows or stops the malfunction of neurons that lead to Alzheimer's symptoms, but can be useful for reducing symptoms.<sup>13-17</sup>

The Alzheimer's agents that are included in this review are listed in Table 1. This review encompasses all dosage forms and strengths. All products with the exception of memantine-donepezil and aducanumab-avwa are available in a generic formulation. This class was last reviewed in November 2020.

**Table 1. Alzheimer's Agents Included in this Review**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
<b>Parasympathomimetic (Cholinergic Agents)</b>			
Donepezil	orally disintegrating tablet, tablet	Aricept <sup>®*</sup>	donepezil, Aricept <sup>®*</sup>
Galantamine	extended-release capsule, solution	Razadyne ER <sup>®*</sup>	galantamine
Rivastigmine	capsule, transdermal patch	Exelon <sup>®*</sup>	rivastigmine
<b>Central Nervous System Agents, Miscellaneous</b>			

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Current PDL Agent(s)
Aducanumab-avwa	injection	Aduhelm®	none
Memantine	extended-release capsule, solution, tablet	Namenda®*, Namenda XR®*	memantine
<b>Combination Products</b>			
Memantine and donepezil	extended-release capsule	Namzaric®	none

\*Generic is available in at least one dosage form or strength.  
PDL=Preferred Drug List.

## II. Evidence-Based Medicine and Current Treatment Guidelines

Current treatment guidelines that incorporate the use of the Alzheimer's agents are summarized in Table 2.

**Table 2. Treatment Guidelines Using the Alzheimer's Agents**

Clinical Guideline	Recommendation(s)
European Federation of Neurological Societies: <b>Guidelines for the Diagnosis and Management of Alzheimer's Disease (2010)</b> <sup>14</sup>	<ul style="list-style-type: none"> <li>• Patients and caregivers should be provided with education and support.</li> <li>• There is insufficient evidence to support the use of any drugs purely for the primary prevention of dementia. Cholinesterase inhibitors, vitamin E, ginkgo and estrogens should not be used as treatments for those with mild cognitive impairment.</li> <li>• In patients with Alzheimer's disease, treatment with cholinesterase inhibitors (donepezil, galantamine, or rivastigmine) should be considered at the time of diagnosis, taking into account expected therapeutic benefits and potential safety issues. Benefits on cognitive and non-cognitive symptoms have been demonstrated in those with mild, moderate and severe disease. Realistic expectations for treatment effects and potential side effects should be discussed with the patient and caregivers.</li> <li>• In patients with moderate to severe Alzheimer's disease, treatment with memantine should be considered taking into account expected therapeutic benefits and potential safety issues. Benefits on cognitive and noncognitive symptoms are apparent, some non-cognitive symptoms (agitation, delusions) may respond better than others. Realistic expectations for treatment effects and potential side effects should be discussed with the patient and caregivers.</li> <li>• Regular patient follow-up should be an integral part of management.</li> <li>• Aspirin should not be used as a treatment for Alzheimer's disease, though it can be used in those with Alzheimer's disease who also have other indications for its use (e.g. to prevent cardiovascular events).</li> <li>• Vitamin E should not be used as a treatment for Alzheimer's disease.</li> <li>• Currently, there is insufficient evidence to support the use of other agents including, anti-inflammatory drugs, nootropics (including piracetam, nicergoline), selegiline, oestrogens, pentoxifylins, or statins in the treatment or prevention of Alzheimer's disease.</li> <li>• Cognitive stimulation or rehabilitation may be considered in patients with mild to moderate Alzheimer's disease.</li> <li>• Management of behavioral and psychological symptoms of dementia should begin with a careful search for triggers and causative factors (i.e. physical illness). Where possible, initial treatment should be non-pharmacological.</li> <li>• Antipsychotics should only be used for moderate or severe behavioral and psychological symptoms of dementia causing significant distress which have either not responded to other treatments (like non-pharmacological measures or cholinesterase inhibitors) or when other treatments are not appropriate. Low dose of atypical agents should be used only after assessment of risk benefit and full discussion with patient (when capacity allows) and caregiver.</li> <li>• Atypical agents have fewer side effects and do not confer a greater risk of stroke or mortality than conventional drugs.</li> </ul>



Clinical Guideline	Recommendation(s)
National Institute for Health and Care Excellence: <b>Dementia: assessment, management and support for people living with dementia and their carers (2018)</b> <sup>15</sup>	<ul style="list-style-type: none"> <li>• Selective serotonin reuptake inhibitors rather than tricyclic antidepressants should be used to treat depression in Alzheimer's disease.</li> </ul> <p><u>Pharmacological management of Alzheimer's disease</u></p> <ul style="list-style-type: none"> <li>• The three acetylcholinesterase (AChE) inhibitors donepezil, galantamine and rivastigmine as monotherapies are recommended as options for managing mild to moderate Alzheimer's disease.</li> <li>• Memantine monotherapy is recommended as an option for managing Alzheimer's disease for people with:             <ul style="list-style-type: none"> <li>○ moderate Alzheimer's disease who are intolerant of or have a contraindication to AChE inhibitors or</li> <li>○ severe Alzheimer's disease.</li> </ul> </li> <li>• For people with an established diagnosis of Alzheimer's disease who are already taking an AChE inhibitor:             <ul style="list-style-type: none"> <li>○ consider memantine in addition to an AChE inhibitor if they have moderate disease</li> <li>○ offer memantine in addition to an AChE inhibitor if they have severe disease.</li> </ul> </li> <li>• Treatment should be under the following conditions:             <ul style="list-style-type: none"> <li>○ For people who are not taking an AChE inhibitor or memantine, prescribers should only start treatment with these on the advice of a clinician who has the necessary knowledge and skills. This could include:                 <ul style="list-style-type: none"> <li>▪ secondary care medical specialists such as psychiatrists, geriatricians and neurologists</li> <li>▪ other healthcare professionals (such as general practitioners (GPs), nurse consultants and advanced nurse practitioners), if they have specialist expertise in diagnosing and treating Alzheimer's disease.</li> </ul> </li> <li>○ Once a decision has been made to start an AChE inhibitor or memantine, the first prescription may be made in primary care.</li> <li>○ For people with an established diagnosis of Alzheimer's disease who are already taking an AChE inhibitor, primary care prescribers may start treatment with memantine without taking advice from a specialist clinician.</li> <li>○ Do not stop AChE inhibitors in people with Alzheimer's disease because of disease severity alone.</li> </ul> </li> <li>• If prescribing an AChE inhibitor (donepezil, galantamine or rivastigmine), treatment should normally be started with the drug with the lowest acquisition cost (taking into account required daily dose and the price per dose once shared care has started). However, an alternative AChE inhibitor could be prescribed if it is considered appropriate when taking into account adverse event profile, expectations about adherence, medical comorbidity, possibility of drug interactions and dosing profiles.</li> <li>• When using assessment scales to determine the severity of Alzheimer's disease, healthcare professionals should take into account any physical, sensory or learning disabilities, or communication difficulties that could affect the results and make any adjustments they consider appropriate. Healthcare professionals should also be mindful of the need to secure equality of access to treatment for patients from different ethnic groups, in particular those from different cultural backgrounds.</li> <li>• When assessing the severity of Alzheimer's disease and the need for treatment, healthcare professionals should not rely solely on cognition scores in circumstances in which it would be inappropriate to do so. These include:             <ul style="list-style-type: none"> <li>○ if the cognition score is not, or is not by itself, a clinically appropriate tool for assessing the severity of that patient's dementia because of the patient's learning difficulties or other disabilities (for example, sensory impairments), linguistic or other communication difficulties or level of education or</li> </ul> </li> </ul>

Clinical Guideline	Recommendation(s)
	<ul style="list-style-type: none"> <li>○ if it is not possible to apply the tool in a language in which the patient is sufficiently fluent for it to be appropriate for assessing the severity of dementia or</li> <li>○ if there are other similar reasons why using a cognition score, or the score alone, would be inappropriate for assessing the severity of dementia.</li> <li>○ In such cases healthcare professionals should determine the need for initiation or continuation of treatment by using another appropriate method of assessment.</li> </ul> <ul style="list-style-type: none"> <li>● Do not offer the following specifically to slow the progress of Alzheimer's disease, except as part of a randomized controlled trial:           <ul style="list-style-type: none"> <li>○ diabetes medicines</li> <li>○ hypertension medicines</li> <li>○ statins</li> <li>○ non-steroidal anti-inflammatory drugs (NSAIDs), including aspirin.</li> </ul> </li> </ul> <p>Pharmacological management of non-Alzheimer's dementia</p> <ul style="list-style-type: none"> <li>● Offer donepezil or rivastigmine to people with mild to moderate dementia with Lewy bodies.</li> <li>● Only consider galantamine for people with mild to moderate dementia with Lewy bodies if donepezil and rivastigmine are not tolerated.</li> <li>● Consider donepezil or rivastigmine for people with severe dementia with Lewy bodies.</li> <li>● Consider memantine for people with dementia with Lewy bodies if AChE inhibitors are not tolerated or are contraindicated.</li> <li>● Only consider AChE inhibitors or memantine for people with vascular dementia if they have suspected comorbid Alzheimer's disease, Parkinson's disease dementia or dementia with Lewy bodies.</li> <li>● Do not offer AChE inhibitors or memantine to people with frontotemporal dementia.</li> <li>● Do not offer AChE inhibitors or memantine to people with cognitive impairment caused by multiple sclerosis.</li> </ul>
<p>American Academy of Neurology:  <b>Practice Guideline Update Summary: Mild Cognitive Impairment (2018)</b><sup>16</sup></p> <p>Reaffirmed January 30, 2021</p>	<p><u>Pharmacologic treatments for patients diagnosed with mild cognitive impairment (MCI)</u></p> <ul style="list-style-type: none"> <li>● Donepezil use over three years is possibly ineffective for reducing the chances of a progression to possible or probably Alzheimer dementia. In patients with MCI, it is unknown whether donepezil slows progression on various cognitive scales.</li> <li>● Galantamine use over 24 months is probably ineffective for reducing progression to dementia.</li> <li>● Rivastigmine use up to 48 months is possibly ineffective for reducing the rate of progression to possible or probable Alzheimer dementia.</li> </ul> <p><u>Recommendations for management of MCI</u></p> <ul style="list-style-type: none"> <li>● For patients diagnosed with MCI, clinicians should wean patients from medications that can contribute to cognitive impairment (where feasible and medically appropriate) and treat modifiable risk factors that may be contributing.</li> <li>● There are no FDA-approved medications for the treatment of MCI. Moreover, there are no high-quality, long-term studies identifying pharmacologic or dietary agents that either improve cognition or delay progression in patients with MCI. For patients diagnosed with MCI, clinicians should counsel the patients and families that there are no pharmacologic or dietary agents currently shown to have symptomatic cognitive benefit in MCI and that no medications are FDA-approved for this purpose.</li> <li>● Studies of cholinesterase inhibitors showed no benefit on cognitive outcomes or reduction in progression from MCI to dementia, although some studies could not exclude an important effect. In addition to lacking efficacy, side effects of cholinesterase inhibitors are common, including gastrointestinal symptoms and</li> </ul>

Clinical Guideline	Recommendation(s)
	<p>cardiac concerns. For patients diagnosed with MCI, clinicians may choose not to offer cholinesterase inhibitors. If clinicians choose to offer cholinesterase inhibitors, they must first discuss with patients the fact that this is an off-label prescription not currently backed by empirical evidence.</p> <ul style="list-style-type: none"> <li>• For patients diagnosed with MCI who are interested in pharmacologic treatment, clinicians may inform these patients of centers or organizations that can connect patients to clinical trials.</li> <li>• For patients diagnosed with MCI, clinicians should recommend regular exercise (twice per week) as part of an overall approach to management.</li> <li>• In patients with MCI, cognitive interventions may be beneficial in improving measures of cognitive function.</li> </ul>
<p>American College of Physicians/American Academy of Family Physicians:  <b>Current Pharmacologic Treatment of Dementia: A Clinical Practice Guideline (2008)</b><sup>17</sup></p>	<ul style="list-style-type: none"> <li>• The decision to initiate therapy should be based on evaluation of benefits and risks associated with an individual patient. All of the drugs have known adverse events, and the decision to manage patients with dementia should balance harms against modest or even no benefit.</li> <li>• Although the evidence shows statistically significant benefits of treatment with some cholinesterase inhibitors and memantine for all kinds of dementia, these benefits, on average, are not clinically significant for cognition and are modest for global assessments. Currently, there is no way to predict which patients might have a clinically important response. The evidence does not support prescribing these medications for every patient with dementia.</li> <li>• Evidence is insufficient to determine the optimal duration of therapy. No evidence demonstrates when it is appropriate to stop the treatment if the patient becomes unresponsive or shows decline in various domains of dementia. If slowing decline is no longer a goal, treatment with memantine or a cholinesterase inhibitor is no longer appropriate.</li> <li>• The evidence is insufficient to compare the effectiveness of different pharmacologic agents for the treatment of dementia. Because few trials compare one drug with another, evidence about effectiveness is insufficient to support the choice of specific drugs for the treatment of dementia. Assessment of the effectiveness of combination therapy is lacking.</li> <li>• Clinicians should base the choice of pharmacologic agents on tolerability, adverse effect profile and ease of use.</li> </ul>
<p>The Movement Disorder Society:  <b>Evidence-Based Medicine Review Update: Treatments for the non-motor symptoms of Parkinson's disease (2019)</b><sup>18</sup></p>	<p><u>Treatment of dementia in Parkinson's disease</u></p> <ul style="list-style-type: none"> <li>• Rivastigmine is efficacious for the treatment of dementia in Parkinson's disease.</li> <li>• There is insufficient evidence for donepezil and galantamine for the treatment of dementia in Parkinson's disease.</li> <li>• Safety conclusions are that the acetylcholinesterase inhibitors donepezil, rivastigmine, and galantamine have an acceptable risk without specialized monitoring.</li> <li>• The practice implications are that rivastigmine is clinically useful for the treatment of dementia in Parkinson's disease, while the practice implications for donepezil and galantamine are that they are both possibly useful for the treatment of dementia in Parkinson's disease.</li> <li>• The practice implications for memantine are that it is investigational for the treatment of dementia in Parkinson's disease.</li> </ul> <p><u>Treatment of non-dementia cognitive impairment in Parkinson's disease</u></p> <ul style="list-style-type: none"> <li>• There is "insufficient evidence" to conclude on the efficacy of rivastigmine or rasagiline for the treatment of cognitive impairment in Parkinson's disease; practice implications are investigational.</li> </ul>

### III. Indications

The Food and Drug Administration (FDA)-approved indications for the Alzheimer's agents are noted in Table 3. While agents within this therapeutic class may have demonstrated positive activity via in vitro trials, the clinical significance of this activity remains unknown until fully demonstrated in well-controlled, peer-reviewed in vivo clinical trials. As such, this review and the recommendations provided are based exclusively upon the results of such clinical trials.

**Table 3. FDA-Approved Indications for the Single Entity Alzheimer's Agents<sup>4</sup>**

Indication	Parasympathomimetic (Cholinergic Agents)			Central Nervous System Agents, Miscellaneous	
	Donepezil	Galantamine	Rivastigmine	Aducanumab-avwa	Memantine
Mild-to-moderate dementia of the Alzheimer's type		✓	✓ †		
Mild, moderate, and severe dementia of the Alzheimer's type	✓		✓ ‡		
Moderate-to-severe dementia of the Alzheimer's type					✓
Mild-to-moderate dementia associated with Parkinson's disease			✓		
Treatment of Alzheimer's disease in patients with mild cognitive impairment or mild dementia stage of disease				✓	

†Capsule and solution.

‡Transdermal patch.

**Table 4. FDA-Approved Indications for the Combination Product Alzheimer's Agents<sup>4</sup>**

Indication	Memantine and Donepezil
Moderate-to-severe dementia of the Alzheimer's type	✓ *

\*In patients stabilized on 10 mg of donepezil hydrochloride once daily.

### IV. Pharmacokinetics

The pharmacokinetic parameters of the Alzheimer's agents are listed in Table 5. Pharmacokinetic properties of the combination products are in line with the properties of their individual components listed below.

**Table 5. Pharmacokinetic Parameters of the Alzheimer's Agents<sup>5</sup>**

Generic Name(s)	Bioavailability (%)	Protein Binding (%)	Metabolism (%)	Excretion (%)	Half-Life
<b>Parasympathomimetic (Cholinergic Agents)</b>					
Donepezil	100	96	Liver (% not reported)	Renal (57) Feces (9 to 15)	70 hours*
Galantamine	90	18	Liver (75)	Renal (95) Feces (5)	7 hours
Rivastigmine	Oral: 36 to 72	40	Liver, extensive Brain, extensive	Renal (>90)	Oral: 1.4 to 1.7 hours Transdermal: 3.0 hours
<b>Central Nervous System Agents, Miscellaneous</b>					
Aducanumab-avwa	Not reported	Not reported	Liver (% not reported)	Not reported	24.8 days
Memantine	Well absorbed	45	Liver, partial	Renal (48)	60 to 80 hours

\* Half-life of 104 hours in subjects over 55 years of age.

## V. Drug Interactions

Major drug interactions with the Alzheimer's agents are listed in Table 6.

**Table 6. Major Drug Interactions with the Alzheimer's Agents<sup>5</sup>**

Generic Name(s)	Interaction	Mechanism
<b>Parasympathomimetic (Cholinergic Agents)</b>		
Donepezil	Azole antifungals	Concomitant use of donepezil, a CYP3A4 substrate that is associated with prolongation of the QT interval, is contraindicated with certain drugs that prolong QT interval and strongly inhibit CYP3A4.
Donepezil	Anticholinergic agents	Concomitant use of a cholinesterase inhibitor, such as donepezil, and an anticholinergic agent may result in interference with the efficacy of both agents.
Donepezil	QT interval prolonging agents	Concurrent use of donepezil and QT interval prolonging agents may result in increased risk of QT-interval prolongation and torsade de pointes.
Donepezil	CYP3A4 inhibitors and inducers	Concurrent use of donepezil and CYP3A4 inhibitors/ inducers may result in increased/decreased donepezil exposure.
Donepezil	Select CYP2D6 inhibitors (clobazam, terbinafine, cinacalcet, peginterferon alfa-2b)	Concurrent use of donepezil and selected CYP2D6 inhibitors may result in increased donepezil exposure.
Donepezil	Seizure threshold lowering agents	Concurrent use of donepezil and seizure threshold lowering agents may result in reduced seizure threshold.
Galantamine	QT interval prolonging agents	Concurrent use of galantamine and QT interval prolonging agents may result in increased risk of QT-interval prolongation and torsade de pointes.
Rivastigmine	Metoclopramide	The concomitant use of metoclopramide and rivastigmine is contraindicated due to potential additive effects of extrapyramidal reactions.
Rivastigmine	Beta blockers	Concomitant use of rivastigmine and beta-blockers, especially cardioselective agents, is not recommended due to additive bradycardic effects resulting in syncope.
<b>Central Nervous System Agents, Miscellaneous</b>		
Memantine	Dextromethorphan, Amantadine, Ketamine	Concurrent use of memantine and selected N-methyl-D-aspartate antagonists may result in increased adverse events of N-methyl-D-aspartate antagonists.
Memantine	Carbonic anhydrase inhibitors	Concurrent use of memantine and carbonic anhydrase inhibitors may result in reduced clearance of memantine due to urinary alkalization.

## VI. Adverse Drug Events

The most common adverse drug events reported with the Alzheimer's agents are listed in Table 7. Adverse drug reactions associated with the combination products are in line with the individual components listed below.<sup>4</sup>

**Table 7. Adverse Drug Events (%) Reported with the Single Entity Alzheimer's Agents<sup>4,6-12</sup>**

Adverse Events	Parasympathomimetic (Cholinergic Agents)			Central Nervous System Agents, Miscellaneous	
	Donepezil	Galantamine	Rivastigmine	Aducanumab-avwa	Memantine
<b>Cardiovascular</b>					
Angina pectoris	-	-	≥1	-	-
Atrial fibrillation	≥1	-	≥1	-	-
Bradycardia	≥1	2	≥1	-	-

Adverse Events	Parasympathomimetic (Cholinergic Agents)			Central Nervous System Agents, Miscellaneous	
	Donepezil	Galantamine	Rivastigmine	Aducanumab-avwa	Memantine
Chest pain	1 to 2	≥1	-	-	-
Heart failure	-	-	≥1	-	≥1
Hemorrhage	2	-	-	-	-
Hypertension	1 to 3	-	3	-	4
Hypotension	≥1	-	≥1	-	-
Myocardial infarction	-	-	≥1	-	-
Palpitation	-	-	≥1	-	-
Peripheral edema	≥1	-	-	-	≥2
Postural hypotension	-	-	≥1	-	-
Syncope	2	2	3	-	≥1
Vasodilation	≥1	-	-	-	-
<b>Central Nervous System</b>					
Abnormal crying	≥1	-	-	-	-
Abnormal dreams	3	-	-	-	-
Aggression	≥1	-	3	-	≥1
Agitation	-	-	≥1	-	≥2
Anxiety	-	-	4 to 5; 3*	-	≥2
Aphasia	≥1	-	-	-	-
Bradykinesia	-	-	≥1	-	-
Cerebrovascular accident	-	-	-	-	≥1
Confusion	2	-	1 to 8	8	6
Convulsion	≥1	-	≥1	-	-
Delusions	≥1	-	-	-	-
Depression	2 to 3	7	1 to 6; 4*	-	≥2
Dizziness	2 to 8	9	6 to 21; 2 to 7*	-	7
Dyskinesia	-	-	≥1	-	-
Emotional lability	2	-	-	-	-
Fatigue	5	5	4 to 9; 2*	-	2
Gait abnormality	-	-	≥1	-	≥2
Hallucination	3	-	4	-	3
Headache	3 to 10	8	4 to 17; 3 to 4*	21	6
Hostility	3	-	-	-	-
Hypokinesia	-	-	-	-	≥1
Insomnia	2 to 14	5	3 to 9; 1 to 4*	-	≥2
Irritability	≥1	-	-	-	-
Malaise	-	≥1	5	-	-
Nervousness	1 to 3	-	-	-	-
Paranoid reaction	-	-	≥1	-	-
Paresthesia	≥1	-	≥1	-	-
Parkinson's disease worsening	-	-	3	-	-
Parkinsonism	-	-	2	-	-
Personality disorder	2	-	-	-	-
Restlessness	≥1	-	≥1	-	-
Somnolence	2	4	4 to 5	-	3
Transient ischemic attack	-	-	≥1	-	≥1
Tremor	≥1	3	4 to 10; ≥1*	-	-
Vertigo	≥1	-	≥1; 0 to 2*	-	≥1
Wandering	≥1	-	-	-	-
<b>Dermatological</b>					
Diaphoresis	≥1	-	4	-	-

Adverse Events	Parasympathomimetic (Cholinergic Agents)			Central Nervous System Agents, Miscellaneous	
	Donepezil	Galantamine	Rivastigmine	Aducanumab-avwa	Memantine
Eczema	3	-	-	-	-
Pruritus	≥1	-	≥1*	-	-
Rash	≥1	-	≥1	-	≥1
Skin ulcer	≥1	-	-	-	-
Urticaria	≥1	-	-	-	-
<b>Gastrointestinal</b>					
Abdominal pain	≥1	5	4 to 13; 2 to 4*	-	-
Anorexia	4 to 8	7 to 9	6 to 17; 3 to 9*	-	≥2
Bloating	≥1	-	-	-	-
Constipation	≥1	-	5; ≥1*	-	5
Diarrhea	5 to 15	6 to 12	7 to 19; 6 to 10*	9	≥2
Dyspepsia	≥1	5	1 to 9	-	-
Epigastric pain	≥1	-	-	-	-
Eructation	-	-	2	-	-
Fecal incontinence	≥1	-	≥1	-	-
Flatulence	-	≥1	4	-	-
Gastritis	-	-	≥1; ≥1*	-	-
Gastrointestinal bleeding	≥1	-	-	-	-
Nausea	3 to 19	13 to 24	29 to 47; 7 to 21*	-	≥2
Toothache	≥1	-	-	-	-
Vomiting	3 to 9	6 to 13	17 to 31; 6 to 19*	-	3
Weight decrease	1 to 3	5 to 7	3; 3 to 8*	-	≥1
<b>Genitourinary</b>					
Cystitis	≥1	-	-	-	-
Frequent urination	2	-	-	-	≥1
Glycosuria	≥1	-	-	-	-
Hematuria	≥1	3	≥1	-	-
Libido increased	≥1	-	-	-	-
Urinary incontinence	2	≥1	≥1*	-	≥2
Urinary tract infection	≥1	8	7; 2*	-	≥2
<b>Laboratory Test Abnormalities</b>					
Alkaline phosphatase increased	≥1	-	-	-	≥1
Creatinine increased	3	-	-	-	-
Hyperlipemia	2	-	-	-	-
Hypokalemia	-	-	≥1	-	-
Lactate dehydrogenase increased	≥1	-	-	-	-
<b>Musculoskeletal</b>					
Arthralgia	-	-	-	-	≥2
Arthritis	1 to 2	-	≥1	-	-
Asthenia	≥1	≥1	2 to 6; 2 to 3*	-	-
Ataxia	≥1	-	≥1	-	≥1
Back pain	3	-	≥1	-	3
Bone fracture	≥1	-	-	-	-
Leg cramps	-	-	≥1	-	-
Muscle cramps	3 to 8	-	-	-	-
Myalgia	-	-	≥1	-	-
Rigors	-	-	≥1	-	-

Adverse Events	Parasympathomimetic (Cholinergic Agents)			Central Nervous System Agents, Miscellaneous	
	Donepezil	Galantamine	Rivastigmine	Aducanumab-avwa	Memantine
<b>Respiratory</b>					
Bronchitis	≥1	-	-		≥2
Cough increased	≥1	-	-		4
Dyspnea	≥1	-	≥1		2
Pharyngitis	≥1	-	-		-
Pneumonia	≥1	-	≥1*		≥1
Respiratory tract infection	-	-	-		≥2
Rhinitis	-	4	4		-
Sore Throat	≥1	-	-		-
<b>Special Senses</b>					
Blurred vision	≥1	-	-		-
Cataract	≥1	-	≥1		≥1
Conjunctivitis	-	-	-		≥1
Eye irritation	≥1	-	-		-
Tinnitus	-	-	≥1		-
<b>Other</b>					
Accident	7 to 13	-	-		-
Accidental trauma	-	-	1 to 10		-
Allergy	-	-	≥1		-
Anemia	-	3	≥1; ≥1*		≥1
ARIA-E				35	
ARIA-H microhemorrhage				19	
ARIA-H superficial siderosis				15	
Dehydration	1 to 2	-	1 to 2; ≥1*		-
Ecchymosis	4 to 5	-	-		-
Edema	≥1	-	≥1		-
Epistaxis	-	-	≥1		-
Fall	-	-	≥1*	15	≥2
Fever	2	≥1	≥1		-
Flu syndrome	≥1	-	3		≥2
Hot flashes	≥1	-	≥1		-
Infection	1 to 11	-	-		-
Inflicted injury	-	-	-		≥2
Influenza	≥1	-	-		-
Pain	3 to 9	-	-		3

ARIA-E=amyloid-related imaging abnormalities-edema, ARIA-H=Amyloid-related imaging abnormalities-hemosiderin deposition

✓ Percent not specified.

- Event not reported or incidence <1%.

\*Transdermal patch.

## VII. Dosing and Administration

The usual dosing regimens for the Alzheimer's agents are listed in Table 8.

**Table 8. Usual Dosing Regimens for the Alzheimer's Agents<sup>4,6-12</sup>**

Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability
<b>Parasympathomimetic (Cholinergic Agents)</b>			
Donepezil	Dementia of the Alzheimer's type (mild to moderate); Tablet and orally disintegrating tablet: initial, 5 mg daily; may increase to 10 mg daily after four to six weeks;	Safety and efficacy not established in the pediatric population.	Orally disintegrating tablet: 5 mg 10 mg



Generic Name(s)	Usual Adult Dose	Usual Pediatric Dose	Availability												
	<p>maintenance, 5 to 10 mg daily</p> <p><u>Dementia of the Alzheimer's type (moderate to severe):</u> Tablet: initial, 5 mg daily; may increase to 10 mg daily after four to six weeks; may increase to 23 mg daily after three months on 10 mg daily dose</p> <p>Orally disintegrating tablet: initial, 5 mg daily; may increase to 10 mg daily after four to six weeks</p>		<p>Tablet: 5 mg 10 mg 23 mg</p>												
Galantamine	<p><u>Mild-to-moderate dementia of the Alzheimer's type:</u> Extended-release capsule: initial, 8 mg daily; maintenance, 16 to 24 mg daily</p> <p>Oral solution: initial, 4 mg twice a day with the morning and evening meals; maintenance: 8 to 12 mg twice a daily</p>	Safety and efficacy not established in the pediatric population.	<p>Extended-release capsule: 8 mg 16 mg 24 mg</p> <p>Solution: 4 mg/mL</p>												
Rivastigmine	<p><u>Mild-to-moderate dementia of the Alzheimer's type:</u> Capsule: initial, 1.5 mg twice daily with the morning and evening meals; maintenance, 3 to 6 mg twice daily</p> <p>Transdermal patch: initial, 4.6 mg/24 hours; maintenance, 9.5 or 13.3 mg/24 hours</p> <p><u>Severe dementia of the Alzheimer's type:</u> Transdermal patch: initial, 4.6 mg/24 hours; maintenance, 13.3 mg/24 hours</p> <p><u>Mild-to-moderate dementia associated with Parkinson's disease:</u> Capsule: initial, 1.5 mg twice daily with the morning and evening meals; maintenance, 3 to 6 mg twice daily</p> <p>Transdermal patch: initial, 4.6 mg/24 hours; maintenance, 9.5 or 13.3 mg/24 hours</p>	Safety and efficacy not established in the pediatric population.	<p>Capsule: 1.5 mg 3 mg 4.5 mg 6 mg</p> <p>Transdermal patch: 4.6 mg/24 hours 9.5 mg/24 hours 13.3 mg/24 hours</p>												
<b>Central Nervous System Agents, Miscellaneous</b>															
Aducanumab-avwa	<p><u>Alzheimer's disease:</u> Solution for IV injection: infuse every 28 days based on the following titration:</p> <table border="1"> <thead> <tr> <th>Infusion (every 28 days)</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>1 mg/kg</td> </tr> <tr> <td>2</td> <td>1 mg/kg</td> </tr> <tr> <td>3</td> <td>3 mg/kg</td> </tr> <tr> <td>4</td> <td>3 mg/kg</td> </tr> <tr> <td>5</td> <td>6 mg/kg</td> </tr> </tbody> </table>	Infusion (every 28 days)	Dose	1	1 mg/kg	2	1 mg/kg	3	3 mg/kg	4	3 mg/kg	5	6 mg/kg	Safety and efficacy not established in the pediatric population.	<p>Injection: 170 mg/1.7 mL 300 mg/3 mL</p>
Infusion (every 28 days)	Dose														
1	1 mg/kg														
2	1 mg/kg														
3	3 mg/kg														
4	3 mg/kg														
5	6 mg/kg														

Generic Name(s)	Usual Adult Dose		Usual Pediatric Dose	Availability
	6	6 mg/kg		
	7 and beyond	10 mg/kg		
Memantine	<p><u>Moderate-to-severe dementia of the Alzheimer's type:</u>            Solution and tablet: initial, 5 mg once daily, increase dose by 5 mg at weekly intervals (twice daily dosing); maintenance, 10 mg twice daily</p> <p>Extended release capsule: initial, 7 mg once daily; maintenance, 28 mg once daily</p>		Safety and efficacy not established in the pediatric population.	<p>Extended release capsule:            7 mg            14 mg            21 mg            28 mg</p> <p>Extended release capsule dose pack:            7 mg (7 count)-14 mg (7 count)-21 mg (7 count)-28 mg (7 count)</p> <p>Solution:            10 mg/5 mL</p> <p>Tablet:            5 mg            10 mg</p> <p>Tablet dose pack:            5 mg (28 count)-10 mg (21 count)</p>
<b>Combination products</b>				
Memantine and donepezil	<p><u>Moderate-to-severe dementia of the Alzheimer's type:</u>            Extended release capsule: patients stabilized on memantine hydrochloride (10 mg twice daily or 28 mg extended-release once daily) and donepezil hydrochloride 10 mg can be switched to 28 mg-10 mg combination capsule, taken once a day in the evening</p>		Safety and efficacy not established in the pediatric population.	<p>Extended release capsule:            7-10 mg            14-10 mg            21-10 mg            28-10 mg</p> <p>Extended release capsule dose pack:            7-10 mg (7 count)-14-10 mg (7 count)-21-10 mg (7 count)-28-10 mg (7 count)</p>

## VIII. Effectiveness

Clinical studies evaluating the safety and efficacy of the Alzheimer's agents are summarized in Table 9.

**Table 9. Comparative Clinical Trials with the Alzheimer's Agents**

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<b>Dementia of Alzheimer's Disease</b>				
Geldmacher et al. <sup>19</sup> (2003)  Donepezil 5 mg/day	OS  Patients with Alzheimer's disease	N=1,115  Variable duration	Primary: Time to nursing home placement  Secondary: Not reported	Primary: Use of donepezil of 5 mg/day or more was associated with significant delays in nursing home placement.  A cumulative dose-response relationship was observed between longer-term sustained donepezil use and delay of nursing home placement.  When donepezil was taken at effective doses for at least nine to 12 months, conservative estimates of the time gained before nursing home placement were 21.4 months for first-dementia-related nursing home placement and 17.5 months for permanent nursing home placement.  Secondary: Not reported
Burns et al. <sup>20</sup> (2007)  Donepezil 5 to 10 mg/day	MC, OL  Patients ≥50 years of age with mild-to-moderate Alzheimer's disease	N=579  132 weeks	Primary: ADAS-cog, CDR-SB, IDDD, QoLS, and adverse events  Secondary: Not reported	Primary: Mean changes in ADAS-cog scores of all patients were improved by approximately two points after six weeks (cumulative week 36) and one point after 12 weeks (cumulative week 42), with improvement compared to the start of OL treatment.  At week 24 (cumulative week 54), mean ADAS-cog scores still showed improvement (approximately 0.5 points) compared to those scores reported at the start of OL treatment. From 24 weeks, ADAS-cog scores declined over the remainder of the study. At the end of 132 weeks of OL treatment (162 weeks total follow-up), the change from DB baseline was 15.6 points for all patients. No difference was seen between patients who had previously received placebo in the DB phase vs those receiving donepezil for the entire treatment period.  CDR-SB scores improved slightly over the first 12 weeks (up to cumulative week 42) of OL treatment and then slowly declined for the

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>remainder of the study period (up to cumulative week 162).</p> <p>Mean IDDD total scores were maintained over the first 24 weeks of OL treatment to within approximately 1 point relative to those at the beginning of this study period. Mean IDDD scores were 138.1 at week 0, 136.9 at week 12, 138.9 at week 24 and 170.8 at week 132 (162 weeks of total follow-up).</p> <p>At the start of the OL extension, QoLS scores were improved compared to baseline, with a mean change of 3.03. The scores remained above the baseline level at weeks six and 12 of OL treatment. At the end of 132 weeks of OL treatment, the decline from the baseline for the DB study was -46.2.</p> <p>Overall, 85% of patients experienced at least one treatment-emergent adverse event. The most common adverse events included diarrhea (12%), nausea (11%), infection (11%) and accidental injury (10%). Nonfatal all-causality and treatment-related serious adverse events were reported for 25 and 7% of patients, respectively.</p> <p>Seventeen patients died during the study or within four weeks after discontinuation of donepezil. The most common causes of death were pneumonia (seven patients) and cerebrovascular accident (two patients). Fifteen deaths were considered unrelated to donepezil. Two deaths, one due to a cerebral hemorrhage diagnosed on day five of treatment and another due to a suspected myocardial infarction on day 55, were considered by the investigators to be possibly related to donepezil.</p> <p>Secondary: Not reported</p>
<p>Hashimoto et al.<sup>21</sup> (2009)</p> <p>Donepezil 5 mg/day</p>	<p>OS, PRO</p> <p>Patients with Alzheimer's disease</p>	<p>N=416</p> <p>12 weeks</p>	<p>Primary: MMSE</p> <p>Secondary: Not reported</p>	<p>Primary: There were significant changes in mean scores on the MMSE (0.9; P&lt;0.01) from baseline to week 12.</p> <p>There was a significant decrease in the personal strain score at week 12 (P=0.002). There was no significant improvement was in role strain.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>There was no significant decrease in the time spent supervising Alzheimer's disease patients.</p> <p>Secondary: Not reported</p>
<p>Homma et al.<sup>22</sup> (2009)</p> <p>Donepezil 10 mg/day</p>	<p>OL</p> <p>Japanese patients ≥50 years of age with severe Alzheimer's disease (modified Hachinski Ischemic Score ≤6, FAST ≥6, MMSE score of 1 to 12)</p>	<p>N=189</p> <p>52 weeks</p>	<p>Primary: SIB, and BEHAVE-AD</p> <p>Secondary: Not reported</p>	<p>Primary: The mean change in SIB scores during the OL study showed improvement until week 24, followed by a decline by week 36. For those patients receiving 52 weeks of treatment, the mean change in SIB from baseline (enrollment in OL study) was -6.1. The mean change in SIB declined more rapidly after 24 weeks.</p> <p>For the BEHAVE-AD, little change was observed during the OL study. The change from baseline to week 24 and week 52 was 0.7 and 0.5, respectively. The level of behavioral symptoms in the study population was low.</p> <p>Overall, 177 patients (93.7%) experienced at least one adverse event. Severe adverse events were reported by 15 patients (7.9%) and serious adverse events were reported by 33 patients (17.5%). The most common adverse events were nasopharyngitis, diarrhea, nausea and vomiting.</p> <p>Secondary: Not reported</p>
<p>Courtney et al.<sup>23</sup> (2004)</p> <p>Donepezil 5 to 10 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, RCT</p> <p>Patients with Alzheimer's disease</p>	<p>N=565</p> <p>156 weeks</p>	<p>Primary: MMSE, BADLS, time to entering institution</p> <p>Secondary: Not reported</p>	<p>Primary: Cognition averaged 0.8 MMSE points better (95% CI, 0.5 to 1.2; P&lt;0.0001) and functionality 1.0 BADLS points better (95% CI, 0.5 to 1.6; P&lt;0.0001) with donepezil over the first two years.</p> <p>No significant benefits were seen with donepezil compared to placebo in institutionalization (42 vs 44% at three years; P=0.4) or progression of disability (58 vs 59% at three years; P=0.4).</p> <p>The RR of entering institutional care in the donepezil group compared to placebo was 0.97 (95% CI, 0.72 to 1.30; P=0.8); the RR of progression of disability or entering institutional care was 0.96 (95% CI, 0.74 to 1.24; P=0.7).</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Similarly, no significant differences were seen between donepezil and placebo in behavioral and psychological symptoms, caregiver psychopathology, adverse events or deaths, or between 5 and 10 mg donepezil.</p> <p>Secondary: Not reported</p>
<p>Sabbagh et al.<sup>24</sup> (2013)</p> <p>Donepezil 23 or 10 mg/day</p>	<p>Post hoc of a 24-week, DB, RCT</p> <p>Patients with moderate to severe Alzheimer's disease (baseline MMSE 0 to 20)</p>	<p>N=</p> <p>Duration not specified</p>	<p>Primary: Cognitive changes in subgroups of patients based on selected baseline and demographic characteristics</p> <p>Secondary: Not reported</p>	<p>Primary: Donepezil 23 mg/day provided statistically significant incremental cognitive benefits over donepezil 10 mg/day irrespective of baseline functional severity, measured by scores on the ADCS-ADL -severe version (P&lt;0.05).</p> <p>When patients were categorized by baseline cognitive severity (MMSE score), significant benefits of donepezil 23 mg/day over 10 mg/day were seen in both subgroups when based on MMSE scores of 0 to 9 vs 10 to 20 (P&lt;0.02 and P&lt;0.01, respectively), and in the more severe subgroup when based on MMSE scores of 0 to 16 vs 17 to 20 (P&lt;0.0001 and P&gt;0.05).</p> <p>Statistically significant incremental cognitive benefits of donepezil 23 mg/day over 10 mg/day were also observed regardless of age, gender, weight, or pre-study donepezil 10mg/day treatment duration (P&lt;0.05).</p> <p>In the multivariate analysis, the only significant interaction was between treatment and baseline MMSE score.</p> <p>Secondary: Not reported</p>
<p>Tariot et al.<sup>25</sup> (2012)</p> <p>Donepezil 23 mg/day</p>	<p>OL</p> <p>Patients with Alzheimer's disease</p>	<p>N=915</p> <p>12 months</p>	<p>Primary: Safety analyses comprised examination of the incidence, severity, and timing of treatment-emergent adverse events;</p>	<p>Primary: In total, 674 patients (74.7%) reported at least one adverse event; in 320 of these patients (47.5%) at least one adverse event was considered to be possibly or probably study drug related.</p> <p>The majority of patients reporting adverse events (81.9%) had adverse events of mild or moderate severity. There were 268 patients (29.7%) who discontinued early, of which 123 (13.6%) were due to adverse events.</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>changes in weight, electrocardiogram, vital signs, and laboratory parameters; and discontinuation due to adverse events all at months three, six, nine, and 12</p> <p>Secondary: Not reported</p>	<p>Patients who had increased donepezil dose from 10 mg/day to 23 mg/day had slightly higher rates of adverse events than patients who were already receiving 23 mg (78.0 and 16.9 vs 72.8 and 14.0%, respectively).</p> <p>The incidence of new adverse events declined rapidly after the first two weeks and remained low throughout the duration of the study.</p> <p>Secondary: Not reported</p>
<p>Winblad et al.<sup>26</sup> (2006)</p> <p><u>RCT</u> Donepezil 10 mg/day</p> <p>vs</p> <p>placebo</p> <p><u>OL</u> Donepezil 5 mg daily for 28 days, then 10 mg/day per clinician's judgment</p>	<p>DB, OL, PC</p> <p>Patients 40 to 90 years of age with a probable or possible diagnosis of Alzheimer's disease</p>	<p>N=286</p> <p>52-week RCT with a 2-year OL extension phase</p>	<p>Primary: GBS</p> <p>Secondary: MMSE, GDS, PDS, NPI</p>	<p>Primary: The GBS total scores indicate that both the continuous-treatment group and delayed-start groups had declined, with the difference between the two groups favoring the continuous-donepezil group, over the three-year period (P=0.056).</p> <p>Secondary: The MMSE declined significantly less in the continuous-treatment group than in the delayed-start group over the course of the study (P=0.004, P=0.057, respectively).</p> <p>GDS declined significantly less over the three-year study period in patients in the continuous-treatment group than in those in the delayed-start group (P=0.0231).</p> <p>There was a trend favoring continuous-donepezil treatment over delayed-start treatment on the PDS, although it was not statistically significant (P=0.091).</p> <p>NPI results showed no significant treatment differences between the groups.</p>
<p>Rogers et al.<sup>27</sup> (1998)</p> <p>Donepezil 5</p>	<p>DB, MC, PC, RCT</p> <p>Patients with mild-to-moderate</p>	<p>N=473</p> <p>24 weeks</p>	<p>Primary: ADAS-Cog, CIBIC</p> <p>Secondary:</p>	<p>Primary: Out of 473 patients, 80% of placebo patients, 85% of 5 mg patients and 68% of 10 mg patients completed the study. Those that discontinued due to adverse effects were 7, 6, and 16% in the placebo, 5 and 10 mg groups,</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day  vs  donepezil 10 mg/day  vs  placebo	Alzheimer's disease		Not reported	respectively.  Primary outcome measure was mean change in scores from baseline to endpoint in the ADAS-Cog. Both donepezil doses were statistically better than placebo (P<0.0001).  Global functioning as measured by the CIBIC plus were statistically better for both donepezil groups compared to placebo at endpoint (P<0.005).  Donepezil 5 and 10 mg treatment showed no statistical difference in improvements.  Secondary: Not reported
Winblad et al. <sup>28</sup> (2006)  Donepezil 10 mg/day  vs  placebo	DB, PC, PG  Patients ≥50 years of age with severe Alzheimer's disease (MMSE score of 1 to 10 and a FAST rating of stage 5 to 7c)	N=248  6 months	Primary: SIB  Secondary: MMSE, NPI, and CGI-I	Primary: At six months, patients assigned donepezil had significantly better mean change from baseline scores than those taking placebo for SIB (P<0.05).  Secondary: CGI-I scores and the mean change from screening scores on the MMSE at six-month follow-up favored donepezil treatment over placebo (all P<0.05).  There was no significant difference between treatment groups on the NPI for the modified intention-to-treat population (P=0.43).
Black et al. <sup>29</sup> (2007)  Donepezil 10 mg/day  vs  placebo	DB, MC, PC, RCT  Patients ≥50 years of age with severe Alzheimer's disease (MMSE score of 1 to 12, modified Hachinski Ischemic score ≤6, and FAST score ≥6)	N=343  24 weeks	Primary: SIB and CIBIC-Plus  Secondary: ADCS-ADL-sev, NPI, MMSE, CBQ, RUSP	Primary: Donepezil was more efficacious when compared to placebo on SIB score change from baseline to endpoint, as well as on CIBIC-Plus score (P<0.05 for all results).  Secondary: On the ADCS-ADL-sev, both the donepezil group and the placebo group declined from baseline, and the treatment difference was NS (P=0.3574).  On the NPI, donepezil was not significantly different from placebo (P=0.4612).



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The donepezil group showed significant improvement from screening to endpoint on the MMSE compared to placebo (P=0.0267).</p> <p>The CBQ stress measure showed no significant change from baseline for either group.</p> <p>The RUSP scores also had low average responses with little movement from baseline and no significant differences.</p>
<p>Homma et al.<sup>30</sup> (2008)</p> <p>Donepezil 5 to 10 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Japanese patients ≥50 years of age with severe Alzheimer's disease (modified Hachinski Ischemic Score ≤6, FAST ≥6, MMSE score of 1 to 12 and diagnosis confirmed by neuroimaging)</p>	<p>N=302</p> <p>24 weeks</p>	<p>Primary: SIB and CIBIC-Plus</p> <p>Secondary: ADCS-ADL-sev and BEHAVE-AD</p>	<p>Primary: Donepezil 5 and 10 mg/day were more effective than placebo on the SIB. At week 24, patients in the donepezil 5 mg/day group had a significant change from baseline of 2.5 points and those in the donepezil 10 mg/day group had a significant change from baseline of 4.7 points. Patients in the placebo group showed significant worsening (-4.2 points) during the course of the study (P&lt;0.001 vs placebo).</p> <p>For the CIBIC-Plus, the analysis was performed on the seven categories of change as well as the three collapsed categories of improved, no change and worsened. In the seven-category analysis, the distribution of CIBIC-Plus scores in the donepezil 10 mg/day group was better than placebo (P=0.003); however, there was no difference with 5 mg/day (P=0.151). In the collapsed-category analysis, the distribution of CIBIC-Plus scores in the donepezil 10 mg/day group was better than placebo (P=0.001); however, there was no difference with 5 mg/day (P=0.129).</p> <p>Secondary: For the ADCS-ADL-sev, there was no significant differences between donepezil and placebo (placebo group, -1.1 points; donepezil 5 mg/day group, -0.1 points; donepezil 10 mg/day group, -0.3 points).</p> <p>For the BEHAVE-AD, there was no significant differences between donepezil and placebo (placebo group, -0.5; donepezil 5 mg/day group, -0.5; donepezil 10 mg/day group, -0.1).</p> <p>Treatment-emergent adverse events were reported by 73.3% of placebo patients, 78.2% of donepezil 5 mg/day patients and 83.3% of donepezil 10 mg/day patients. There was no significant difference in adverse events</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>between the donepezil groups and the placebo group. The most common adverse events reported are consistent with the known cholinergic side effects of donepezil. Serious adverse events were reported by 15 placebo patients (14.3%), 12 donepezil 5 mg/day patients (11.9%) and 10 donepezil 10 mg/day patients (10.4%).</p> <p>Five patients died during the treatment period. The causes of death were acute pneumonia (placebo group), acute myocardial infarction (donepezil 5 mg/day group), suspected stomach cancer (donepezil 5 mg/day group; the patient died 80 days after discontinuation), vomit-induced tracheal occlusion (donepezil 10 mg/day group; the patient died seven days after completion) and arrhythmia (donepezil 10 mg/day group).</p>
<p>Birks et al.<sup>31</sup> (2006)</p> <p>Donepezil 5 to 10 mg/day</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients with Alzheimer's disease</p>	<p>N=5,796 (24 trials)</p> <p>12 to 60 weeks</p>	<p>Primary:            ADAS-Cog, MMSE, CIBIC-Plus, ADL, withdrawals and adverse events</p> <p>Secondary:            Not reported</p>	<p>Primary:</p> <p>A significant difference was seen on the ADAS-Cog scale for patients treated with donepezil 5 mg at 24 weeks (WMD, -2.02 points; 95% CI, -2.77 to -1.26; P&lt;0.00001) and 10 mg at 24 weeks (WMD, -2.81 points; 95% CI, -3.55 to -2.06; P&lt;0.00001).</p> <p>A significant difference was seen on the MMSE for patients treated with donepezil 10 mg/day as compared to placebo at 52 weeks (WMD, 1.84 points; 95% CI, 0.53 to 3.15; P=0.006).</p> <p>Global Clinical State, CIBIC-Plus scores showed significant benefit in patients treated with donepezil 5 and 10 mg/day (OR, 2.38; 95% CI, 1.78 to 3.19; P&lt;0.00001 and OR, 1.82; 95% CI, 1.42 to 2.35; P&lt;0.00001).</p> <p>Improvements were seen in ADL scores for patients in the donepezil group over those in the placebo group (P&lt;0.01 for all scales used).</p> <p>Significantly more patients treated with donepezil 10 mg/day withdrew from treatment (24 vs 20%; P=0.003); however, there was no difference in withdrawal rates between the 5 mg/day and placebo group (P=0.56). Adverse events that occurred significantly more frequently in both the 5 and 10 mg/day treatment groups as compared to placebo are: anorexia, diarrhea, and muscle cramps.</p> <p>Secondary:</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Wallin et al.<sup>32</sup> (2007)</p> <p>Donepezil 5 to 10 mg/day</p> <p>vs</p> <p>historical data</p>	<p>MC, PRO</p> <p>Patients ≥40 years of age with probable Alzheimer's disease</p>	<p>N=435</p> <p>3 years</p>	<p>Primary: MMSE, ADAS-Cog, CIBIC, IADL</p> <p>Secondary: Not reported</p>	<p>Not reported</p> <p>Primary: For the MMSE, patients had a mean score of 22.0 at baseline and 19.1 at 36 months. After 36 months of donepezil treatment, the mean decline was 3.8 points (95% CI, 3.0 to 4.7).</p> <p>For ADAS-Cog, patients had a mean score of 20.7 at baseline and 26.1 at 36 months. After 36 months, the mean increase was 8.2 points (95% CI, 6.4 to 10.0). A modeling equation predicts an increase in ADAS-Cog to be 4 to 9 points in 12 months without treatment. Scores for the treatment group were significantly better than predicted scores for non-treatment (95% CI, 14.5 to 16.6).</p> <p>For CIBIC, at two months, 34% of patients were considered improved, 59% unchanged and 7% were worse. At six months, 28% of patients were considered improved, 46% unchanged and 26% were worse. At 12 months, 20% of patients were considered improved, 29% unchanged and 51% were worse. At 36 months, 30% of patients were considered improved or unchanged.</p> <p>The IADL change from baseline at six months was 1.01, at 12 months 2.19, and at 36 months 6.18.</p> <p>Secondary: Not reported</p>
<p>Farlow et al.<sup>33</sup> (2010)</p> <p>Donepezil 10 mg/day</p> <p>vs</p> <p>donepezil 23 mg/day</p>	<p>DB, MC, RCT</p> <p>Patients 45 to 90 years of age with moderate-to-severe Alzheimer's disease who took donepezil 10 mg/day ≥12 weeks</p>	<p>N=1,467</p> <p>24 weeks</p>	<p>Primary: Efficacy as measured by SIB-cognition and CIBIC-global function rating; tolerability</p> <p>Secondary: Not reported</p>	<p>Primary: After 24 weeks, the change in SIB-cognition score was significantly greater with donepezil 23 mg/day compared to donepezil 10 mg/day (2.6 vs 0.4, respectively; P&lt;0.001).</p> <p>There was no significant different in CIBIC score with donepezil 23 mg/day compared to donepezil 10 mg/day (4.23 vs 4.29, respectively).</p> <p>In a post-hoc analysis, the least square mean changes in SIB score and CIBIC treatment effect at end point were greater with donepezil 23 mg/day compared to donepezil 10 mg/day in patients with more advanced Alzheimer's disease compared to less impaired patients (SIB, 1.6 vs -1.5,</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>respectively; P&lt;0.001; CIBIC, 4.31 vs 4.42; P=0.028).</p> <p>Treatment emergent adverse events were reported in 73.7% of patients who received donepezil 23 mg/day and in 63.7% of patients who received donepezil 10 mg/day.</p> <p>Adverse events were reported as follows with donepezil 23 mg/day: mild (30.8%), moderate (34.5%), and severe (8.4%). The most common treatment emergent adverse events were nausea (6.1%), vomiting (5%) and diarrhea (3.2%). Severe treatment emergent adverse events that were reported included nausea (0.9%), dizziness (0.7%) and vomiting (0.6%).</p> <p>Adverse events were reported as follows with donepezil 10 mg/day: mild (31.2%), moderate (25.3%), and severe (7.2%). The most common treatment emergent adverse events were nausea (1.9%), vomiting (0.8%) and diarrhea (1.5%). Severe treatment emergent adverse events that were reported included nausea (0.2%) and dizziness (0.2%).</p> <p>Secondary: Not reported</p>
<p>Ferris et al.<sup>34</sup> (2011)</p> <p>Donepezil 10 mg/day</p> <p>vs</p> <p>donepezil 23 mg/day</p>	<p>DB, MC, RCT (post-hoc analysis)</p> <p>Patients 45 to 90 years of age with moderate-to-severe Alzheimer's disease who took donepezil 10 mg/day ≥12 weeks</p>	<p>N=1,467</p> <p>24 weeks</p>	<p>Primary: SIB-Language scale and 21-item SIB-derived language scale</p> <p>Secondary: Correlation of SIB-Language scale and SIB-derived language scale with ADCS-ADL-sev, CIBIC-plus/CIBIC-plus, and MMSE</p>	<p>Primary: At week 24, there was an improvement in language noted with donepezil 23 mg/day compared to a decline in language function with donepezil 10 mg/day (SIB-Language scale treatment difference, 0.8; P=0.0013, SIB-derived language scale treatment difference, 0.8; P=0.0009).</p> <p>Secondary: At week 24, SIB-Language scale and SIB-derived language scale scores were moderately correlated with scores on the ADCS-ADL-sev and CIBIC-plus. Results were similar in both moderate (MMSE, 17 to 20) and severe (MMSE, 0 to 16) Alzheimer's disease patients.</p>
<p>Farlow et al.<sup>35</sup> (2011)</p>	<p>DB, MC, RCT (post-hoc analysis)</p>	<p>N=1,434</p>	<p>Primary: Safety and</p>	<p>Primary: Of the 963 patients receiving donepezil 23 mg/day and 471 patients</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Donepezil 10 mg/day</p> <p>vs</p> <p>donepezil 23 mg/day</p>	<p>Patients 45 to 90 years of age with moderate-to-severe Alzheimer's disease who took donepezil 10 mg/day <math>\geq</math>12 weeks</p>	<p>24 weeks</p>	<p>tolerability</p> <p>Secondary: Not reported</p>	<p>receiving donepezil 10 mg/day, a total of 71.1 and 84.7% completed the study, respectively.</p> <p>The most common adverse events causing early discontinuation were higher in the donepezil 23 mg/day group compared to the donepezil 10 mg/day group (18.6 vs 7.9%, respectively). Adverse events that contributed the most to the discontinuations were vomiting (2.9 vs 0.4%, respectively), nausea (1.9 vs 0.4%, respectively), diarrhea (1.7 vs 0.4%, respectively), and dizziness (1.1 and 0%, respectively).</p> <p>The most common adverse events with donepezil 23 mg/day compared to donepezil 10 mg/day were nausea (11.8 vs 3.4%, respectively), vomiting (9.2 vs 2.5%, respectively) and diarrhea (8.3 vs 5.3%, respectively).</p> <p>Serious adverse events occurred in 8.3% of patients receiving donepezil 23 mg/day and in 9.6% of patients receiving donepezil 10 mg/day. These included urinary tract infection (0.6 vs 0.4%, respectively), fall (0.6 vs 0.4%, respectively), pneumonia (0.3 vs 0.6%, respectively), syncope (0.2 vs 1.1%, respectively), aggression (0.2 vs 0.8%, respectively), and confusional state (0.1 vs 0.6%, respectively).</p> <p>Secondary: Not reported</p>
<p>Doody et al.<sup>36</sup> (2012)</p> <p>Donepezil 23 mg/day</p> <p>vs</p> <p>donepezil 10 mg/day</p> <p>Patients were allowed to also take memantine.</p>	<p>DB, MC</p> <p>Patients with moderate-to-severe Alzheimer's disease</p>	<p>N=not specified</p> <p>24 weeks</p>	<p>Primary: Efficacy and safety</p> <p>Secondary: Not reported</p>	<p>Primary: At week 24, donepezil 23 mg/day provided significant cognitive benefits over 10 mg/day (<math>P&lt;0.01</math>) on the SIB, with or without concomitant memantine.</p> <p>The higher dose showed no benefit on the global function, MMSE or ADL measures in either memantine subgroup.</p> <p>Rates of treatment-emergent adverse events were higher for donepezil 23 mg/day with memantine (80.7%) than 23 mg/day without memantine (69.7%) or 10 mg/day with/without memantine (66.7/62.0%); across all treatment groups, most events were mild/moderate in severity. Individual rates of serious adverse events were low (<math>&lt;1.0\%</math>), regardless of concomitant memantine use.</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Raskind et al. <sup>37</sup> (2004)  Galantamine 24 mg/day	OL  Patients with mild-to-moderate Alzheimer's disease	N=194  36 months	Primary: ADAS-Cog, adverse events  Secondary: Not reported	Primary: Patients treated continuously with galantamine for 36 months increased a mean of 10.2±0.9 points on the ADAS-Cog. This was a substantially smaller cognitive decline (approximately 50%) than that predicted for the placebo group.  Patients discontinuing galantamine therapy before 36 months had declined at a similar rate before discontinuation as those completing 36 months of treatment.  Almost 80% of patients who received galantamine for 36 months seemed to demonstrate cognitive benefits compared to those predicted for untreated patients.  Secondary: Not reported
Rockwood et al. <sup>38</sup> (2008)  Galantamine 24 mg/day	MC, OL  Patients with Alzheimer's disease who had received galantamine treatment for up to 36 months	N=240  Up to 48 months	Primary: ADAS-Cog, DAD, adverse events  Secondary: Not reported	Primary: Mean ADAS-Cog worsened from 22.6±8.6 at baseline to 31.3±13.1 at 48 months.  DAD worsened from 73.4±18.1 at baseline to 36.1±29.0 at 48 months.  Fifty one patients withdrew from the study.  Secondary: Not reported
Wallin et al. <sup>39</sup> (2011)  Galantamine 24 mg/day	MC, OL, PRO  Patients with Alzheimer's disease and no previous cholinesterase inhibitor therapy	N=280  36 months	Primary: MMSE, ADAS-cog, IADL, CIBIC  Secondary: Subgroup analysis by K-means cluster analysis	Primary: From baseline to 36 months, MMSE decreased from 23.3 to 21.74. The MMSE score was significantly better at two months (P<0.001) and at six months (P=0.006) compared to baseline, and was stable at 12 months (P=0.616) compared to baseline. The total mean decline in MMSE score from baseline after three years of treatment was 2.6  From baseline to 36 months, ADAS-cog increased from 16.85 to 19.39.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The total change in ADAS-cog score after three years of treatment was 5.6 points above baseline values.</p> <p>The ADAS-cog scores at six months were not different from baseline (P=0.248), but deteriorated after that.</p> <p>Mean IADL scores demonstrated deteriorated at all time points compared to baseline (12.76 to 17.13).</p> <p>According to CIBIC scores at two months, 93% of patients remaining in the study were “improved or unchanged”, at months six, 12, 24, and 36; 81, 69, 50, and 41% of the patients were “improved or unchanged”, respectively.</p> <p>Secondary: Cluster analysis identified two response clusters. Cluster 1 included patients with low ability in ADAS-cog and IADL scores at baseline. These patients were older and less educated, but responded better at six months compared to cluster two patients. Cluster 2 patients included better ADAS-cog and IADL scores at baseline. Cluster 2 patients had a higher frequency of the APOE ε4 allele.</p>
<p>Brody et al.<sup>40</sup> (2006)</p> <p>Galantamine 2 to 50 mg/day</p>	<p>OL, OS, PRO</p> <p>Patients diagnosed with mild-to-moderately severe dementia</p>	<p>N=345 ITT N= 229 PP</p> <p>6 month follow-up</p>	<p>Primary: MMSE, ADAS-Cog, CIBIC-Plus, IADL</p> <p>Secondary: Not reported</p>	<p>Primary: For the MMSE 65% of PP patients had an increased score at the three-month assessment as compared to baseline with an overall 92% response rate. 70% of PP patients had an increased score at the six-month assessment as compared to baseline with an overall 91% response rate. 44% of ITT patients had an increased score at the six-month assessment as compared to baseline (P values were not reported).</p> <p>For ADAS-Cog at 6 months, 86% of the PP patients and 33% of the ITT patients had a decrease in ADAS-Cog score. P value was not reported.</p> <p>For CIBIC-Plus at three months, 91% of PP patients were considered responders by their physicians; 28% were unchanged, 38% were minimally improved, 22% were much improved, 4% were very much improved (P values not reported). For CIBIC-Plus at six months, 86% of PP patients were considered responders by their physicians; 20% were</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>unchanged, 26% were minimally improved, 32% were much improved, 7% were very much improved. In the ITT patients, 54 % were classified as responders at six months (P values not reported).</p> <p>Most PP patients had no change in IADL scores at three and six months (P value not reported).</p> <p>Most PP patients had no change in behavior scores at three and six months (P value not reported).</p> <p>Secondary: Not reported</p>
<p>Richarz et al.<sup>41</sup> (2014)</p> <p>Galantamine 8 to 24 mg/day</p>	<p>OL, PRO</p> <p>Patients ≥45 years of age with mild to moderate Alzheimer's disease</p>	<p>N=75 (36 months)</p> <p>N=159 (24 months)</p> <p>N=269 (6 months)</p> <p>Up to 36 months</p>	<p>Primary: ADAS-cog/11</p> <p>Secondary: Bayer-ADL, NPI, CGI-C, adverse events</p>	<p>Primary: Mean ADAS-cog score improved significantly during the first six months, with improvement maintained until month 12. During follow-up, mean ADAS-cog score returned to baseline levels between months 18 and 24; after 36 months, it had deteriorated (increased) by 2.87 ± 11.07 points.</p> <p>Secondary: Mean NPI score improved significantly in the first 12 months and worsened thereafter. In the 36-month sample, patient self-rated Bayer ADL scores remained stable until 24 months of treatment; then, a significant deterioration had occurred; a significant deterioration from baseline in caregivers' Bayer ADL scores occurred after month 12. After six months of treatment, 84% of the patients who completed the six-month observation period were considered to be improved or unchanged compared with baseline on the CGI-C. In the 36-month sample, the corresponding value was 54%.</p> <p>In the 36-month sample, 54 patients (72%) reported at least one treatment-emergent adverse event throughout the treatment period, with most events occurring during the first two years of treatment.</p>
<p>Cummings et al.<sup>42</sup> (2004)</p> <p>Galantamine 8 to 24 mg/day</p>	<p>DB, PC, RCT</p> <p>Patients with mild-moderate Alzheimer's disease</p>	<p>N=978</p> <p>21 weeks</p>	<p>Primary: NPI, caregiver distress related to patients' behavior</p>	<p>Primary: NPI scores worsened with placebo, whereas patients treated with 16 or 24 mg/day of galantamine had no change in NPI scores.</p> <p>Behavioral improvement in patients symptomatic at baseline ranged from</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  placebo			Secondary: Not reported	29 to 48%. Changes were evident in patients receiving 16 and 24 mg/day of galantamine.  High-dose galantamine was associated with a significant reduction in caregiver distress.  Secondary: Not reported
Scarpini et al. <sup>43</sup> (2011)  <u>Phase 1</u> Galantamine 8 to 16 mg/day  <u>Phase 2</u> Galantamine 16 mg/day  vs  placebo	<u>Phase 1</u> MC, OL  <u>Phase 2</u> DB, MC, RCT  Mild to moderate Alzheimer's disease in patients ≥50 years of age (MMSE, 11 to 24)	N=393  36 months	Primary: ADAS-cog/11 deterioration ≥4 points  Secondary: CIBIC-plus, adverse events	<u>Phase 1</u> Primary: Cognitive functions improved significantly on the ADAS-cog/11 scale with galantamine treatment at month seven relative to baseline (from 24.1 to 22.9, difference, -1.2; 95% CI, -2.3 to -0.1; P<0.01). Scores were similar to baseline values at the end of the OL phase at month 12 (mean score at baseline, 24.1; mean score at month 12, 24.7; 95% CI, -0.5 to 1.7, P=0.16).  Secondary: CIBIC-plus score improved in 34.3%, was unchanged in 30.9%, and worsened in 34.9% of patients when compared to baseline.  A total of 50.4% of patients reported adverse events, of which the most common was gastrointestinal disorders (21.3%), nervous system disorders (9.8%), and psychiatric disorders (19.7%). Serious adverse events were reported in 12.2%.  <u>Phase 2</u> Primary: Patients receiving placebo were more likely to discontinue therapy prematurely compared to galantamine for any reason (HR, 1.76; 95% CI, 1.10 to 2.81; P=0.02) or lack of efficacy (HR, 1.80; 95% CI, 1.02 to 3.18; P=0.04). No significant difference was observed by ADAS-cog >4 between the groups (HR, 1.66; 95% CI, 0.78 to 3.54; P=0.19).  Secondary: There were no significant differences between the treatment groups concerning mean values of the CIBIC-plus scale.

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>A total of 34.1% of patients receiving galantamine and 27% of patients receiving placebo experienced adverse events. The most common adverse events were nervous system disorders (6.6%) and psychiatric disorders (5.3%). Serious adverse events were reported in 14.5% of galantamine-treated patients compared to 6.3% of patients in the placebo group.</p>
<p>Kavanagh et al.<sup>44</sup> (2011)</p> <p>Galantamine 16 to 24 mg/day</p> <p>vs</p> <p>placebo</p>	<p>OL, RCT</p> <p>Patients with mild-to-moderate Alzheimer's disease</p>	<p>N=3,523 (5 trials)</p> <p>5 to 6 months</p>	<p>Primary: Changes from baseline in ADAS-Cog 11 at trial endpoint (two to five months after reaching maintenance doses)</p> <p>Secondary: Not reported</p>	<p>Primary: The proportion of patients who met criteria for "improved", "stable", or "non-rapid decline" at trial endpoint were 45.8, 59.5, and 87.6%, respectively with galantamine compared to 27.2, 37.1, and 67.7%, respectively with placebo.</p> <p>Changes in ADAS-Cog 11 scores with galantamine were -4.9, -4.7, and -2.9 points, respectively, for "improved", "stable" and "non-rapid decline" compared to -3.6, -3.4, and -1.2, respectively with placebo.</p> <p>Patients receiving galantamine who were reported to be "improved" or "stable" experienced improvement in ADAS-Cog 11 scores until 18 months after starting treatment, and attenuated deterioration thereafter. For galantamine-treated patients exhibiting "non-rapid decline", mean ADAS-Cog 11 score returned to baseline after approximately 12 months.</p> <p>Secondary: Not reported</p>
<p>Burns et al.<sup>45</sup> (2009)</p> <p>Galantamine 24 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients 40 to 95 years of age with severe dementia of the Alzheimer type or probable Alzheimer's disease (MMSE, 5 to 12 points)</p>	<p>N=407</p> <p>6 months</p>	<p>Primary: SIB, MDS-ADL, and adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: In the completer analysis, the mean total SIB score of the galantamine group increased to 69.1 points at week 26. The mean SIB score in the placebo group decreased to 66.9. The between group least squares mean difference was 4.36 (95% CI, 1.3 to 7.5; P=0.006).</p> <p>In the completer analysis, the mean total MDS-ADL self-performance score worsened in both groups: scores at week 26 were 13.0 points in the galantamine group and 13.6 points in the placebo group. The between-group least squares mean difference was -0.41 points (95% CI, -1.3 to 0.5; P=0.383).</p> <p>In the LOCF analysis, the mean SIB score in the galantamine group</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>increased to 69.3 points. In the placebo group, the mean SIB score decreased by 3.2 points. The between-group least squares mean difference was 5.02 points (95% CI, 2.17 to 7.86; P=0.0006).</p> <p>In the LOCF analysis, the mean total seven-item MDS-ADL self-performance score in the galantamine group worsened at endpoint to 13.1 points and to 14.0 points in the placebo group. Changes from baseline in the seven-item MDS-ADL self-performance score were 1.3 points and 1.7 points, respectively. The between-group least squares mean difference was -0.50 (95% CI, -1.39 to 0.39; P=0.394).</p> <p>Significant between-group differences were seen in the galantamine group for memory (P=0.006), praxis (P=0.010), and visuospatial ability (P=0.002). There were no significant differences in language (P=0.064) or attention (P=0.075).</p> <p>Scores for all eleven-item MDS-ADL self-performance subscales worsened in both treatment arms. The deterioration in the subscale score for locomotion on unit was significantly less in the galantamine group (P=0.021).</p> <p>During the study, 88% of patients who received galantamine and 89% who received placebo had at least one adverse event. The most common adverse events in both treatment groups were urinary tract infections, vomiting, diarrhea, nausea, and falls.</p> <p>Secondary: Not reported</p>
<p>Raskind et al.<sup>46</sup> (2004)</p> <p>Galantamine 24 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, RCT</p> <p>Patients with mild-moderate Alzheimer's disease</p>	<p>N=194</p> <p>36 months</p>	<p>Primary: ADAS-Cog, adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: Patients treated continuously with galantamine for 36 months increased a mean of 10.2±0.9 points on the ADAS-Cog. This was a substantially smaller cognitive decline (approximately 50%) than that predicted for the placebo group.</p> <p>Patients discontinuing galantamine therapy before 36 months had declined at a similar rate before discontinuation as those completing 36 months of treatment.</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Almost 80% of patients who received galantamine for 36 months seemed to demonstrate cognitive benefits compared to those predicted for untreated patients.</p> <p>Secondary: Not reported</p>
<p>Wilcock et al.<sup>47</sup> (2000)</p> <p>Galantamine 24 mg/day</p> <p>vs</p> <p>galantamine 32 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB</p> <p>Patients with mild-moderate Alzheimer's disease</p>	<p>N=653</p> <p>6 months</p>	<p>Primary: ADAS-Cog, adverse events</p> <p>Secondary: Not reported</p>	<p>Primary: Both doses of galantamine were statistically better than placebo in the mean change in ADAS-Cog from baseline to endpoint (P&lt;0.0001).</p> <p>Patients taking galantamine 24 mg had a -0.5 point mean change on the ADAS-Cog scale, while the 32 mg group had a -0.8 change. This compares to a +2.4 change for the placebo group. Statistical comparisons between the 24 mg group and the 32 mg group were not conducted.</p> <p>Discontinuations due to adverse events were 9, 14 and 22% in the placebo, 24 and 32 mg dose groups, respectively.</p> <p>Secondary: Not reported</p>
<p>Dunbar et al.<sup>48</sup> (2006)</p> <p>Galantamine IR 8 to 16 or 24 mg/day</p> <p>vs</p> <p>galantamine ER 8 to 16 or 24 mg/day</p> <p>vs</p> <p>placebo</p>	<p>Post hoc analysis, DB, MC, PC, RCT</p> <p>Patients with mild-to-moderate probable Alzheimer's disease</p>	<p>N=965</p> <p>7 months</p>	<p>Primary: Nausea and vomiting</p> <p>Secondary: Not reported</p>	<p>Primary: Nausea reports were as follows: 16.9% of the galantamine ER group, 13.8% of galantamine IR group and 5.0% of placebo group.</p> <p>Vomiting reports were as follows: 6.6% of the galantamine ER groups, 8.6% of the galantamine IR group and 2.2% of the placebo group.</p> <p>During dose titration, the area under the curve of daily percentage of patients reporting nausea or vomiting was significantly higher in the galantamine IR group compared to placebo (320.9 vs 102.9; P=0.01) but for galantamine ER vs placebo and galantamine ER vs galantamine IR no significant differences were seen ([173.5 vs 102.9; P=NS], [320.9 vs 173.5; P=NS]).</p> <p>The mean daily nausea rate and the mean daily vomiting rate for galantamine ER and galantamine IR were not significantly different but</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>when both were compared to placebo, significance was seen (P&lt;0.05).</p> <p>The galantamine IR had a greater mean percentage of days with nausea compared to galantamine ER (38 vs 18.4%; P=0.014) while there was no significance for both galantamine groups compared to placebo.</p> <p>Secondary: Not reported</p>
<p>Brody et al.<sup>49</sup> (2005)</p> <p>Galantamine IR 8 to 16 or 24 mg/day</p> <p>vs</p> <p>galantamine ER 8 to 16 or 24 mg/day</p> <p>vs</p> <p>placebo</p>	<p>AC, DB, MC, PC, PG, RCT</p> <p>Patients with mild-to-moderate probable Alzheimer's disease</p>	<p>N=971</p> <p>6 months</p>	<p>Primary: ADAS-cog/11, CIBIC-Plus</p> <p>Secondary: ADCS-ADL, NPI, ADAS-cog/13, nonmemory ADAS-cog/ memory, ADAS-Cog</p>	<p>Primary: Compared to placebo, galantamine was significantly more effective with improvement from baseline in ADAS-cog/11 scores (mean change, 1.3 and -1.4, respectively; P&lt;0.001; 95% CI, -3.74 to -1.68; LOCF mean change, 1.2 and -1.3, respectively; P&lt;0.001; 95% CI, -3.34 to -1.49).</p> <p>Galantamine also showed similar results when compared to placebo (OC mean change, -1.8 and 1.3, respectively; P&lt;0.001; 95% CI, -4.17 to -2.08; LOCF mean change, -1.6 and 1.2, respectively; P&lt;0.01; 95% CI, -3.70 to -1.86).</p> <p>Secondary: ADCS-ADL scores were significantly improved in the galantamine group vs placebo (P=0.003; 95% CI, 0.85 to 4.03; LOCF; P&lt;0.001; 95% CI, 1.09 to 3.91).</p> <p>In galantamine groups vs placebo, NPI scores were not statistically significant but instead numerically significant (P=0.451; 95% CI, -2.77 to 1.23; LOCF; P=0.941; 95% CI, -1.85 to 1.82), (OC; P&lt;0.205; 95% CI, -3.31 to 0.71; LOCF; P&lt;0.102; 95% CI, -3.42 to 0.23).</p> <p>Statistical significance was found in cognition improvement from baseline for both galantamine groups compared to placebo based on ADAS-cog/13, non-memory ADAS-Cog, and memory ADAS-Cog scores.</p>
<p>Loy et al.<sup>50</sup> (2006)</p> <p>Galantamine 8 to 36 mg/day</p>	<p>MA (10 trials)</p> <p>Patients diagnosed with mild cognitive impairment or</p>	<p>N=6,805</p> <p>12 weeks-2 years</p>	<p>Primary: CIBIC-plus, ADAS-Cog, ADCS-ADL, DAD, NPI</p>	<p>Primary: Statistically significant difference was seen on the global rating scales for patients treated with galantamine, at all durations and all doses but 8 mg/day (P values varied).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo	Alzheimer's disease		Secondary: Not reported	<p>Statistically significant difference was seen on the ADAS-Cog scale for patients treated with galantamine at all doses, with greater effect at six months than three months (P values varied).</p> <p>When reported, ADCS-ADL, DAD, and NPI scores for patients treated with galantamine were significantly improved over those in the placebo group (P values not reported).</p> <p>Secondary: Not reported</p>
Herrmann et al. <sup>51</sup> (2011)  Memantine 20 mg/day	OL  Patients with moderate-to-severe Alzheimer's disease	N=31  3 months	<p>Primary NPI-NH change in agitation and aggression subscale, CGI-C scale, caregiver impact, and effect on nursing burden measured by M- NCAS</p> <p>Secondary: Caregiver distress subscale of the NPI- NH, changes in psychotropic medications</p>	<p>Primary: There was a significant decrease in the NPI-NH agitation/aggression subscale score with memantine (P=0.014).</p> <p>According to the CGI-C scores, 48% of patients were improved (much improved or minimally improved). A total of 52% of patients did not benefit from treatment (no change, minimally worse or much worse).</p> <p>There was a significant decrease in the M-NCAS total score (P=0.005), as well as decreases on the attitude (P=0.009) and strain (P=0.013) subscales with memantine therapy.</p> <p>Secondary: The NPI-NH subscale score decreased significantly with memantine therapy (P=0.009).</p> <p>Psychotropic medications were available in 28 patients, with 64.3% receiving at least one dose during the study. Lorazepam was the most commonly used psychotropic (P=0.046). Overall, seven patients decreased psychotropic medication use during the study, while three increased usage; Most remained the same for psychotropic usage.</p>
Bakchine et al. <sup>52</sup> (2007)  Memantine 20 mg/day	DB, PC  Patients with mild- to-moderate Alzheimer's disease	N=470  24 weeks	<p>Primary: ADAS-COG and CIBIC-plus</p> <p>Secondary: Not reported</p>	<p>Primary: Patients in the memantine group showed a statistically significant improvement relative to placebo in ADAS-COG and CIBIC-plus at weeks 12 and 18. There was no significant difference between the groups at week 24.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo				Secondary: Not reported
Reisberg et al. <sup>53</sup> (2003)  Memantine 20 mg/day  vs  placebo	DB, PG  Patients with moderate-to-severe Alzheimer's disease	N=252  28 weeks	Primary: CIBIC-Plus and ADCS-ADL  Secondary: SIB	Primary: A significantly greater effect was observed in the memantine group compared to the placebo group on the ADCS-ADL (P=0.03).  There was a significant difference in favor of memantine at week 28 on the CIBIC-Plus using the observed-cases analysis (mean score, 4.7 placebo vs 4.4, memantine; P=0.03), and a numerical difference at study endpoint in favor of memantine using the last-observed-carried-forward analysis (mean score, 4.8 placebo vs 4.5 memantine; P=0.06).  Secondary: Memantine patients showed significantly less cognitive decline on the SIB total score compared to placebo-treated patients over the 28-week study period (P=0.002).
Winblad et al. <sup>54</sup> (1999)  Memantine 10 mg/day  vs  placebo	DB, PC  Patients in Latvia with severe dementia, either Alzheimer's disease or vascular dementia	N=166  12 weeks	Primary: CGI-C and BGP  Secondary: Safety	Primary: Significantly greater improvement was observed in the memantine group compared to the placebo group on the BGP and the CGI-C (P<0.016 and P<0.001, respectively).  Separate analyses of the Alzheimer's disease population alone also yielded statistically significant results in favor of patients receiving memantine, by either the last-observed-carried-forward analysis or the observed-cases analysis on both outcome measures.  At study endpoint, memantine patients showed significantly greater functional improvement compared to patients who received placebo, at study endpoint (P=0.012).  Secondary: No significant differences in safety were found between the groups.
Winblad et al. <sup>55</sup> (2007)  Memantine 20	MA  Four studies: memantine as	N=1,826 in subgroup with moderate-to-	Primary: CIBIC-Plus, SIB, ADAS-Cog, ADCS-ADL, NPI	Primary: There was a statistically significant advantage for the memantine group over the placebo group in all 4 efficacy domains: CIBIC-Plus or global status (P<0.001), SIB or ADAS-Cog status (P<0.001), ADCS-ADL

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
mg/day vs placebo	monotherapy, 2 studies of memantine vs placebo in patients already taking an acetylcholinesterase inhibitor; patients diagnosed with moderate-to-severe Alzheimer's disease	severe Alzheimer's disease  24 to 28 weeks	Secondary: Not reported	(P<0.001) and NPI (P=0.03).  Secondary: Not reported
Wilkinson et al. <sup>56</sup> (2007)  Memantine 20 mg/day  vs  placebo	MA  Patients diagnosed with moderate-to-severe Alzheimer's disease	N=1,826  24 to 28 weeks	Primary: ADAS-Cog, SIB, CIBIC-Pus, ADCS-ADL  Secondary: Not reported	Primary: Significantly more patients in the placebo group (21%) had marked clinical worsening, as demonstrated by deteriorating scores, than in the memantine group (11%; P<0.001).  Significantly more patients in the placebo group (28%) compared to the memantine group (18%) had documentation of worsening in any outcome measure (P<0.001).  Secondary: Not reported
McShane et al. <sup>57</sup> (2006)  Memantine 10 to 30 mg/day  vs  placebo	MA (12 trials)  Patients diagnosed with mild-to-moderate, moderate-to-severe and mild-to-moderate vascular dementia	N=3,731 (15 trials)  Variable duration	Primary: CIBIC-Plus, SIB, ADAS-Cog, ADCS-ADL, NPI  Secondary: Not reported	Primary: Significant improvement at six months was seen for patients with mild-to-moderate dementia treated with memantine on the ADAS-Cog scale (P=0.03); however, there was no significant difference seen for behavior and ADL scales.  Significant improvement at six months was seen for patients with moderate-to-severe dementia treated with memantine for the following scales: CIBIC-Plus (P<0.00001), SIB (P<0.00001), ADCS-ADL (P=0.003) and NPI (P=0.004).  Patients with vascular dementia treated with memantine had significant improvement in cognition scores and behavior scores but no significant change in global rating scales (ADAS-Cog; P=0.0002, NPI; P=0.03).  Secondary:



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Not reported
Grossberg et al. <sup>58</sup> (2013)  Memantine extended-release 28 mg once daily  vs  placebo	DB, MC, RCT  Outpatients with Alzheimer's disease (MMSE scores of three to 14) who were receiving stable, ongoing cholinesterase inhibitor treatment	N=677  24 weeks	Primary: Baseline-to-endpoint score change on the SIB and the endpoint score on the CIBIC-Plus  Secondary: Baseline-to-endpoint score change on the ADCS-ADL19; additional parameters included the baseline-to-endpoint score changes on the NPI and verbal fluency test	Not reported  Primary: At 24 weeks memantine-treated patients significantly outperformed placebo-treated patients on the SIB (2.6; 95% CI, 1.0 to 4.2; P=0.001) and CIBIC-Plus (P=0.008).  Secondary: At 24 weeks memantine-treated patients significantly outperformed placebo-treated patients on the NPI (P=0.005), and verbal fluency test (P=0.004); the effect did not achieve significance on ADCS-ADL19 (P=0.177).  Adverse events with a frequency of >5.0 % that were more prevalent in the memantine group were headache (5.6 vs 5.1 %) and diarrhea (5.0 vs 3.9 %).
Grossberg et al. <sup>59</sup> (2018)  Memantine extended-release 28 mg once daily  vs  placebo	Post-hoc analysis of DB, RCT  Outpatients with Alzheimer's disease (MMSE scores of three to 14) who were receiving stable, ongoing cholinesterase inhibitor (ChEI) treatment	N=677  24 weeks	Primary: Comparing patients receiving memantine ER/ cholinesterase inhibitor (ChEI) to placebo/ChEI for time to onset of response and if the response was maintained (achieving improvement at weeks eight, 12, or 18 and maintaining through	Primary: Greater percentages of memantine ER/ChEI patients achieved an early response that was maintained on SIB, NPI, and CIBIC-Plus (P<0.05) versus placebo/ChEI. Greater percentages of memantine ER/ChEI-treated patients achieved and maintained a clinically notable response on ADL/NPI, SIB/ADL/NPI, and SIB/ADL/CIBIC-Plus, compared with placebo/ChEI (P<0.05). Memantine ER results in early, maintained improvement in patients with moderate to severe Alzheimer's disease concurrently taking ChEIs, compared with cholinesterase treatment alone.  Secondary: When comparing memantine ER/ChEI-treated versus placebo/ChEI-treated responders for all possible combinations of two, three, or four efficacy measures, a greater proportion of memantine ER/ChEI patients showed no decline and clinically notable response versus ChEI alone. The difference between treatments for patients who showed no decline did not

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			endpoint/week 24)  Secondary: Comparing percentages of patients for all possible combinations of two to four assessments with either no decline or clinically notable response	reach statistical significance; the combination of efficacy outcomes with the greatest difference was SIB/CIBIC-Plus (P=0.0541).
Hager et al. <sup>60</sup> (2016)  Galantamine  vs  placebo  Memantine was taken at baseline and throughout the study by 24.5% of galantamine-treated patients and 24.0% of placebo-treated patients	Post-hoc analysis of DB, PC, PRO, RCT  Patients with mild-to-moderate Alzheimer's disease or mixed dementia stratified by the presence or absence of concomitant memantine	N=2,045  2 years	Primary: Mortality and efficacy parameters including MMSE scores, DAD scores, and nursing home placement  Secondary: Not reported	Primary: In memantine users, mortality rates were not reduced by galantamine (HR, 1.25; 95% CI, 0.63 to 2.46) as they were in nonusers (HR, 0.33; 95% CI, 0.18 to 0.61). Mortality rates in the galantamine-treated groups, compared with placebo, were lower in patient groups with $\geq$ median age and higher MMSE score (18 to 26).  In memantine users, galantamine did not reduce MMSE decline at any time point. In contrast, in memantine nonusers the galantamine group showed reduced decline in MMSE scores as compared with the placebo group at all time points, with a numerical increase in the effect size over time (P>0.05 for all comparisons).  Examination of DAD scores at month 24 demonstrated a benefit in galantamine-treated memantine nonusers, with attenuation of this benefit in the memantine user group across the range of baseline MMSE scores.  In memantine users, the risk of new nursing home admission during year one was higher in the galantamine group than in the placebo group (3.70; 95% CI, 1.04 to 13.23; P=0.03). In memantine nonusers, the risk of nursing home placement tended to be lower in galantamine-treated patients than in placebo-treated patients in year two (RR, 0.19; 95% CI, 0.02 to 1.57; P=0.08). The cumulative numerical percentages of nursing home placements were 5.0% and 18.8% in memantine users on placebo

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>and galantamine, respectively, and 5.0% and 1.8% in memantine nonusers on placebo and galantamine.</p> <p>Overall, the beneficial effects of galantamine at two years post treatment were not observed in patients who had been placed on background memantine.</p> <p>Secondary: Not reported</p>
<p>Burns et al.<sup>61</sup> (2004)</p> <p>Rivastigmine</p>	<p>RETRO</p> <p>Patients with moderately severe Alzheimer's disease/dementia</p>	<p>N=2,126</p> <p>3 trials, each 6 months</p>	<p>Primary: Effectiveness</p> <p>Secondary: Not reported</p>	<p>Primary: Mean ADAS-Cog score declined by 6.3 points in the placebo group and increased by 0.2 points in the rivastigmine group (P&lt;0.001).</p> <p>Clinical benefits were also observed with the MMSE, the six-item PDS, and items of the BEHAV-AD assessed efficacy.</p> <p>Rivastigmine showed the same pattern of adverse events as in other studies, but the RR of dropping out due to adverse events was lower than in subjects with milder Alzheimer's disease.</p> <p>Secondary: Not reported</p>
<p>Dantoine et al.<sup>62</sup> (2006)</p> <p>Rivastigmine 3 to 12 mg/day</p> <p>Addition of memantine 5 to 20 mg/day was allowed for non-responders of rivastigmine at the end of week 16.</p>	<p>MC, OL</p> <p>Patients at least 50 years of age with probable Alzheimer's disease according to criteria of DSM-IV, baseline scores of &lt;18 for MMSE or scores of &gt;4 on GDS, previously treated for at least 6 months prior with donepezil 5 to 10</p>	<p>N=202</p> <p>16 weeks of rivastigmine monotherapy (Phase 1)</p> <p>Additional 12 weeks of rivastigmine and memantine combination therapy for non-</p>	<p>Primary: MMSE</p> <p>Secondary: MMSE, Mini-Zarit inventory, NPI, Ten-point Clock-drawing Test, D-KEFS verbal fluency test, CGI-C</p>	<p>Primary: Based on MMSE scores, 46.3% of patients improved or stabilized on rivastigmine monotherapy at the end of Phase 1.</p> <p>For those patients previously on donepezil or galantamine, responder rates were also similar (46.6 and 46.4%).</p> <p>At the end of Phase 2 with combination therapy of rivastigmine and memantine, according to MMSE scores, 77.9% of patients improved or stabilized.</p> <p>Patients switching to combination therapy from galantamine responded more significantly than those who switched from donepezil (84.2 vs 72.3%; P=0.047).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
	mg/day or galantamine 16 to 24 mg/day and considered not stabilized, current stabilized medications allowed	responders of rivastigmine monotherapy (Phase 2)  Total 28 weeks		<p>Secondary: According to CGI-C data, no change or improvement was seen in 76.5% of patients who completed the study at the end of Phase 1.</p> <p>For the 82.6% who worsened from baseline at the end of Phase 1, 81.4% improved or had no change at the end of Phase 2 with the addition of memantine on the CGI-C.</p> <p>At the end of Phase 1, MMSE and NPI showed significant improvements (P&lt;0.001 and P&lt;0.05, respectively) while there was no change from baseline for Ten-point Clock-drawing Test and D-KEFS verbal fluency test scores and the Mini-Zarit interview.</p> <p>At the end of Phase 2, D-KEFS verbal fluency test, Mini-Zarit, and especially MMSE scores showed significant improvement (P&lt;0.05, P&lt;0.001 and P&lt;0.001, respectively).</p>
<p>Olin et al.<sup>63</sup> (2010)</p> <p>Rivastigmine 6 to 12 mg/day and memantine 20 mg/day</p>	<p>MC, OL, PRO</p> <p>Patients ≥50 years of age with moderate-to-severe Alzheimer's disease (MMSE ≥10 to ≤20)</p>	<p>N=116</p> <p>26 weeks</p>	<p>Primary: Safety and tolerability</p> <p>Secondary: ADCS-CGIC, ADCS-ADL measured</p>	<p>Primary: Nausea and vomiting occurred in 26.7 and 10.3% of patients, respectively. Most cases were mild with few severe cases reported (2.6 and 2.6%, respectively).</p> <p>At least one treatment-emergent adverse event was experienced by 81.9% of patients. The most common adverse events were nausea (26.7%), dizziness (11.2%), vomiting (10.3%), and diarrhea (10.3%).</p> <p>No patients exhibited clinically significant ECG abnormalities.</p> <p>Secondary: At week 26, 59% of patients experienced no decline in MMSE total score from baseline. The mean change from baseline in MMSE total score was 0.7.</p> <p>At week 26, there was no change in global ADCS-CGIC scores.</p> <p>Patient and caregiver assessed mental/cognitive state, behavior and functioning severity scores were maintained to a similar extent throughout the study.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>The mean overall rating on the ADCS-CGIC was 4.0. At week 26, 64.5% of patients were considered unchanged or improved.</p> <p>The mean ADAS-ADL scores significantly declined by -2.9.</p> <p>At week 26, cognition, behavior and global functioning were unchanged or improved in 63.2, 71.1 and 77.6% of patients respectively.</p>
<p>Gauthier et al.<sup>64</sup> (2010)</p> <p>Rivastigmine 3 to 12 mg/day</p>	<p>MC, OL, OS, PRO</p> <p>Patients with mild-moderate Alzheimer's disease</p>	<p>N=3,800</p> <p>12 months</p>	<p>Primary: Physician-assessed abbreviated CGI-C, MMSE, psychotropic medication use</p> <p>Secondary: Not reported</p>	<p>Primary:</p> <p>At six months, the proportion of patients who were reported as being improved vs no change vs deteriorating were 46.4 vs 44.9 vs 8.8% for attention; 42.8 vs 50.0 vs 7.2% for apathy; 41.1 vs 49.5 vs 9.4% for anxiety; 33.8 vs 68.4 vs 7.7% for agitation; 35.1 vs 54.8 vs 10.1% for irritability; and 30.8 vs 63.8 vs 5.4% for sleep disturbance.</p> <p>At 12 months, the proportion of patients who were reported as being improved vs no change vs deteriorating were 47.9 vs 41.0 vs 11.1% for attention; 44.1 vs 46.7 vs 9.2% for apathy; 41.8 vs 47.3 vs 10.9% for anxiety; 33.5 vs 57.6 vs 8.9% for agitation; 33.8 vs 56.4 vs 9.8% for irritability; and 29.7 vs 64.7 vs 5.6% for sleep disturbance.</p> <p>Overall, CGI-C at six and 12 months demonstrated a larger percentage of patients with improvement vs deterioration. At six months, 54% of patients overall demonstrated no change. At 12 months, 52% of patients overall demonstrated no change.</p> <p>MMSE scores were 20.8 at baseline, 21.5 after three months, 21.3 after six months, and 21.3 after 12 months.</p> <p>At baseline, 61.3% of patients were not taking a psychotropic medication. At six months, the proportion of patients not taking any psychotropic medications increased to 70.8%; at 12 months, it was 84.7%.</p>
<p>Birks et al.<sup>65</sup> (2000)</p> <p>Rivastigmine 6 to 12 mg/day</p>	<p>MA (8 trials)</p> <p>Patients diagnosed with Alzheimer's disease</p>	<p>N=3,660</p> <p>12 to 52 weeks</p>	<p>Primary: ADAS-Cog, ADL, adverse events</p> <p>Secondary:</p>	<p>Primary:</p> <p>Statistically significant differences were seen in patients treated with rivastigmine at doses of 6 to 12 mg/day as compared to placebo for the following outcomes: ADAS-Cog (WMD, -2.09; 95% CI, -2.65 to -1.54) and ADL (WMD, -2.15; 95% CI, -3.16 to -1.13).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			Not reported	<p>At 26 weeks, 55% of patient had severe dementia in the rivastigmine group as compared to 59% in the placebo group (OR, 0.78; 95% CI, 0.64 to 0.94).</p> <p>Adverse events (nausea, vomiting, diarrhea, anorexia, headache, syncope, abdominal pain and dizziness) were reported significantly more frequently in the rivastigmine group than with placebo.</p> <p>Secondary: Not reported</p>
<p>Birks et al.<sup>66</sup> (2009)</p> <p>Rivastigmine</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients diagnosed with probable Alzheimer's disease</p>	<p>N=4,775 (9 trials)</p> <p>Variable duration</p>	<p>Primary: Cognitive function, global impression, activities of daily living, behavioral disturbance, withdrawal rates, and incidence of adverse effects</p> <p>Secondary: Not reported</p>	<p>Primary: <u>Cognitive function</u></p> <p>The meta-analysis, using WMD, demonstrated benefit on cognitive function as measured by ADAS-Cog test scores for rivastigmine compared to placebo as follows: rivastigmine 1 to 4 mg/day at 18 weeks (WMD, -1.07; 95% CI, -1.66 to -0.48; P=0.0004) and 26 weeks (WMD, -0.84; 95% CI, -1.48 to -0.19; P=0.01); rivastigmine 6 to 12 mg/day at 12 weeks (WMD, -1.49; 95% CI, -1.96 to -1.01; P&lt;0.00001), 18 weeks (WMD, -1.79; 95% CI, -2.30 to -1.29; P&lt;0.00001) and 26 weeks (WMD, -1.99; 95% CI, -2.49 to -1.50; P&lt;0.00001).</p> <p>An additional analysis of ADAS-Cog dichotomized into those showing less than four points improvement and those showing four or more points improvement at 26 weeks shows benefit for cognitive function for the 6 to 12 mg daily of rivastigmine compared to placebo (83% did not show four points improvement compared to 89%; OR, 0.6; 95% CI, 0.4 to 0.8). There was no difference for the 1 to 4 mg/day dose compared to placebo (88% did not show four points improvement compared to 90%; OR, 0.84; 95% CI, 0.60 to 1.19).</p> <p>MMSE shows similar results in favor of rivastigmine at 26 weeks compared to placebo as follows: rivastigmine 1 to 4 mg/day at 26 weeks (WMD, 0.43; 95% CI, 0.08 to 0.78; P=0.02) and rivastigmine 6 to 12 mg/day at 26 weeks (WMD, 0.82; 95% CI, 0.56 to 1.08; P&lt;0.00001).</p> <p>One study used the SIB, which shows benefit associated with higher dose</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>rivastigmine compared to placebo at 26 weeks (WMD, 4.53; 95% CI, 0.47 to 8.59; P=0.03).</p> <p><u>Global assessment</u>            Using the CIBIC-Plus scale or the ADCS-CGIC scale, there were benefits associated with rivastigmine compared to placebo as follows: rivastigmine 6 to 12 mg/day at 12 weeks (OR, 0.74; 95% CI, 0.60 to 0.92; P=0.008), 18 weeks (OR, 0.79; 95% CI, 0.64 to 0.98; P=0.03) and at 26 weeks (OR, 0.66; 95% CI, 0.55 to 0.79; P&lt;0.00001); rivastigmine 1 to 4 mg/day at 26 weeks (OR, 0.71; 95% CI, 0.55 to 0.93; P=0.01).</p> <p>Using GDS, there were benefits associated with rivastigmine 6 to 12 mg/day compared to placebo (55% showed the worse condition compared to 59%; OR, 0.78; 95% CI, 0.64 to 0.94; P=0.01) but not with 1 to 4 mg daily rivastigmine compared to placebo.</p> <p><u>ADL</u>            The PDS showed an improvement associated with rivastigmine compared to placebo as follows: rivastigmine 6 to 12 mg/day at 12 weeks (WMD, 1.08; 95% CI, 0.19 to 1.98; P=0.02), 18 weeks (WMD, 1.90; 95% CI, 0.93 to 2.88; P=0.0001), and 26 weeks (WMD, 2.15; 95% CI, 1.13 to 3.16; P&lt;0.0001). One study assessing ADL using the ADCS-ADL scale and showed benefit for rivastigmine 6 to 12 mg/day at 24 weeks (WMD, 1.80; 95% CI, 0.20 to 3.40; P=0.03).</p> <p><u>Behavioral disturbance</u>            There was no difference between rivastigmine and placebo in behavioral disturbance found in two studies using the neuropsychiatric instrument (NPI-10, and NPI-12).</p> <p><u>Withdrawals before the end of treatment</u>            There were no significant differences in withdrawal rates with rivastigmine 1 to 4 mg/day and placebo at 12, 18 and 26 weeks.</p> <p>There were significant differences in withdrawal rates for the higher dose group in favor of placebo as follows: rivastigmine 6 to 12 mg/day at 12 weeks (OR, 2.60; 95% CI, 1.19 to 5.68; P=0.02), 18 weeks (OR, 4.02;</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>95% CI, 1.31 to 12.32; P=0.01), and 26 weeks (OR, 2.19; 95% CI, 1.83 to 2.63; P&lt;0.00001).</p> <p><u>Adverse events</u> There were no significant differences in the numbers of patients with at least one adverse event between the lower dose rivastigmine (1 to 4 mg/day) and placebo groups. There were significant differences between the higher dose rivastigmine (6 to 12 mg/day) and placebo groups in favor of placebo by the end of the titration period (OR, 2.96; 95% CI, 2.39 to 3.68; P&lt;0.00001) and by 26 weeks (OR, 2.49; 95% CI, 2.05 to 3.02; P&lt;0.00001).</p> <p>There were no significant differences in the numbers of patients with at least one severe adverse event between the lower dose rivastigmine (1 to 4 mg/day) and placebo groups. There were significant differences between the higher dose rivastigmine (6 to 12 mg daily) and placebo groups in favor of the placebo group for the titration period (OR, 1.88; 95% CI, 1.39 to 2.55; P&lt;0.0001).</p> <p>There were significant differences, in favor of placebo, for the rivastigmine 6 to 12 mg/day group by the end of the titration period, and by 26 weeks for the number of patients suffering nausea, vomiting, diarrhea, anorexia, headache, syncope, abdominal pain and dizziness. There were significant differences in favor of placebo, for the rivastigmine 1 to 4 mg/day group by the end of the titration period and by 26 weeks for the number of patients suffering nausea, vomiting, diarrhea, and anorexia.</p> <p>Secondary: Not reported</p>
<p>Rosler et al.<sup>67</sup> (1999)</p> <p>Rivastigmine 1 to 4 mg/day</p> <p>vs</p>	<p>DB, MC, PC, RCT</p> <p>Patients 50 to 85 years of age and not able to bear children, all patients met criteria for Alzheimer's type</p>	<p>N=725</p> <p>Dose titration over the first 12 weeks with a subsequent assessment</p>	<p>Primary: Improvements in cognitive function and overall clinical status measured by the ADAS-Cog, CIBIC, PDS, MMSE and GDS</p>	<p>Primary: Significant improvement in cognitive function assessed by the ADAS-Cog was observed with the higher dose group by <math>\geq 4</math> points compared to placebo (P&lt;0.05).</p> <p>At week 26, significantly more patients in both rivastigmine groups had improved in global function as assessed by the CIBIC compared to those in the placebo group (P&lt;0.05).</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>rivastigmine 6 to 12 mg/day</p> <p>vs</p> <p>placebo</p>	<p>dementia as described in the DSM-IV and criteria for probable Alzheimer's disease</p>	<p>period of 14 weeks, total of 26 weeks</p>	<p>Secondary: Safety and tolerability</p>	<p>Mean scores on the PDS improved from baseline in the higher dose group but fell in the placebo group (P&lt;0.05).</p> <p>At week 26, mean scores in the MMSE and the GDS significantly improved in patients receiving rivastigmine 6 to 12 mg/day (P&lt;0.05).</p> <p>Secondary: Discontinuation rates for any reason were significantly higher in the higher dose group than in the lower dose or placebo group (33% vs 14%).</p> <p>Adverse events related to treatment including nausea, vomiting, diarrhea, abdominal pain and anorexia, were generally mild and occurred most frequently during the dose escalation phase (23% in higher dose group, 7% in lower dose group and 7% in placebo group).</p>
<p>Articus et al.<sup>68</sup> (2011)</p> <p>Rivastigmine patch 9.5 mg/24 hours</p>	<p>MC, OL</p> <p>Patients with Alzheimer's disease</p>	<p>N=208</p> <p>24 weeks</p>	<p>Primary: Proportion of patients treated with rivastigmine for ≥8 weeks at week 24</p> <p>Secondary: Tolerability, week 24 MMSE, ADCS-CGIC, ADCS-ADL, ADCPQ, Zarit Burden Interview Score</p>	<p>Primary: In the ITT population, 80.8% of patients (95% CI, 75.0 to 86.5) were treated for at least eight weeks with rivastigmine. A total of 74.2% of patients (95% CI, 67.8 to 80.5) were treated for at least eight weeks and completed the study.</p> <p>A total of 74.2% of patients treated rivastigmine patch were able to reach and maintain the maximum dose for at least eight weeks. The most common adverse events being nausea (10.1%), erythema (8.7%), pruritus (8.2%), and vomiting (7.2%).</p> <p>Secondary: The most common adverse events were nausea (10.1%), erythema (8.7%), pruritus (8.2%), vomiting (7.2%), diarrhea (4.3%) and agitation (4.3%).</p> <p>At week 24, improvements were seen on: MMSE (1.3), and ADCS-ADL (1.3).</p> <p>At week 24, improvements in ADCS-CGIC were demonstrated in 34.6% of patients as assessed by patients, and in 29.7% of patients as assessed by the caregiver.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Grossberg et al.<sup>69</sup> (2009)</p> <p>Rivastigmine patch 9.5 mg/24 hours to 17.4 mg/24 hours</p>	<p>OL</p> <p>Patients 50 to 85 years of age with Alzheimer's disease (MMSE scores 10 to 20)</p>	<p>N=870</p> <p>28 weeks (weeks 25 to 52 of open-label extension)</p>	<p>Primary: Safety and tolerability</p> <p>Secondary: ADAS-cog</p>	<p>ADCPQ scores improved 18.5 points, and Zarit Burden Interview Score improved slightly at each visit until week 24 (-0.4).</p> <p>Primary: During the first four weeks of the open-label extension, patients formerly randomized to rivastigmine treatment (capsule or patch) reported fewer adverse events than those formerly randomized to placebo (<math>\leq 15.2</math> vs <math>28.2\%</math>). This prior exposure effect was noted for nausea (<math>\leq 2.5</math> vs <math>8.5\%</math>) and vomiting (<math>\leq 1.9</math> vs <math>6.0\%</math>).</p> <p>A total of 57.6% of patients reported adverse events during the OL extension (weeks 25 to 52), with nausea and vomiting being reported most frequently (15.7 and 14.3%, respectively).</p> <p>During the OL extension, over 90% of all patients experienced “no, slight, or mild” skin irritation as their most severe application-site reaction. The symptoms that were most commonly reported as moderate or severe were erythema and pruritus (7.7 and 5.6%, respectively).</p> <p>Serious adverse events occurred in 1.0% of patients during the first four weeks of the OL extension phase (weeks 25 to 28) and 9.4% of patients during the full open-label extension phase (weeks 25 to 52). The most common serious adverse events were gastrointestinal disorders (2.0%), infections and infestations (2.0%), cardiac disorders (1.7%), and nervous system disorders (1.5%).</p> <p>Eight deaths occurred during the OL extension phase and a further two occurred during the 30-day follow-up period. The causes of death were most commonly cardiac disorders (n=5) and nervous system disorders (n=3). None were considered treatment related.</p> <p>Secondary: Patients previously randomized to placebo who were switched to the 9.5 mg/24 hour rivastigmine patch during the OL extension experienced a 1.3-point increase in their ADAS-cog scores during weeks 24 to 40. There was no overall change in ADAS-cog score at week 40 compared to baseline (95% CI, -1.4 to 0.6). The increase in ADAS-cog score was not sustained beyond week 40.</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Patients receiving rivastigmine treatment for the entire study (weeks 0 to 52) showed a deterioration of 0.3 points (95% CI, -0.4 to 0.9) on the ADAS-cog at week 52. Those receiving placebo for weeks 0 to 24, followed by the patch, showed a deterioration of 0.9 points [95% CI, -0.4 to 2.1).
Gauthier et al. <sup>70</sup> (2013)  Rivastigmine transdermal patch 4.6 mg/24 hours or 9.5 mg/24 hours, once daily	OS  Patients with Alzheimer's disease with MMSE score of 10 to 26 and GDS score of 4 to 6	N=1,204  18 months	Primary: Change in MMSE from baseline to 18 months  Secondary: Change in MMSE at six and 12 months and change in GDS, assessment of patient ability, overall patient assessment rating, caregiver-reported compliance and treatment satisfaction at six, 12, and 18 months	Primary: Over 18 months of treatment there were no clinically significant changes in MMSE.  Secondary: Over 18 months of treatment there were no clinically significant changes in GDS.  The majority of patients showed improvement or no change in GDS, assessment of patient ability and overall patient assessment rating over 18 months.  The proportion with reported improvement in GDS, assessment of patient ability and overall patient assessment rating was higher than the proportion that deteriorated. Compliance improved from baseline to 18 months and for 88.2% of patient's caregivers preferred the transdermal patch to oral medications.
Sadowsky et al. <sup>71</sup> (2010)  <u>US13 and US18</u> Rivastigmine capsules 3 to 12 mg/day  <u>US38</u> Rivastigmine patch 4.6 mg/24 hours for 5 weeks, then rivastigmine patch	<u>US13 and US18</u> PRO, MC, OL  <u>US38</u> RCT, MC, OL  Patients ≥49 years of age with a diagnosis of dementia of the Alzheimer type (MMSE ≥8 to ≤26 or MMSE >10 to	N=592  25 to 26 weeks	Primary: Safety and tolerability	Primary: In US13 and US18, 67.7% of patients completed the studies and 32.3% of patients withdrew due to adverse events (59.8%), unsatisfactory treatment effect (15.9%), withdrawal of consent (15%), and loss to follow-up (6.5%). The remaining 2.7% of patients discontinued due to protocol deviation, administrative problem, or death.  In US13 and US18, the most frequently reported adverse events (AEs) were nausea (32.9%), vomiting (24.1%), dizziness (11.8%), weight loss (9.1%) agitation (7.9%), fall (7.9%) and confused state (7.9%). Serious AE's were reported in 6% of patients and included pneumonia (1.8%), syncope (1.2%), dehydration (1.2%) and vomiting (1.2%).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
9.5 mg/24 hours for 20 weeks	≤24) who showed a poor response to donepezil			<p>In US38, 67.4% of patients completed the study. The primary reasons for not completing the study were adverse events (44.7%), withdrawal of consent (29.4%), unsatisfactory treatment effect (10.6%), protocol deviation (7.1%), and loss to follow-up (3.5%). The remaining 4.7% of patients discontinued due to administrative problems, abnormal test procedure, or death.</p> <p>In US38, 70.5% of patients reported at least 1 AE. More patients in the immediate-switch group (73.3%) experienced at least one AE during the study than in the delayed-switch group (67.7%). The most common adverse events were application site reaction (15.3%), and agitation (6.9%). The most common serious AEs reported were syncope (1.1%), dehydration (0.8%) and pneumonia (0.4%).</p> <p>Discontinuation due to AE (14.6%) was the most common reason for patients not completing the extension phase in both immediate- and delayed-switch groups; the differences between the groups were NS. Discontinuations occurred for the following reasons: application site reaction (4.2%), disease progression (2.3%), and agitation (1.5%). Discontinuation due to gastrointestinal AEs was lower for the rivastigmine patch compared to the capsules.</p>
<p>Cummings et al.<sup>72</sup> (2012)</p> <p>10 cm<sup>2</sup> rivastigmine patch (9.5 mg/24 hours)</p> <p>vs</p> <p>15 cm<sup>2</sup> rivastigmine patch (13.3 mg/24 hours)</p>	<p>DB, PG, RCT</p> <p>Patients 50 to 85 years of age with MMSE scores of 10 to 24 diagnosed with Alzheimer's disease, all patients were required to be living with someone or to be in daily contact with a caregiver</p>	<p>N=567</p> <p>48 weeks</p>	<p>Primary: ADCS-IADL scale and ADAS-cog</p> <p>Secondary: Time to functional decline on the ADCS-IADL, change in the Trail Making Test parts A and B, and change in the NPI-10, and the NPI-caregiver distress scale.</p>	<p>Primary: The 13.3 mg/24 hours patch was statistically superior to the 9.5 mg/24 hours patch on the ADCS-IADL scale from week 16 (P=0.025) onwards including week 48 (P = 0.002), and ADAS-cog at week 24 (P= 0.027), but not at week 48 (P = 0.227).</p> <p>Secondary: Functional decline on the ADCS-IADL tended to occur later in the 13.3 mg/24 h patch group than in the 9.5 mg/24 hours patch group, but the observed difference did not reach significance.</p> <p>Proportion of patients with functional decline was 77.0% in the 13.3 mg/24 hours patch group compared to 81.2% with the 9.5 mg/24 hours patch Group. The difference was not statistically significant.</p> <p>Patients in the 13.3 mg/24 hours patch group had smaller increases in time</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>to complete the Trail Making Test parts A at weeks 24 and 48 compared to those in the 9.5 mg/24 hours patch group, but the observed difference did not reach significance.</p> <p>Differences were not significantly different in changes in the change in the 10-item (NPI-10), and the NPI-caregiver distress scale.</p> <p>The most frequently reported adverse events by primary system organ class were gastrointestinal disorders (29.3 vs. 19.1%, 13.3 and 9.5 mg/24 hours patch, respectively), psychiatric disorders (25.4 vs. 21.6%, respectively) and nervous system disorders (21.4 vs. 18.4%, respectively). Skin and subcutaneous tissue disorders were less frequently observed with the 13.3 mg/24 hours than the 9.5 mg/24 hours patch (2.1 vs 6%).</p>
<p>Cummings et al.<sup>73</sup> (2010)</p> <p>Rivastigmine patch 9.5 mg/24 hours</p> <p>vs</p> <p>rivastigmine patch 17.4 mg/24 hours</p> <p>vs</p> <p>placebo</p>	<p>DB, PC, PRO, RCT</p> <p>Patients 50 to 85 years of age with mild-to-moderate Alzheimer's disease</p>	<p>N=1,195</p> <p>24 to 52 weeks</p>	<p>Primary: Tolerability at 24 weeks</p> <p>Secondary: Patients skin condition at the application site at 28 weeks</p>	<p>Primary: No serious skin reactions were reported in either the 24 or 28 week phases of the study.</p> <p>During the 24 week period, 574 patients wearing an active patch and 579 patients wearing a placebo patch underwent at least one assessment of application-site skin condition. Of patients on the 9.5 mg/24 hour patch, erythema and pruritus were the most commonly reported reactions (moderate in 7.6% of patients and severe in 6.7% of patients). A total of 89.6% of patients in the patch group had "no, slight, or mild" signs and symptoms for their most severe application site reaction.</p> <p>Secondary: A total of 870 patients entered the 28 week phase of the study and received rivastigmine 9.5 mg/24 hours patch.</p> <p>Overall, the skin tolerability profile was similar to the DB phase. A total of 91.5% of patients experienced "no, slight, or mild" symptoms as their most severe application site reaction, with erythema and pruritus being the most common finding. A total of 3.7% of patients discontinued treatment due to skin reactions during the open-label extension, and there was no increase in the severity of skin reaction noted.</p>
<p>Molinuevo et al.<sup>74</sup> (2012)</p>	<p>MC, OS, PRO</p>	<p>N=649</p>	<p>Primary: Adherence rates</p>	<p>Primary: At baseline, 0.6% of patients were taking <math>\geq 80\%</math> of their medication as</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
Rivastigmine patch 9.5 mg/24 hours  vs  rivastigmine 3 to 12 mg/day	Patients with mild-to-moderate Alzheimer's disease	6 months	Secondary: Strategies followed by a physician to improve adherence and reasons for nonadherence reported by patients	prescribed. At three and six months, 77 and 88.1%, respectively, were noted to be taking more than 80% of their medication as prescribed (P<0.0001 vs baseline). The proportion of adherent patients at three months was 73.6% and at six months was 85.9% (P<0.0001).  Secondary: Modification of Alzheimer's disease treatment was the only intervention that substantially improved adherence at three months (P<0.0001). At the six month visit, psychoeducation was the only effective strategy that reached statistical significance (P<0.0001).  The most common reasons for nonadherence include forgetfulness (56.4%), avoidance of adverse events (30.7%), and refusal of treatment (25.3%).
Boada et al. <sup>75</sup> (2013)  Rivastigmine transdermal patch  vs  rivastigmine capsules	OL  Patients treated with rivastigmine	N=1,078  Duration not specified	Primary: Patient satisfaction (Treatment Satisfaction with Medicines and the Morisky-Green questionnaires)  Secondary: Not reported	Primary: Satisfaction reported was greater with transdermal than oral rivastigmine: mean+standard deviation of the total Treatment Satisfaction with Medicines score, 72.5+14.1 vs 65.2+12.5; P<0.001.  The proportion of adherent patients was greater with transdermal than with oral rivastigmine (65.0 vs 41.4%; P<0.001).  Satisfaction, in turn, was significantly greater in adherent cases than in nonadherent cases.  Secondary: Not reported
Blesa González et al. <sup>76</sup> (2011)  Rivastigmine 6 to 12 mg/day (RO)  vs  rivastigmine patch	MC, OL, RCT  Patients ≥60 years of age with mild-to-moderate Alzheimer's disease who were previously treated with oral rivastigmine	N=142  3 months	Primary: Gastrointestinal adverse events  Secondary: Overall tolerance, local tolerance for those patients on patches, satisfaction level, and cognitive	Primary: Gastrointestinal adverse events were reported in <5% of patients receiving patches (4.7% in RPT and 4.3% in RP) vs 6.1% in RO patients. No statistical significance was reached (P=0.8667). Gastrointestinal adverse events were noted in 11 cases, two in RPT patients, six in RP patients, and three in the RO patients (P=0.3067).  Secondary: Overall tolerability did not reveal any significant differences among the groups (P=0.8239).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
titrated to 9.5 mg/24 hours (RPT)  vs  rivastigmine patch 9.5 mg/24 hours (RP)			state by MMSE	<p>Local tolerability revealed skin or subcutaneous tissue adverse events reported in 11.6% of patients in the RPT group vs 17% of patients in the RP group (P=0.4055). All skin adverse events were reported as slight or moderate intensity.</p> <p>RP was defined by 72% of patients as very easy to use, while RO was considered very easy to use by 30% of patients (P=0.0005). In RP patients, 67% considered it very easy to follow compared to 19% of RO patients (&lt;0.0001). A total of 72% of RP patients confirmed the treatment never interfered with their daily lives vs 40% of the RO group (P=0.0085). Overall satisfaction comparisons revealed that in RP patients, 60% were very satisfied vs 14% in RO patients (P&lt;0.0001).</p> <p>MMSE did not demonstrate significant differences among treatment groups when compared at one and three month visits.</p>
Winblad et al. <sup>77</sup> (2007)  Rivastigmine patch 9.5 mg/24 hours  vs  rivastigmine patch 17.4 mg/24 hours  vs  rivastigmine 12 mg/day  vs  placebo	DD, PC, RCT  Patients 50 to 85 years of age with MMSE scores of 10 to 20 diagnosed with Alzheimer's disease, all patients were required to be living with someone or to be in daily contact with a caregiver	N=1,195  Dose titration in 4-week intervals over 16 weeks and maintained at their highest well-tolerated dose for a further 8 weeks, total of 24 weeks	Primary: ADAS-Cog subscale (assess orientation, memory, language, visuospatial and praxis function), ADCS-CGIC (assess single global rating)  Secondary: ADCS-ADL, MMSE, NPI, Ten Point Clock-drawing Test, and Trail-making Test part A	Primary: Patients in all rivastigmine groups (patch and capsule) showed significant improvements compared to placebo at week 24 with respect to ADAS-Cog and the ADCS-CGIC (all P<0.05 vs placebo).  Secondary: All rivastigmine groups (patch and capsule) showed statistically significant benefits over placebo on the ADCS-ADL, MMSE and Trail-making Test part A (all P<0.05 vs placebo).  Statistically significant treatment effects were not attained on the NPI or Ten Point Clock-drawing Test (P value not reported).
Winblad, Kawata et al. <sup>78</sup>	DB, DD, PC	N=1,059	Primary: ADCPQ	Primary: At 8 weeks, general preference was seen for the patch:

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2007)</p> <p>10 cm<sup>2</sup> rivastigmine patch (9.5 mg/24 hours)</p> <p>vs</p> <p>20 cm<sup>2</sup> rivastigmine patch (17.4 mg/24 hours)</p> <p>vs</p> <p>rivastigmine 6 mg capsules twice daily</p> <p>vs</p> <p>placebo</p>	<p>ACs included different size rivastigmine patches and rivastigmine capsules</p>	<p>24 week</p>	<p>Secondary: Not reported</p>	<p>68% of caregivers preferred the patch over capsule form (P&lt;0.0001). 70% of caregivers preferred the patch due to ease of schedule (P&lt;0.0001). 55% of caregivers preferred the patch due to ease of use (P=0.0008).</p> <p>At 24 weeks, general preference was seen for the patch: 72% of caregivers preferred the patch over capsule form (P&lt;0.0001). 74% of caregivers preferred the patch due to ease of schedule (P&lt;0.0001). 64% of caregivers preferred the patch due to ease of use (P&lt;0.0001). Caregivers preferred the patch over capsule dosage form, regardless of size of patch (P&lt;0.0001).</p> <p>At 8 weeks, caregivers indicated greater satisfaction overall (P&lt;0.0001), greater satisfaction with administration (P&lt;0.0001), less interference with daily life with the patch than the capsule (P&lt;0.01).</p> <p>Secondary: Not reported</p>
<p>Winblad et al.<sup>79</sup> (2007)</p> <p>Rivastigmine patch 9.5 mg/24 hours</p> <p>vs</p> <p>rivastigmine patch 17.4 mg/24 hours</p> <p>vs</p> <p>rivastigmine 12 mg/day</p>	<p>DB, DD, MC, PG</p> <p>Women or men 50 to 85 years of age with a diagnosis of dementia of the Alzheimer's type according to the DSM-IV, and probable Alzheimer's disease</p>	<p>N=1,195</p> <p>24 weeks</p>	<p>Primary: ADAS-Cog, ADCS-CGIC</p> <p>Secondary: ADCS-ADL scale; NPI for behavior and psychiatric symptoms; MMSE for cognition; Ten Point Clock-drawing Test for assessment of visuospatial and executive functions; Trail Making Test</p>	<p>Primary: Patients receiving rivastigmine patches or capsules showed significant benefits compared to placebo at week 24 on the ADAS-Cog subscale (P&lt;0.05 vs placebo for all rivastigmine groups).</p> <p>Treatment differences on the ADCS-CGIC were statistically significant for the 10 cm<sup>2</sup> patch and capsule group (all P&lt;0.05 vs placebo). The 20 cm<sup>2</sup> patch did not achieve statistical significance compared to placebo in the analysis (P=0.054).</p> <p>Secondary: Rivastigmine patches and capsule provided statistically significant benefits over placebo on the ADCS-ADL, MMSE and Trail-making Test A (all P&lt;0.05 vs placebo).</p> <p>Changes from baseline on the NPI, NPI-distress subscale, and Ten-point</p>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo			Part A for assessment of attention, visual tracking and motor processing speed	Clock-drawing Test in the rivastigmine groups were not significantly different from those in the placebo groups (all P>0.05).
Blesa et al. <sup>80</sup> (2007)  Rivastigmine patch 9.5 mg/24 hours  vs  rivastigmine patch 17.4 mg/24 hours  vs  rivastigmine 12 mg/day  vs  placebo	DB, DD, PC  ACs included different size rivastigmine patches and rivastigmine capsules, caregiver preference based on data generated during the IDEAL trial (Winblad et al)	N=1,059  24 week	Primary: ADCPQ  Secondary: Not reported	Primary: At 8 weeks, general preference was seen for the patch: 68% of caregivers preferred the patch over capsule form (P<0.0001). 70% of caregivers preferred the patch due to ease of schedule (P<0.0001). 55% of caregivers preferred the patch due to ease of use (P=0.0008).  At 24 weeks, general preference was seen for the patch: 72% of caregivers preferred the patch over capsule form (P<0.0001). 74% of caregivers preferred the patch due to ease of schedule (P<0.0001). 64% of caregivers preferred the patch due to ease of use (P<0.0001). Caregivers preferred the patch over capsule dosage form, regardless of size of patch (P<0.0001).  At eight weeks, caregivers indicated greater satisfaction overall (P<0.0001), greater satisfaction with administration (P<0.0001), less interference with daily life with the patch than the capsule (P<0.01).  Secondary: Not reported
Farlow et al. <sup>81</sup> (2011)  Rivastigmine patch 9.5 mg/24 hours  vs  rivastigmine patch 17.4 mg/24 hours  vs	RETRO  Patients with mild-to-severe Alzheimer's disease	N=1,050  24 weeks	Primary: ADAS-cog, ADCS-CGIC, and ADCS-ADL  Secondary: Not reported	Primary: In patients with moderate disease, there was a significant improvement on ADAS-cog scores with the rivastigmine 17.4 mg/24 hour patch (P=0.0009) and rivastigmine capsule (P=0.0128).  For patients with moderately severe disease, there was a significant improvement in ADAS-cog scores with the rivastigmine 17.4 mg/24 hour patch (P=0.006), rivastigmine 9.5 mg/24 hour patch (P=0.0163), and rivastigmine capsule (P=0.0071) compared to placebo.  For patients with severe disease, there was a significant improvement on ADCS-CGIC scores with the rivastigmine 9.5 mg/24 hour patch (P=0.037) and rivastigmine capsule (P=0.0073) compared to placebo.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
rivastigmine 12 mg/day  vs  placebo				<p>For patients with moderately severe disease, there was a significant improvement on ADCS-CGIC scores with the rivastigmine 17.4 mg/24 hour patch (P=0.043) and rivastigmine 9.5 mg/24 hour patch (P=0.0116) compared to placebo.</p> <p>Significant improvement on ADCS-CGIC scores were seen with the rivastigmine 17.4 mg/24 hour patch in patients with moderate disease (P=0.03) and mild to moderate disease (P=0.0455) compared to placebo.</p> <p>For patients with moderately severe disease, there was a significant improvement on ADCS-ADL scores with the rivastigmine 17.4 mg/24 hour patch (P=0.0211) compared to placebo.</p> <p>For patients with moderate disease, there was a significant improvement on ADCS-ADL scores with the rivastigmine 17.4 mg/24 hour patch (P=0.0194) and rivastigmine capsule (P=0.0077) compared to placebo.</p> <p>There was no significant difference in ADCS-ADL scores among the treatment groups in patients with severe AD.</p>
Choi et al. <sup>82</sup> (2011)  Rivastigmine patch 4.6 mg/24 hours for 4 weeks, then rivastigmine patch 9.5 mg/24 hours for 4 weeks, then rivastigmine patch 9.5 mg/24 hours and memantine 5 mg/day titrated to 20 mg/day  vs	MC, OL, RCT  Patients with mild-to-moderate Alzheimer's disease	N=172  24 weeks	Primary: Tolerability  Secondary: Efficacy as measured by CMAI-K, ADAS-cog, K-MMSE, FAB, CGA-NPI, ADCS-ADL and CDR-SB scores	Primary: The incidence of adverse events (53.4 vs 50.6%) and discontinuation due to adverse events (6.8 vs 4.8%) was not different between patients with and without memantine, respectively.  The most common adverse events were skin irritation in both treatment groups (42 vs 34.9%; P=0.71), but discontinuation was rare (4.5 vs 2.4%; P=0.74).  Secondary: CMAI-K scores favored rivastigmine monotherapy vs combination therapy at the end of treatment (P=0.01). Changes in other efficacy measures (ADAS-cog, K-MMSE, FAB, CGA-NPI, ADCS-ADL and CDR-SB) were not significantly different.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>rivastigmine patch 9.5 mg/24 hours</p> <p>Farlow et al.<sup>83</sup> (2010)</p> <p>Rivastigmine patch 9.5 mg/24 hours and memantine</p> <p>vs</p> <p>rivastigmine patch 9.5 mg/24 hours</p>	<p>OL, RCT</p> <p>Patients ≥50 years of age with mild-to-moderate Alzheimer's disease who had been receiving donepezil for at least 6 months and at a stable dose of 5-10 mg/day for a minimum of 3 months</p>	<p>N=261</p> <p>25 weeks</p>	<p>Primary: Safety and tolerability of rivastigmine transdermal patch, with or without concomitant memantine</p> <p>Secondary: Changes in cognition, global functioning and activities of daily living measured by MMSE and ADCS-ADL using the CGIC</p>	<p>Primary: The incidences of adverse events (73.3 vs 67.5%) and serious adverse events (10.4 vs 7.1%) were both slightly higher in patients receiving concomitant memantine, but the differences were NS (95% CIs, -5.2 to 16.9 and -3.6 to 10.1 for adverse events and serious adverse events, respectively).</p> <p>The most frequent adverse events in the combination therapy group and the rivastigmine monotherapy group were application site reactions (17.5 vs 13.5%, respectively) and agitation (5.9 vs 7.9%, respectively).</p> <p>Secondary: Concomitant memantine was associated with no significant changes in efficacy, as assessed by CGIC and MMSE scores. Global functioning remained unchanged or improved (CGIC rating ≤4) in 57.7 and 67.2% of patients with memantine and patients without memantine, respectively (P=0.604).</p> <p>ADCS-ADL scores deteriorated from baseline in both groups, with significant worsening in patients receiving memantine compared to those not receiving memantine (mean change from baseline rivastigmine and memantine vs rivastigmine monotherapy: -5.3 vs -2.0; P=0.043).</p>
<p>Harry et al.<sup>84</sup> (2005)</p> <p>Donepezil with doses ranging from 5 to 10 mg/day</p> <p>or</p> <p>galantamine with doses ranging from 8 to 36 mg/day</p>	<p>MA</p> <p>Patients with mild-to-moderate Alzheimer's disease, and without diagnosis of any other psychiatric or neurological disorder</p>	<p>N=3,353</p> <p>3 donepezil studies</p> <p>5 galantamine studies</p> <p>Duration varied</p>	<p>Primary: ADAS-Cog or MMSE</p> <p>Secondary: Not reported</p>	<p>Primary: The majority of patients showed no difference compared to placebo.</p> <p>There was no significant difference in efficacy between the groups.</p> <p>Secondary: Not reported</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo				
<p>Wilcock et al.<sup>85</sup> (2003)</p> <p>Donepezil 10 mg/day</p> <p>vs</p> <p>galantamine 24 mg/day</p>	<p>MC, PG, RCT</p> <p>Patients with Alzheimer's disease</p>	<p>N=182</p> <p>52 weeks</p>	<p>Primary: BrADL</p> <p>Secondary: MMSE, ADAS-Cog, NPI</p>	<p>Primary: BrADL total score showed no significant difference between treatment groups in mean change from baseline to week 52.</p> <p>Secondary: Galantamine patients' scores on the MMSE at week 52 did not differ significantly from baseline, whereas donepezil patients' scores deteriorated significantly from baseline (P&lt;0.0005). The between group difference in MMSE change did not reach statistical significance.</p> <p>In the ADAS-Cog analysis, between group differences for the total population were NS, whereas galantamine treated patients with MMSE scores of 12 to 18 demonstrated an increase (worsening) in the ADAS-Cog score of 1.61+/-0.80 vs baseline, compared to an increase of 4.08+/-0.84 for patients treated with donepezil.</p> <p>More caregivers of patients receiving galantamine reported reductions in burden compared to donepezil.</p> <p>Changes from baseline in NPI were similar for both treatments.</p>
<p>Jones et al.<sup>86</sup> (2004)</p> <p>Donepezil 10 mg/day</p> <p>vs</p> <p>galantamine 12 mg twice daily</p>	<p>OL, RCT</p> <p>Patients with Alzheimer's disease</p>	<p>N=120</p> <p>12 weeks</p>	<p>Primary: Ease of use and tolerability, ADAS-Cog, effects on cognition and activities of daily living</p> <p>Secondary: Not reported</p>	<p>Primary: Physicians and caregivers reported statistically significant greater satisfaction/ ease of use with donepezil compared to galantamine at weeks four and 12.</p> <p>Significantly greater improvements in cognition were observed for donepezil vs galantamine on the ADAS-Cog at week 12 and at endpoint.</p> <p>Activities of daily living improved significantly in the donepezil group compared to the galantamine group at weeks four and 12 (P&lt;0.05).</p> <p>Forty-six percent of galantamine patients reported gastrointestinal adverse events vs 25% of donepezil patients.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Modrego et al. <sup>87</sup> (2010)  Donepezil 10 mg/day  vs  memantine 20 mg/day	PG, RCT, SB  Patients with mild-to-moderate Alzheimer's disease	N=63  6 months	Primary: ADAS-cog, NPI, DAD, changes in N-acetylaspartate metabolite levels  Secondary: Not reported	Primary: There were no significant differences in the clinical scales with donepezil and memantine (donepezil: ADAS-cog, -0.12; P=NS, NPI, -0.04; P=NS, DAD, 6.67; P=0.014) (memantine: ADAS-cog, -1.37; P=NS, NPI, 1.25; P=NS, DAD, 4.46; P=NS). More patients worsened than improved on either drug.  Daily living activities decreased by 4.4% in the memantine group and 6.6% in the donepezil group (P=0.6).  At baseline, N-acetylaspartate/Cr ratio in the PCG correlated significantly with the ADAS-cog (P=0.02) and MEC (P=0.02). The N-acetylaspartate/Cr ratio correlated with the baseline ADAS-cog (P=0.02) in the left temporal lobe.  At week 24, the PCG was the only area where the correlation was significant. The patients who improved in the ADAS-cog showed increases in the N-acetylaspartate/Cr ratios (P=0.004). None of the baseline metabolite levels predicted response to treatment in any of the examined areas.  Secondary: Not reported
Rozankovic et al. <sup>88</sup> (2021) [Abstract only]  Donepezil  vs  Memantine	OL, RCT  Patients with moderate Alzheimer's disease	N=85  6 months	Primary: NPI at baseline to six months  Secondary: Not reported	Primary: The NPI total score improved from baseline to month six in both groups (P<0.0001).  Analyses of the NPI subdomains revealed that both donepezil treatment and memantine treatment produced statistically significant improvement in all of the NPI domains except euphoria and apathy, for which no improvement was observed after memantine treatment.  Secondary: Not reported
Wilkinson et al. <sup>89</sup>	OL, RCT	N=111	Primary:	Primary:

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>(2002)</p> <p>Donepezil 10 mg/day</p> <p>vs</p> <p>rivastigmine 6 mg twice daily</p>	<p>Patients with mild-to-moderate Alzheimer's disease</p>	<p>12 weeks</p>	<p>ADAS-Cog, tolerability</p> <p>Secondary: Not reported</p>	<p>More patients taking donepezil completed the study (89.3%) compared to the rivastigmine group (69.1%; P=0.009).</p> <p>10.7% of the donepezil group and 21.8% of the rivastigmine group discontinued treatment due to adverse events.</p> <p>87.5% of the donepezil patients and 47.3% of the rivastigmine patients remained on the maximum approved dose of each drug at the last study visit.</p> <p>Both groups showed comparable improvements in ADAS-Cog administered at weeks four and 12.</p> <p>Secondary: Not reported</p>
<p>Van Puyvelde et al.<sup>90</sup> (2011)</p> <p>Galantamine</p> <p>vs</p> <p>donepezil or rivastigmine (safety control group)</p>	<p>MC, OS, PRO</p> <p>Patients with mild-to-moderate Alzheimer's disease</p>	<p>N=128</p> <p>6 months</p>	<p>Primary: Safety, patients and caregiver satisfaction, global impression as reported by the physician</p> <p>Secondary; Not reported</p>	<p>Primary: Adverse events were similar among both treatment groups (galantamine, 34%; SCG, 34.4%). The incidence of serious (12 events) and severe (15 events) adverse events with galantamine was similar to the SCG group (serious: galantamine 9.3% vs safety control group 9.7%); severe: galantamine 11.3% vs safety control group 12.9%.</p> <p>A total of 84.5% of patients treated with galantamine continued their treatment after six months.</p> <p>Patients receiving galantamine reported their condition as improved (49%), unchanged (47%) and worsened (4%).</p> <p>Caregivers rated global evaluation as better (37%), unchanged (41%) and worse (22%) with galantamine.</p> <p>Physicians rated global clinical impression of change as better (46%), unchanged (34%) and worse (20%) with galantamine.</p> <p>Measurements of cognition and behavior remained stable. The appreciation of physicians and caregivers corresponded well (P&lt;0.001).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Tariot et al. <sup>91</sup> (2004)  Memantine 20 mg/day  vs  donepezil	DB, MC, PC, RCT  Patients with moderate-to-severe Alzheimer's disease who received stable doses of donepezil	N=404  24 weeks	Primary: SIB, ADCS-ADL, CIBIC-Plus, BGP  Secondary: Not reported	Secondary: Not reported  Primary: A significantly greater therapeutic effect was observed in the memantine group than in the placebo group on the ADCS-ADL, SIB and CIBIC-Plus.  Patients receiving memantine in combination with donepezil demonstrated significantly less decline in ADCS-ADL scores compared to patients receiving donepezil-placebo over the 24-week study period (P=0.02).  Patients receiving memantine showed significantly less cognitive decline in SIB scores compared to patients receiving placebo. Therapy with memantine-donepezil resulted in sustained cognitive performance above baseline compared to the progressive decline seen with the donepezil-placebo treatment.  The change in total mean scores favored memantine vs placebo for the CIBIC-Plus (possible score range was 1-7), 4.41 vs 4.66, respectively (P=0.03).  Treatment discontinuations due to adverse events for memantine vs placebo were 7.4% of the patients compared to 12.4%.  Secondary: Not reported
Bullock et al. <sup>92</sup> (2005)  Rivastigmine 3 to 12 mg/day  vs  donepezil 5 to 10 mg/day	DB, MC, RCT  Patients 50 to 85 years of age with moderate to moderately-severe Alzheimer's disease (MMSE score 10-20)	N=994  24 months	Primary: SIB  Secondary: GDS, ADCS-ADL, MMSE, NPI	Primary: Donepezil-treated patients declined 9.91 points from baseline on the SIB as compared to rivastigmine-treated patients, who declined by 9.30 points (P=NS).  Secondary: Rivastigmine was more effective than donepezil on the ADCS-ADL, on which there was a between-treatment difference of 2.1 points after two years (P=0.007), and greater efficacy on the GDS (P=0.049). There were no significant differences in MMSE and NPI between the treatment groups.

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				More patients receiving rivastigmine reported 'any adverse event' compared to those receiving donepezil during the titration phase (82.0 and 64.7%, respectively). Adverse events were higher with rivastigmine during the titration phase and included nausea (32.9 vs 15.2%) and vomiting (27.9 vs 5.8%). In the maintenance phase, adverse event rates in the two groups were similar (78.7% for the rivastigmine group and 76.9% for the donepezil group). Premature discontinuations due to adverse events were higher in the rivastigmine group during the titration phase (14.1 vs 7.0% for donepezil) but similar in the maintenance phase (17.9 vs 14.1% for donepezil).
Mossello et al. <sup>93</sup> (2004)  Donepezil 5 to 10 mg/day  vs  galantamine 16 to 24 mg/day  vs  rivastigmine 6 to 12 mg/day	OL, OS  Patients with mild-to-moderate Alzheimer's disease	N=407  9 months	Primary: MMSE, ADL and IADL  Secondary: Not reported	Primary: There were no differences amongst the three groups in regard to any of the outcome measures (galantamine was not included in the MMSE comparison due to the small number of treated patients).  Discontinuation due to adverse effects was lower in those patients on donepezil (3%) vs rivastigmine (17%; P=0.01) and vs galantamine (21%; P=0.01).  Secondary: Not reported
Aguglia et al. <sup>94</sup> (2004)  Donepezil  vs  galantamine  vs  rivastigmine	OL  Patients with Alzheimer's disease	N=242  6 months	Primary: MMSE, ADAS-Cog, ADL and IADL  Secondary: Not reported	Primary: There were no statistical differences on changes in the MMSE, ADAS-Cog, ADL or IADL measures amongst the three groups.  There were no differences on changes in the IADL measure among the three groups.  In the ADL measure, donepezil and galantamine patients showed a decrease while there was no change for rivastigmine patients.  Rivastigmine showed a small numerical advantage (but not statistically) compared to donepezil and galantamine on the ADAS-Cog.



Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				Secondary: Not reported
Lopez-Pousa et al. <sup>95</sup> (2005)  Donepezil  vs  galantamine  vs  rivastigmine  vs  historical controls	OL, PRO  Patients with mild-to-moderate Alzheimer's disease	N=147  6 months	Primary: MMSE  Secondary: Not reported	Primary: All three treatment groups had better MMSE scores compared to control (donepezil; P<0.001, galantamine; P<0.01, and rivastigmine; P<0.03).  There were no statistical differences between the groups on measures of cognitive decline (via MMSE).  Secondary: Not reported
Rodda et al. <sup>96</sup> (2009)  Donepezil 5 to 10 mg/day  vs  galantamine 8 to 24 mg/day  vs  rivastigmine 9 to 17.4 mg/day	RETRO  Patients with Alzheimer's disease being treated with donepezil, rivastigmine or galantamine monotherapy	N=6,110  12 to 170 weeks	Primary: NPI  Secondary: Not reported	Primary: Three of the 14 studies reviewed reported statistically significant improvement in overall NPI score or in the agitation/aggression item of the NPI only. One study demonstrated a significant difference in NPI score between groups randomized to either continuation or discontinuation of donepezil (placebo following an initial OL treatment phase. Of these four positive studies, two specified a minimum level of behavioral disturbance at baseline and used behavioral scores as a primary outcome.  Secondary: Not reported
Howard et al. <sup>97</sup> (2012)	DB, MC, RCT	N=295	Primary: Standardized Mini-	Primary: Mean donepezil vs placebo Standardized Mini-Mental State Examination

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Donepezil 10 mg/day</p> <p>vs</p> <p>memantine 20 mg/day</p> <p>vs</p> <p>donepezil 10 mg/day and memantine 20 mg/day</p> <p>vs</p> <p>placebo</p>	<p>Community-based patients with moderate-to-severe Alzheimer's disease who were taking donepezil 10 mg/day for <math>\geq 3</math> months</p>	<p>52 weeks</p>	<p>Mental State Examination and BADLS scores</p> <p>Secondary: NPI, caregiver health status assessed by General Health Questionnaire 12</p>	<p>scores were higher with donepezil (better cognitive function) by an average of 1.9 points (95% CI, 1.3 to 2.5; <math>P &lt; 0.001</math>) and BADLS scores were lower (less functional impairment) by 3.0 points (95% CI, 1.3 to 2.5; <math>P &lt; 0.001</math>). Both outcomes demonstrated significant heterogeneity in treatment efficacy over time (<math>P = 0.002</math> and <math>P = 0.004</math>, respectively), with less benefit apparent at the six week assessment than at later time points. From six weeks onward, differences were roughly parallel.</p> <p>Mean donepezil+memantine vs placebo+memantine Standardized Mini-Mental State Examination scores were higher with donepezil by an average of 1.2 points (95% CI, 0.6 to 1.8; <math>P &lt; 0.001</math>) and BADLS scores were lower by 1.8 points (95% CI, 0.3 to 2.8; <math>P &lt; 0.001</math>). Both outcomes were smaller than the minimum clinically important difference. Interactions of memantine therapy with visit were NS. Both donepezil and memantine demonstrated benefits on both Standardized Mini-Mental State Examination and BADLS larger in the absence of other agents alone, though statistically insignificant (<math>P = 0.14</math> and <math>P = 0.09</math>, respectively).</p> <p>No significant benefits were seen adding memantine to donepezil on Standardized Mini-Mental State Examination scores (0.8 points higher with memantine and placebo; 95% CI, -0.1 to 1.6; <math>P = 0.07</math>) or BADLS scores (0.5 points lower with memantine than placebo; 95% CI, 2.2 to 1.2; <math>P = 0.57</math>).</p> <p>Secondary: NPI scores were lower for patients on memantine compared to placebo, indicating fewer behavioral and psychological symptoms by 4.0 points (99% CI, 0.6 to 7.4; <math>P = 0.002</math>).</p> <p>No observable NPI differences noted with continuation, as compared to discontinuation of donepezil therapy (2.3 points lower with continuation; 95% CI, -1.1 to 5.7; <math>P = 0.08</math>). Donepezil+memantine vs donepezil demonstrated a lower NPI score by 5.1 points (99% CI, 0.3 to 9.8; <math>P = 0.006</math>).</p> <p>Continuation of donepezil and donepezil+memantine compared to the placebo and memantine + placebo demonstrated larger average decreases</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				(indicating fewer psychological symptoms) across trial visits in General Health Questionnaire 12 scores for caregiver health status. There was a 0.5 point larger decrease with continuation vs discontinuation of donepezil (99% CI, -0.01 to 1.0; P=0.01) and 0.5 point larger decrease with memantine vs placebo (95% CI, -0.1 to 0.9; P=0.03), though significance was not reached to allow for multiple secondary outcomes.
Porsteinsson et al. <sup>98</sup> (2008)  Memantine 20 mg/day plus cholinesterase inhibitor  vs  cholinesterase inhibitor plus placebo	PC, R  Patients with probable Alzheimer's disease, MMSE scores between 10 to 22, concurrently taking a cholinesterase inhibitor	N=433  24 weeks	Primary: ADAS-cog, CIBIC-Plus  Secondary: ADCS-ADL, NPI, MMSE	Primary: No significant difference in ADAS-cog and CIBIC-Plus was found between memantine and placebo.  Secondary: No significant difference in ADCS-ADL, NPI or MMSE was found between memantine and placebo.
Cumming et al. <sup>99</sup> (2006)  Memantine 20 mg/day plus donepezil  vs  donepezil	DB, PC, PG, PRO  Patients with moderate-to-severe Alzheimer's disease who received stable doses of donepezil	N=404  24 weeks	Primary: NPI  Secondary: Not reported	Primary: NPI scores significantly favored the memantine group at 12 weeks and at 24 weeks. At week 12, NPI scores increased (worsening behavior) 1.7 points in the placebo group and decreased 2.5 points in the memantine group (P<0.001). At week 24, NPI scores increased 3.7 points (worsening behavior) in the placebo groups and the memantine group returned to baseline (P=0.002).  Fewer patients developed delusions in the memantine treatment group than the placebo group (P=0.011).  Secondary: Not reported
Maidment et al. <sup>100</sup>  Memantine 20 mg daily	MA  Patients with probable	N=1,750  Duration varied	Primary: NPI  Secondary:	Primary: Compared to the placebo group patients receiving memantine improved by 1.99 on the NPI scale (95% CI, -0.08 to -3.91; P=0.041).

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs  placebo  or  memantine 20 mg daily in combination with a cholinesterase inhibitor (doses varied)  vs  placebo in combination with a cholinesterase inhibitor (doses varied)	Alzheimer's disease		Not reported	Secondary: Not reported
Wilkinson et al. <sup>101</sup> (2009)  Cholinesterase inhibitors (donepezil 5 or 10 mg/day)  vs  placebo	MA  Patients with mild-to-moderate Alzheimer's disease	N=906 (3 trials)  24 weeks	Primary: MMSE  Secondary: Not reported	Primary: A significantly greater percentage of placebo patients than donepezil-treated patients met the specified criteria for all three definitions of clinical worsening. The OR for clinical worsening were significantly reduced for donepezil-treated patients compared to placebo patients (P<0.0001 for all definitions).  Among patients meeting criteria for clinical worsening, mean declines in MMSE scores were greater for placebo than donepezil-treated patients.  This outcome was also apparent when milder (MMSE, 18 to 26) and more moderate (MMSE, 10 to 17) subgroups were analyzed separately.  Secondary: Not reported
Feldman et al. <sup>102</sup>	OS, PRO	N=548	Primary:	Primary:

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
(2009)  Cholinesterase inhibitors	Alzheimer's disease patients with and without cerebrovascular disease	7 years	Time to nursing home placement  Secondary: Identify factors noted to reduce risk of NHP, including measurement of DAD and MMSE	The overall median time to permanent institutional admission was 42.4 months (95% CI, 38.0 to 48.0 months).  Secondary: Factors noted to reduce the risk of being admitted to a nursing home included higher baseline DAD and MMSE scores, Alzheimer's disease diagnosis, living with caregiver, country, and treatment duration (P<0.05).  Each year of treatment demonstrated a reduced risk of nursing home admission (galantamine, -31%, other cholinesterase inhibitors, -29%).
Trinh et al. <sup>103</sup> (2003)  Cholinesterase inhibitors  vs  placebo	MA  Trials included outpatients with mild or moderate Alzheimer's disease who were treated for at least one month with a cholinesterase inhibitor	29 trials  Duration varied	Primary: NPI, ADAS-noncog, ADL and IADL  Secondary: Not reported	Primary: Cholinesterase inhibitors improved the NPI statistically better than placebo (95% CI, 0.87 to 2.57).  Cholinesterase inhibitors improved the ADAS-noncog measure numerically but not statistically compared to placebo (95% CI, 0.0 to 0.05).  Cholinesterase inhibitors improved ADL numerically but not significantly better than placebo (95% CI, 0.0 to 0.19).  Cholinesterase inhibitors improved IADL statistically compared to placebo (95% CI, 0.01 to 0.17).  Secondary: Not reported
Lanctot et al. <sup>104</sup> (2003)  Cholinesterase inhibitors  vs  placebo	MA  Adult patients diagnosed with Alzheimer's disease	N=7,954  16 trials that varied in duration	Primary: Global responders, using CGI-C, CIBIC, adverse, events, dropouts  Secondary: Not reported	Primary: For cholinesterase inhibitors the pooled mean proportion of global responders was in excess by 9% when compared to the placebo treatment (9%; 95% CI, 6 to 12).  In the cholinesterase inhibitor treatment groups the rates of adverse events, dropout for any reason and dropout because of adverse events were higher compared to the placebo treatment groups (8%; 95% CI, 5 to 11; 8%; 95% CI, 5 to 11; and 7%; 95% CI, 3 to 10).  The number needed to treat for one additional patient to benefit was 7

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>(95% CI, 6 to 9) for stabilization or better, 12 (95% CI, 9 to 16) for minimal improvement or better and 42 (95% CI, 26 to 114) for marked improvement.</p> <p>The number needed to treat for one additional patient to experience an adverse event was 12 (95% CI, 10 to 18).</p> <p>Secondary: Not reported</p>
<p>Birks et al.<sup>105</sup> (2006)</p> <p>Cholinesterase inhibitors</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients diagnosed with mild, moderate or severe dementia due to Alzheimer's disease</p>	<p>N=7,298</p> <p>Minimum 6 months</p>	<p>Primary: CIBIC-Plus, GBS, GDS, ADAS-Cog, MMSE, SIB, NPI, ADL scored by PDS and DAD</p> <p>Secondary: Withdrawals prior to six months, adverse events</p>	<p><i>Cholinesterase inhibitor vs placebo (12 trials)</i></p> <p>Primary: Significant benefit was seen in CIBIC-Plus for patients treated with a cholinesterase inhibitor over placebo; more patients were scored as "showed improvement" than "showed decline/no change" (OR, 1.56; 95% CI, 1.32 to 1.85; P&lt;0.00001); eight studies.</p> <p>No significant difference was seen in GBS between the cholinesterase inhibitor and placebo groups at one year (P value not reported): one trial.</p> <p>Significant improvement in ADAS-Cog was found for patients treated with donepezil, galantamine, or rivastigmine over placebo (WMD, -2.66; 95% CI, -3.02 to -2.31; P&lt;0.00001): 10 studies.</p> <p>Significant benefit was seen in MMSE for patients treated with a cholinesterase inhibitor over placebo (WMD, 1.37; 95% CI, 1.13 to 1.61; P&lt;0.00001): nine studies.</p> <p>Significant benefit was seen in ADL-PDS and DAD for patients treated with a cholinesterase inhibitor over placebo (WMD, 2.40; 95% CI, 1.55 to 3.37; P&lt;0.00001 for PDS; and WMD, 4.39; 95% CI, 1.96 to 6.81; P=0.0004 for DAD).</p> <p>Significant benefit was seen in NPI for patients treated with a cholinesterase inhibitor over placebo (WMD, -2.44; 95% CI, -4.12 to -0.76; P=0.004).</p> <p>Secondary:</p>

Alzheimer's Agents  
AHFS Classes 120400 and 289200

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Significantly more patients treated with a cholinesterase inhibitor (29%) withdrew prior to six months than those in the placebo groups (18%; <math>P &lt; 0.00001</math>).</p> <p>Adverse events that occurred significantly more frequently in the cholinesterase inhibitor group than the placebo group, from pooled data from at least 6 trials included: abdominal pain, anorexia, dizziness, diarrhea, headache (<math>P &lt; 0.0001</math>), insomnia (<math>P = 0.007</math>), nausea, vomiting (<math>P &lt; 0.00001</math> unless noted).</p> <p><i>Donepezil vs rivastigmine (one trial)</i>  <b>Primary:</b>            There was no statistically significant difference between the treatment groups for cognitive function, ADL scales, behavior disturbances and global assessment (P values not reported).</p> <p><b>Secondary:</b>            Significantly fewer patients in the donepezil group withdrew from treatment after 2 years than in the rivastigmine group (OR, 0.64; 95% CI, 0.50 to 0.83; <math>P = 0.0006</math>).</p> <p>Adverse events that occurred significantly more frequently at 12-16 weeks of treatment in the rivastigmine group than in the donepezil group included: nausea (<math>P &lt; 0.00001</math>), vomiting (<math>P &lt; 0.00001</math>), falls (<math>P = 0.01</math>), hypertension (<math>P = 0.01</math>), anorexia (<math>P = 0.0005</math>) and weight loss (<math>P = 0.001</math>), and after 16 weeks to 2 years of treatment: nausea (<math>P = 0.0002</math>), vomiting (<math>P &lt; 0.00001</math>) and anorexia (<math>P = 0.02</math>).</p> <p>No significant difference between treatment groups for serious adverse events was noted (P value not reported).</p>
Hansen et al. <sup>106</sup> (2008)  Cholinesterase inhibitors	MA  Patients with Alzheimer's disease	26 trials  Variable duration	<b>Primary:</b> Cognition (ADAS-cog), function, behavior (NPI), global assessment of change (CIBIC+ and CGI-C)	<b>Primary:</b> <u>Cognition (14 studies)</u> The pooled WMD in change between active treatment and placebo was -2.67 (95% CI -3.28 to -2.06) for donepezil, -2.76 (95% CI -3.17 to -2.34) for galantamine, and -3.01 (95% CI -3.80 to -2.21) for rivastigmine.  <u>Function (14 studies)</u>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			Secondary: Not reported	<p>The pooled standardized mean difference between active treatment and placebo was 0.31 (95% CI, 0.21 to 0.40) for donepezil, 0.27 (95% CI, 0.18 to 0.36) for galantamine, and 0.26 (95% CI, 0.11 to 0.40) for rivastigmine.</p> <p><u>Behavior (seven studies)</u> The pooled WMD in NPI score between active treatment and placebo was -4.3 (95% CI, -5.95 to -2.65) for donepezil and -1.44 (95% CI, -2.39 to -0.48) for galantamine.</p> <p><u>Global assessment of change (nine studies)</u> The pooled RR of responding for active treatment compared to placebo was 1.88 (95% CI, 1.50 to 2.34) for donepezil, 1.15 (95% CI, 0.96 to 1.39) for galantamine, and 1.64 (95% CI, 1.29 to 2.09) for rivastigmine.</p> <p>Secondary: Not reported</p>
Kim et al. <sup>107</sup> (2011)  Cholinesterase inhibitors	MA  Cognitively impaired older adults	54 trials  Variable duration	Primary: Falls, syncope, fracture and accidental injury reported  Secondary: Not reported	<p>Primary: Cholinesterase inhibitors usage was associated with the greatest risk of syncope compared to placebo (OR, 1.53; 95% CI, 1.02 to 2.30), but not with any other events: falls (OR, 0.88; 95% CI, 0.74 to 1.04); fracture (OR, 1.39; 95% CI, 0.75 to 2.56); accidental injury (OR, 1.13; 95% CI, 0.87 to 1.45).</p> <p>Memantine was associated with fewer fractures (OR, 0.21; 95% CI, 0.05 to 0.85), but not with other events: falls (OR, 0.92; 95% CI, 0.72 to 1.18), syncope (OR, 1.04; 95% CI, 0.35 to 3.04); accidental injury (OR, 0.80; 95% CI, 0.56 to 1.12).</p> <p>There were no differential effects noted according to type and severity of cognitive impairment, residential status, or length of follow-up.</p>
<b>Alzheimer's Disease</b>				
Haeberlein SB et al. <sup>108</sup> (2022) EMERGE & ENGAGE	Two DB, MC, PC, RCT  Patients 50 to 85 years of age who met clinical criteria	N=1,638 (EMERGE) and 1,647 (ENGAGE)  78 weeks	Primary: Change from baseline to week 78 on CDR-SB  Secondary:	Primary: <i>Note: An independent data monitoring committee reviewed the unblinded results of the interim analysis and made the recommendation to the sponsor to terminate the studies, which occurred on March 21, 2019. Following this futility announcement (March 21, 2019), all dosing stopped at the study sites, and data collection and data cleaning continued. The</i>



Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
<p>Aducanumab low-dose (titrated to target dose of 3 mg/kg for ApoE ε4+ or 6 mg/kg for ApoE ε4-) IV every four weeks</p> <p>vs</p> <p>aducanumab high-dose (titrated to a target dose of 6 mg/kg for ApoE ε4+ or 10 mg/kg for ApoE ε4-) IV every four weeks</p> <p>vs</p> <p>placebo</p>	<p>for MCI due to Alzheimer's disease or mild Alzheimer's disease dementia, with amyloid pathology confirmed by visual assessment of amyloid PET, MMSE score of 24 to 30, and CDR global score of 0.5.</p>		<p>MMSE, ADAS-Cog13, ADCS-ADL-MCI, biomarker endpoints and safety assessments</p>	<p>prespecified futility methodology used pooled data from EMERGE and ENGAGE to predict the future unobserved treatment effect. The individual study results using the prespecified primary efficacy analysis methods on the futility data set showed a -18% treatment difference on the CDR-SB, favoring high-dose aducanumab in EMERGE, and a 15% treatment difference on CDR-SB, favoring placebo in ENGAGE. This refuted an assumption of conditional power (i.e., that the treatment effect in the two studies would be similar), so conditional power was then recalculated using the data from each of the two studies to predict the future unobserved treatment effect, and this non-pooled analysis yielded estimates of 59% and 0% on the primary endpoint for the high-dose groups in EMERGE and ENGAGE, respectively.</p> <p>High-dose aducanumab in EMERGE demonstrated a difference of -0.39 vs placebo in the mean change from baseline in CDR-SB score at week 78 (95% CI, -0.69 to -0.09; P=0.012), a 22% reduction in decline. Results from the low-dose aducanumab arm were not statistically significant compared to placebo on the primary endpoint.</p> <p>In ENGAGE, the difference in mean change from baseline in CDR-SB scores at week 78 was 0.03 with high-dose aducanumab vs placebo (95% CI, -0.26 to 0.33; P=0.833), an increase of 2%. Results from the low-dose aducanumab arm were not statistically significant compared to placebo on the primary endpoint.</p> <p>Secondary:        In EMERGE, the high-dose aducanumab arm also showed less decline vs placebo on each of the three specified secondary endpoints: MMSE (difference of 0.6 vs placebo in mean change from baseline; -18%; P=0.049), ADAS-Cog13 (difference of -1.40 vs placebo in mean change from baseline; -27%; P=0.010), and ADCS-ADL-MCI (difference of 1.7 vs placebo in mean change from baseline; -40%; P&lt;0.001) at week 78.</p> <p>Results from three out of four endpoints from the low-dose aducanumab arm were numerically intermediate between those of the high-dose and placebo arms but not statistically significant vs placebo on any secondary endpoint.</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>In ENGAGE, in the high-dose aducanumab arm, the differences in mean change from baseline at week 78 vs placebo on the MMSE (-0.1; 3%; P=0.811), ADAS-Cog13 (-0.59; -11%; P=0.258), and ADCS-ADL-MCI (0.7; -18%; P=0.151) were not statistically significant. Results from the low-dose aducanumab arm were not statistically significant vs placebo on any of the secondary endpoints and were consistent with those from EMERGE.</p> <p>Amyloid PET substudies assessed 488 and 585 patients in EMERGE and ENGAGE, respectively. These substudies showed a dose- and time-dependent reduction in amyloid PET SUVR in both EMERGE and ENGAGE. At week 78, the difference in adjusted mean change from baseline between high-dose aducanumab and placebo was -0.278 (95% CI, -0.306 to -0.250; P&lt;0.0001) for EMERGE and -0.232 (95% CI, -0.256 to -0.208; P&lt;0.0001) for ENGAGE. For the high-dose aducanumab arm, the reduction in adjusted mean change from baseline in amyloid PET SUVR in ENGAGE was 16.5% less than that in EMERGE at week 78. The adjusted mean changes from baseline in amyloid PET SUVR for low-dose aducanumab arms were similar between EMERGE and ENGAGE at week 78.</p> <p>The incidence of adverse events was similar across dose groups in both studies. Adverse events with an incidence &gt;10% in any dose group were ARIA-E, headache, brain microhemorrhages, nasopharyngitis, fall, localized superficial siderosis and dizziness. The incidence of ARIA-E was higher in the high-dose groups compared with low-dose groups (35% vs 26%, respectively, in EMERGE and 36% vs 26%, respectively, in ENGAGE).</p>
<b>Dementia of Parkinson's Disease</b>				
Emre et al. <sup>109</sup> (2004)  Rivastigmine 3 to 12 mg/day; average dose 8.6 mg/day	DB, MC, PC, RCT  Patients at least 50 years of age with mild-to-moderate dementia developed 2 years after the	N=541  Dose titration over the first 16 weeks with a subsequent	Primary: ADAS-Cog, ADCS-CGIC  Secondary: ADCS-ADL, NPI-10, MMSE, CDR	Primary: Patients who were receiving rivastigmine had significant improvement of 2.1 points in the 70-point ADAS-Cog scores vs worsening of 0.7 point in the placebo group from baseline (P<0.001).  19.8% of patients in the rivastigmine group and 14.5% in the placebo group clinically improved in the ADCS-CGIC scores. 13% of patients in

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
vs placebo	diagnosis of Parkinson's disease	assessment period of 8 weeks  Total of 24 weeks	power of attention tests, D-KEFS verbal fluency test, Ten Point Clock-drawing Test	the rivastigmine group and 23.1% in the placebo group clinically worsened in the ADCS-CGIC scores (P=0.007).  Secondary: All secondary outcomes were significantly better in the rivastigmine group compared to placebo, as reflected by the changes in the ADCS-ADL score (P=0.02), NPI-10 (P=0.02), MMSE (P=0.03), CDR power of attention tests (P=0.009), D-KEFS verbal fluency test (P<0.001), and the Ten Point Clock-drawing Test (P=0.02).
Wesnes et al. <sup>110</sup> (2005)  Rivastigmine 3 to 12 mg/day, average dose 8.6 mg/day  vs placebo	DB, MC, PC, RCT  Patients at least 50 years old with Parkinson's disease	N=487  24 weeks	Primary: Power of attention, continuity of attention, cognitive reaction time, reaction time variability  Secondary: Not reported	Primary: At week 16, there was no statistical significance from baseline scores between rivastigmine and placebo for power of attention (P=0.11) but there was a significance at week 24 (P<0.01).  By week 16, there was a significant improvement with continuity of attention (P=0.001) compared to placebo and this parameter continued to improve at week 24 (P=0.0001).  Cognitive reaction time showed significant improvement by the end of week 24 (P<0.001) vs week 16 (P=0.064) but declined with placebo.  Reaction time variability continued to show improvement over placebo from week 16 (P<0.05) to week 24 (P<0.001).  Secondary: Not reported
Schmitt et al. <sup>111</sup> (2010)  Rivastigmine 3 to 12 mg/day  vs placebo	DB, MC, PC, RCT  Patients with Parkinson's disease dementia	N=541  24 weeks	Primary: Executive function as assessed by D-KEFS measures  Secondary: Not reported	Primary: Rivastigmine was associated with significantly more correct responses, fewer set loss errors, and more total responses made (within time available), compared to placebo (all P<0.05). There was no significant difference in total repetition errors (P=0.57).  Rivastigmine was associated with a significantly higher Card Sorting recognition description score than placebo (P=0.03). Word reading errors, word comprehension, and sort recognition errors were NS. There were significantly more correct substitutions on the Symbol Digit Modalities Test compared to placebo (P=0.02).

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Rivastigmine was associated with significantly fewer self-corrected errors on the Color-Word Interference inhibition/switching subtest compared to placebo (P=0.049). Treatment differences in numbers of correct responses were near statistical significance (P=0.050). Other treatment differences in this battery of executive function tests were not statistically significant.</p> <p>Secondary: Not reported</p>
<p>Olin et al.<sup>112</sup> (2010)</p> <p>Rivastigmine 3 to 12 mg/day</p> <p>vs</p> <p>placebo</p>	<p>DB, MC, PC, RCT</p> <p>Patients ≥50 years of age with Parkinson's disease dementia</p>	<p>N=541</p> <p>24 weeks</p>	<p>Primary: Tolerability and efficacy as measured by ADCS-ADL</p> <p>Secondary: Not reported</p>	<p>Primary: A total of 75.8% of patients completed the study (rivastigmine, 72.7% vs placebo, 82.1%). The primary reasons for discontinuation were adverse events (17.1% for rivastigmine vs 7.8% for placebo) and withdrawal of consent (5.8% rivastigmine vs 1.1% placebo).</p> <p>At 24 weeks, rivastigmine was associated with significantly less deterioration compared to placebo based on ADCS-ADL total scores (-1.1 vs -3.6, respectively; P=0.023). Similar improvements were seen with rivastigmine compared to placebo on the basic ADCS-ADL subscale (-0.5 vs -1.7, respectively; P=0.025), and on high level function ADLs (0.1 vs -1.0; P=0.017). No other measures were significantly different among the treatment groups.</p> <p>Secondary: Not reported</p>
<p>Maidment et al.<sup>113</sup> (2006)</p> <p>Rivastigmine 3 to 12 mg/day</p> <p>vs</p> <p>placebo</p>	<p>MA</p> <p>Patients diagnosed with mild-to-moderately severe dementia, which developed at least 2 years after Parkinson's disease was diagnosed</p>	<p>N=541 (1 study)</p> <p>24 weeks</p>	<p>Primary: ADAS-Cog, ADCS-CGIC</p> <p>Secondary: MMSE, ADCS-ADL, NPI, CDR, D-KEFS, Ten Point Clock-drawing Test, UPDRS, adverse events</p>	<p>Primary: Significant improvement in ADAS-Cog was found for patients treated with rivastigmine over placebo (WMD, -2.80; 95% CI, -4.26 to -1.34; P=0.0002).</p> <p>Results in ADCS-CGIC significantly favored patients treated with rivastigmine over placebo (WMD, -0.50; 95% CI, -0.77 to -0.23; P=0.0004). 19.8% of rivastigmine patients experienced "clinically meaningful (moderate or marked) improvement" compared to 14.5% of the placebo group; 13.0% of rivastigmine patients experienced "clinically meaningful worsening" compared to 23.1% in the placebo group (P values not reported).</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
				<p>Secondary: Results for MMSE significantly favored patients treated with rivastigmine over placebo (WMD, 1.00; 95% CI, 0.33 to 1.67; P=0.003).</p> <p>Results for ADCS-ADL significantly favored patients treated with rivastigmine over placebo (WMD, 2.50; 95% CI, 0.43 to 4.57; P=0.02).</p> <p>Results for NPI significantly favored patients treated with rivastigmine over placebo (WMD, -2.00; 95% CI, -3.91 to -0.09; P=0.04).</p> <p>For CDR no statistically significant difference was found (P=0.25).</p> <p>For D-KEFS, results significantly favored patients treated with rivastigmine over placebo (WMD, 2.80; 95% CI, 1.47 to 4.13; P&lt;0.0001).</p> <p>Full UPDRS was not reported. No statistically significant difference was found for motor score, including tremor (P=0.83 and P=0.84).</p> <p>Significantly more patients in the rivastigmine group than the placebo group experienced one or more adverse events (P=0.0006). Adverse events included: nausea, vomiting, tremor, and dizziness.</p> <p>Significantly more patients treated with rivastigmine withdrew from treatment for any reason than those treated with placebo (P=0.02).</p>
<p>Emre et al.<sup>114</sup> (2014)</p> <p>Rivastigmine capsules 6 mg twice daily</p> <p>vs</p> <p>rivastigmine patch 9.5 mg/24 hours</p>	<p>OL, PRO, RCT</p> <p>Patients 50 to 85 years of age with mild-to-moderately severe Parkinson disease dementia, which developed at least one year after Parkinson's disease was diagnosed</p>	<p>N=583</p> <p>76 weeks</p>	<p>Primary: Incidence of predefined adverse events due to worsening Parkinson's disease motor symptoms (tremor, rigidity, bradykinesia, and falls) and discontinuation rate due to predefined</p>	<p>Primary: The incidence of adverse effects due to worsening motor symptoms in the capsule groups was 36.1% (95% CI, 30.6 to 41.8), with tremor the most commonly reported (24.5%; 95% CI, 19.7 to 29.8). Overall, 4.4% (95% CI, 2.4 to 7.4) of capsule-treated patients discontinued due to worsening motor symptoms.</p> <p>Secondary: The incidence of adverse effects due to worsening motor symptoms in the patch group (31.9%; 95% CI, 26.6 to 37.7) was similar to capsules. Fewer patients experienced tremor with patch (9.7%; 95% CI, 6.6 to 13.7) compared to capsules. The incidences of bradykinesia, rigidity, and fall</p>

Study and Drug Regimen	Study Design and Demographics	Study Size and Study Duration	End Points	Results
			<p>potential adverse effects with capsules</p> <p>Secondary: Same as primary but with patch</p> <p>Efficacy: Mattis Dementia Rating Scale, ADCS-ADL, NPI-10</p>	<p>were similar between groups. The incidence of discontinuation due to worsening of motor symptoms was 2.4% (95% CI, 1.0 to 4.9) with patch.</p> <p>Efficacy: Improvements on Mattis Dementia Rating Scale and NPI-10 from baseline were observed in both groups at weeks 24 and 52. Deterioration in ADCS-ADL score from baseline was observed in both groups at all time points. The size of initial improvement on the Mattis Dementia Rating Scale and NPI-10 gradually declined in both groups; decline was greater in the patch group. In the overall population, there was a statistically significant difference in favor of capsules compared with patch at weeks 24 to 76 for MDRS; weeks 52 and 76 for ADCS-ADL; and weeks 24 and 76 for NPI-10.</p>

Study abbreviations: AC=active control, CI=confidence interval, DB=double blind, DD=double dummy, ER=extended release, HR=hazard ratio, IR=immediate release, ITT=intent to treat, LOCF=last observation carried forward, MA=meta-analysis, MC=multicenter, NS=not significant, OL=open label, OR=odds ratio, OS=observational study, PC=placebo controlled, PG=parallel group, PP=per protocol, PRO=prospective, RCT=randomized controlled trial, RETRO=retrospective, RR=relative risk, SB=Single-blind, WMD=weighted mean difference

Efficacy Measures Key: ADAS-Cog=Alzheimer's Disease Assessment Scale-Cognitive subscale, ADAS-cog/10=10-item cognitive subscale of the Alzheimer's Disease Assessment Scale, ADAS-cog/11=11-item cognitive subscale of the Alzheimer's Disease Assessment Scale, ADAS-cog/13=13-item cognitive subscale of the Alzheimer's Disease Assessment Scale, ADAS-cog/memory=Alzheimer's Disease Assessment Scale-Cognitive/Memory, ADAS-noncog=Alzheimer Disease Assessment Scale-Noncognitive, ADCPQ=Alzheimer's Disease Caregiver Preference Questionnaire, ADCS-ADL=Alzheimer's Disease Cooperative Study-Activities of Daily Living scale, **ADCS-ADL-MCI= Alzheimer's Disease Cooperative Study-Activities of Daily Living-Mild Cognitive Impairment scale**, ADCS-ADL-sev=Alzheimer Disease Cooperative Study-Activities of Daily Living-severe version, ADCS-CGIC=Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change, ADL=Activity of Daily Living, **ApoE=apolipoprotein E**, **ARIA-E=amyloid related imaging abnormalities-edema**, BADLS=Bristol Activities of Daily Living Scale, BEHAV-AD=Behavioral Pathology in Alzheimer's Disease Rating Scale, BGP=Behavioral Rating Scale for Geriatric Patients, BrADL=Bristol Activities of Daily Living Scale, CBQ=Caregiver Burden Questionnaire, CDR=Cognitive Drug Research, CDR-SB=Clinical Dementia Rating Scale-Sum of Boxes, CGA-NPI=Caregiver-Administered Neuropsychiatric Inventory, CGI-C=Clinical Global Impression of Change, CGI-I=Clinical Global Impression of Improvement scale, CIBIC=Clinician Interview-Based Impression of Change Scale, CIBIC-Plus=Clinician's Interview-Based Impression of Change Plus Caregiver Input, CMAI-K=Cohen Mansfield Agitation Inventory-Korean type, DAD=Disability Assessment, D-KEFS=Delis-Kaplan Executive Function System, ECG=electrocardiogram, FAB=Frontal Assessment Battery, FAST=Functional Assessment Staging, GBS=Gottfried-Bråne-Steen scale, GDS=Global Deterioration Scale, IADL=Instrumental Activity of Daily Living, IDDD=Interview for Deterioration in Daily Functioning Activities in Dementia, K-MMSE=Korean Mini-Mental Status Exam, MDS-ADL=Minimum Data Set-Activities of Daily Living, MMSE=Mini-Mental Status Exam, M-NCAS=Modified Nursing Care Assessment Scale, NPI=Neuropsychiatric Inventory, NPI-10=10-item Neuropsychiatric Inventory, QOL=quality of life, QoLS=Quality of Life Scale, PDS=Progressive Deterioration Scale, **PET=positron emission tomography**, RUSP=Resource Utilization for Severe Alzheimer Disease Patients, SIB=Severe Impairment Battery, **SUVR=standardized uptake value ratio**, UPDRS=Unified Parkinson's Disease Rating Scale

## Additional Evidence

### Dose Simplification

A search of Medline and PubMed did not reveal data pertinent to this topic.

### Stable Therapy

The cholinesterase inhibitors exhibit similar pharmacologic properties, and evidence from comparative studies support a switch strategy when patients are intolerant to one drug or when a therapeutic dose cannot be reached.<sup>115</sup> Gauthier et al. reported that when switched from donepezil to rivastigmine, approximately 50% of those who had adverse events or a lack of efficacy with donepezil tolerated or responded well to rivastigmine.<sup>116</sup> Wilkinson et al. found no difference in tolerability when patients were switched from donepezil to galantamine using either a four-day washout period or a seven-day washout period.<sup>117</sup> Sadowsky et al. evaluated immediate switch (no washout) or delayed switch (seven-day washout) from oral donepezil to transdermal rivastigmine following a four-week treatment period with donepezil.<sup>118</sup> The authors found that the rates of discontinuation due to any reason or adverse events were similar between the treatment groups. They concluded that both switch strategies were safe and well tolerated. Sakka et al. evaluated patients with moderate-to-severe Alzheimer's disease who were switched to donepezil after experiencing a treatment failure or intolerance with memantine.<sup>119</sup> The authors concluded that donepezil was effective and well tolerated in patients who discontinued memantine monotherapy, including those patients with previous exposure to cholinesterase inhibitors. A post-hoc analysis of five-month trial data with galantamine demonstrated that patients had similar efficacy outcomes, whether or not they had received prior anticholinesterase therapy, suggesting that a previous failure did not predict response to galantamine.<sup>120</sup>

### Impact on Physician Visits

Fillenbaum et al. evaluated the frequency of outpatient visits for patients with Alzheimer's disease.<sup>121</sup> Outpatient visit ranged from 81 to 95% and was not related to the stage of dementia or institutional status. Leibson et al. demonstrated that the onset of Alzheimer's disease is not associated with greater use of acute care services, nor is the high use of nursing home care offset by fewer emergency room or hospital encounters.<sup>122</sup> Clark et al. evaluated a telephone intervention program where healthcare professionals work with patients and caregivers to determine resources within the family of an Alzheimer's patient.<sup>123</sup> Alzheimer's patients in the program felt less embarrassed and isolated because of their memory problems and reported less problems coping with their disease. Intervention patients with more severe impairment had fewer physician visits, were less likely to have an emergency room visit or hospital admission, and had decreased depression and strain. Wimo et al. demonstrated that the use of memantine in patients with moderate-to-severe Alzheimer's disease was associated with less total caregiver time compared to placebo.<sup>124</sup> There were also fewer patients institutionalized at week 28 in the memantine group compared to placebo.

## IX. Cost

A "relative cost index" is provided below as a comparison of the average cost per prescription for medications within this American Hospital Formulary Service (AHFS) drug class. To differentiate the average cost per prescription from one product to another, a specific number of '\$' signs from one to five is assigned to each medication. Assignment of relative cost values is based upon current Alabama Medicaid prescription claims history and the average cost per prescription as paid at the retail pharmacy level. For brand or generic products with little or no recent utilization data, the average cost per prescription is calculated by using the Alabama Medicaid average acquisition cost (AAC) and the standard daily dosing per product labeling. Please note that the relative cost index does not factor in additional cost offsets available to the Alabama Medicaid program via pharmaceutical manufacturer rebating.

The relative cost index scale for this class is as follows:

Relative Cost Index Scale	
\$	\$0-\$30 per Rx
\$\$	\$31-\$50 per Rx
\$\$\$	\$51-\$100 per Rx
\$\$\$\$	\$101-\$200 per Rx
\$\$\$\$\$	Over \$200 per Rx

Rx=prescription

**Table 10. Relative Cost of the Alzheimer's Agents**

Generic Name(s)	Formulation(s)	Example Brand Name(s)	Brand Cost	Generic Cost
<b>Parasympathomimetic (Cholinergic Agents)</b>				
Donepezil	orally disintegrating tablet, tablet	Aricept®*	\$\$\$\$\$	\$
Galantamine	extended-release capsule, solution	Razadyne® ER*	\$\$\$\$\$	\$\$
Rivastigmine	capsule, solution, transdermal patch	Exelon®*	\$\$\$\$\$	\$\$\$
<b>Central Nervous System Agents, Miscellaneous</b>				
Aducanumab-avwa	injection	Aduhelm®	\$\$\$\$\$	N/A
Memantine	extended-release capsule, solution, tablet	Namenda®*, Namenda® XR*	\$\$\$\$\$	\$
<b>Combination Products</b>				
Memantine and donepezil	extended-release capsule	Namzaric®	\$\$\$\$\$	N/A

\*Generic is available in at least one dosage form or strength.  
 N/A=Not available.

## X. Conclusions

The cholinesterase inhibitors are approved for the treatment of mild-to-moderate Alzheimer's disease. Donepezil is also approved for the treatment of severe disease. The N-Methyl-D-aspartate (NMDA) receptor antagonist, memantine, has only been approved for the treatment of moderate-to-severe Alzheimer's disease. Although these agents provide symptomatic benefit, they have not been shown to delay the progression of neurodegeneration. Aducanumab-avwa is an amyloid beta-directed antibody and is the first disease modifying therapy approved for the treatment of Alzheimer's disease. Aducanumab received FDA approval based on a surrogate endpoint (reduction in amyloid-β plaques) and has not yet been shown to provide a clinical benefit. All products with the exception of memantine-donepezil and aducanumab-avwa are available in a generic formulation.

There are several guidelines which discuss the role of these agents in the management of Alzheimer's disease.<sup>14-18</sup> The primary goal of treatment is to delay the progression of symptoms and preserve functional ability. The use of a cholinesterase inhibitor may lead to modest improvements in some patients; therefore, it is appropriate to offer a trial of one of these agents for patients with mild-to-moderate disease.<sup>17</sup> In patients with Alzheimer's disease, treatment with cholinesterase inhibitors (donepezil, galantamine, or rivastigmine) should be considered at the time of diagnosis, taking into account expected therapeutic benefits and potential safety issues. In patients with moderate to severe Alzheimer's disease, treatment with memantine should be considered taking into account expected therapeutic benefits and potential safety issues.<sup>14-15</sup> Guidelines do not give preference to one agent over another and have not been updated to include the place in therapy for aducanumab-avwa. Clinicians should base the treatment decision on tolerability, adverse events, and ease of use.<sup>14-18</sup>

Numerous clinical trials have evaluated the efficacy and safety of the cholinesterase inhibitors and memantine. Several outcomes have been assessed (using more than 40 different instruments), including cognition, global function, behavior, and quality of life. There is consistent evidence from well-designed studies that donepezil, galantamine, rivastigmine, and memantine positively affect cognition and global function, although the improvements are modest. The findings are less consistent for other outcomes, including behavior and quality of life. In most cases, the duration of these clinical trials was less than one year. Thus, there is insufficient evidence to determine the optimal duration of therapy.<sup>15</sup>

Two randomized clinical trials, EMERGE and ENGAGE, evaluated the efficacy of aducanumab in patients with confirmed presence of amyloid pathology based on positron emission tomography (PET) scan who met clinical criteria for mild cognitive impairment due to Alzheimer's disease or mild Alzheimer's disease dementia. Both studies were terminated early after an interim analysis for futility, which occurred when approximately 50% of the pooled patients in both studies completed week 78. The primary endpoint of change from baseline to week 78 on the CDR-SB, an integrated scale that assesses both function and cognition, was met in EMERGE but not in



ENGAGE. Amyloid PET substudies showed a dose- and time-dependent reduction in amyloid PET standardized uptake value ratio in both studies. The high-dose aducanumab group in EMERGE demonstrated a difference of -0.39 vs placebo in the mean change from baseline in CDR-SB score at week 78 (95% CI, -0.69 to -0.09; P=0.012), a 22% reduction in decline. Results from the low-dose aducanumab arm were not statistically significant compared to placebo on this same measure. In ENGAGE, the difference in mean change from baseline in CDR-SB scores at week 78 was 0.03 with high-dose aducanumab vs placebo (95% CI, -0.26 to 0.33; P=0.833), an increase of 2%. Results from the low-dose aducanumab arm were not statistically significant compared to placebo on the same measure.<sup>108</sup> There are relatively few studies that directly compare the efficacy and safety of the Alzheimer's agents. Most of the trials have compared active treatment to placebo or no treatment. The studies also differ with regards to design, patient population, and treatment duration, which make it difficult to directly compare the results.<sup>19-114</sup>

The Centers for Medicare & Medicaid Services (CMS) completed further evaluation of aducanumab (Aduhelm<sup>®</sup>), including stakeholder input. After extensive evaluation and review due to the limited clinical efficacy outcomes data, beyond that seen with surrogate marker evaluation, compared to placebo in clinical trials and the significant safety concerns, CMS published the final national coverage determination for aducanumab (Aduhelm<sup>®</sup>) on April 7, 2022. The final Medicare national coverage determination indicates coverage of FDA-approved monoclonal antibody therapies directed against amyloid for the treatment of Alzheimer's disease when provided in accordance with the extensive coverage criteria outlined for coverage with evidence development for patients who have a clinical diagnosis of mild cognitive impairment due to Alzheimer's disease or mild Alzheimer's disease dementia, both with confirmed presence of amyloid beta pathology consistent with Alzheimer's disease. The guidance indicates that monoclonal antibodies directed against amyloid that are approved by FDA for the treatment of Alzheimer's disease based upon evidence of efficacy from a change in a surrogate endpoint (e.g., amyloid reduction), considered as reasonably likely to predict clinical benefit, may be covered in a randomized controlled trial conducted under an investigational new drug application. Additionally, monoclonal antibodies directed against amyloid that are approved by FDA for the treatment of Alzheimer's disease based upon evidence of efficacy from a direct measure of clinical benefit may be covered in CMS-approved prospective comparative studies, that meet specific protocol and analysis requirements outlined by CMS.<sup>125</sup>

There is insufficient evidence to support that one brand Alzheimer's agent is safer or more efficacious than another. Formulations without a generic alternative should be managed through the medical justification portion of the prior authorization process.

Therefore, all brand Alzheimer's agents within the class reviewed are comparable to each other and to the generic products in the class (if applicable) and offer no significant clinical advantage over other alternatives in general use.

## **XI. Recommendations**

No brand Alzheimer's agent is recommended for preferred status. Alabama Medicaid should accept cost proposals from manufacturers to determine the most cost effective products and possibly designate one or more preferred brands.

Aducanumab-avwa should not be placed in preferred status regardless of cost.

## XII. References

1. American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association, 2013.
2. Wolk DA, Dickerson BC. Clinical features and diagnosis of Alzheimer disease. In: UpToDate [database on the internet]. Waltham (MA); 2022 [cited 2022 Jan]. Available from: <http://www.uptodate.com/utd/index.do>.
3. Press D, Buss SS. Treatment of Alzheimer dementia. In: UpToDate [database on the internet]. Waltham (MA): UpToDate; 2022 [cited 2022 Jan]. Available from: <http://www.uptodate.com/utd/index.do>.
4. Lexicomp Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.; 2022 [cited 2022 Jan]. Available from: <http://online.lexi.com>. Subscription required to view..
5. Micromedex® Healthcare Series [database on the Internet]. Greenwood Village (CO): Thomson Reuters (Healthcare) Inc.; Updated periodically [cited 2022 Jan]. Available from: <http://www.thomsonhc.com/>.
6. Aricept® [package insert]. Woodcliff Lake, NJ: Eisai, Inc.; December 2021.
7. Razadyne® [package insert]. Titusville, NJ: Janssen Pharmaceuticals, Inc.; August 2021.
8. Exelon® Patch [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; Jun 2020.
9. Namenda® [package insert]. Madison, NJ: Allergan USA, Inc.; November 2018.
10. Namenda® XR [package insert]. Madison, NJ: Allergan USA, Inc.; November 2019.
11. Namzaric® [package insert]. St. Louis, MO: Forest Pharmaceuticals, Inc.; Jan 2019.
12. Aduhelm® [package insert]. Cambridge (MA): Biogen, Inc.; 2021 Jul.
13. Rabins PV, Blacker D, Rovner BW, Rummans T, Schneider LS, Tariot PN, et al. American Psychiatric Association practice guideline for the treatment of patients with Alzheimer's disease and other dementias. Second edition. *Am J Psychiatry*. 2007 Dec;164(12 Suppl):5-56
14. Hort J, O'Brien JT, Gainotti G, et al. EFNS Scientist Panel on Dementia. EFNS guidelines for the diagnosis and management of Alzheimer's disease. *Eur J Neurol*. 2010;17(10):1236-48.
15. National Institute for Health and Care Excellence. Dementia: assessment, management and support for people living with dementia and their carers. National Institute for Health and Care Excellence (NICE); June 2018. Available at: <https://www.nice.org.uk/guidance/ng97>. Accessed June 2019.
16. Petersen RC, Lopez O, Armstrong MJ, et al. Practice guideline update summary: Mild cognitive impairment: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90(3):126-135. doi:10.1212/WNL.0000000000004826.
17. Qaseem A, Snow V, Cross J, et al. American College of Physicians/American Academy of Family Physicians Panel on Dementia. Current pharmacologic treatment of dementia: a clinical practice guideline from the American College of Physicians and the American Academy of Family Physicians. *Ann Intern Med*. 2008;148:370-8.
18. Seppi K, Chaudhuri KR, Coelho M, Fox SH, Katzenschlager R, et al. The Movement Disorder Society Evidence-Based Medicine Review Update: Treatments for the non-motor symptoms of Parkinson's disease. *Mov Disord*. 2018 Dec. Available at <https://www.movementdisorders.org/MDS/Resources/Publications-Reviews/EBM-Reviews.htm>.
19. Geldmacher DS, Provenzano G, McRae T, Mastey V, Ieni JR. Donepezil is associated with delayed nursing home placement in patients with Alzheimer's disease. *J Am Geriatr Soc*. 2003 Jul;51(7):937-44.
20. Burns A, Gauthier S, Perdomo C, et al. Efficacy and safety of donepezil over 3 years: an open-label, multicentre study in patients with Alzheimer's disease. *Int J Geriatr Psychiatry*. 2007;22:806-12.
21. Hashimoto M, Yatabe Y, Kaneda K, et al. Impact of donepezil hydrochloride on the care burden of family caregivers of patients with Alzheimer's disease. *Psychogeriatrics*. 2009;9:196-203.
22. Homma A, Imai Y, Tago H, et al. Long-term safety and efficacy of donepezil in patients with severe Alzheimer's disease: results from a 52-week, open-label, multicenter, extension study in Japan. *Dement Geriatr Cogn Disord*. 2009;27:232-9.
23. Courtney C, Farrell D, Gray R, Hills R, Lynch L, Sellwood E, Edwards S, Hardyman W, Raftery J, Crome P, Lendon C, Shaw H, Bentham P; AD2000 Collaborative Group. Long-term donepezil treatment in 565 patients with Alzheimer's disease (AD2000): randomised double-blind trial. *Lancet*. 2004 Jun 26;363(9427):2105-15.
24. Sabbagh M, Cummings J, Christensen D, Doody R, Farlow M, Liu L, Mackell J, Fain R. Evaluating the cognitive effects of donepezil 23 mg/d in moderate and severe Alzheimer's disease: analysis of effects of baseline features on treatment response. *BMC Geriatrics*. 2013;13:56.
25. Tariot P, Salloway S, Yardley J, Mackell J, Moline M. Long-term safety and tolerability of donepezil 23 mg in patients with moderate to severe Alzheimer's disease. *BMC Research Notes*. 2012;5:283.

26. Winblad B, Wimo A, Engedal K, Soininen H, Verhey F, Waldemar G, et al. 3-year study of donepezil therapy in Alzheimer's disease: effects of early and continuous therapy. *Dement Geriatr Cogn Disord*. 2006;21(5-6):353-63.
27. Rogers SL, Farlow MR, Doody RS, Mohs R, Friedhoff LT. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. Donepezil Study Group. *Neurology*. 1998 Jan;50(1):136-45.
28. Winblad B, Kilander L, Eriksson S, Minthon L, Batsman S, Wetterholm AL, Jansson-Blixt C, et al; Severe Alzheimer's Disease Study Group. Donepezil in patients with severe Alzheimer's disease: double-blind, parallel-group, placebo-controlled study. *Lancet*. 2006 Apr 1;367(9516):1057-65.
29. Black SE, Doody R, Li H, McRae T, Jambor KM, et al. Donepezil preserves cognition and global function in patients with severe Alzheimer disease. *Neurology*. 2007;69:459-69.
30. Homma A, Imai Y, Tago H, et al. Donepezil treatment of patients with severe Alzheimer's disease in a Japanese population: results from a 24-week, double-blind, placebo-controlled, randomized trial. *Dement Geriatr Cogn Disord*. 2008;25:399-407.
31. Birks J, Harvey RD. Donepezil for dementia due to Alzheimer's disease. *Cochrane Database Syst Rev*. 2006 Jan 25;(1):CD001190. doi: 10.1002/14651858.CD001190.pub2.
32. Wallin A et al. Donepezil in Alzheimer's disease: what to expect after 3 years of treatment in a routine clinical setting. *Dementia and Geriatric Cognitive Disorders*. 2007;23:150-60.
33. Farlow MR, Salloway S, Tariot PN, et al. Effectiveness and tolerability of high-dose (23 mg/d) vs standard-dose (10 mg/d) donepezil in moderate to severe Alzheimer's disease: A 24-week, randomized, double-blind study. *Clin Ther*. 2010;32:1234-51.
34. Ferris SH, Schmitt FA, Saxton J, et al. Analyzing the impact of 23 mg/day donepezil on language dysfunction in moderate to severe Alzheimer's disease. *Alzheimers Res Ther*. 2011;3:22.
35. Farlow M, Veloso F, Moline M, et al. Safety and tolerability of donepezil 23 mg in moderate to severe Alzheimer's disease. *BMC Neurol*. 2011;11:57.
36. Doody RS, Geldmacher DS, Farlow MR, Sun Y, Moline M, Mackell J. Efficacy and safety of donepezil 23 mg vs donepezil 10 mg for moderate-to-severe Alzheimer's disease: a subgroup analysis in patients already taking or not taking concomitant memantine. *Dementia & Geriatric Cognitive Disorders*. 2012;3(2-3):164-73.
37. Raskind MA, Peskind ER, Truyen L, Kershaw P, Damaraju CV. The cognitive benefits of galantamine are sustained for at least 36 months: a long-term extension trial. *Arch Neurol*. 2004 Feb;61(2):252-6.
38. Rockwood K, Dai D, Mitnitski A, et al. Patterns of decline and evidence of subgroups in patients with Alzheimer's disease taking galantamine for up to 48 months. *Int J Geriatr Psychiatry*. 2008;23:207-14.
39. Wallin AK, Wattmo C, Minthon L. Galantamine treatment in Alzheimer's disease: response and long-term outcome in a routine clinical setting. *Neuropsychiatr Dis Treat*. 2011;7:565-76.
40. Brodaty H et al. A Naturalistic Study of Galantamine for Alzheimer's disease. *CNS Drugs*. 2006;20(11):935-46.
41. Richarz U, Gaudig M, Rettig K, Schauble B. Galantamine treatment in outpatients with mild Alzheimer's disease. *Acta Neurol Scand*. 2014 Jun;129(6):382-92.
42. Cummings JL, Schneider L, Tariot PN, Kershaw PR, Yuan W. Reduction of behavioral disturbances and caregiver distress by galantamine in patients with Alzheimer's disease. *Am J Psychiatry*. 2004 Mar;161(3):532-8.
43. Scarpini E, Bruno G, Zappalà G, et al. Cessation vs continuation of galantamine treatment after 12 months of therapy in patients with Alzheimer's disease: a randomized, double blind, placebo controlled withdrawal trial. *J Alzheimers Dis*. 2011;26:211-20.
44. Kavanagh S, Howe I, Brashear HR, et al. Long-term response to galantamine in relation to short-term efficacy data: pooled analysis in patients with mild to moderate Alzheimer's disease. *Curr Alzheimer Res*. 2011;8:175-86.
45. Burns A, Bernabei R, Bullock R, et al. Safety and efficacy of galantamine (Reminyl) in severe Alzheimer's disease (the SERAD study): a randomised, placebo-controlled, double-blind trial. *Lancet Neurol*. 2009;8:39-47.
46. Raskind MA, Peskind ER, Truyen L, Kershaw P, Damaraju CV. The cognitive benefits of galantamine are sustained for at least 36 months: a long-term extension trial. *Arch Neurol*. 2004 Feb;61(2):252-6.
47. Wilcock GK, Lilienfeld S, Gaens E. Efficacy and safety of galantamine in patients with mild-to-moderate Alzheimer's disease: multicentre randomised controlled trial. Galantamine International-1 Study Group. *BMJ*. 2000 Dec 9;321(7274):1445-9.
48. Dunbar F, Zhu Y, Brashear H. Post hoc comparison of daily rates of nausea and vomiting with once- and twice-daily galantamine from a double-blind, placebo-controlled, parallel-group, 6-month study. *Clin Ther*. 2006 Mar;28(3):365-72.

49. Brodaty H, Corey-Bloom J, Potocnik F, Truyen L, Gold M, et al. Galantamine prolonged-release formulation in the treatment of mild-to-moderate Alzheimer's disease. *Dementia and Geriatric Cognitive Disorders*. 2005;20:120-32.
50. Loy C, Schneider L. Galantamine for Alzheimer's disease and mild cognitive impairment. *Cochrane Database Syst Rev*. 2006, Issue 1. Art.No.:CD001747. DOI: 10.1002/14651858.CD001747.pub3.
51. Herrmann N, Cappell J, Eryavec GM, Lanctôt KL. Changes in nursing burden following memantine for agitation and aggression in long-term care residents with moderate to severe Alzheimer's disease: an open-label pilot study. *CNS Drugs*. 2011;25:425-33.
52. Bakchine S, Loft H. Memantine treatment in patients with mild to moderate Alzheimer's disease: results of a randomised, double-blind, placebo-controlled 6-month study. *J Alzheimers Dis*. 2007;11:471-9.
53. Reisberg B, Doody R, Stoffler A, Schmitt F, Ferris S, Mobius HJ; Memantine Study Group. Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med*. 2003 Apr 3;348(14):1333-41.
54. Winblad B, Poritis N. Memantine in severe dementia: results of the 9M-Best Study (benefit and efficacy in severely demented patients during treatment with memantine). *Int J Geriatr Psychiatry*. 1999;14(2):135-46.
55. Winblad B, Jones RW, Wirth Y, Stoffler A, Mobius HJ. Memantine in moderate-to-severe Alzheimer's disease: a meta-analysis of randomised clinical trials. *Dement Geriatr Cogn Disord*. 2007;24(1):20-7.
56. Wilkinson D and Andersen HF. Analysis of the effect of memantine in reducing the worsening of clinical symptoms in patients with moderate-to-severe Alzheimer's disease. *Dementia and Geriatric Cognitive Disorders*. 2007;24:138-45.
57. McShane R, Areosa Sastre A, Minakaran N. Memantine for dementia. *Cochrane Database of Systematic Reviews* 2006, Issue 2. Art.No.:CD003154. DOI: 10.1002/14651858.CD003154.pub5.
58. Grossberg GT, Manes F, Allegri RF, Gutierrez-Robledo LM, Gloger S, Xie L, Jia XD, Pejovic V, Miller ML, Perhach JL, Graham SM. The safety, tolerability, and efficacy of once-daily memantine (28 mg): a multinational, randomized, double-blind, placebo-controlled trial in patients with moderate-to-severe Alzheimer's disease taking cholinesterase inhibitors. *CNS Drugs*. 2013 Jun;27(6):469-78.
59. Grossberg GT, Alva G, Hendrix S, Ellison N, Kane MC, Edwards J. Memantine ER Maintains Patient Response in Moderate to Severe Alzheimer's Disease: Post Hoc Analyses From a Randomized, Controlled, Clinical Trial of Patients Treated With Cholinesterase Inhibitors. *Alzheimer Dis Assoc Disord*. 2018;32(3):173-178. doi:10.1097/WAD.0000000000000261.
60. Hager K, Baseman AS, Nye JS, Brashear HR, Han J, Sano M, et al. Effect of concomitant use of memantine on mortality and efficacy outcomes of galantamine-treated patients with Alzheimer's disease: post-hoc analysis of a randomized placebo-controlled study. *Alzheimers Res Ther*. 2016 Nov 15;8(1):47.
61. Burns A, Spiegel R, Quarg P. Efficacy of rivastigmine in subjects with moderately severe Alzheimer's disease. *Int J Geriatr Psychiatry*. 2004 Mar;19(3):243-9
62. Dantoine T, Auriacombe S, Sarazin M, et al. Rivastigmine monotherapy and combination therapy with memantine in patients with moderately severe Alzheimer's disease who failed to benefit from previous cholinesterase inhibitor treatment. *International Journal of Clinical Practice*. 2006;60(1):110-8.
63. Olin JT, Bhatnagar V, Reyes P, et al. Safety and tolerability of rivastigmine capsule with memantine in patients with probable Alzheimer's disease: a 26-week, open-label, prospective trial (Study ENA713B US32). *Int J Geriatr Psychiatry*. 2010;25:419-26.
64. Gauthier S, Juby A, Dalziel W, Réhel B, Schecter R; EXPLORE investigators. Effects of rivastigmine on common symptomatology of Alzheimer's disease (EXPLORE). *Curr Med Res Opin*. 2010;26:1149-60.
65. Birks J, Grimely EJ, Iakovidou V, Tsolaki M. Rivastigmine for Alzheimer's disease. *Cochrane Database Syst Rev*. 2000, Issue 4. Art. No.: CD001191. DOI: 10.1002/14651858.CD001191.
66. Birks J, Grimley Evans J, Iakovidou V, et al. Rivastigmine for Alzheimer's disease. *Cochrane Database Syst Rev* 2009; (2); Cochrane AN: CD001191.
67. Rosler M, Anand R, Cicin-Sain A, Gauthier S, et al. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomized controlled trial. *BMJ*. 1999;318:633-40.
68. Articus K, Baier M, Tracik F, et al. A 24-week, multicentre, open evaluation of the clinical effectiveness of the rivastigmine patch in patients with probable Alzheimer's disease. *Int J Clin Pract*. 2011;65:790-6.
69. Grossberg G, Sadowsky C, Fröstl H, et al. Safety and tolerability of the rivastigmine patch: results of a 28-week open-label extension. *Alzheimer Dis Assoc Disord*. 2009;23:158-64.
70. Gauthier S, Robillard A, Cohen S, Black S, Sampalis J, Colizza D, de Takacs F, Schecter R. EMBRACE investigators. Real-life effectiveness and tolerability of the rivastigmine transdermal patch in patients with mild-to-moderate Alzheimer's disease: the EMBRACE study. *Current Medical Research & Opinion*. 2013 Aug;29(8):989-1000.

71. Sadowsky CH, Farlow MR, Meng X, Olin JT. Safety and tolerability of rivastigmine transdermal patch compared to rivastigmine capsules in patients switched from donepezil: data from three clinical trials. *Int J Clin Pract.* 2010;64:188-93.
72. Cummings J, Froelich L, Black SE, Bakchine S, Bellelli G, Molinuevo JL, et al. Randomized, double-blind, parallel-group, 48-week study for efficacy and safety of a higher-dose rivastigmine patch (15 vs. 10 cm<sup>2</sup>) in Alzheimer's disease. *Dement Geriatr Cogn Disord.* 2012;33(5):341-53.
73. Cummings JL, Farlow MR, Meng X, et al. Rivastigmine transdermal patch skin tolerability: results of a 1-year clinical trial in patients with mild-to-moderate Alzheimer's disease. *Clin Drug Investig.* 2010;30:41-9.
74. Molinuevo JL, Arranz FJ. Impact of transdermal drug delivery on treatment adherence in patients with Alzheimer's disease. *Expert Rev Neurother.* 2012;12:31-7.
75. Boada M, Arranz FJ. Transdermal is better than oral: observational research of the satisfaction of caregivers of patients with Alzheimer's disease treated with rivastigmine. *Dementia & Geriatric Cognitive Disorders.* 2013;35(1-2):23-33.
76. Blesa González R, Boada Rovira M, Martínez Parra C, et al. Evaluation of the convenience of changing the rivastigmine administration route in patients with Alzheimer disease. *Neurologia.* 2011;26:262-71.
77. Winblad B, Grossberg G, Frolich L, Farlow M, et al. IDEAL: a 6 month, double-blind, placebo-controlled study of the first skin patch for Alzheimer disease. *Neurology.* 2007;69(Suppl1):S14-S22.
78. Winblad B, Kawata AK, Beusterien KM, et al. Caregiver preference for rivastigmine patch relative to capsules for treatment of probable Alzheimer's disease. *Int J Geriatr Psychiatry.* 2007 May;22(5):485-91.
79. Winblad B, Cummings J, Andreasen N, Grossberg G, et al. A six-month double-blind, randomized, placebo-controlled study of a transdermal patch in Alzheimer's disease-rivastigmine patch vs capsule. *International Journal of Geriatric Psychiatry.* 2007;22:456-67.
80. Blesa R, et al. Caregiver preference for rivastigmine patches vs capsules for the treatment of Alzheimer disease. *Neurology.* 2007;69(Suppl 1):S23-S28.
81. Farlow MR, Grossberg GT, Meng X, et al. Rivastigmine transdermal patch and capsule in Alzheimer's disease: influence of disease stage on response to therapy. *Int J Geriatr Psychiatry.* 2011 Dec;26(12):1236-43.
82. Choi SH, Park KW, Na DL, et al.; Expect Study Group. Tolerability and efficacy of memantine add-on therapy to rivastigmine transdermal patches in mild to moderate Alzheimer's disease: a multicenter, randomized, open-label, parallel-group study. *Curr Med Res Opin.* 2011;27:1375-83.
83. Farlow MR, Alva G, Meng X, Olin JT. A 25-week, open-label trial investigating rivastigmine transdermal patches with concomitant memantine in mild-to-moderate Alzheimer's disease: a post hoc analysis. *Curr Med Res Opin.* 2010;26:263-9.
84. Harry RD, Zakzanis KK. A comparison of donepezil and galantamine in the treatment of cognitive symptoms of Alzheimer's disease: a meta-analysis. *Hum Psychopharmacol.* 2005 Apr;20(3):183-7.
85. Wilcock G, Howe I, Coles H, Lilienfeld S, Truyen L, Zhu Y, Bullock R, Kershaw P; GAL-GBR-2 Study Group. A long-term comparison of galantamine and donepezil in the treatment of Alzheimer's disease. *Drugs Aging.* 2003;20(10):777-89.
86. Jones RW, Soininen H, Hager K, Aarsland D, Passmore P, Murthy A, Zhang R, Bahra R. A multinational, randomised, 12-week study comparing the effects of donepezil and galantamine in patients with mild-to-moderate Alzheimer's disease. *Int J Geriatr Psychiatry.* 2004 Jan;19(1):58-67.
87. Modrego PJ, Fayed N, Errea JM, et al. Memantine vs donepezil in mild to moderate Alzheimer's disease: a randomized trial with magnetic resonance spectroscopy. *Eur J Neurol.* 2010;17:405-12.
88. Rozankovic PB, Rozankovic M, Badzak J, Stojic M, Sporis IS. Impact of Donepezil and Memantine on Behavioral and Psychological Symptoms of Alzheimer Disease: Six-month Open-label Study. *Cogn Behav Neurol.* 2021 Dec 2;34(4):288-294.
89. Wilkinson DG, Passmore AP, Bullock R, Hopker SW, Smith R, Potocnik FC, Maud CM, Engelbrecht I, Hock C, Ieni JR, Bahra RS. A multinational, randomised, 12-week, comparative study of donepezil and rivastigmine in patients with mild-to-moderate Alzheimer's disease. *Int J Clin Pract.* 2002;56(6):441-6.
90. Van Puyvelde K, Mets T; RODOS Study Group. Galantamine (Reminyl) once daily outcome and satisfaction survey (RODOS) in mild to moderate Alzheimer's disease: a study in a real life population. *Geriatr Gerontol Int.* 2011;11:256-61.
91. Tariot PN, Farlow MR, Grossberg GT, Graham SM, McDonald S, Gergel I; Memantine Study Group. Memantine treatment in patients with moderate-to-severe Alzheimer disease already receiving donepezil: a randomized controlled trial. *JAMA.* 2004 Jan 21;291(3):317-24.
92. Bullock R, Touchon J, Bergman H, et al. Rivastigmine and donepezil treatment in moderate to moderately-severe Alzheimer's disease over a 2-year period. *Curr Med Res Opin.* 2005;21:1317-1327.

93. Mossello E, Tonon E, Caleri V, Tilli S, Cantini C, Cavallini MC, Bencini F, Mecacci R, Marini M, Bardelli F, Sarcone E, Razzi E, Biagini CA, Masotti G. Effectiveness and safety of cholinesterase inhibitors in elderly subjects with Alzheimer's disease: a "real world" study. *Arch Gerontol Geriatr Suppl.* 2004;(9):297-307.
94. Aguglia E, Onor ML, Saina M, Maso E. An open-label, comparative study of rivastigmine, donepezil and galantamine in a real-world setting. *Curr Med Res Opin.* 2004 Nov;20(11):1747-52.
95. Lopez-Pousa S, Turon-Estrada A, Garre-Olmo J, Pericot-Nierga I, Lozano-Gallego M, Vilalta-Franch M, Hernandez-Ferrandiz M, Morante-Munoz V, Isern-Vila A, Gelada-Battle E, Majo-Llopert J. Differential efficacy of treatment with acetylcholinesterase inhibitors in patients with mild and moderate Alzheimer's disease over a 6-month period. *Dement Geriatr Cogn Disord.* 2005;19(4):189-95.
96. Rodda J, Morgan S, Walker Z. Are cholinesterase inhibitors effective in the management of the behavioral and psychological symptoms of dementia in Alzheimer's disease? A systematic review of randomized, placebo-controlled trials of donepezil, rivastigmine and galantamine. *Int Psychogeriatr* 2009;21:813-24.
97. Howard R, McShane R, Lindesay J, et al. Donepezil and memantine for moderate-to-severe Alzheimer's disease. *N Engl J Med.* 2012;366:893-903.
98. Porsteinsson AP, Grossberg GT, Mintzer J, et al. Memantine treatment in patients with mild to moderate Alzheimer's disease already receiving a cholinesterase inhibitor: a randomized, double-blind, placebo-controlled trial. *Curr Alzheimer Res.* 2008;5:83-9.
99. Cumming JL, Schneider E, Tariot P, et al. Behavioral effects of memantine in Alzheimer disease patients receiving donepezil treatment. *Neurology.* 2006;67:57-63.
100. Maidment ID, Fox CG, Boustani M, et al. Efficacy of memantine on behavioral and psychological symptoms related to dementia: a systematic meta-analysis. *Ann Pharmacother.* 2008;42:32-8.
101. Wilkinson D, Schindler R, Schwam E, et al. Effectiveness of donepezil in reducing clinical worsening in patients with mild-to-moderate Alzheimer's disease. *Dement Geriatr Cogn Disord.* 2009;28:244-51.
102. Feldman HH, Pirttila T, Dartigues JF, et al. Treatment with galantamine and time to nursing home placement in Alzheimer's disease patients with and without cerebrovascular disease. *Int J Geriatr Psychiatry.* 2009;24:479-88.
103. Trinh NH, Hoblyn J, Mohanty S, Yaffe K. Efficacy of cholinesterase inhibitors in the treatment of neuropsychiatric symptoms and functional impairment in Alzheimer disease: a meta-analysis. *JAMA.* 2003 Jan 8;289(2):210-6.
104. Lanctôt K, Herrmann N, Yau K, Khan L, Liu B et al. Efficacy and safety of cholinesterase inhibitors in Alzheimer's disease: a meta-analysis. *CMAJ.* 2003 Sept 16;169(6):169.
105. Birks J. Cholinesterase inhibitors for Alzheimer's disease. *Cochrane Database Syst Rev.* 2006, Issue 1. Art.No.: CD005593. DOI: 10.1002/14651858.CD005593.
106. Hansen RA, Gartlehner G, Webb AP, et al. Efficacy and safety of donepezil, galantamine, and rivastigmine for the treatment of Alzheimer's disease: a systematic review and meta-analysis. *Clin Interv Aging.* 2008;3:211-25.
107. Kim DH, Brown RT, Ding EL, et al. Dementia medications and risk of falls, syncope, and related adverse events: meta-analysis of randomized controlled trials. *J Am Geriatr Soc.* 2011 Jun;59(6):1019-31.
108. Haeblerlein SB, Aisen PS, Barkhof F, Chalkias S, Chen T, Cohen S et al. Two randomized phase 3 studies of aducanumab in early Alzheimer's Disease. *J Prev Alz Dis* 2022; Published online March 18, 2022, <http://dx.doi.org/10.14283/jpad.2022.30>.
109. Emre M, Aarsland D, Albanese A, et al. Rivastigmine for dementia associated with Parkinson's disease. *NEJM.* 2004;351(24):2509-18.
110. Wesnes KA, McKeith I, Edgar C, Emre M, Lane R. Benefits of rivastigmine on attention in dementia associated with Parkinson disease. *Neurology.* 2005;65:1654-6.
111. Schmitt FA, Farlow MR, Meng X, et al. Efficacy of rivastigmine on executive function in patients with Parkinson's disease dementia. *CNS Neurosci Ther* 2010;16:330-6.
112. Olin JT, Aarsland D, Meng X. Rivastigmine in the treatment of dementia associated with Parkinson's disease: effects on activities of daily living. *Dement Geriatr Cogn Disord.* 2010;29(6):510-5.
113. Maidment I, Fox C, Boustani M. Cholinesterase inhibitors for Parkinson's disease dementia. *Cochrane Database Syst Rev.* 2006, Issue 1. Art. No.: CD004747. DOI: 10.1002/14651858.CD004747.pub2.
114. Emre M, Poewe W, De Deyn PP, et al. Long-term safety of rivastigmine in parkinson disease dementia: an open-label, randomized study. *Clin Neuropharmacol.* 2014 Jan-Feb;37(1):9-16.
115. Scarpini E, Scheltens P, Feldman H. Treatment of Alzheimer's disease: current status and new perspectives. *Lancet Neurol.* 2003 Sep;2(9):539-47.

116. Gauthier S, Emre M, Farlow MR, Bullock R, Grossberg GT, Potkin SG. Strategies for continued successful treatment of Alzheimer's disease: switching cholinesterase inhibitors. *Curr Med Res Opin* 2003;19(8):707-14.
117. Wilkinson DG, Howe I. Switching from donepezil to galantamine: a double-blind study of two wash-out periods. *Int J Geriatr Psychiatry*. 2005 May;20(5):489-91.
118. Sadowsky CH, Dengiz A, Olin JT, et al. Switching from donepezil tablets to rivastigmine transdermal patch in Alzheimer's disease. *Am J Alzheimers Dis Other Dement*. 2009;24:267-75.
119. Sakka P, Tsolaki M, Hort J, et al. Effectiveness of open-label donepezil treatment in patients with Alzheimer's disease discontinuing memantine monotherapy. *Curr Med Res Opin*. 2007;23:3153-65.
120. Emre M. Switching cholinesterase inhibitors in patients with Alzheimer's disease. *Int J Clin Pract Suppl*. 2002 Jun;(127):64-72.
121. Fillenbaum G, Heyman A, Peterson BL, Pieper CF, Weiman AL. Use and cost of outpatient visits of AD patients: CERAD XXII. *Neurology*. 2001 Jun 26;56(12):1706-11.
122. Leibson C, Owens T, O'Brien P, Waring S, Tangalos E, Hanson V, Plevak M, Kokmen E. Use of physician and acute care services by persons with and without Alzheimer's disease: a population-based comparison. *J Am Geriatr Soc*. 1999 Jul;47(7):864-9.
123. Clark PA, Bass DM, Looman WJ, McCarthy CA, Eckert S. Outcomes for patients with dementia from the Cleveland Alzheimer's Managed Care Demonstration. *Aging Ment Health*. 2004 Jan;8(1):40-51.
124. Wimo A, Winblad B, Stoffler A, Wirth Y, Mobius HJ. Resource utilization and cost analysis of memantine in patients with moderate-to-severe Alzheimer's disease. *Pharmacoeconomics*. 2003;21(5):327-40.
125. Centers for Medicare & Medicaid Services. Medicare Coverage Policy for Monoclonal Antibodies Directed Against Amyloid for the Treatment of Alzheimer's Disease [Fact Sheet available online]. Baltimore (MD): Centers for Medicare & Medicaid Services; April 7, 2022 [cited 2022 Apr 12]. Available at: <https://www.cms.gov/newsroom/fact-sheets/medicare-coverage-policy-monoclonal-antibodies-directed-against-amyloid-treatment-alzheimers-disease>.