## **ACHN RISK-STRATIFICATION ALGORITHM - GENERAL**

Step 1: Use the following criteria to risk stratify the recipient (Within the most recent 90 days at least two per risk level needed)

### **High Risk**

#### **Risk Stratification Data:**

One or more Core \*\*\*SDoH Three or more ED visits Delivering mother has history of Any readmission

spontaneous abortions Delivering mother aged 21 or younger Delivering mother aged 35 or older

Delivering mother enrolled 2<sup>nd</sup>

trimester or greater

No prenatal care for previous pregnancy

Inappropriate birth spacing

Domestic violence (still residing with partner)

Substance Misuse Depression

#### Clinical:

\*\* One unstable behavioral health diagnosis Two or more stable behavioral health

diagnoses

Three or more active chronic conditions.

Active cancer

Pregnancy ended within previous 3 months In-home delivery within previous 3 months

#### **Medium Risk**

#### **Risk Stratification Data:**

Moderate claims history One or more Primary SDoH History of substance use/misuse in past year

Language /communication barriers that interfere with recipient's ability to understand

## **Utilization:**

**Utilization:** 

One hospitalization related to

Admission to substance use

treatment facility in past 30 days

chronic conditions and/or pregnancy

One to two ED visits related to chronic conditions

Any hospitalization within 12 months post-partum

Tobacco products user

## Clinical:

\* One stable behavioral health diagnosis. Fewer than three active chronic conditions

Any hospitalization History of cancer

Delivering mother history of STDs/STIs

## Low Risk

## **Risk Stratification Data:**

**Utilization**:

Clinical:

Low claims history No SDoH

No ED visits No hospitalizations

No chronic health conditions At least one maintenance medication

Two or more doctor visits per year for at least 3 months

Step 2: Use objective and subjective data to assign a risk-stratification level for the recipient. Clinical judgement and subjective data must be thoroughly documented in HIMS.

#### Key

\*Stable Behavioral Diagnoses:

\*\*Unstable Behavioral Diagnoses:

Includes medication adherence, no mental health crisis episodes in past 3 mos.

Includes medication non-compliance, 1 missed refill within 3 mos. period, Patient admission of missed medication for 3 or more consecutive days, behavioral

changes (missed work/tardiness), mental health crisis in past 3 mos.

# **Claims History Thresholds:**

**Significant Claims History** 

**Moderate Claims History** 

**Low Claims History** 

Is 1% to 6% of the ACHN-eligible population in terms of cost.

Is 7% to 13% of the ACHN-eligible population in terms of cost.

Is 14% to 20% of the ACHN -eligible population in terms of cost.

\*\*\*SDoH

Is the abbreviation for Social Determinant of Health.

The four Agency-recognized Core SDoHs and their ICD-10 Z-Codes are:

Z59.41 **Food Insecurity** (no food or insufficient supply of food)

Z59.0 Housing Instability (specifically homeless)

Z59.82 Lack of Transportation (no private owned transportation or access to public transit)

Interpersonal Violence (domestic violence) (currently living in an unsafe environment and subjected to partner violence)